



Title	Effect of chlorination on anti-acetylcholinesterase activity of organophosphorus insecticide solutions and contributions of the parent insecticides and their oxons to the activity
Author(s)	Matsushita, Taku; Fujita, Yuki; Omori, Kei; Huang, Yuxiang; Matsui, Yoshihiko; Shirasaki, Nobutaka
Citation	Chemosphere, 261, 127743 <a href="https://doi.org/10.1016/j.chemosphere.2020.127743">https://doi.org/10.1016/j.chemosphere.2020.127743</a>
Issue Date	2020-12
Doc URL	<a href="http://hdl.handle.net/2115/86348">http://hdl.handle.net/2115/86348</a>
Rights	© <2020>. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Type	article (author version)
File Information	matsushita-chemosphere_2020.pdf



[Instructions for use](#)

# Effect of chlorination on anti-acetylcholinesterase activity of organophosphorus insecticide solutions and contributions of the parent insecticides and their oxons to the activity

Taku Matsushita<sup>a\*</sup>, Yuki Fujita<sup>b</sup>, Kei Omori<sup>b</sup>, Yuxiang Huang<sup>b</sup>,  
Yoshihiko Matsui<sup>a</sup>, and Nobutaka Shirasaki<sup>a</sup>

<sup>a</sup>Faculty of Engineering, Hokkaido University, N13W8, Sapporo 060-8628, Japan

<sup>b</sup>Graduate School of Engineering, Hokkaido University, N13W8, Sapporo 060-8628, Japan

\*Corresponding author: taku-m@eng.hokudai.ac.jp; +81-11-706-7279

## Abstract

Organophosphorus insecticides are known to be partly transformed to their respective oxons during the chlorination step of drinking water treatment. For most organophosphorus insecticides, the toxicological endpoint for determining acceptable daily intake levels is inhibition of acetylcholinesterase (AChE). Like the parent insecticides, oxons also inhibit AChE, so the presence of oxons in drinking water is also evaluated. However, no attention is paid to the possible presence of transformation products (TPs) other than oxons. In the present study, we determined whether the anti-AChE activity observed for chlorinated solutions of the organophosphorus insecticides malathion and methidathion could be solely attributed to the parent compounds and their oxons. Upon chlorination, both malathion and methidathion were immediately transformed to their oxons; the maximum transformation ratios were 60% and 30%, respectively, indicating that at least 40% and 70% of these compounds were transformed into other TPs. Before chlorination, malathion- and methidathion-containing solutions exhibited little to no anti-AChE activity, but the solutions showed strong activity after chlorination. The contributions of the parent insecticides and their oxons to the activities of the chlorinated samples were calculated from the concentrations of the compounds in the samples and dose–response curves for chemical standards of the compounds. For both the malathion-containing solution and the methidathion-containing solution, the calculated anti-AChE activities were almost the same as the observed activities at every chlorination time. This suggests that the observed activities could be attributed solely to the parent insecticides and their oxons, indicating that other TPs need not be considered.

**Keywords:** Diazinon; dimethoate; drinking water; malathion; methidathion; toxicity.

## 1. Introduction

Around the globe, large amounts of pesticides have been applied to agricultural fields to control crop-damaging insects and weeds. Organophosphorus insecticides are among the most commonly used

pesticides because of their low cost, ready availability, wide-ranging efficacy, and ability to control many pest species (Tankiewicz et al. 2010). In 2017, worldwide sales of organophosphorus insecticides amounted to approximately 2.5 billion USD, second only to the sales of nicotinamides (including neonicotinoid insecticides) at approximately 3.5 billion USD (Phillips 2019). However, organophosphorus insecticides are toxic to humans. The toxicity is induced by inhibition of acetylcholinesterase (AChE) in synapses and on the membranes of red blood cells, which results in accumulation of acetylcholine (ACh) and overstimulation of ACh receptors in synapses of the autonomic nervous system and neuromuscular junctions (Eddleston et al. 2008). According to the Joint FAO/WHO Meeting on Pesticide Residues (WHO 2020), USEPA (2017) and the Food Safety Commission of Japan (FSCJ 2020), the toxicological endpoint for determining the acceptable daily intake levels and chronic reference doses of most organophosphorus insecticides is inhibition of AChE.

After being applied to agricultural fields, organophosphorus insecticides flow into rivers during rain events (Wee and Aris 2017, Derbalah et al. 2019, Xu et al. 2020); and the resulting contaminated river water can be drawn into downstream drinking water treatment plants. Because organophosphorus insecticides are quite difficult to be removed by conventional drinking water treatment processes consisting of coagulation, sedimentation, and sand filtration (Stackelberg et al. 2007, Ormad et al. 2008, Matsushita et al. 2018), these compounds are present in the treated water when it is disinfected with chlorine prior to release into the distribution system. Upon exposure of the insecticides to free chlorine, oxidation of the P=S bond to a P=O bond results in transformation of some of the molecules into their oxons (Duirk et al. 2009, Kamel et al. 2009, Li et al. 2016). Under the circumstances that transformation products (TPs) produced in the environment and water treatments recently attracts increasing attention worldwide (Yin et al. 2017, Kiefer et al. 2019, Zahn et al. 2019), attention is paid not only for the organophosphorus insecticides but also for their oxons in drinking waters, because the oxons inhibit AChE activity as well as their parent insecticides do (Eddleston et al. 2008).

The European Union Drinking Water Directive sets the maximum allowable concentration for any individual pesticide at 0.1 µg/L, and this value applies not only to the active substance but also to its relevant TPs (European Commission 2018). For this purpose, a TP is defined as a compound that is formed either in organisms or in the environment and that “has intrinsic properties comparable to the parent substance in terms of its biological target activity, or...poses a higher or comparable risk to organisms than the parent substance or...has certain toxicological properties that are considered unacceptable” (EN 2009). The oxons of organophosphorus insecticides are included in the regulation because they have the same mode of action as their parent insecticides and are produced and detected in environmental water as well (Duttgupta et al. 2020). According to the Japanese drinking water quality guideline, the oxons of 10 organophosphorus insecticides (butamifos, chlorpyrifos, diazinon, EPN (*O*-ethyl *O*-(4-nitrophenyl) phenylphosphonothioate), fenitrothion, fenthion, isofenphos, isoxathion, malathion, and prothiofos) and the parent insecticides must be quantified in finished water.

However, the percentage of the parent insecticide that is transformed into the oxon, the transformation ratio, is usually less than 100% (Onodera et al. 1995, Li et al. 2016), which means that the remainder of the insecticide is transformed into other TPs. In fact, Magara et al. (1994) reported that chlorination of diazinon produces not only diazinon-oxon but also diethyl phosphate, dichloroacetic acid, and trichloroacetic acid. Even though non-oxon TPs may inhibit AChE, these TPs are unlikely to be taken into account in the European Union regulation and are not taken into account in the Japanese drinking water quality guideline (exceptions include the sulfoxide, sulfone, oxon-sulfoxide, and oxon-sulfone of fenthion, because they are known to inhibit AChE (Tahara et al. 2008)).

Oxons are generated not only by chlorination but also by other water treatment processes, such as ozonation (Wu et al. 2009), UV irradiation (Zamy et al. 2004, Čolović et al. 2010), UV irradiation in the presence of H<sub>2</sub>O<sub>2</sub> (Li et al. 2015), and UV irradiation in the presence of TiO<sub>2</sub> (Bavcon Kralj et al. 2007, Calza et al. 2008). Bavcon Kralj et al. (2007) reported that AChE was inhibited by a UV/TiO<sub>2</sub>-treated malathion-containing solution and implied that the inhibition was due to the oxon of malathion, on the basis of the fact that the anti-AChE activity of the solution depended on the oxon concentration. However, these investigators did not determine whether the pure oxon at the concentrations present in the irradiated samples showed the same inhibitory activity against AChE as did the irradiated samples. Onodera et al. (1995) concluded that AChE inhibition by a chlorinated EPN-containing solution was due to the oxon of EPN, but no evidence was provided for this conclusion. Only Čolović et al. (2010) did the necessary quantitative comparison; specifically, they compared anti-AChE activity by a UV-treated diazinon-containing solution with anti-AChE activity by a chemical standard of diazinon-oxon at the same concentration as that in the UV-treated sample, and they concluded that the anti-AChE activity observed at the beginning of the treatment (after irradiation for 5 min) could be attributed to diazinon-oxon in the UV-treated sample. To our knowledge, no one has quantitatively compared observed anti-AChE activity with anti-AChE activity thought to be derived from an oxon generated during chlorination, leaving it unclear as to whether the toxicities of compounds other than parent insecticides and their oxons in drinking waters after chlorination should be considered.

Accordingly, the objectives of the present study were to investigate whether the activities of insecticides and their oxons could fully explain anti-AChE activity observed upon chlorination of solutions containing organophosphorus insecticides. First, we investigated anti-AChE activities by four organophosphorus insecticides (malathion, methidathion, diazinon, and dimethoate) and their respective oxons by using chemical standards. On the basis of the resulting data, we selected two insecticides (malathion and methidathion) for additional experiments. Solutions containing each insecticide were separately treated with free chlorine, and samples were withdrawn at predetermined intervals. The concentrations of insecticides and their oxons in the samples were determined, and their anti-AChE activities were measured. Finally, the contributions of the parent insecticides and their oxons to observed activity were evaluated by combining their concentrations in the chlorinated

samples with dose–response curves generated with chemical standards. In this way, we were able to clarify whether it is reasonable to consider only parent insecticides and their oxons when assessing contamination of drinking water.

## 2. Materials and Methods

### 2.1 Chemicals

Chemical standards of malathion, methidathion, diazinon, dimethoate, and their respective oxons (Fig. S1) were purchased from Fujifilm Wako Pure Chemical Corporation (Osaka, Japan) and were used as received. To avoid any effect of organic solvents on the experimental results, stock solutions of the insecticides (malathion, 29  $\mu\text{M}$ ; methidathion, 230  $\mu\text{M}$ ; diazinon, 38  $\mu\text{M}$ ; dimethoate, 30  $\mu\text{M}$ ) and oxons (malathion-oxon, diazinon-oxon, and dimethoate-oxon, 100  $\mu\text{M}$ ; methidathion-oxon, 1000  $\mu\text{M}$ ) were prepared in a Milli-Q water-based (Milli-Q Advantage, Millipore Co., Bedford, MA, USA) phosphate buffer (10 mM, pH 7.0). AChE (derived from human erythrocytes) was purchased from Merck KGaA (Darmstadt, Germany). ACh and choline (Ch) were purchased from Fujifilm Wako Pure Chemical Corporation.

### 2.2 Anti-acetylcholinesterase activity assay

An *in vitro* anti-AChE activity assay was conducted with commercially available AChE and ACh according to Liu et al. (2014) with some modification as follows. Phosphate buffer (1 mM, pH 7.4), prepared by dissolving  $\text{Na}_2\text{HPO}_4$  and  $\text{NaH}_2\text{PO}_4$  in Milli-Q water, was supplemented with NaCl at a concentration of 150 mM. Hereafter, this buffer is referred to as the assay buffer. A working solution of AChE was prepared by adding AChE (240 units/L) to the assay buffer, and a working solution of ACh (120  $\mu\text{M}$ ) was prepared by adding ACh to the assay buffer.

Each sample solution was diluted to the desired concentration with the assay buffer, and then 285  $\mu\text{L}$  of the diluted sample was poured into at least three wells of an ice-cooled 96-well microplate. Each sample was supplemented with 7.5  $\mu\text{L}$  of the AChE working solution and then preincubated at 37  $^\circ\text{C}$  for 30 min, during which inhibition of the AChE by compounds in the sample was expected to occur. After the preincubation, the 96-well plates were cooled on ice, and then each sample was supplemented with 7.5  $\mu\text{L}$  of the ACh working solution on ice and then incubated at 37  $^\circ\text{C}$  for 120 min, during which time release of Ch from ACh by uninhibited AChE was expected to occur. After the incubation, an aliquot (200  $\mu\text{L}$ ) of each sample was mixed with 200  $\mu\text{L}$  of acetonitrile to inactivate the AChE. The concentration of Ch in the final solution was measured by liquid chromatography–mass spectrometry (LC-MS). A control (prepared with the assay buffer instead of the diluted sample) and a blank (prepared with the assay buffer instead of both the diluted sample and the AChE working solution) were also subjected to the assay procedure.

Anti-AChE activity by each sample was calculated by means of Eq. (1):

$$\text{Anti-AChE activity} = \frac{[\text{Ch}]_{\text{control}} - [\text{Ch}]_{\text{sample}}}{[\text{Ch}]_{\text{control}} - [\text{Ch}]_{\text{blank}}} \quad (1)$$

where  $[\text{Ch}]_{\text{control}}$ ,  $[\text{Ch}]_{\text{sample}}$ , and  $[\text{Ch}]_{\text{blank}}$  are the Ch concentrations in the control, the sample, and the blank, respectively.

### 2.3 Batch chlorination experiments

Malathion and methidathion were separately dissolved in 10 mM phosphate buffer (pH 7.0) at concentrations of 28 and 230  $\mu\text{M}$ , respectively, in glass vials. For chlorination, these solutions were supplemented with sodium hypochlorite at mol- $\text{Cl}_2$ /mol-C ratios of 1 and 3 for malathion and methidathion, respectively, so that chlorine was present throughout the procedure; and the supplemented samples were magnetically stirred for 5 min. Then the vials were tightly sealed with screw caps and Parafilm to prevent volatilization and allowed to stand at 20 °C in the dark for 168 h. During this period, aliquots were withdrawn at predetermined intervals, quenched with sodium sulfite to remove residual chlorine, and subjected to the anti-AChE activity assay (after 200-fold dilution with the assay buffer). In addition, the concentrations of the parent insecticides and their oxons in the aliquots were determined by LC-MS.

### 2.4 Quantification of the concentrations of choline, parent insecticides, and oxon

The concentration of Ch was measured with a hybrid quadrupole-orbitrap mass spectrometer (Q-Exactive, Thermo Fisher Scientific, Inc., Waltham, MA, USA) coupled with an LC system (UltiMate 3000, Thermo Fisher Scientific). A 5- $\mu\text{L}$  sample of each solution was assayed by LC on a 100 mm  $\times$  2.1 mm Cortecs UPLC HILIC column (1.6- $\mu\text{m}$  particle size, Waters Corporation, Milford, MA, USA). The mobile phase was a binary gradient of 100 mM ammonium formate in Milli-Q water (solvent A) and 100% acetonitrile (solvent B) at a flow rate of 200  $\mu\text{L}/\text{min}$  as follows: begin with 5% B (v/v), increase linearly to 60% B over a period of 0.75 min, hold at that percentage for 0.25 min, decrease linearly to 30% B over a period of 0.25 min, hold at that percentage for 1 min, and then decrease linearly to 5% B over a period of 3.75 min. The mass spectrometer was operated in electrospray-ionization mode (positive) with a spray voltage of 3.2 kV. The temperatures of the capillary heater and the electrospray-ionization-probe heater were 220 and 450 °C, respectively. The flow rates of the sheath gas, auxiliary gas, and sweep gas were 50, 15, and 0 units, respectively. The S-lens ratio frequency level was set to 57. The concentration of Ch was quantified in selected-ion-monitoring mode ( $m/z$  104.1070) at a resolution of 70,000. Detection limit of Ch was 5 nM.

The concentrations of insecticides and oxons were measured with the same apparatus used to measure Ch concentration. A 5- $\mu\text{L}$  sample of each solution was assayed by LC on a 50 mm  $\times$  2.1 mm Hypersil Gold column (1.9- $\mu\text{m}$  particle size, Thermo Fisher Scientific). The mobile phase was a binary gradient of 2 mM ammonium formate in Milli-Q water (solvent A) and 100% methanol (solvent B) at a flow rate of 200  $\mu\text{L}/\text{min}$  as follows: 1% B (v/v) for 1.5 min, increase linearly to 60% B over a period of 1 min, increase linearly to 99% B over a period of 5.5 min, hold at that percentage

for 1.5 min, decrease linearly to 1% B over a period of 0.5 min, and then hold at that percentage for 2.0 min. The mass spectrometer was operated in electrospray-ionization mode (positive) with a spray voltage of 3.2 kV. The temperatures of the capillary heater and the electrospray-ionization-probe heater were 220 and 450 °C, respectively. The flow rates of the sheath gas, auxiliary gas, and sweep gas were 50, 15, and 0 units, respectively. The S-lens ratio frequency level was set to 57. The concentrations of malathion, malathion-oxon, methidathion, and methidathion-oxon were quantified in selected-ion-monitoring mode ( $m/z$  331.0433, 315.0662, 302.9691, and 286.9920, respectively) at a resolution of 70,000. Detection limits of malathion, malathion-oxon, methidathion, and methidathion-oxon were 3, 2, 0.3, and 2 nM, respectively.

### 3. Results and Discussion

#### 3.1 Comparison of anti-acetylcholinesterase activities of insecticides and oxons

First, we assayed the anti-AChE activities of chemical standards of the four parent insecticides and their respective oxons (Fig. 1). Among the insecticides, only malathion showed relatively strong anti-AChE activity; methidathion and diazinon showed only slight activity, and dimethoate showed no activity at all in the tested concentration range. In contrast, all the oxons showed dose-dependent anti-AChE activity, and in all cases the oxons were more active than the respective parent insecticides. To compare the activities of the compounds, we calculated median half maximal inhibitory concentrations ( $IC_{50}$ ) by using logistic regression analysis (Table 1). The  $IC_{50}$  values of the parent insecticides were more than 3 orders of magnitude higher than those of their respective oxons, indicating that the oxons were >1,000 times as active as the parent insecticides. This tendency is consistent with published data obtained by means of *in vitro* assays (Tahara et al. 2005, Krstić et al. 2008, Čolović et al. 2010, Čolović et al. 2011), although the  $IC_{50}$  values were different among literature probably due to the differences in experimental conditions as shown in Table S1. The activities of the compounds differed considerably and decreased in the order malathion-oxon > diazinon-oxon  $\approx$  methidathion-oxon  $\gg$  dimethoate-oxon  $\approx$  malathion > methidathion  $\approx$  diazinon > dimethoate. Notably, the activity of dimethoate-oxon was as low as that of malathion, even though the former was an oxon.

#### 3.2 Transformation of malathion and methidathion to their oxons during chlorination

We investigated the effects of chlorination on the activities of solutions of malathion and methidathion because of the relatively high anti-AChE activities of the oxons of these two insecticides. During chlorination, malathion immediately reacted with free chlorine (Fig. 2a). The malathion almost completely disappeared within 15 min. At that point, the malathion-oxon concentration was equivalent to 60% of the initial malathion concentration, and the concentration of the oxon then gradually decreased with chlorination time. The transformation of malathion to its oxon during chlorination that we observed is consistent with that previously reported (Duirk et al. 2009, Li et al. 2016).

Likewise, methidathion reacted with free chlorine immediately, disappearing almost completely within 10 min, and methidathion-oxon was produced by the reaction, as has previously been reported (Duirk et al. 2009, Kamel et al. 2009). However, the ratio of the oxon concentration to the initial methidathion concentration (30%) was half the corresponding ratio for the malathion oxon. The lower transformation ratio for methidathion may have been due partly to differences in the initial insecticide and free chlorine concentrations: both concentrations were necessarily higher for methidathion than for malathion because the anti-AChE activity of methidathion-oxon was lower than that of malathion-oxon. This possibility was confirmed by an experiment showing that when the initial concentrations of methidathion and free chlorine were decreased to 33 nM (from 230  $\mu$ M) and to 1 mg-Cl<sub>2</sub>/L, the ratio increased to 83% (Fig. S2).

### 3.3 Effect of chlorination on anti-acetylcholinesterase activity by insecticide-containing solutions

Before chlorination, the malathion-containing solution slightly inhibited AChE (Fig. 3a). Upon exposure to free chlorine, the anti-AChE activity of the solution rapidly increased and then gradually decreased with chlorination time. In contrast, the methidathion-containing solution showed no activity before chlorination (Fig. 3b), but upon exposure to free chlorine, the solution showed high activity; then the activity rapidly decreased, and no activity was observed after 24 h of chlorination. Our results are consistent with previously reported chlorination-induced increases in the anti-AChE activity of solutions containing other organophosphorus insecticides, such as EPN (Onodera et al. 1995) and fenthion (Tahara et al. 2008). Our findings clearly show that both malathion and methidathion were converted to their respective TPs and that some of the TPs showed anti-AChE activity. During chlorination, the parent insecticides were transformed to their oxons (Fig. 2), and the oxons were more active than the parent insecticides (Fig. 1). Accordingly, transformation of the insecticides to their oxons probably contributed to the increase in activity.

However, as mentioned above, only 60% of the initial malathion and 30% of the initial methidathion were transformed to their oxons, which means that the remainder of the initial amounts of the two insecticides were transformed into TPs other than oxons. Duirk et al. (2009) reported that organophosphorus insecticides are oxidized to their oxons by HOCl but are hydrolyzed by OCl<sup>-</sup>. In fact, we detected dimethyl phosphate, a hydrolyte of methidathion-oxon, during chlorination of the methidathion-containing solution (Fig. S3). This compound may also have been produced from malathion, judging from the fact that both insecticides have an *O,O*-dimethyl dithiophosphate group (Fig. S1). Therefore, we assayed the anti-AChE activity of a chemical standard of dimethyl phosphate; the assay showed that this TP had low anti-AChE activity (Fig. S4) and thus that it did not contribute to the observed activity of the chlorinated solutions. Nevertheless, other, as-yet-unidentified TPs with anti-AChE activity may have contributed to the observed activity.

### 3.4 Contributions of parent insecticides and their oxons to anti-acetylcholinesterase activity by

### *chlorinated insecticide-containing solutions*

Because the dose–response curves for the activities of the parent insecticides and their oxons were not linear (Fig. 1), the inhibition was not dose-additive; that is, the activity of a solution containing both the insecticide and its oxon could not be calculated simply by summing the individual activities of the insecticide and its oxon. To estimate the total anti-AChE activity induced by the parent insecticides and their oxons in the samples, a mixture containing the same concentrations of the parent insecticides and their oxons as those in the chlorination samples should be prepared and then subjected to the anti-AChE activity assay (Krstić et al. 2008, Čolović et al. 2011). However, before chlorination (chlorination time = 0 h), the solutions contained only the parent insecticides and no oxons; whereas after chlorination, the solutions contained only the oxons and no parent insecticides. Therefore, we could determine the anti-AChE activity of the parent insecticide from the data for the sample before chlorination, and we could determine the activity of the oxon from the data for the sample after chlorination. For the unchlorinated solutions, we calculated the activity of the parent insecticide from its concentration in the initial sample (Fig. 2) and the dose–response curve (Fig. 1), being sure to take into account the dilution factor in the anti-AChE activity assay (200-fold). For the solutions after chlorination, we used the same procedure to calculate the activity of the oxon at every chlorination time.

Comparison of the observed and calculated anti-AChE activities (Fig. 4) revealed that the observed activity of the malathion-containing solution before chlorination was roughly the same as the activity calculated as described above, indicating that the observed activity of the unchlorinated solution was attributable malathion. After treatment with free chlorine, the observed activity was almost the same as the calculated activity at every chlorination time, which clearly indicates that the observed activity was due to malathion-oxon and that the other, as-yet-unidentified TPs did not inhibit AChE. As mentioned in the introduction, the Japanese drinking water quality guideline requires both malathion and its oxon to be measured. Our current findings indicate that this guideline is reasonable for regulating malathion in drinking water.

Likewise, the lack of anti-AChE activity of the methidathion solution before chlorination is consistent with the activity calculated from the methidathion concentration in the sample and the dose–response curve (Fig. 4b). After treatment with free chlorine, the observed and calculated activities were still almost the same at every chlorination time. This comparison clearly indicates that the activity of the chlorinated solution was due only to methidathion-oxon and not to any other TPs. The current Japanese drinking water quality guideline does not require quantification of the oxon of methidathion, even though methidathion must be quantified; the reason for the difference between the two compounds is unclear. Our results suggest the oxon of methidathion should be quantified.

Krstić et al. (2008) reported that antagonistic inhibition effects were observed when AChE was simultaneously exposed to malathion-oxon and isomalathion, both of which had anti-AChE activity,

under relatively high concentrations. Therefore, in the present study, some unidentified TPs having anti-AChE activity might have been produced, but their contributions to the observed anti-AChE activity might not have been recognized because of the antagonistic effect with oxons having strong anti-AChE activity.

The crop protection market has changed over the past two decades. Specifically, Europe and North America decreased their shares in 2018, and the shares attributable to Asia and Central and South America increased. The increase was driven in large part by developing countries in the latter regions (Phillips 2019). Explosive population growth in these developing countries will increase the need for food production and thus will most likely further increase the market shares attributable to Asia and Central and South America, and African countries may follow them. Our findings revealed that the anti-AChE activities of malathion- and methidathion-containing solutions after chlorination are due solely to the parent insecticides and their oxons. Accordingly, if developing countries already regulate or plan to regulate malathion and methidathion, we recommend that the oxons also be regulated.

#### **4. Conclusions**

1. The anti-AChE activities of the oxons of the four insecticides evaluated in this study were >1,000 times the activities of their respective parent insecticides. Anti-AChE activity decreased in the order malathion-oxon > diazinon-oxon  $\approx$  methidathion-oxon  $\gg$  dimethoate-oxon  $\approx$  malathion > methidathion  $\approx$  diazinon > dimethoate.
2. Both malathion and methidathion immediately reacted with free chlorine, and their respective oxons were produced. The concentrations of the oxons decreased with increasing chlorination time.
3. Solutions of malathion and methidathion showed little or no anti-AChE activity before chlorination. However, chlorination resulted in a rapid increase in the activities of the solutions, but thereafter the activities decreased with increasing chlorination time.
4. The activities of the chlorinated malathion- and methidathion-containing solutions were attributable solely to the parent insecticides and their respective oxons. No other TPs were shown to inhibit AChE.
5. If any countries have already incorporated or plan to incorporate malathion and methidathion into regulatory guidelines, the oxons of these two insecticides should also be included.

#### **Acknowledgments**

This research was supported in part by a Health and Labor Sciences Research Grant (no. 19LA0501) from the Ministry of Health, Labor, and Welfare of Japan and by a grant from the Fuso Innovative Technology Fund.

## References

- Bavcon Kralj, M., Černigoj, U., Franko, M. and Trebše, P. (2007) Comparison of photocatalysis and photolysis of malathion, isomalathion, malaoxon, and commercial malathion—Products and toxicity studies. *Water Research* 41(19), 4504-4514.
- Calza, P., Massolino, C. and Pelizzetti, E. (2008) Light induced transformations of selected organophosphorus pesticides on titanium dioxide: Pathways and by-products evaluation using LC–MS technique. *Journal of Photochemistry and Photobiology A: Chemistry* 199(1), 42-49.
- Čolović, M., Krstić, D., Petrović, S., Leskovac, A., Joksić, G., Savić, J., Franko, M., Trebše, P. and Vasić, V. (2010) Toxic effects of diazinon and its photodegradation products. *Toxicology Letters* 193(1), 9-18.
- Čolović, M.B., Krstić, D.Z., Ušćumlić, G.S. and Vasić, V.M. (2011) Single and simultaneous exposure of acetylcholinesterase to diazinon, chlorpyrifos and their photodegradation products. *Pesticide Biochemistry and Physiology* 100(1), 16-22.
- Derbalah, A., Chidya, R., Jadoon, W. and Sakugawa, H. (2019) Temporal trends in organophosphorus pesticides use and concentrations in river water in Japan, and risk assessment. *Journal of Environmental Sciences* 79, 135-152.
- Duirk, S.E., Desetto, L.M. and Davis, G.M. (2009) Transformation of Organophosphorus Pesticides in the Presence of Aqueous Chlorine: Kinetics, Pathways, and Structure–Activity Relationships. *Environmental Science & Technology* 43(7), 2335-2340.
- Duttagupta, S., Mukherjee, A., Bhattacharya, A. and Bhattacharya, J. (2020) Wide exposure of persistent organic pollutants (PoPs) in natural waters and sediments of the densely populated Western Bengal basin, India. *Science of The Total Environment* 717, 137187.
- Eddleston, M., Buckley, N.A., Eyer, P. and Dawson, A.H. (2008) Management of acute organophosphorus pesticide poisoning. *The Lancet* 371(9612), 597-607.
- EN (2009) Regulation (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC. *Official Journal of the European Union* 52, 1-50.
- European Commission (2018) Annexes to the Proposal for a Directive of the European Parliament and of the Council on the quality of water intended for human consumption (recast).
- FSCJ (2020) Risk assessment reports for pesticides, [http://www.fsc.go.jp/english/evaluationreports/agrichemicals1\\_e1.html](http://www.fsc.go.jp/english/evaluationreports/agrichemicals1_e1.html), visited on 9/April/2020.
- Kamel, A., Byrne, C., Vigo, C., Ferrario, J., Stafford, C., Verdin, G., Siegelman, F., Knizner, S. and Hetrick, J. (2009) Oxidation of selected organophosphate pesticides during chlorination of simulated drinking water. *Water Research* 43(2), 522-534.
- Kiefer, K., Müller, A., Singer, H. and Hollender, J. (2019) New relevant pesticide transformation products in groundwater detected using target and suspect screening for agricultural and urban micropollutants with LC-HRMS. *Water Research* 165, 114972.

- Krstić, D.Z., Čolović, M., Bavcon kralj, M., Franko, M., Krinulović, K., Trebše, P. and Vasić, V. (2008) Inhibition of AChE by malathion and some structurally similar compounds. *Journal of Enzyme Inhibition and Medicinal Chemistry* 23(4), 562-573.
- Li, W., Liu, Y., Duan, J., van Leeuwen, J. and Saint, C.P. (2015) UV and UV/H<sub>2</sub>O<sub>2</sub> treatment of diazinon and its influence on disinfection byproduct formation following chlorination. *Chemical Engineering Journal* 274, 39-49.
- Li, W., Wu, R., Duan, J., Saint, C.P. and van Leeuwen, J. (2016) Impact of prechlorination on organophosphorus pesticides during drinking water treatment: Removal and transformation to toxic oxon byproducts. *Water Research* 105, 1-10.
- Liu, W., Yang, Y., Cheng, X., Gong, C., Li, S., He, D., Yang, L., Wang, Z. and Wang, C. (2014) Rapid and sensitive detection of the inhibitive activities of acetyl- and butyryl-cholinesterases inhibitors by UPLC–ESI-MS/MS. *Journal of Pharmaceutical and Biomedical Analysis* 94, 215-220.
- Magara, Y., Aizawa, T., Matsumoto, N. and Souna, F. (1994) Degradation of pesticides by chlorination during water purification. *Water Science & Technology* 30(7), 119-128.
- Matsushita, T., Morimoto, A., Kuriyama, T., Matsumoto, E., Matsui, Y., Shirasaki, N., Kondo, T., Takanashi, H. and Kameya, T. (2018) Removals of pesticides and pesticide transformation products during drinking water treatment processes and their impact on mutagen formation potential after chlorination. *Water Research* 138, 67-76.
- Onodera, S., Hirose, Y. and Ishikura, S. (1995) Changes in Genetic and Enzymatic Inhibiting Effects of Organophosphorus Pesticides during Aqueous Chlorination. *Journal of Environmental Chemistry* 5(1), 65-71.
- Ormad, M.P., Miguel, N., Claver, A., Matesanz, J.M. and Ovelheiro, J.L. (2008) Pesticides removal in the process of drinking water production. *Chemosphere* 71(1), 97-106.
- Phillips, M.W.A. (2019) Agrochemical industry development, trends in R&D and the impact of regulation. *Pest Management Science*.
- Stackelberg, P.E., Gibs, J., Furlong, E.T., Meyer, M.T., Zaugg, S.D. and Lippincott, R.L. (2007) Efficiency of conventional drinking-water-treatment processes in removal of pharmaceuticals and other organic compounds. *Science of The Total Environment* 377(2), 255-272.
- Tahara, M., Kubota, R., Nakazawa, H., Tokunaga, H. and Nishimura, T. (2005) Use of cholinesterase activity as an indicator for the effects of combinations of organophosphorus pesticides in water from environmental sources. *Water Research* 39(20), 5112-5118.
- Tahara, M., Kubota, R., Tokunaga, H., Nakazawa, H. and Nishimura, T. (2008) The behaviour and cholinesterase inhibitory activity of fenthion and its products by light and chlorination. *Journal of Water Supply: Research and Technology-Aqua* 57(3), 143-151.
- Tankiewicz, M., Fenik, J. and Biziuk, M. (2010) Determination of organophosphorus and organonitrogen pesticides in water samples. *TrAC Trends in Analytical Chemistry* 29(9), 1050-1063.
- USEPA (2017) Human health benchmarks for pesticides: update 2017 technical document.
- Wee, S.Y. and Aris, A.Z. (2017) Ecological risk estimation of organophosphorus pesticides in riverine

ecosystems. *Chemosphere* 188, 575-581.

WHO (2020) Inventory of evaluations performed by the Joint Meeting on Pesticide Residues (JMPR), <https://apps.who.int/pesticide-residues-jmpr-database>, visited on 9/April/2020.

Wu, J., Lan, C. and Chan, G.Y.S. (2009) Organophosphorus pesticide ozonation and formation of oxon intermediates. *Chemosphere* 76(9), 1308-1314.

Xu, L., Granger, C., Dong, H., Mao, Y., Duan, S., Li, J. and Qiang, Z. (2020) Occurrences of 29 pesticides in the Huangpu River, China: Highest ecological risk identified in Shanghai metropolitan area. *Chemosphere* 251, 126411.

Yin, L., Wang, B., Yuan, H., Deng, S., Huang, J., Wang, Y. and Yu, G. (2017) Pay special attention to the transformation products of PPCPs in environment. *Emerging Contaminants* 3(2), 69-75.

Zahn, D., Mucha, P., Zilles, V., Touffet, A., Gallard, H., Knepper, T.P. and Frömel, T. (2019) Identification of potentially mobile and persistent transformation products of REACH-registered chemicals and their occurrence in surface waters. *Water Research* 150, 86-96.

Zamy, C., Mazellier, P. and Legube, B. (2004) Phototransformation of selected organophosphorus pesticides in dilute aqueous solutions. *Water Research* 38(9), 2305-2314.




Fig. 1 Concentration dependence of AChE inhibition by pesticides (white triangles) and their oxons (black circles).




Fig. 2 Effect of chlorination time on the concentrations of (a) malathion and (b) methidathion (white triangles) and their respective oxons (black circles) relative to the initial concentrations of the parent insecticides. The initial concentrations of malathion and methidathion were 28 and 230  $\mu\text{M}$ , respectively. The concentrations were measured at 0, 0.25, 0.5, 1, 3, 12, 24, 48, 72, 96, 120, and 168 h for malathion, and at 0, 0.17, 1, 3, 6, 9, 12, 24, 48, 72, and 168 h for methidathion. Error bars indicate standard deviations of three measurements.




Fig. 3 Effect of chlorination time on AChE inhibition by (a) malathion and (b) methidathion. Error bars indicate standard deviations of three measurements.




Fig. 4 Comparison of chlorination time dependence of (gray) observed anti-AChE activities of chlorinated solutions and (black) activities calculated from insecticide/oxon concentrations and the dose-response curves for (a) malathion and (b) methidathion. Error bars indicate standard deviations of three measurements.

Supplementary information for:

**Effect of chlorination on anti-acetylcholinesterase activity of organophosphorus insecticide solutions and contributions of the parent insecticides and their oxons to the activity**

Taku Matsushita<sup>a\*</sup>, Yuki Fujita<sup>b</sup>, Kei Omori<sup>b</sup>, Yuxiang Huang<sup>b</sup>,  
Yoshihiko Matsui<sup>a</sup>, and Nobutaka Shirasaki<sup>a</sup>

<sup>a</sup>Faculty of Engineering, Hokkaido University, N13W8, Sapporo 060-8628, Japan

<sup>b</sup>Graduate School of Engineering, Hokkaido University, N13W8, Sapporo 060-8628, Japan

\*Corresponding author: taku-m@eng.hokudai.ac.jp; +81-11-706-7279




Fig. S1 Target pesticides and their respective oxons.

Table S1 Differences in experimental conditions for anti-ChE activity assay among published literature.

	Incubation time	Enzyme	Substrate
Krstić et al. (2008)	not described	Bovine erythrocyte AChE	Acetylthiocholine iodide
Tahara et al. (2005)	7 min	Butyrylcholinesterase	5-Methyl-2-thenoylthiocholine iodide
Čolović et al. (2010)	20 min	Human erythrocyte AChE	Acetylthiocholine iodide
Čolović et al. (2011)	20 min	Electric eel AChE	Acetylthiocholine iodide
This study	120 min	Human erythrocyte AChE	Acetylcholine chloride




Fig. S2 Effect of chlorination time on the methidathion (white triangles) and methidathion-oxon (black circles) concentrations relative to the initial methidathion concentration (33 nM). The initial free chlorine concentration was 1 mg-Cl<sub>2</sub>/L.




Fig. S3 Effect of chlorination time on the dimethyl phosphate concentration relative to the initial concentration of methidathion. Error bars indicate standard deviations of three measurements.




Fig. S4 AChE inhibition activity of dimethyl phosphate.