



Title	A new brackish tanaidacean, <i>Sinelobus kisui</i> sp. nov. (Crustacea, Peracarida, Tanaidacea), from Japan, with a key to <i>Sinelobus</i> species and barcode information from two loci
Author(s)	Hirano, Kyoko; Kakui, Keiichi
Citation	Zoosystematics and Evolution, 98(2), 245-256 https://doi.org/10.3897/zse.98.84818
Issue Date	2022-06-29
Doc URL	http://hdl.handle.net/2115/86485
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	article
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	2022_Hirano_Kakui_Sinelobus.pdf ()



[Instructions for use](#)

A new brackish tanaidacean, *Sinelobus kisui* sp. nov. (Crustacea, Peracarida, Tanaidacea), from Japan, with a key to *Sinelobus* species and barcode information from two loci

Kyoko Hirano¹, Keiichi Kakui²

¹ Department of Biological Sciences, School of Science, Hokkaido University, Sapporo 060-0810, Japan

² Department of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

<https://zoobank.org/506932DF-85C7-491E-AA6E-C832996FC66D>

Corresponding author: Keiichi Kakui (kakui@eis.hokudai.ac.jp)

Academic editor: Luiz F. Andrade ♦ Received 28 April 2022 ♦ Accepted 10 June 2022 ♦ Published 29 June 2022

Abstract

We describe the new brackish tanaidid species *Sinelobus kisui* sp. nov. from Hagi, Yamaguchi, Japan. *Sinelobus kisui* is similar to *S. barretti* and *S. vanhaareni* in having antennal article 2 with one outer distal seta, the dorsodistal crotchet on pereopods 2 and 3 carpi shorter than half propodus length, and pereopodal carpi 2–6 with five distal crotchets, but differs from them in having (1) the inner of two ventro-subdistal circumplumose setae on the maxillipedal endite longer than the outer; (2) the maxillipedal endite with one mid-inner spiniform seta; (3) the pereopod-1 propodus with one middle setulate seta; and (4) the pleopod-1 protopod lacking inner plumose setae. Our study confirmed that character states of the chelipeds in strongly dimorphic males are useful in *Sinelobus* taxonomy. We determined partial sequences for the cytochrome *c* oxidase subunit I (COI; *cox1*) and 18S rRNA (18S) genes in *S. kisui* for future DNA barcoding and phylogenetic analyses. Morphological and/or molecular data reveal that *S. kisui* also occurs in Kagawa and Osaka, Japan. A key to species in *Sinelobus* is provided.

Key Words

Integrative taxonomy, intraspecific variation, non-marine, Tanaididae, tube dweller

Introduction

The brackish genus *Sinelobus* Sieg, 1980 is one of 20 genera in Tanaididae Nobili, 1906. After Sieg's (1980) revision of Tanaididae, the genus was regarded as monotypic, containing only the single cosmopolitan species *Sinelobus stanfordi* (Richardson, 1901), originally described from Clipperton Island in the eastern Pacific Ocean. Since 2008, however, five additional species have been described: *Sinelobus barretti* Edgar, 2008 (Tasmania), *Sinelobus bathykolpos* Bamber, 2014 (Hong Kong), *Sinelobus pinkenba* Bamber, 2008 (Australia), *Sinelobus stromatoliticus* Rishworth, Perissinotto & Błażewicz, 2018 (South Africa), and *Sinelobus vanhaareni* Bamber, 2014 (Netherlands) (Fig. 1a).

Sinelobus tanaidaceans are benthic tube-dwellers, lack a planktonic larval stage, and inhabit brackish water (note:

there are some records from freshwater environments; e.g., Stephensen 1936), suggesting that they have low dispersibility, especially across oceans. It is thus likely that this genus contains many locally endemic species that have not yet been detected. They have been reported from various localities in Japan, either identified as *S. stanfordi* or not identified to species level (e.g., Stephensen 1936, Ariyama and Ohtani 1990, Kakui 2022) (Fig. 1b).

We investigated *Sinelobus* tanaidaceans collected from Hagi, Yamaguchi, Japan and concluded they represent an undescribed species. Here we describe this species as new, present partial sequences for the mitochondrial cytochrome *c* oxidase subunit I (COI) and nuclear 18S rRNA (18S) genes for future DNA barcoding and phylogenetic analyses, and provide a key to the species in *Sinelobus*.

Materials and methods

Sinelobus specimens were collected among tubular *Ulva* algae in a brackish stream (no salinity data) at Hagi, Yamaguchi, Japan on 22 May 2014 (Fig. 1). Some individuals were maintained in a small aquarium (20 °C; 14 h light/10 h dark; 3‰ salinity; fed porphyridized dry feed for crayfish [JAN code 4971618829092; Kyorin, Japan] every 3 days), and one female and one male hatched in the aquarium were photographed live to document their body pigmentation pattern. All other individuals were fixed and preserved in 70–99% ethanol. We also obtained *Sinelobus* specimens from two other Japanese localities, Kagawa (collected by Yoshihiro Hayashi) and Osaka (collected by Michio Ohtani). The Kagawa individuals (= *Sinelobus* sp. 1 sensu

Kakui et al. [2011]) came from the mouth of the Aya River on 2 June 2008 and were fixed in 99% ethanol. The Osaka individuals were collected at Station 4 in the Shin Yodo River on 19 February 1989 and fixed in 10% formalin (see Ariyama and Ohtani [1990] for additional details). The methods used for dissection, preparation of slides, light microscopy, and drawing were as described by Kakui and Angsupanich (2012). The specimens studied were deposited in the Invertebrate Collection of the Hokkaido University Museum (ICHUM), Sapporo, Japan, and the Osaka Museum of Natural History (OMNH), Osaka, Japan.

Orientation and terminology here follow Larsen (2003), except that the term “plumose sensory seta(e)” (PSS; Bird 2011) is used instead of “broom seta(e),” and the term “protopod” is used instead of “basal article” for pleopods




Figure 1. Sampling sites and type localities for *Sinelobus* species. **a.** Map showing the type localities for seven *Sinelobus* species, including *Sinelobus kisui* sp. nov.; **b.** Map showing the sampling site at Hagi, Yamaguchi and previously reported sites for *Sinelobus* individuals in Japan (*, Kochi, where *Sinelobus* sp. 2 sensu Kakui et al. [2011] was collected); **c.** Location of the sampling site for *Sinelobus kisui* in Hagi; **d.** Photograph showing the point of sampling in a brackish stream in Hagi. Maps were generated by using GMT6 (a, b, Wessel et al. 2019) or were based on GSI Maps (c. Geospatial Information Authority of Japan 2022). Sources and raw data for the plots are available in the Figshare repository (Kakui 2022).

and uropods. Three other setal terms used are: “crotchet(s)” (Bird 2019), “flattened denticulate seta(e)” (Edgar 2008), and “step-tipped plumose seta(e)” (Kakui et al. 2010). Body length (BL) was measured from the base of the antennules to the tip of the pleotelson, and body width at the widest portion of the cephalothorax (CW, cephalothorax width). Appendages were measured in the holotype and allotype specimens. Measurements were made axially with ImageJ (Rasband 2022) from digital images: dorsally on the body, antennules, antennae, and uropods; laterally on the pereopods and pleopods. Unless otherwise noted, all measurements in the text are in millimeters.

DNA was extracted from three Yamaguchi specimens by using a NucleoSpin Tissue XS Kit (Macherey-Nagel, Germany). For PCR amplification, primers used for COI were LCO1490 (Folmer et al. 1994) and HCOoutout (Prendini et al. 2005); these two and HCO2198 (Folmer et al. 1994) were used in cycle sequencing. Amplification primers for 18S were 18S-a1F and 18S-a9R (Kakui et al. 2011); seven primers (18S-b3F, 18S-b4F, 18S-b4R, 18S-b6F, 18S-a6R, 18S-b8R, and 18S-b8F; Kakui et al. 2011, Kakui and Shimada 2017) were used in cycle sequencing. PCR amplification conditions for COI with TaKaRa Ex Taq DNA polymerase (TaKaRa Bio, Japan) were 94 °C for 1 min; 35 cycles of 98 °C for 10 s, 42 °C for 30 s, and 72 °C for 1 min; and 72 °C for 2 min. Amplification conditions for 18S with KOD FX Neo (Toyobo, Japan) were 94 °C for 2 min; 45 cycles of 98 °C for 10 s, 52 °C for 30 s, and 68 °C for 75 s; and 68 °C for 3 min.

Nucleotide sequences were determined by direct sequencing with a Big Dye Terminator Kit ver. 3.1 and a 3730 DNA Analyzer (Life Technologies, USA). Fragments were concatenated by using MEGA7 (Kumar et al. 2016). Two previously determined, unpublished COI sequences from two Kagawa *Sinelobus* individuals were included in this study; the method used to determine these sequences was described by Okamoto et al. (2020). The sequences we determined were deposited in the International Nucleotide Sequence Database (INSD) through the DNA Data Bank of Japan. Kimura (1980) 2-parameter (K2P) distances among *Sinelobus* COI sequences were calculated with MEGA7.

Results

Taxonomy

Family Tanaididae Nobili, 1906

Genus *Sinelobus* Sieg, 1980

[New Japanese name: *Kisui-tanaisu-zoku*]

Remarks. *Sinelobus* differs from confamilial genera in the following combination of characters: pereonite 1 not extended anterolaterally; four free pleonites; dorso-transverse plumose setal row on pleonites 1 and 2 present but incomplete (absent in mid-dorsal region); antennal article 4 without distal setal tuft; antennal article

5 at least 0.6 times length of article 4; labial outer lobe without terminal process; and prominent sexual dimorphism (Sieg 1980, Bamber 2005, Bamber and Boxshall 2006, Chim and Tong 2019).

Sinelobus kisui sp. nov.

<https://zoobank.org/FA0362B0-1A05-41B3-AC84-DA36C622D3BA>

[Japanese name: *Kisui-tanaisu*]

Figs 2–6

Sinelobus stanfordi: Ariyama and Ohtani 1990, 25–26, figs 2–5; Kakui 2016, 610 (not *Sinelobus stanfordi* Richardson, 1901).

Sinelobus sp. 1: Kakui et al. 2011, 751; Kakui et al. 2012, 128–129, fig. 1D.

Sinelobus sp.: Kaji et al. 2016, 3–8, figs 1–5; Kakui et al. 2021, 3.

Diagnosis. Antennal article 2 with outer distal simple seta. Maxillipedal endite with one mid-inner and two dorsodistal spiniform setae; and two ventro-subdistal circumplumose setae, inner one longer than outer. Pereopod 1 with carpus bearing two dorsal and two ventral simple setae; propodus with middle setulate seta. Dorsodistal crotchet on carpi of pereopods 2 and 3 shorter than half propodus length. Pereopod 3 basis with ventral PSS. Pereopods 2–6 with carpus bearing five distal crotchets. Pleopod 1 protopod without inner plumose setae. In strongly sexually dimorphic males, chelipedal merus with three ventral processes; angle of dorsal margin of fixed finger to distal margin of palm about 90°.

Etymology. The specific name is derived from the Japanese noun *kisui* (brackish water), referring to the habitat of this species, and also referable to the Japanese name *Kisui-tanaisu*, proposed by Ariyama and Ohtani (1990) for this species in the Shin Yodo River, Osaka.

Material examined. Holotype: Female, ICHUM8301 (BL 4.00, CW 0.67), Hagi, Yamaguchi, Japan (34°26.168'N, 131°25.140'E) (type locality), among tubular *Ulva* algae, brackish stream, 22 May 2014, coll. by Keiichi Kakui. **Allotype:** Male, ICHUM8302 (BL 3.17, CW 0.62), same collection data as for holotype. **Paratypes:** Five females (ICHUM8303, BL 3.06, CW 0.56; ICHUM8304, BL 2.91, CW 0.52; ICHUM8305, CW 0.51, INSD accession number LC705431 [COI]; ICHUM8306, BL 3.18, CW 0.52; ICHUM8312, BL 2.10, CW 0.42, INSD accession numbers LC705430 [18S], LC705432 [COI]). Six males (ICHUM8307, BL 3.04, CW 0.56; ICHUM8308, BL 2.85, CW 0.56, INSD accession number LC705433 [COI]; ICHUM8309, BL 2.89, CW 0.52; ICHUM8310, BL 3.36, CW 0.61; ICHUM8311, BL 2.42, CW 0.46; ICHUM8313, BL 2.46, CW 0.53). Collection data for ICHUM8303–8311 same as for holotype. ICHUM8312 and 8313 hatched in an aquarium in Kakui laboratory, Sapporo, descendants of individuals collected from type locality on 22 May 2014.

Other material: One female, ICHUM8314 (BL 2.79, CW 0.51; INSD accession number LC664100 [COI]), Aya River, Sakaide, Kagawa, Japan (approximate coordinates



Figure 2. *Sinelobus kisui* sp. nov., paratype female (a. ICHUM8312) and male (b. ICHUM8313), dorsal views, fresh specimens. Scale bars: 0.5 mm.

34°19.693'N, 133°52.499'E), 2 June 2008, coll. by Yoshihiro Hayashi; one male, ICHUM4031 (INSD accession numbers AB618192 [18S], LC664101 [COI]), same collection data as for ICHUM8314; one male, OMNH-Ar12459 (BL 2.97, CW 0.67), Station 4, Shin Yodo River, Osaka, Osaka, Japan

(approximate coordinates: 34°41.447'N, 135°26.613'E), under rocks, 19 February 1989, coll. by Michio Ohtani.

Description of females (based on holotype unless noted otherwise). Body (Figs 2a, 3a, b) 5.96 times as long as wide, with brown pigmentation (retained in eth-




Figure 3. *Sinelobus kisui* sp. nov. **a, b.** Body, female holotype, dorsal (**a**) and left (**b**) views; **c, d.** Body, male allotype, dorsal (**c**) and left (**d**) views. Scale bars: 1 mm.

anol). Cephalothorax 0.19 times BL, tapering anteriorly, with mid-lateral pair of simple setae and pair of simple setae posterior to eyes. Dorsal pigmentation pattern on carapace: anterior region dark; subanterior region without dark pigmentation; posterolateral region with two dark diagonal bands; reticulate pattern of pigmentation elsewhere (often faint). Pereonites 1–6 with length ratio 1.00:1.71:2.07:2.50:2.58:2.20; all with one or two pairs of simple setae. Pereonites 1–6 width-to-length ratio 0.30, 0.51, 0.63, 0.82, 0.86, and 0.80, respectively. Pleo-

nites 1–3 with pair of dorsal simple seta; pleonites 1 and 2 with dorso-transverse plumose setal row (but incomplete, absent in dorsal region); pleonites 2 and 3 with lateral plumose setal row. Pleonite 4 with several pairs of simple setae. Pleotelson with seven pairs of simple setae.

Antennule (Fig. 4a) 0.65 times as long as cephalothorax; articles 1–4 with length ratio 1.00:0.37:0.33:0.03. Article 1 with two outer distal and one inner distal simple setae, and several proximal and distal PSS. Article 2 with three outer distal and two inner distal simple setae, and




Figure 4. *Sinelobus kisui* sp. nov. **a, c, e–k, m, n.** Female holotype; **b, d.** Male allotype; **l.** Female paratype (ICHUM8304); **a, b.** Right (**a**) and left (**b**) antennules, dorsal views; **c, d.** Left antenna, dorsal (**c**) and outer (**d**) views; **e.** Labrum, lateral view; **f, g.** Left mandible, posterior (**f**) and inner (**g**) views; **h.** Right mandible, inner view; **i.** Right molar; **j.** Labium; **k.** Right maxillule, palp broken; **l.** Left maxilla; **m.** Maxillipeds, dorsal view (right palp omitted; outer portion of right endite bent inward); **n.** Left epignath. Scale bars: 0.1 mm.

several distal PSS. Article 3 with two distal simple setae and several distal PSS. Article 4 with tuft of distal simple setae, distal PSS, and two aesthetascs.

Antenna (Fig. 4c) 0.84 times as long as antennule; articles 1–6 with length ratio 1.00:3.00:1.08:3.29:2.37:0.34. Articles 1 and 3 naked. Article 2 with one outer distal, one ventrodiscal, and one inner distal simple setae. Article 4 with distal simple seta and distal PSS. Article 5 with three distal simple setae and three distal PSS. Article 6 with tuft of distal simple setae.

Labrum (Fig. 4e) setulate distally. Mandibles (Fig. 4f–i) with well-developed molar process bearing many small teeth on masticatory surface; left mandible (Fig. 4f, g) with weakly denticulate incisor, wide denticulate lacinia mobilis, and bifurcate serrate accessory seta; right mandible (Fig. 4h, i) with seven-toothed incisor, peg-like lacinia mobilis, and bifurcate serrate accessory seta. Labium (Fig. 4j) with distally setulate inner lobe; outer lobes smaller. Maxillule (Fig. 4k) endite bearing eight distal spiniform setae and outer subdistal setation;

palp broken. Maxilla (ICHUM8304; Fig. 4l) with small serrations on outer distal margins.

Maxilliped (Fig. 4m) with coxa bearing simple seta (not illustrated). Basis with ventrodistal simple seta. Endite with outer distal tufts of fine setae, one mid-inner and two dorsodistal spiniform setae; and two ventro-subdistal circumplumose setae, inner one longer than outer; distal region setulate. Palp article 1 with outer distal serrations; article 2 with one outer distal and four inner simple setae, and inner distal serrate spiniform seta; article 3 with four inner distal simple setae and six inner plumose setae; article 4 with one mid-outer and two outer-subdistal simple setae, two inner plumose setae, and five distal serrate setae. Epignath (Fig. 4n) with kidney-shaped lobe, margins finely setulate; terminal seta setulate.

Cheliped (Fig. 5a, b) with triangular articulation to cephalothorax via sclerite (Fig. 3b). Basis slightly longer than wide, with one outer dorsal and one ventrodistal simple setae. Merus with one outer dorsal and one ventral simple setae. Carpus 1.46 times as long as wide, with one dorsal, two dorsodistal and two ventral simple setae. Propodus 0.77 times as long as carpus; palm with two outer and one inner simple setae at insertion of dactylus; fixed finger with one ventral, four outer subdistal, and two inner subdistal simple setae, and dorso-subdistal broad lamellar expansion and triangular claw. Dactylus as long as fixed finger, with inner simple seta, row of ventral spiniform setae, and bifurcate serrate seta; unguis triangular, curved ventrally.

Pereopod 1 (Fig. 5e, f) length 0.25 times BL, with length ratio of basis, merus, carpus, propodus, and dactylus-unguis 1.00:0.28:0.36:0.51:0.40. Coxa with dorsal simple seta. Basis cylindrical, narrow, 4.61 times as long as wide, with dorso-subproximal simple seta and dorso-subproximal PSS. Merus with one ventrodistal simple seta. Carpus with two dorsal and two ventrodistal simple setae. Propodus with one dorsodistal and two ventrodistal simple setae, middle PSS, and middle setulate seta. Dactylus naked; unguis shorter than dactylus, naked.

Pereopod 2 (Fig. 5g–i) with length ratio of articles from basis to propodus 1.00:0.53:0.34:0.46. Coxa with dorsal simple seta (not illustrated). Basis cylindrical, narrow, 3.17 times as long as wide, with ventrodistal simple seta and dorso-subproximal PSS. Merus with ventrodistal simple seta and ventrodistal crotchet (Fig. 5h). Carpus with dorsodistal simple seta and six distal crotchets (dorsodistal one shorter than half propodus length). Propodus with two dorsodistal and one ventrodistal simple setae and mid-dorsal PSS. Dactylus (ICHUM8303; Fig. 5i) naked; unguis shorter than dactylus, naked.

Pereopod 3 (ICHUM8303; Fig. 5j) with length ratio of articles from basis to dactylus-unguis 1.00:0.39:0.34:0.36:0.37; similar to pereopod 2, except basis with ventral PSS, carpus with five distal crotchets, and propodus with dorsodistal simple seta.

Pereopod 4 (ICHUM8303; Fig. 5k) with length ratio of articles from basis to dactylus-unguis 1.00:0.29:0.32:0.43:0.32. Coxa with simple seta. Basis 3.21 times as long as wide, with ventrodistal simple seta, and two dorso-subproximal

and two ventro-subdistal PSS. Merus with outer distal simple seta and two ventrodistal crotchets. Carpus with inner dorsodistal simple seta and five distal crotchets. Propodus with outer distal simple seta, dorso-subdistal PSS, and one outer dorsodistal and one inner dorsodistal setulate setae. Dactylus-unguis fused to form claw, strongly arcuate, with inner and outer rows of ventral spines.

Pereopod 5 (Fig. 5l) with length ratio of articles from basis to dactylus-unguis 1.00:0.43:0.30:0.51:0.35; similar to pereopod 4.

Pereopod 6 (Fig. 5m, n) with length ratio of articles from basis to dactylus-unguis 1.00:0.40:0.33:0.58:0.39; similar to pereopod 4 except basis without ventro-subdistal PSS and propodus with five inner distal flattened denticulate setae and inner distal biserrate seta.

Pleopod 1 (Fig. 6a) with protopod bearing six outer plumose setae; endopod with one inner and 14 outer plumose setae, and outer distal step-tipped plumose seta; exopod 1.42 times as long as endopod, with 29 outer plumose setae. Pleopod 2 (Fig. 6b) with protopod bearing six outer plumose setae; one inner plumose seta present on right protopod but absent on left; endopod with one inner and 13 outer plumose setae, and outer distal step-tipped plumose seta; exopod 1.37 times as long as endopod, with 30 outer plumose setae. Pleopod 3 (Fig. 6c) smaller than pleopods 1 and 2, with protopod bearing three outer plumose setae; endopod with one inner and 11 outer plumose setae, and outer distal step-tipped plumose seta; exopod 1.36 times as long as endopod, with 22 outer plumose setae.

Uropod (ICHUM8303; Fig. 6g) with four articles (protopod and triarticulate ramus). Protopod with four distal simple setae. Ramus article 1 naked; article 2 longest, with one middle and two distal simple setae and two distal PSS; article 3 with one middle and four distal simple setae.

Description of males (based on allotype). Body (Figs 2b, 3c, d) similar to female except for cephalothorax shape: cephalothorax strongly narrowed anteriorly.

Antennule (Fig. 4b) 0.93 times as long as cephalothorax; articles 1–4 with length ratio of 1.00:0.33:0.27:0.02. Article 1 with two outer distal and two inner distal simple setae, and several proximal and distal PSS. Article 2 with three outer distal and two inner distal simple setae, and several distal PSS. Article 3 with two distal simple setae and several distal PSS. Article 4 with tuft of distal simple setae, distal PSS, and three aesthetascs.

Antenna (Fig. 4d) 0.84 times as long as antennule; articles 1–6 with length ratio 1.00:2.47:1.59:3.43:1.93:0.19. Articles 1 and 3 naked. Article 2 with one outer distal, one ventrodistal, and one inner distal simple setae. Article 4 with distal simple seta and distal PSS. Article 5 with three distal simple setae and three distal PSS. Article 6 with tuft of distal simple setae and several distal PSS.

Labrum, mandibles, labium, maxillule, maxilla, maxilliped, and epignath similar to those of female.

Cheliped (Fig. 5c, d) with triangular articulation to cephalothorax via sclerite (Fig. 3d). Basis as long as wide, with one outer dorsal and one ventrodistal simple setae. Merus with one outer dorsal and one ventral simple




Figure 5. *Sinelobus kisui* sp. nov. **a, b, e–g, l–n.** Female holotype; **c, d.** Male allotype; **h–k.** Female paratype (ICHUM8303); **a, c.** Left (**a**) and right (**c**) chelipeds, outer views; **b.** Dactylus–unguis of cheliped and bifurcate serrate seta, inner view; **d.** Merus and carpus of cheliped, inner views; **e, g, j–m.** Left (**e, g, k–m**) and right (**j**) pereopods 1–6, inner (**e, k**) and outer (**g, j, l, m**) views; **f, i, n.** Distal part of right (**f**) and left (**i, n**) pereopods 1, 2, and 6, outer (**f, i**) and inner (**n**) views; **h.** Crotchet. White and black arrowheads, ventral processes on merus and carpus, respectively; gray arrowhead, inner broad thickening on merus; gray and black arrows, ventroproximal and mid-ventral processes on propodus, respectively. Scale bars: 0.1 mm (**a–g, i–m**); 0.05 mm (**h, n**).

setae, inner broad thickening (Fig. 5d, gray arrowhead), and three ventral processes (Fig. 5c, d, white arrowheads). Carpus 1.22 times as long as wide, with one dorsal, two dorsodistal, and two ventral simple setae, and three ventral processes (Fig. 5c, d, black arrowheads). Chela 1.15 times as long and 1.24 times as wide as carpus; ventral margin with middle simple seta and one proximal and one middle projection (Fig. 5c, gray and black arrows, respectively). Palm with two outer and one inner simple setae at insertion of dactylus. Fixed finger with four outer subdistal and two inner subdistal simple setae, dorso-subdistal triangular lamellar expansion, and triangular claw; base of fixed finger distant (greater than dactylus width) from base of dactylus; angle of dorsal margin of fixed finger to distal margin of palm about 90°. Dactylus strongly curved ventrally, with inner simple seta, row of ventral spiniform setae, and bifurcate serrate seta; unguis triangular.

Pereopods 1–6 more slender than but similar to those of females, with differences in setal number (see Suppl. material 1: Table S1 for details).

Pleopods (Fig. 6d–f) similar to those of female, with differences in setal number (see Suppl. material 1: Table S1 for details).

Uropod similar to that of female.

Variation. In addition to the holotype and allotype, two female (ICHUM8303, ICHUM8304) and one male (ICHUM8307) paratypes of *Sinelobus kisui* sp. nov. were dissected, and selected characters were observed for the

antennule, antenna, maxilliped, cheliped, pereopods 1–6, and pleopods 1–3. The raw morphological data we obtained are in Suppl. material 1: Table S1. All specimens shared the same state for the following selected characters: (1) antennule with two (females) or three (males) aesthetascs; (2) antennal article 2 with one outer distal, one ventrodistal, and one inner distal simple setae; (3) maxillipedal endite with one mid-inner and two distal spiniform setae, and two ventro-subdistal circumplumose setae, inner one longer than outer; (4) pereopod 1 with carpus bearing two dorsal and two ventral simple setae and propodus bearing one middle setulate seta; (5) dorsodistal crotchet on carpi of pereopods 2 and 3 shorter than half propodus length; (6) pereopod 3 basis with one ventral PSS; (7) pleopod 1 protopod lacking inner plumose setae. Two strongly sexually dimorphic males (allotype and paratype ICHUM8307) shared all states observed in the chelipeds.

The following setae varied in number among specimens (selected characters only are presented; ranges are in parentheses): ventrodistal simple setae on basis of pereopods 4–6 (1–2); inner distal flattened denticulate setae on pereopod 6 propodus (4–5); outer plumose setae on protopods of pleopods 1 and 2 (4–6) and on protopod of pleopod 3 (3–4); outer plumose setae on endopods of pleopods 1 and 2 (8–14) and on endopod of pleopod 3 (8–11); outer plumose setae on exopods of pleopods 1 and 2 (21–30) and on exopod of pleopod 3 (20–22). In the holotype individual, the right pleopod-2 protopod bears




Figure 6. *Sinelobus kisui* sp. nov. a–c. Female holotype; d–f. Male allotype; g. Female paratype (ICHUM8303); a–f. Right (a–c) and left (d–f) pleopods 1 (a, d), 2 (b, e), and 3 (e, f) (all setal ornamentation omitted); g. Right uropod. Scale bars: 0.1 mm.

one inner plumose seta (Fig. 6b) but the left one lacks it. This seta was not found on the corresponding article in the other four individuals, suggesting that its presence in the one case observed may be abnormal. We observed six distal crotchets on carpi of left pereopods 2 and 3 in the holotype individual, but all other pereopods we observed bore five, indicating the former state may be abnormal.

Morphology of individuals from Kagawa and Osaka. One female from Kagawa (ICHUM8314) and one strongly sexually dimorphic male from Osaka (OMNH-Ar12459) were dissected and observed. In the former individual, all diagnostic character states were observed, except for those found only in strongly sexually dimorphic males. The latter individual was whitish, having lost its pigmentation (but see Ariyama and Ohtani 1990), and showed all diagnostic character states.

Genetic information. One partial 18S (INSD accession number LC705430; 1920 bp long) and three COI (LC705431–LC705433; 815 bp long, encoding 271 amino acids) sequences were determined from Yamaguchi individuals. The 18S sequence was identical to that from a *Sinelobus* sp. 1 sensu Kakui et al. (2011) individual from Kagawa (AB618192, Kakui et al. 2011) but was 4.6% divergent in K2P distance from an 18S sequence from *Sinelobus* sp. 2 sensu Kakui et al. (2011) (AB618193, Kakui et al. 2011). For COI, K2P distances among three individuals from Yamaguchi were 0.0–0.4%; those between three Yamaguchi individuals and two Kagawa individuals (LC664100, LC664101; 658 bp long, encoding 219 amino acids) were 0.3–1.5%; and that between two Kagawa sequences was 1.5%.

Discussion

Sinelobus kisui sp. nov. is similar to *S. barretti* and *S. vanhaareni* in having antennal article 2 with one outer distal seta, the carpi of pereopods 2 and 3 with a dorsodistal crotchet shorter than half the propodus length, and the carpi of pereopods 2–6 with five distal crotchets (Table 1).

Sinelobus kisui differs from *S. barretti* in having (1) the inner of the two ventro-subdistal circumplumose setae on the maxillipedal endite longer than the outer; (2) the pereopod-1 carpus with two dorsal and two ventral simple setae; (3) the pereopod-1 propodus with a middle

setulate seta; and (4) the pleopod-1 protopod lacking inner plumose setae.

Sinelobus kisui differs from *S. vanhaareni* in having (1) the maxillipedal endite with one mid-inner seta; (2) the pereopod-1 carpus with two dorsal and two ventral simple setae; (3) the pereopod-3 basis with a ventral PSS; and (4) the pleopod-1 protopod lacking inner plumose setae. In addition, their strongly sexually dimorphic (SSD) males differ in the number of ventral processes on the chelipedal merus (three in *S. kisui*; two in *S. vanhaareni*). The two species share a state that is uncommon in *Sinelobus*, which is that the pleopod-3 protopod lacks inner setae, although *S. vanhaareni* bears a single inner plumose seta on the protopods of pleopods 1 and 2 (van Haaren and Soors 2009, Bamber 2014).

Our study confirms that character states of the chelipeds of SSD males (e.g., the number of ventral processes on the merus and carpus; the angle between the dorsal margin of the fixed finger and the distal margin of the palm; Table 1) are useful in *Sinelobus* taxonomy. Here we regard males having chelipeds in which the base of the fixed finger is distant (greater than dactylus width) from the base of the dactylus as SSD males. By this definition, *S. barretti* and *S. pinkenba* lack information on SSD males. Differences among *Sinelobus* species are not large, and we recommend that the chelipeds of SSD males be observed in future studies dealing with *Sinelobus* taxonomy.

In addition to individuals from the type locality (Yamaguchi), we observed *Sinelobus* individuals collected from Kagawa and Osaka and conclude that they are conspecific with Yamaguchi individuals, i.e., they are *S. kisui*. The K2P distances between the Yamaguchi and Kagawa COI sequences (0.3–1.5%) were within the range previously attributed to intraspecific variation in confamilial tanaidacean species: up to 1.1% in *Hexapleomera ulsana* Wi, Jeong & Kang, 2018 (Wi et al. 2018); up to 1.5% in *Zeuxo ezoensis* Okamoto, Oya & Kakui, 2020 (Okamoto et al. 2020). The fact that the 18S sequences were identical between the two populations supports the conclusion of conspecificity from our morphological observations. The species identity of *Sinelobus* tanaidaceans reported from other Japanese localities (cf. Fig. 1b) must be carefully investigated. Kakui et al. (2011) detected at least one other species molecularly, *Sinelobus* sp. 2, from Kochi, which shares a coastline with Kagawa (Fig. 1b).

Key to the seven species currently placed in *Sinelobus*

- | | | |
|---|---|---|
| 1 | Pleopod 1 protopod with inner plumose seta | 2 |
| – | Pleopod 1 protopod without inner plumose seta | <i>S. kisui</i> sp. nov. |
| 2 | Carpus of pereopods 2–6 with four distal crotchets | <i>S. bathykolpos</i> Bamber, 2014 |
| – | Carpus of pereopods 2–6 with five distal crotchets | 3 |
| 3 | Dorsodistal crotchet on carpi of pereopods 2 and 3 longer than about half propodus length | <i>S. stromatoliticus</i> Rishworth, Perissinotto & Błażewicz, 2018 |
| – | Dorsodistal crotchet on carpi of pereopods 2 and 3 shorter than half propodus length | 4 |
| 4 | Antennal article 2 with three or more ventrodiscal simple setae | 5 |
| – | Antennal article 2 with one ventrodiscal simple seta | 6 |

- 5 Pereopod 1 propodus with middle seta..... *S. pinkenba* Bamber, 2008
- Pereopod 1 propodus without middle seta *S. stanfordi* (Richardson, 1901)
- 6 Maxillipedal endite with mid-inner spiniform seta; pereopod 1 propodus without middle seta..... *S. barretti* Edgar, 2008
- Maxillipedal endite without mid-inner spiniform seta; pereopod 1 propodus with middle seta..... *S. vanhaareni* Bamber, 2014

Table 1. Comparison of selected characters for seven *Sinelobus* species. L, length; Mxp, maxilliped; nd, no data; Pr, pereopod; PSS, plumose sensory seta; SSD, strongly sexually dimorphic.

	<i>S. kisui</i> sp. nov.	<i>S. barretti</i>	<i>S. bathykolpos</i>	<i>S. pinkenba</i>	<i>S. stanfordi</i>	<i>S. stromatoliticus</i>	<i>S. vanhaareni</i>
Antennal article 2: number of outer distal setae	1	1	1	3	4	1	1
Mxp endite: mid-inner spiniform seta	Present	Present	Absent	Absent	Absent	Absent	Absent
Mxp endite: L of two circumplumose setae	Inner > outer	Outer > inner	Inner > outer	Inner > outer	Inner > outer	Inner > outer	Inner > outer
Chelipedal merus (SSD male): number of ventral processes	3	nd	1	nd	0	2	2
Chela (SSD male): angle of dorsal margin of fixed finger to distal margin of palm	≈ 90°	nd	≈ 50°	nd	≈ 90°	≈ 70°	≈ 90°
Pr1 carpus: number of simple setae (dorsal, ventral)	2, 2	1, 1	1, 1	1, 1	1, 1	1, 1	2, 1
Pr1 propodus: middle seta	Present	Absent	Present	Present	Absent	Present	Present
Pr2–3 carpi: relative L of dorsodistal crotch- et to propodus L	< 0.5	< 0.5	< 0.5	< 0.5	< 0.5	≈ 0.5	< 0.5
Pr3 basis: ventral PSS	Present	Present	Absent	Absent	Absent	Absent	Absent
Pr2–6 carpi: number of distal crotchets	5 (rarely 6)	5	4	5	5	5	5
Pleopod 1 protopod: number of inner plumose seta	0	1	1	1	1	1	1
Pleopod 3 protopod: number of inner plumose seta	0	1	1	1?	1	1	0
Source	This study	Edgar (2008)	Bamber (2014)	Bamber (2008)	Sieg (1980)	Rishworth et al. (2018)	Bamber (2014)

Acknowledgements

We thank Susumu Ohtsuka for help in sampling; Yoshihiro Hayashi and Hiroyuki Ariyama for providing *Sinelobus* samples from Kagawa and Osaka, respectively; Chizue Hiruta for help in rearing tanaidaceans; Kazuhiro Kogame for comments on our results; and Matthew H. Dick for reviewing the manuscript and editing the English. This study was supported in part by a KAKENHI grant (JP19K06800) to KK from the Japan Society for the Promotion of Science (JSPS).

References

Ariyama H, Ohtani M (1990) Record of tanaidacean Crustacea *Sinelobus stanfordi* (Richardson) from the Yodo River in Osaka. Nanki Seibutu 32: 23–26. [in Japanese with English abstract]

Bamber RN (2005) The tanaidaceans (Arthropoda: Crustacea: Peracarida: Tanaidacea) of Esperance, Western Australia, Australia. In: Wells FE, Walker DI, Kendrick GA (Eds) The Marine Flora and Fauna of Esperance, Western Australia. Western Australian Museum, Perth, 613–727.

Bamber RN (2008) Tanaidaceans (Crustacea: Peracarida: Tanaidacea) from Moreton Bay, Queensland. Memoirs of the Queensland Museum. Nature 54: 143–217.

Bamber RN (2014) Two new species of *Sinelobus* Sieg, 1980 (Crustacea: Tanaidacea: Tanaididae), and a correction to the higher taxonomic nomenclature. Journal of Natural History 48(33–34): 2049–2068. <https://doi.org/10.1080/00222933.2014.897767>

Bamber RN, Boxshall GA (2006) A new genus and species of the Langitainae (Crustacea: Peracarida: Tanaidacea: Tanaididae) bearing a new

genus and species of nicothoid parasite (Crustacea: Copepoda: Siphonostomatoida: Nicothoidae) from the New Caledonia Slope. Species Diversity: 11(2): 137–148. <https://doi.org/10.12782/specdiv.11.137>

Bird GJ (2011) Paratanaoidean tanaidaceans (Crustacea: Peracarida) from littoral and shallow sublittoral habitats in New Zealand, with descriptions of three new genera and seven new species. Zootaxa 2891(1): 1–62. <https://doi.org/10.11646/zootaxa.2891.1.1>

Bird GJ (2019) Tanaidacea (Crustacea: Peracarida) from the Southern French Polynesia Expedition, 2014. I. Tanaidomorpha. Zootaxa 4548(1): 1–75. <https://doi.org/10.11646/zootaxa.4548.1.1>

Chim CK, Tong SJW (2019) *Xenosinelobus balanocolus*, a new tanaidid genus and species (Crustacea: Peracarida: Tanaidacea) from barnacles on intertidal rocky shores and seawalls in the Singapore Strait. Zootaxa 4629(3): 413–427. <https://doi.org/10.11646/zootaxa.4629.3.9>

Edgar GJ (2008) Shallow water Tanaidae (Crustacea: Tanaidacea) of Australia. Zootaxa 1836(1): 1–92. <https://doi.org/10.11646/zootaxa.1836.1.1>

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3: 294–299.

Geospatial Information Authority of Japan (2022) GSI Maps. <https://maps.gsi.go.jp/> [Accessed 28 March 2022]

Kaji T, Kakui K, Miyazaki N, Miura K, Palmer AR (2016) Mesoscale morphology at nanoscale resolution: Serial block-face scanning electron microscopy reveals fine 3D detail of a novel silk spinneret system in a tube-building tanaid crustacean. Frontiers in Zoology 13(1): e14. <https://doi.org/10.1186/s12983-016-0146-0>

Kakui K (2016) Review of the taxonomy, diversity, ecology, and other biological aspects of order Tanaidacea from Japan and sur-

- rounding waters. In: Motokawa M, Kajihara H (Eds) *Species Diversity of Animals in Japan*, Springer, Berlin, 603–627. https://doi.org/10.1007/978-4-431-56432-4_23
- Kakui K (2022) Occurrence records of Japanese *Sinelobus* tanaidaceans and their published sources and online sources ver. 20220321. Figshare. <https://doi.org/10.6084/m9.figshare.19561966> [Accessed 11 June 2022]
- Kakui K, Angsupanich S (2012) *Birdotanaïs songkhlaensis*, a new genus and species of Nototanaididae (Crustacea: Tanaidacea) from Thailand. *The Raffles Bulletin of Zoology* 60: 421–432.
- Kakui K, Shimada D (2017) A new species of *Tanaopsis* (Crustacea: Tanaidacea) from Japan, with remarks on the functions of serial ridges and grooves on the appendages. *Zootaxa* 4282(2): 324–336. <https://doi.org/10.11646/zootaxa.4282.2.6>
- Kakui K, Kajihara H, Mawatari SF (2010) A new species of *Nesotanaïs* Shiino, 1968 (Crustacea, Tanaidacea) from Japan, with a key to species and a note on male chelipeds. *ZooKeys* 33: 1–17. <https://doi.org/10.3897/zookeys.33.296>
- Kakui K, Katoh T, Hiruta SF, Kobayashi N, Kajihara H (2011) Molecular systematics of Tanaidacea (Crustacea: Peracarida) based on 18S sequence data, with an amendment of suborder/superfamily-level classification. *Zoological Science* 28(10): 749–757. <https://doi.org/10.2108/zsj.28.749>
- Kakui K, Kobayashi N, Kajihara H (2012) Phylogenetic position of *Arctotanaïs* in the suborder Tanaidomorpha (Peracarida: Tanaidacea). *Journal of Crustacean Biology* 32(1): 127–139. <https://doi.org/10.1163/193724011X615406>
- Kakui K, Fleming JF, Mori M, Fujiwara Y, Arakawa K (2021) Comprehensive transcriptome sequencing of Tanaidacea with proteomic evidences for their silk. *Genome Biology and Evolution* 13: evab281. <https://doi.org/10.1093/gbe/evab281>
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16(2): 111–120. <https://doi.org/10.1007/BF01731581>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33(7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Larsen K (2003) Proposed new standardized anatomical terminology for the Tanaidacea (Peracarida). *Journal of Crustacean Biology* 23(3): 644–661. <https://doi.org/10.1651/C-2363>
- Nobili G (1906) Diagnoses préliminaires de Crustacés, Décapodes et Isopodes nouveaux recueillis par M. le Dr. G. Seurat aux îles Touamotou. *Bulletin du Muséum d'Histoire Naturelle* 12: 256–270. <https://doi.org/10.5962/bhl.part.4097>
- Okamoto N, Oya Y, Kakui K (2020) A new species of *Zeuxo* (Crustacea: Peracarida: Tanaidacea) from Japan, with remarks on carapace pigmentation as a potentially useful taxonomic character. *Marine Biology Research* 16(5): 411–422. <https://doi.org/10.1080/17451000.2020.1766693>
- Prendini L, Weygoldt P, Wheeler WC (2005) Systematics of the *Damon variegatus* group of African whip spiders (Chelicerata: Amblypygi): Evidence from behaviour, morphology and DNA. *Organisms, Diversity & Evolution* 5(3): 203–236. <https://doi.org/10.1016/j.ode.2004.12.004>
- Rasband WS (2022) ImageJ v1.53p. <https://imagej.nih.gov/ij/index.html>. [Accessed 24 March 2022]
- Richardson H (1901) Papers from the Hopkins Stanford Galapagos Expedition, 1898–1899. *Proceedings of the Washington Academy of Sciences* 3: 565–568. <https://doi.org/10.5962/bhl.part.26343>
- Rishworth GM, Perissinotto R, Błażewicz M (2018) *Sinelobus stromatoliticus* sp. nov. (Peracarida: Tanaidacea) found within extant peritidal stromatolites. *Marine Biodiversity* 49(2): 783–794. <https://doi.org/10.1007/s12526-018-0851-3> [Printed version was published in 2019]
- Sieg J (1980) *Taxonomische Monographie der Tanaidae Dana, 1849* (Crustacea, Tanaidacea). *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 537: 1–267.
- Stephensen K (1936) A tanaid (*Tanaïs stanfordi* Richardson) found in fresh-water in the Kurile Islands, with taxonomic remarks on the genus *Tanaïs* sensu lat. (*Tanaïs* Audouin et Milne-Edwards 1829, and *Anatanais* Nordenstam 1930). *Annotationes Zoologicae Japonenses* 15: 361–373. <https://www.doi.org/10.34434/za000395>
- Van Haaren T, Soors J (2009) *Sinelobus stanfordi* (Richardson, 1901): A new crustacean invader in Europe. *Aquatic Invasions* 4(4): 703–711. <https://doi.org/10.3391/ai.2009.4.4.20>
- Wessel P, Luis JF, Uieda L, Scharroo R, Wobbe F, Smith WHF, Tian D (2019) The Generic Mapping Tools Version 6. *Geochemistry Geophysics Geosystems* 20(11): 5556–5564. <https://doi.org/10.1029/2019GC008515>
- Wi JH, Jeong M-K, Kang C-K (2018) A new species of the genus *Hexapleomera* Dudich, 1931 from the South Korea, with molecular evidence (Crustacea, Tanaidacea, Tanaididae). *ZooKeys* 739: 1–28. <https://doi.org/10.3897/zookeys.739.21580>

Supplementary material 1

Table S1

Authors: Kyoko Hirano, Keiichi Kakui

Data type: Xlsx file.

Explanation note: Variation in characters among dissected specimens of *Sinelobus kisui* sp. nov.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zse.98.84818.suppl1>