



Title	Collagen XVII deficiency alters epidermal patterning
Author(s)	Wang, Yunan; Kitahata, Hiroyuki; Kosumi, Hideyuki; Watanabe, Mika; Fujimura, Yu; Takashima, Shota; Osada, Shin-ichi; Hirose, Tomonori; Nishie, Wataru; Nagayama, Masaharu; Shimizu, Hiroshi; Natsuga, Ken
Citation	Laboratory investigation, 102(6), 581-588 https://doi.org/10.1038/s41374-022-00738-2
Issue Date	2022-06-01
Doc URL	http://hdl.handle.net/2115/86545
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Lab Invest 102 581-588.pdf



[Instructions for use](#)

Collagen XVII deficiency alters epidermal patterning

Yunan Wang¹, Hiroyuki Kitahata², Hideyuki Kosumi¹, Mika Watanabe^{1,3}, Yu Fujimura¹, Shota Takashima¹, Shin-Ichi Osada⁴, Tomonori Hirose⁵, Wataru Nishie¹, Masaharu Nagayama⁶, Hiroshi Shimizu¹, Ken Natsuga^{1*}

¹Department of Dermatology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo, Japan

²Department of Physics, Graduate School of Science, Chiba University, Chiba, Japan

³Department of Life Sciences and Systems Biology, Molecular Biotechnology Centre, University of Turin, Turin, Italy

⁴Department of Dermatology, Nippon Medical School, Tokyo, Japan

⁵Department of Molecular Biology, Yokohama City University Graduate School of Medical Science, Yokohama, Japan

⁶Research Institute for Electronic Science, Hokkaido University, Sapporo, Japan

Short title

COL17 and epidermal patterning

*Correspondence and reprint requests to:

Ken Natsuga, M.D., Ph.D.

Department of Dermatology, Hokkaido University Faculty of Medicine and

Graduate School of Medicine

North 15 West 7, Sapporo 060-8638, Japan

Telephone: +81-11-716-1161, ext. 5962

E-mail: natsuga@med.hokudai.ac.jp

Abstract

Vertebrates exhibit patterned epidermis, exemplified by scales/interscales in mice tails and grooves/ridges on the human skin surface (microtopography).

Although the role of spatiotemporal regulation of stem cells (SCs) has been implicated in this process, the mechanism underlying the development of such epidermal patterns is poorly understood. Here, we show that collagen XVII

(COL17), a niche for epidermal SCs, helps stabilize epidermal patterns. Gene

knockout and rescue experiments revealed that COL17 maintains the width of

the murine tail scale epidermis independently of epidermal cell polarity. Skin

regeneration after wounding was associated with slender scale epidermis,

which was alleviated by overexpression of human COL17. COL17-negative skin

in human junctional epidermolysis bullosa showed a distinct epidermal pattern

from COL17-positive skin that resulted from revertant mosaicism. These results

demonstrate that COL17 contributes to defining mouse tail scale shapes and

human skin microtopography. Our study sheds light on the role of the SC niche

in tissue pattern formation.

Introduction

The skin is the body's outermost organ and is composed of the multilayered epithelium, the epidermis, and the underlying dermis. The epidermis serves as a physical barrier to pathogens and prevents water leakage from the body ¹.

The epidermis is maintained by a fine-tuned balance between the proliferation and differentiation of epidermal stem cells (SCs), which reside in the epidermal basal layer ². Epidermal SCs need niche proteins such as integrins and collagen XVII (COL17) for their proper function ^{3,4}. Functional loss of these proteins leads to transient hyperproliferation of the developing epidermis due to disturbed SC maintenance ⁵⁻⁷.

Vertebrates have distinct skin patterns. In some, the patterns are visible through melanin distribution in the skin (e.g., zebra and tiger stripes); in others, the allocation of skin components forms patterns (e.g., human microtopography, hair follicles (HFs), and fish scales). Murine tail skin serves as a robust model for examining epidermal pattern formation ⁸. The tail epidermis consists of scale (parakeratotic) and interscale (orthokeratotic) areas, which are arranged alternately. These scale and interscale areas are distinguished by the expression of keratin 31 (K31) and keratin 10 (K10), respectively ⁹. Label-

retaining and lineage-tracing experiments have revealed that K10+ interscale epidermis is slow-cycling, whereas K31+ scale areas are fast-cycling. Two distinct stem cell populations (Dlx1+ and Slc1a3+) give rise to interscale and scale epidermis, respectively ¹⁰, although it is unclear how these cell populations are arranged into scale/interscale patterns.

The expression of epidermal SC niche proteins, including integrins and COL17, shows alternate patterns in the human epidermis, where their expression is enriched in the epidermis facing the dermal protrusion but not in the epidermal rete ridges ¹¹⁻¹³. Conversely, the scale/interscale patterns are absent in β 1 integrin-null tail epidermis ¹⁴. These previous studies suggest the involvement of SC niche proteins in epidermal pattern formation. However, whether these SC niche proteins indeed regulate the epidermal patterns and the mechanisms underlying such regulation are unknown.

Here, we demonstrate that COL17, an SC niche protein ^{7,15-17}, helps in the formation of proper epidermal patterns in mice and humans. Interestingly, disturbed epidermal patterning through COL17 deletion is independent of aberrant epidermal cell polarity, but could involve wound-related skin changes.

Material and methods

Animals

C57BL/6 (wild-type, WT) mice were purchased from Clea (Tokyo, Japan).

Col17a1^{-/-}, K14-hCOL17 (h: human, a courtesy gift from Prof. Kim B

Yancey), hCOL17⁺; *Col17a1*^{-/-}, K5-Cre;*aPKCλ*^{ΔE5/ΔE5} (*aPKCλ* eKO), and

Prkcz^{-/-} (*aPKCζ* KO) were generated as previously described¹⁸⁻²¹.

aPKCλ^{ΔE5/ΔE5} and *Prkcz*^{+/-} mice were used as *aPKCλ* eKO and *aPKCζ* KO

controls, respectively. *aPKCλ* eKO and *aPKCλ*^{ΔE5/ΔE5} littermates were

generated by mating *aPKCλ* eKO and *aPKCλ*^{ΔE5/ΔE5} mice. *aPKCζ* KO and

Prkcz^{+/-} littermates were generated by mating *aPKCζ* KO and *Prkcz*^{+/-}

mice. Littermate *Col17a1*^{+/-} or *Col17a1*^{+/+} mice were used as *Col17a1*^{-/-}

or hCOL17⁺;*Col17a1*^{-/-} control. *Col17a1*^{-/-}, *Col17a1*^{+/-}, and *Col17a1*^{+/+}

littermates were generated by mating *Col17a1*^{+/-} female and male mice.

hCOL17⁺;*Col17a1*^{-/-}, *Col17a1*^{+/-}, and *Col17a1*^{+/+} littermates were

generated by mating hCOL17⁺;*Col17a1*^{+/-} and *Col17a1*^{+/-} mice. K14-

hCOL17 and WT littermates were generated by mating K14-hCOL17 and

WT mice. Female mice were used for the wound-healing experiments on

WT mice. Otherwise, sex-matched mice were used in each experiment.

The institutional review board of the Hokkaido University Graduate School of Medicine approved all animal studies described below.

Cell Culture

hTERT-immortalized human primary keratinocytes (KerCT; ATCC, Manassas, VA, USA), and spontaneously transformed murine keratinocytes (PAM212)²² were cultured in serum-free keratinocyte growth medium (KGM; Lonza). The cells were transfected with 10 μ M of human *COL17A1* siRNA, murine *Col17a1* siRNA or the control (Mock) (Silencer Select siRNAs, Thermo Fisher Scientific, Waltham, Massachusetts, USA) using Lipofectamine 2000 (Thermo Fisher Scientific) and Opti-MEM (Thermo Fisher Scientific). The cells were analyzed at 48 or 72 h after the knockdown procedure.

Antibodies

The following antibodies were used: polyclonal anti-K31 (Progen, hHa1), polyclonal anti-K10 (BioLegend, Poly19054), polyclonal anti-K6 (BioLegend, Poly19057), monoclonal anti-cytoplasmic COL17 (Abcam, Cambridge, MA, USA, ab186415/EPR14758).

Quantitative RT-PCR (qRT-PCR)

RNA was isolated from the tail epidermis or cultured cells using the RNeasy Mini kit (QIAGEN, Hilden, Germany), and cDNA was synthesized using the SuperScript III First-Strand Synthesis System (Thermo Fisher Scientific, Waltham, MA, USA). qRT-PCR was carried out using the designated primers and fast SYBR Green (Thermo Fisher Scientific) in a STEP-One Plus sequence detection system (Applied Biosystems, Waltham, MA, USA).

The following primers were used for the analysis: Forward primer and reverse primer are: murine *Krt6a*, CACGTTAAGAAGCAGTGTGCC and GCTCTGAGCACGGGATTCT; murine *Krt6b*, AGGAGTGCAGGTTGAATGGTG and AAAAAGAGAAGCGAGAGGACACA; murine *Krt16*, TCCCAGCTCAGCATGAAAG and GAGCTGTGGATATTCTCGCCA; murine *Krt17*, AGACAGAGAACCGCTACTGC and CGGGTGGTCACAGGTTCTTTT; murine *Col17a1*, GATGGCACTGAAGTCACCGA and TATCCATTGCTGGTGCTCCC; murine *Cyc1*, ATCGTTTCGAGCTAGGCATGG and GCCGGGAAAGTAAGGGTTGA; human *KRT17*, CAGAGAACCGCTACTGCGTG and

GTCACCGGTTCTTTCTTGTACTG; human *KRT16*,
GCTCAGCATGAAAGCATCCC and GACCTCGCGGGAAGAATAGG; human
KRT6A, AGTGCAGGCTGAATGGCGAA and TGGGACCGAGAGCTAGCAGA;
human *KRT6B*, TTCATCGACAAGGTGCGGT and
CAGCTCCGAGTCCAGACGAC; human *COL17A1*,
TCAACCAGAGGACGGAGTCA and TCGACTCCCCTTGAGCAAAC; human
RNA18SN1 (18S), GGCGCCCCCTCGATGCTCTTAG and
GCTCGGGCCTGCTTTGAACACTCT.

Immunofluorescence staining

Paraffin sections were deparaffinized and boiled in citrate or EDTA buffer for 20 min in a microwave oven. Frozen sections or cultured cells were fixed in 4% paraformaldehyde (PFA) for 10 min at room temperature (RT), or cold acetone or used without fixation. After washing with PBS, sections were treated with blocking buffer (0.5% fish skin gelatine, 5% goat serum, 4% BSA in PBS) for 1 h. The samples were incubated with primary antibodies at 4 °C overnight and were subsequently incubated with secondary antibodies conjugated with Alexa fluor 488, Alexa fluor 647, or FITC at RT for 1 h. Nuclei were stained with 4',6-

diamidino-2-phenylindole (DAPI). Images were obtained using confocal microscopy (FV1000, Olympus, Tokyo, Japan; LSM-710, Zeiss, Germany) or fluorescence microscopy (BZ-9000, Keyence, Osaka, Japan).

Whole mount staining

Tail skin was incubated in 5 mM EDTA/PBS on a shaker at 37 °C for 4 h to separate the epidermis from the dermis. Epidermal sheets were fixed in 4% paraformaldehyde (PFA) for 1h at RT. After blocking, epidermal sheets were incubated with primary antibodies overnight at RT, and then washed in 0.2% Tween/PBS. Samples were subsequently incubated with secondary antibodies. After washing, epidermal sheets were mounted on glass slides in Mowiol solution. The images of whole mount stained samples were obtained using FV1000 confocal laser scanning microscope (Olympus, Tokyo, Japan) or BZ9000 fluorescence microscope (Keyence, Osaka, Japan). The size and shape of the scales near the midline of the tail epidermis were analyzed using ImageJ (NIH, Bethesda, MD, USA). When the scales of the littermates were compared, the area, length, and width of the scales were normalized to the

whole tail equivalents to exclude the effects of the organismal size of each mouse.

Wound healing experiments

The surface of the tail skin (epidermis and papillary dermis, approximately 5 × 4 mm in size) was removed using a scalpel from 1-month-old (1MO) WT mice or K14-hCOL17 mice to produce superficial skin wounds (**Supplementary Fig. 5A**). The wounded skin was collected and analyzed when the healing process was complete (typically 4–6 weeks after wounding) and at a later time point (3 months after wounding).

Junctional epidermolysis bullosa (JEB) skin analysis

Photographs of the skin of a JEB patient²³ who was compound heterozygous for c.1179del (p.Ala394Leufs*9) and c.4159C>T (p.Gln1387*) in *COL17A1* (NM_000494.4) were taken by TG-5 (Olympus). Three revertant mosaicism spots and three adjacent diseased skin areas from the upper arm were further analyzed as described below. Although the fingerprints have been absent in other JEB patients with *COL17A1* p.Arg1303Gln mutations²⁴⁻²⁶, the patient in

our study maintained his fingerprints. This discrepancy is probably due to the difference of the *COL17A* mutations. The institutional review board of the Hokkaido University Graduate School of Medicine approved all human studies described above (ID: 13-043). The study was conducted according to the principles of the Declaration of Helsinki. The participant provided written informed consent.

Quantification of the skin microtopography

We selected several regions like the one surrounded by the red circle in the left panels of **Supplementary Fig. 6A and 6B** for Diseased skin and Reverted skin. We converted the image inside each circle to a grayscale image and calculated the two-dimensional autocorrelation function ²⁷ (right panels in **Supplementary Fig. 6A and 6B**). Using the two-dimensional autocorrelation function, the characteristic direction of the epidermal pattern was detected by determining the direction in which the autocorrelation in the range of a distance less than 1 mm is the largest. The one-dimensional autocorrelation function was calculated (**Supplementary Fig. 6C, 6D**) in the direction perpendicular to the characteristic direction, represented by the red lines in the right panels in **Supplementary**

Fig. 6A and 6B. The peak height (Δ) of the one-dimensional autocorrelation function in the range of distance less than 1 mm was adopted to quantify the regularity of the pattern. In the case where no peak was detected, the peak height was set to zero. The above-mentioned image analyses were performed with ImageJ NIH (Bethesda, Maryland, USA; <https://imagej.nih.gov/ij/>) by preparing a plug-in.

Statistical analysis

Statistical analysis was performed using GraphPad Prism (GraphPad Software, La Jolla, CA, USA). p-values were determined using Welch's t-test, Student's t-test, or Mann-Whitney test. p-values are indicated as * $0.01 < p < 0.05$, ** $0.001 < p < 0.01$, *** $0.0001 < p < 0.001$, **** $p < 0.0001$. The values are shown as violin plots. Violin plots show median (dashed line) and quartiles (dotted line).

RESULTS

COL17 deficiency alters scale shape in the tail skin

We first characterized the scale/interscale patterning of *Col17a1*^{-/-} ¹⁹ tail epidermis. We selected the time point of 1MO, when the scales became mature ⁸ (**Supplementary Fig. 1**). Immunofluorescence of the tail sections showed K10+ interscale and K31+ scale alternate patterns in both *Col17a1*^{-/-} and controls (**Fig. 1A; Supplementary Fig. 2A**). We then examined the scale shape by whole skin imaging. We defined the length and width of a scale as the diameter of the anterior-posterior (AP) and lateral-medial (LM) axes, respectively (**Fig. 1B**). Since the length and width of whole tail samples varied among the mice (**Supplementary Table 1**), we normalized the scale length/width by dividing them by each mouse tail length/width to exclude the effects of organismal size in the analysis. Whole-mount imaging showed that the scale size was smaller and the shape was more slender (shorter on the LM axis) in the *Col17a1*^{-/-} tail epidermis than in the littermate controls (**Fig. 1C-1F, Supplementary Fig. 2B**). The shorter scale width in *Col17a1*^{-/-} explains the smaller size of *Col17a1*^{-/-} scales because their length was comparable to that of the controls (**Fig. 1G, 1H**). Although the basal cell number of maximum diameter on the AP axis was comparable between

Col17a1^{-/-} and control scales, the cell number on the LM axis was smaller in *Col17a1*^{-/-} scales than in control scales (**Supplementary Fig. 2C**), implying the altered alignment of the basal cells in *Col17a1*^{-/-} scales. As the C57BL/6 (WT) scale shape was wider at P14 (postnatal day 14) than at 1MO (**Fig. 1I**), the slender scale shape in *Col17a1*^{-/-} mice was not due to the delayed development of the mice. *Col17a1*^{-/-} tails were generally smaller but not more slender than those of controls (**Supplementary Table 1**), excluding the involvement of organismal proportions in *Col17a1*^{-/-} slender scales. Transgenic rescue by the expression of hCOL17 under the keratin 14 (K14) promoter in *Col17a1*^{-/-} mice¹⁹ reversed the slender scale phenotype (**Fig. 2A-2C**). These data indicate that COL17 helps to define the scale proportion.

aPKC deregulation does not phenocopy *Col17a1*^{-/-} scale shape

Atypical protein kinase C (aPKC) is a key regulator of epithelial polarity²⁸, and the epidermis expresses two aPKC isoforms (aPKC λ and aPKC ζ)²⁹. The ablation of aPKC λ in the epidermis (K5-Cre;*aPKC* $\lambda^{\Delta E5/\Delta E5}$, aPKC λ eKO) and *Col17a1*^{-/-} mice share premature aging phenotypes such as gray hair and hair loss^{15,19,21,28}. This phenotypic similarity has been proposed to be a

consequence of the interaction between COL17 and the aPKC complex³⁰. We asked if the destabilized aPKC accounts for the slender scale shape of *Col17a1*^{-/-} mice (**Fig. 3A-3F**). Whole-mount imaging showed that the scale size was smaller in aPKC λ eKO than in littermate controls, but its scale proportion (width/length ratio) was wider than that of controls (**Fig. 3A, 3C, 3D**), which is in contrast to the slender scales of *Col17a1*^{-/-} mice (**Fig. 2**). aPKC ζ knockout (aPKC ζ KO, *Prkcz*^{-/-}) mice, which have no apparent skin phenotype²⁰, showed slightly larger scales, while the width/length ratio did not exhibit significant change (**Fig. 3B, 3E, 3F**). These results indicate that the aberrant cell polarity in *Col17a1*^{-/-} epidermis^{17,30} is not involved in altering the scale shape.

Expression of wound-induced keratins is pronounced in *Col17a1*^{-/-} tail epidermis

One of the factors that may affect scale shape is the cytoskeleton of epidermal keratinocytes. Keratin 6, 16, and 17 (K6/K16/K17) are the well-known keratins expressed upon physical injury. We reasoned that these wound-induced keratins might be enriched in *Col17a1*^{-/-} epidermis at steady state because COL17 deficiency leads to epidermolysis bullosa in humans³¹ and shows skin

fragility in mice ¹⁹. Quantitative PCR revealed that the gene expression of these wound-induced keratins was higher in *Col17a1*^{-/-} tail epidermis (**Fig. 4A**). In addition to gene expression, immunofluorescence analyses showed ectopic K6 expression in *Col17a1*^{-/-} tail epidermis at 1MO. K6 expression in *Col17a1*^{-/-} tail epidermis was also observed during the developmental stages (P1 and P14; **Fig. 4B-4D**). These data indicate that keratin profiles are skewed towards the wound-induced subsets in *Col17a1*^{-/-} epidermis. In contrast, human or murine cultured keratinocytes knocked down for *COL17A1* or *Col17a1* did not result in the expression of wound-induced keratins at the mRNA or protein level (**Supplementary Fig. 3-4**), suggesting that this phenotype is dependent on the in vivo setting. Rather, *KRT16* and *Krt6b* expression was reduced in human and murine knockdown experiments, respectively, for unknown reasons.

Scale shape becomes slender after skin regeneration, and COL17 overexpression rescues the phenotype

The expression of wound-induced keratins in *Col17a1*^{-/-} epidermis led us to ask whether wounding itself alters the tail scale shape upon skin regeneration (**Fig. 5A-5F**). The regenerated tail epidermis (4 to 6 weeks after wounding) exhibited

a more slender scale shape than the non-lesional areas (**Fig. 5A, 5C, 5D**), recapitulating the *Col17a1*^{-/-} scale (**Fig. 2**). The slender scale phenotype in the regenerated epidermis was not reversed 3 months after wounding (**Supplementary Fig. 5B-5D**). To see that additive COL17 prevents the alteration of the scale shape in the regenerated epidermis, we utilized K14-hCOL17 transgenic mice, which overexpress hCOL17 under the K14 promoter. The scale shape of the regenerated K14-hCOL17 skin was not as slender as that of the regenerated WT skin (**Fig. 5B, 5E, 5F**). These results suggest that COL17 prevents wound-induced scale deformation.

COL17 influences human skin microtopography

We finally asked whether the presence or absence of COL17 also affects epidermal patterning in humans. Although scale/interscale epidermal patterns in mouse tails are not conserved, skin surface patterns consisting of grooves and ridges are visible in humans. We took advantage of the revertant mosaicism in epidermolysis bullosa (EB), in which the mutated genes are corrected spontaneously^{32,33}. We compared COL17-negative (diseased) and COL17-positive (revertant) skin from a junctional EB (JEB) patient with *COL17A1*

mutations²³ (**Fig. 6A-6C, Supplementary Fig. 6**). The diseased skin surface appeared coarse, while the revertant skin was smooth (**Fig. 6B**). We calculated the autocorrelation functions of these skin images to quantify the skin microtopography and found that the diseased skin shows a distinct pattern from the revertant skin (**Fig. 6D**). These findings demonstrate that COL17 is a deterministic factor of epidermal patterning in mice and humans.

Discussion

COL17 is a hemidesmosomal protein that anchors basal keratinocytes to the dermis. COL17 stabilizes hemidesmosomes by binding to various basement membrane zone proteins including BP230^{34,35}, α 6 integrin³⁶, β 4 integrin^{35,37-39}, plectin^{35,40}, laminin-332^{25,41}, and type IV collagen^{42,43}. As a consequence, COL17 deficiency results in epidermolysis bullosa^{19,31}. Recently, COL17 has also been highlighted as an SC niche protein of HFs and epidermis, and its deficiency destabilizes epithelial SC maintenance^{7,15-17}. Our study provides new insights into COL17 biology. *Col17a1*^{-/-} mice have slender tail scales and are characterized by the expression of wound-induced keratins in the epidermis. In line with this, the regenerated epidermis after wounding shows slender tail scales. Human COL17 overexpression reverses the alteration of scale shapes upon wounding (**Fig. 7**).

Epidermal cell polarity regulates symmetrical and asymmetrical cell division of basal keratinocytes⁴⁴⁻⁴⁶ and is regulated by the aPKC complex⁴⁷⁻⁴⁹. Increased asymmetrical cell division in aPKC λ eKO epidermis²⁸ and possible stem cell depletion explain the smaller scales in aPKC λ eKO mice (**Fig. 3**). Although COL17 interacts with the aPKC complex³⁰ and helps maintain

epidermal cell polarity^{17,30}, *Col17a1*^{-/-} did not exhibit proportionally small scales as seen in aPKC λ eKO mice. This phenotypical difference indicates that the slender scale phenotype in *Col17a1*^{-/-} mice is independent of aberrant cell polarity.

A limitation of our study is that it does not explain which contributes more to the altered epidermal patterns in *Col17a1*^{-/-} mice: epidermal SC instability or weakened epidermal-dermal adhesion. As the HF abnormality of *Col17a1*^{-/-} mice becomes apparent at 3 months old¹⁵, it is not very likely that HFs are involved in the phenotypes of epidermal patterns at 1 month old in our study. Furthermore, the hard palate in the oral mucosa, where HFs are absent, shows an epithelial pattern of fast- and slow-cycling stem cells⁵⁰, recapitulating the epidermal pattern of the tail skin. Thus, HFs might not be essential for epithelial pattern formation.

Among various skin patterns, fingerprints, also called dermatoglyphics, are the most well-characterized in humans. Fingerprints show an alternate pattern of epidermal ridges and grooves. Loss of fingerprints has been described in EB patients, including Kindler syndrome^{51,52} and junctional EB with *COL17A1* p.Arg1303Gln mutation²⁴⁻²⁶. Our study has also demonstrated that

the presence of COL17 alters human skin microtopography (**Fig. 6**). These facts corroborate the role of COL17 in epidermal patterning and highlight COL17 as a therapeutic target for wound-induced skin deformations.

Human skin microtopography is not identical to mouse tail scale patterns because humans lost their tails after evolutionarily branching off from other primates. It has not been determined whether human skin ridges or grooves correspond to the fast- or slow-cycling areas that characterize mouse tails. However, the data that COL17 absence disturbs proper skin patterning in both humans and mice point to the involvement of COL17 in regulating skin surface texture across species.

Wound repair in mouse back skin requires wound contraction⁵³; however, the contribution of wound contraction has been regarded as minimal in the tail skin⁵⁴. Recent studies suggest that two-thirds of tail wound healing is due to epithelial regeneration, while wound contraction explains the remainder⁵⁵. We believe that wound contraction does not play a major role in the slender scale phenotype in the regenerated tail skin (**Fig. 5**) for the following reasons: 1) the scale shape in the regenerated skin would become proportionally small rather than slender if the wound contraction affected this phenotype, and 2) the

slender scale phenotype in the regenerated skin is rescued by K14-driven human COL17 overexpression, which might not influence wound contraction because the transgene expression is confined to the epidermis. The slender scales in *Col17a1*^{-/-} mice most likely represent wound-related skin changes that involve the expression of wound-induced keratins in *Col17a1*^{-/-} epidermis (**Fig. 4**). However, the mechanisms by which wound-related skin changes affect epidermal patterning need further investigation.

In conclusion, our study highlights the unrecognized role of COL17 in epidermal patterning. We propose that COL17 modulation can be utilized to prevent epidermal deformation upon wounding.

Acknowledgments

We thank Ms. Megumi Takehara for their technical assistance. We also thank Professor Kim B Yancey for providing K14-hCOL17 transgenic mice.

Ethics Approval and Consent to Participate

The institutional review board of the Hokkaido University Graduate School of Medicine approved all human studies described above (ID: 13-043). The study was conducted according to the principles of the Declaration of Helsinki. The participant provided written informed consent.

Author contributions

Y. W. performed the experiments, analyzed the data, interpreted the results, and wrote the manuscript. K.H. analyzed the data, interpreted the results, and wrote the manuscript. H.K., M.W., Y.F., S.T., SI.O., T.H., and N.W. performed the experiments and analyzed the data. M.N. and S.H. interpreted the results and supervised the study. K.N. designed the experiments, analyzed the data, interpreted the results, wrote the manuscript and supervised the study.

Funding

This work was funded by AMED to KN (20ek0109380h0003), the Lydia O'Leary Memorial Pias Dermatological Foundation to WN, JST CREST (JPMJCR15D2) to MN, and Five-star Alliance in NJRC Master. & Dev to HK/MN.

Data Availability Statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest

The authors declare that no conflicts of interest exist.

References

- 1 Natsuga K. Epidermal barriers. *Cold Spring Harb Perspect Med* **4**, a018218 (2014).
- 2 Moreci RS, Lechler T. Epidermal structure and differentiation. *Curr Biol* **30**, R144-R149 (2020).
- 3 Natsuga K, Watanabe M, Nishie W, Shimizu H. Life before and beyond blistering: the role of collagen XVII in epidermal physiology. *Exp Dermatol* **28**, 1135-1141 (2019).
- 4 Watt FM, Fujiwara H. Cell-extracellular matrix interactions in normal and diseased skin. *Cold Spring Harb Perspect Biol* **3**, a005124 (2011).
- 5 Brakebusch C, Grose R, Quondamatteo F, Ramirez A, Jorcano JL, Pirro A *et al.* Skin and hair follicle integrity is crucially dependent on β 1 integrin expression on keratinocytes. *EMBO J* **19**, 3990-4003 (2000).
- 6 Niculescu C, Ganguli-Indra G, Pfister V, Dupé V, Messaddeq N, De Arcangelis A *et al.* Conditional ablation of integrin alpha-6 in mouse epidermis leads to skin fragility and inflammation. *Eur J Cell Biol* **90**, 270-277 (2011).

- 7 Watanabe M, Natsuga K, Nishie W, Kobayashi Y, Donati G, Suzuki S *et al.* Type XVII collagen coordinates proliferation in the interfollicular epidermis. *Elife* **6**, e26635 (2017).
- 8 Dekoninck S, Hannezo E, Sifrim A, Miroshnikova YA, Aragona M, Malfait M *et al.* Defining the Design Principles of Skin Epidermis Postnatal Growth. *Cell* (2020).
- 9 Gomez C, Chua W, Miremadi A, Quist S, Headon DJ, Watt FM. The interfollicular epidermis of adult mouse tail comprises two distinct cell lineages that are differentially regulated by Wnt, Edaradd, and Lrig1. *Stem Cell Rep* **1**, 19-27 (2013).
- 10 Sada A, Jacob F, Leung E, Wang S, White BS, Shalloway D *et al.* Defining the cellular lineage hierarchy in the interfollicular epidermis of adult skin. *Nat Cell Biol* **18**, 619-631 (2016).
- 11 Jones PH, Harper S, Watt FM. Stem cell patterning and fate in human epidermis. *Cell* **80**, 83-93 (1995).
- 12 Kobayashi Y, Yasugahira Y, Kitahata H, Watanabe M, Natsuga K, Nagayama M. Interplay between epidermal stem cell dynamics and dermal deformation. *Npj Comput Mater* **4**, 1-9 (2018).

- 13 Wang S, Drummond ML, Guerrero-Juarez CF, Tarapore E, MacLean AL, Stabell AR *et al.* Single cell transcriptomics of human epidermis identifies basal stem cell transition states. *Nat Commun* **11**, 1-14 (2020).
- 14 López-Rovira T, Silva-Vargas V, Watt FM. Different consequences of β 1 integrin deletion in neonatal and adult mouse epidermis reveal a context-dependent role of integrins in regulating proliferation, differentiation, and intercellular communication. *J Invest Dermatol* **125**, 1215-1227 (2005).
- 15 Tanimura S, Tadokoro Y, Inomata K, Binh NT, Nishie W, Yamazaki S *et al.* Hair follicle stem cells provide a functional niche for melanocyte stem cells. *Cell Stem Cell* **8**, 177-187 (2011).
- 16 Matsumura H, Mohri Y, Binh NT, Morinaga H, Fukuda M, Ito M *et al.* Hair follicle aging is driven by transepidermal elimination of stem cells via COL17A1 proteolysis. *Science* **351** (2016).
- 17 Liu N, Matsumura H, Kato T, Ichinose S, Takada A, Namiki T *et al.* Stem cell competition orchestrates skin homeostasis and ageing. *Nature* **568**, 344-350 (2019).

- 18 Hirate Y, Hirahara S, Inoue K, Suzuki A, Alarcon VB, Akimoto K *et al.*
Polarity-dependent distribution of angiotensin II receptor type 1 localizes Hippo signaling in preimplantation embryos. *Curr Biol* **23**, 1181-1194 (2013).
- 19 Nishie W, Sawamura D, Goto M, Ito K, Shibaki A, McMillan JR *et al.*
Humanization of autoantigen. *Nat Med* **13**, 378-383 (2007).
- 20 Noguchi N, Hirose T, Suzuki T, Kagaya M, Chida K, Ohno S *et al.*
Atypical protein kinase C isoforms differentially regulate directional keratinocyte migration during wound healing. *J Dermatol Sci* **93**, 101-108 (2019).
- 21 Osada S-I, Minematsu N, Oda F, Akimoto K, Kawana S, Ohno S.
Atypical protein kinase C isoform, aPKC λ , is essential for maintaining hair follicle stem cell quiescence. *J Invest Dermatol* **135**, 2584-2592 (2015).
- 22 Yuspa SH, Hawley-Nelson P, Koehler B, Stanley JR. A Survey of Transformation Markers in Differentiating Epidermal Cell Lines in Culture. *Cancer Res* **40**, 4694-4703 (1980).

- 23 Nakamura H, Sawamura D, Goto M, Nakamura H, Kida M, Ariga T *et al.* Analysis of the COL17A1 in non-Herlitz junctional epidermolysis bullosa and amelogenesis imperfecta. *Int J Mol Med* **18**, 333-337 (2006).
- 24 Yuen W, Pas H, Sinke R, Jonkman M. Junctional epidermolysis bullosa of late onset explained by mutations in COL17A1. *Br J Dermatol* **164**, 1280-1284 (2011).
- 25 Nishimura M, Nishie W, Shirafuji Y, Shinkuma S, Natsuga K, Nakamura H *et al.* Extracellular cleavage of collagen XVII is essential for correct cutaneous basement membrane formation. *Hum Mol Genet* **25**, 328-339 (2016).
- 26 Stouthamer A, Nieboer C, Van der Waal R, Jonkman M. Normal expression of the 19-DEJ-1 epitope in two siblings with late-onset junctional epidermolysis bullosa. *Br J Dermatol* **144**, 1054-1057 (2001).
- 27 Okada M, Sumino Y, Ito H, Kitahata H. Spontaneous deformation and fission of oil droplets on an aqueous surfactant solution. *Phys Rev E* **102**, 042603 (2020).
- 28 Niessen MT, Scott J, Zielinski JG, Vorhagen S, Sotiropoulou PA, Blanpain C *et al.* aPKC λ controls epidermal homeostasis and stem cell

- fate through regulation of division orientation. *J Cell Biol* **202**, 887-900 (2013).
- 29 Helfrich I, Schmitz A, Zigrino P, Michels C, Haase I, le Bivic A *et al.* Role of aPKC isoforms and their binding partners Par3 and Par6 in epidermal barrier formation. *J Invest Dermatol* **127**, 782-791 (2007).
- 30 Watanabe M, Kosumi H, Osada SI, Takashima S, Wang YA, Nishie W *et al.* Type XVII collagen interacts with the aPKC-PAR complex and maintains epidermal cell polarity. *Exp Dermatol* **30**, 62-67 (2021).
- 31 McGrath JA, Gatalica B, Christiano AM, Owaribe K, McMillan JR, Eady RA *et al.* Mutations in the 180-kD bullous pemphigoid antigen (BPAG2), a hemidesmosomal transmembrane collagen (COL17A1), in generalized atrophic benign epidermolysis bullosa. *Nat Genet* **11**, 83-86 (1995).
- 32 Jonkman MF, Pasmooij AM. Revertant mosaicism—patchwork in the skin. *N Engl J Med* **360**, 1680-1682 (2009).
- 33 Nomura T. Recombination-induced revertant mosaicism in ichthyosis with confetti and loricrin keratoderma. *J Dermatol Sci* **97**, 94-100 (2020).
- 34 Hopkinson SB, Jones JC. The N terminus of the transmembrane protein BP180 interacts with the N-terminal domain of BP230, thereby mediating

- keratin cytoskeleton anchorage to the cell surface at the site of the hemidesmosome. *Mol Biol Cell* **11**, 277-286 (2000).
- 35 Koster J, Geerts D, Favre B, Borradori L, Sonnenberg A. Analysis of the interactions between BP180, BP230, plectin and the integrin $\alpha 6\beta 4$ important for hemidesmosome assembly. *J Cell Sci* **116**, 387-399 (2003).
- 36 Hopkinson SB, Baker SE, Jones J. Molecular genetic studies of a human epidermal autoantigen (the 180-kD bullous pemphigoid antigen/BP180): identification of functionally important sequences within the BP180 molecule and evidence for an interaction between BP180 and alpha 6 integrin. *J Cell Biol* **130**, 117-125 (1995).
- 37 Aho S, Uitto J. Direct interaction between the intracellular domains of bullous pemphigoid antigen 2 (BP180) and $\beta 4$ integrin, hemidesmosomal components of basal keratinocytes. *Biochem Biophys Res Commun* **243**, 694-699 (1998).
- 38 Schaapveld RQ, Borradori L, Geerts D, Van Leusden MR, Kuikman I, Nievers MG *et al.* Hemidesmosome formation is initiated by the $\beta 4$ integrin subunit, requires complex formation of $\beta 4$ and HD1/plectin, and

- involves a direct interaction between $\beta 4$ and the bullous pemphigoid antigen 180. *J Cell Biol* **142**, 271-284 (1998).
- 39 Hamill KJ, Hopkinson SB, Jonkman MF, Jones JC. Type XVII collagen regulates lamellipod stability, cell motility, and signaling to Rac1 by targeting bullous pemphigoid antigen 1e to $\alpha 6\beta 4$ integrin. *J Biol Chem* **286**, 26768-26780 (2011).
- 40 Natsuga K, Nishie W, Nishimura M, Shinkuma S, Watanabe M, Izumi K *et al.* Loss of interaction between plectin and type XVII collagen results in epidermolysis bullosa simplex. *Hum Mutat* **38**, 1666-1670 (2017).
- 41 Tasanen K, Tunggal L, Chometon G, Bruckner-Tuderman L, Aumailley M. Keratinocytes from patients lacking collagen XVII display a migratory phenotype. *Am J Pathol* **164**, 2027-2038 (2004).
- 42 Nishie W, Kiritsi D, Nyström A, Hofmann SC, Bruckner-Tuderman L. Dynamic interactions of epidermal collagen XVII with the extracellular matrix: laminin 332 as a major binding partner. *Am J Pathol* **179**, 829-837 (2011).
- 43 Kamaguchi M, Iwata H. The diagnosis and blistering mechanisms of mucous membrane pemphigoid. *Front Immunol* **10**, 34 (2019).

- 44 Morrow A, Underwood J, Seldin L, Hinnant T, Lechler T. Regulated spindle orientation buffers tissue growth in the epidermis. *Elife* **8**, e48482 (2019).
- 45 Muroyama A, Lechler T. Polarity and stratification of the epidermis. *Semin Cell Dev Biol* **23**, 890-896 (2012).
- 46 Poulson ND, Lechler T. Asymmetric cell divisions in the epidermis. *Int Rev Cell Mol Biol* **295**, 199-232 (2012).
- 47 Niessen MT, Iden S, Niessen CM. The in vivo function of mammalian cell and tissue polarity regulators—how to shape and maintain the epidermal barrier. *J Cell Sci* **125**, 3501-3510 (2012).
- 48 Tellkamp F, Vorhagen S, Niessen CM. Epidermal polarity genes in health and disease. *Cold Spring Harb Perspect Med* **4**, a015255 (2014).
- 49 Vorhagen S, Niessen CM. Mammalian aPKC/Par polarity complex mediated regulation of epithelial division orientation and cell fate. *Exp Cell Res* **328**, 296-302 (2014).
- 50 Byrd KM, Piehl NC, Patel JH, Huh WJ, Sequeira I, Lough KJ *et al.* Heterogeneity within Stratified Epithelial Stem Cell Populations Maintains

- the Oral Mucosa in Response to Physiological Stress. *Cell Stem Cell* **25**, 814-829 e816 (2019).
- 51 Has C, Burger B, Volz A, Kohlhase J, Bruckner-Tuderman L, Itin P. Mild clinical phenotype of Kindler syndrome associated with late diagnosis and skin cancer. *Dermatology* **221**, 309-312 (2010).
- 52 de Almeida Jr HL, Isaacsson H, Guarenti IM, e Silva RM, de Castro LAS. IMAGES IN DERMATOLOGY. *An Bras Dermatol* **90**, 581-584 (2015).
- 53 Grada A, Mervis J, Falanga V. Research techniques made simple: animal models of wound healing. *J Invest Dermatol* **138**, 2095-2105. e2091 (2018).
- 54 Falanga V, Schraye D, Cha J, Butmarc J, Carson P, Roberts AB *et al.* Full-thickness wounding of the mouse tail as a model for delayed wound healing: accelerated wound closure in Smad3 knock-out mice. *Wound Repair Regen* **12**, 320-326 (2004).
- 55 Aragona M, Dekoninck S, Rulands S, Lenglez S, Mascré G, Simons BD *et al.* Defining stem cell dynamics and migration during wound healing in mouse skin epidermis. *Nat Commun* **8**, 1-14 (2017).

Figure legends

Fig. 1. Slender tail scales of *Col17a1*^{-/-} mice.

(A) Images showing K31 and K10 staining of *Col17a1*^{-/-} and control tail skin samples at 1MO (n=4). Scale bar: 500 μ m. (B) Schematic of distribution of scales in mouse tail epidermis. (C) K31 whole mount staining images of *Col17a1*^{-/-} and littermate control tail epidermis at 1MO (n=3). Scale bar: 500 μ m. (D) Phase-contrast images of *Col17a1*^{-/-} and littermate control tail epidermis at 1MO (n=4). Scale bar: 500 μ m. (E-H) Quantification of the size and shape of tail scales. Scale area (E), width/length (F), length (G), and width (H) of *Col17a1*^{-/-} and littermate control tail scales at 1MO are shown (n=298 scales from three control and 413 scales from three *Col17a1*^{-/-} mice). (I) Width/length ratio of tail scales in P14 and 1MO WT mice (n=148 scales from three P14 WT mice and 123 scales from three 1MO WT mice). ****p<0.0001, Welch's t-test.

Fig. 2. Restoration of the scale shape in *Col17a1*^{-/-} tail epidermis by human COL17 overexpression.

(A) Whole mount phase-contrast imaging of hCOL17⁺; *Col17a1*^{-/-} and littermate control (hCOL17⁻; *Col17a1*^{+/+} or hCOL17⁻; *Col17a1*^{+/-}) tail epidermis at 1MO (n=3). Scale bar: 500 μ m. (B, C) Quantification of the size and shape of tail scales. Scale area (B) and width/length (C) of hCOL17⁺; *Col17a1*^{-/-} and littermate control tail scales at 1MO are shown (n=266 scales from three control and 359 scales from three hCOL17⁺; *Col17a1*^{-/-} mice). The raw data used for **Fig. 2C** are shown in **Supplementary Table 2**. *0.01 < p < 0.05, Welch's t-test.

Fig. 3. Small, but not slender, tail scales of aPKC λ eKO mice.

(A) Whole mount phase-contrast imaging of aPKC λ eKO and littermate control tail epidermis (n=3). Scale bar: 500 μ m. (B) Whole mount phase-contrast imaging of aPKC ζ KO and littermate control tail epidermis (n=3). Scale bar: 500 μ m. (C-F) Quantification of the size and shape of tail scales. Scale area and width/length of aPKC λ eKO (C, D) and aPKC ζ KO (E, F) tail scales at 1MO are shown (n=226 scales from three control mice and 357 scales from three aPKC λ eKO mice, n=300 scales from three control mice and 317 scales from three aPKC ζ KO mice). The raw data used for **Fig. 3F** are shown in **Supplementary Table 3**. ****p<0.0001, *0.01<p<0.05, Welch's t-test.

Fig. 4. Expression of wound-induced keratins in *Col17a1*^{-/-} tail epidermis.

(A) Gene expression of *Krt6a*, *Krt6b*, *Krt16*, and *Krt17* in the tail epidermis of *Col17a1*^{-/-} and littermate control at 1MO (n = 7). *0.01 < p < 0.05, Student's t-test.

(B-D) K6 staining of *Col17a1*^{-/-} and littermate control at P1 (B, n = 3), P14 (C, n = 3), and 1MO (D, n = 3). Scale bar: 100 μ m.

Fig. 5. Slender tail scales after skin regeneration and COL17 transgenic rescue.

(A-B) Whole mount phase-contrast imaging of WT (A) and K14-hCOL17 (B) tail epidermis 4-6 weeks after wounding (n=4 (WT) and 6 (K14-hCOL17), respectively). Scale bar: 500 μ m. (C-F) Quantification of the size and shape of tail scales. Scale area and width/length of WT (C, D) and K14-hCOL17 (E, F) tail scales after skin regeneration are shown (n=542 scales from unwounded areas and 467 scales from regenerated areas from four WT mice, n=424 scales from unwounded areas and 440 scales from regenerated areas from six K14-hCOL17 mice). ****p<0.0001, Welch's t-test.

Fig. 6. Distinct epidermal patterns of COL17-positive and negative human skin.

(A) The upper arm of a junctional epidermolysis bullosa patient who is compound heterozygous for c.1179del (p.Ala394Leufs*9) and c.4159C>T (p.Gln1387*) in *COL17A1* (NM_000494.4). The revertant skin areas are circled by dotted lines. (B) Representative images of the diseased and revertant skin. Scale bar: 3 mm. These images are exemplified in **Supplementary Fig. 6**. (C) COL17 labeling of the diseased and revertant skin. Scale bar: 100 μ m. (D) Quantification of skin microtopography (n=75 spots (radius: 3 mm) from three skin areas, respectively). The degree of regularity on the epidermal pattern was quantified by the peak height of the autocorrelation function (**Supplementary Fig. 6**). ***0.0001<p<0.001, Mann-Whitney test.

Fig. 7. COL17-related epidermal patterning.

A graphical abstract of the study. COL17 deficiency or wounding leads to slender scale epidermis, which is reversed by overexpression of human COL17 in the epidermis.






Figure 1. Slender tail scales of *Col17a1-/-* mice.

A


A
↑
↓
P



Control

hCOL17+;*Col17a1*^{-/-}

B



C




Figure 2. Restoration of the scale shape in *Col17a1*^{-/-} tail epidermis by human COL17 overexpression








Figure 3. Small, but not slender, tail scales of aPKC λ eKO mice.

A ↔ P

A



B

Control

Col17a1^{-/-}

P1

C

P14

D

1MO




Figure 4. Expression of wound-induced keratins in *Col17a1^{-/-}* tail epidermis.




Figure 5. Slender tail scales after skin regeneration and COL17 transgenic rescue.




Figure 6. Distinct epidermal patterns of COL17-positive and negative human skin.

Mouse tail epidermis

- Wound-related?
- Independent of aPKC




Figure 7. COL17-related epidermal patterning