



Title	The use of digestate from animal wastes and its impacts on the soil microbiome and nutrient dynamics in agricultural soils
Author(s)	MADEGWA, Yvonne Musavi
Citation	北海道大学. 博士(農学) 甲第14653号
Issue Date	2021-09-24
DOI	10.14943/doctoral.k14653
Doc URL	<a href="http://hdl.handle.net/2115/86916">http://hdl.handle.net/2115/86916</a>
Type	theses (doctoral)
File Information	madegwa_yvonne_musavi.pdf



[Instructions for use](#)

**The use of digestate from animal wastes and its impacts on the soil microbiome and  
nutrient dynamics in agricultural soils**

**(家畜糞尿由来消化液の利用が農地土壤微生物叢と栄養素ダイナミクスに及ぼす影響  
の評価)**

**北海道大学 大学院農学院**

**共生基盤学専攻 博士後期課程**

**Yvonne Musavi Madegwa**

## **Abstract**

Abstract of a thesis submitted in partial fulfilment of the requirements for the Degree of Doctor of Philosophy

### **The use of digestate from animal wastes and its impacts on the soil microbiome and nutrient dynamics in agricultural soils**

Yvonne Musavi Madegwa

Digestate is the semi liquid byproduct of the biogas production process. Due to its basic pH and high nutrient content (i.e inorganic nitrogen, phosphorous, organic carbon), digestate use as a fertilizer has been increasing. Additionally, to improve its efficiency as a fertilizer, digestate can be separated into solid and liquid fractions using a flocculant. However, the effect of digestate and its separated fractions on the soil microbiome and nutrient dynamics in agricultural soils is not clear. Therefore, this study, involving fieldwork and incubation experiments was conducted to address this research need. The field work involved sampling soils from farmers' fields in Kamishihoro town in Hokkaido to determine the effect of land use, seasons and fertilizer application on the soil microbiome and related functions. The second experiment was an incubation to determine the effect of soil pH on microbial stability against the application of digestate from dairy wastes. The third experiment was an incubation to analyze the effect of solid-liquid separation of digestate on soil nutrient dynamics and Japanese mustard spinach yields in different soils.

The field study was conducted in an Andosol (volcanic soil) dominated agricultural region in cool temperate climate to determine the effect of land use (cropland, grassland), season (spring, summer) and fertilizer (digestate) on soil microorganisms and related functions. Soils were sampled from farmers' fields, DNA extracted and sequenced targeting 16S rRNA region. In result, land use had a significant effect on beta diversity and evenness with higher values recorded in cropland than grassland. However, grassland had a higher number of unique operational taxonomic units (OTUs) (10303) compared to cropland (5112). In cropland, season had a significant effect on beta diversity, evenness, OTU numbers and Shannon index with higher values recorded in summer compared to spring. Based on predicted soil functions, nitrogenase (*nifH*) had significantly higher values in cropland-summer while nitrite reductase (*nirK*) and ammonia

monooxygenase (*amoA*) were significantly higher in cropland-spring. In grassland, season had a significant effect on beta diversity only. These results indicate that grassland microorganisms were stable and more resistant to seasonal changes than cropland, suggesting that conventional tillage practices have a negative effect on soil microbial stability. Additionally, grassland-spring (7059) had a higher number of unique OTUs than grassland-summer (2597). Based on predicted soil functions, *nifH* was significantly higher in grassland-spring while *nirK* and *amoA* were significantly higher in grassland-summer. These results indicate that the impact of seasons on soil microorganisms' distribution and abundance in cropland and grassland may directly affect soil functions.

The first incubation experiment was performed to understand the effect of lime application (pH = 6.5 and 5.5 for the soils with and without lime, respectively) and fertilizer (digestate, urea and control) on the soil microbial community structures, stability and gene functions. Soils were sampled weekly after the application of fertilizers for a month. For microbial community analysis, DNA was extracted and sequenced targeting 16S rRNA region. For gene abundances (i.e 16S rRNA, ammonia oxidizing archaea (AOA), ammonia oxidizing bacteria (AOB), nitrous oxide reductase (*nosZ*) and nitrite reductase (*nirS*), quantitative PCR was conducted. In result, the relative abundance of *Actinobacteria* was influenced more strongly by digestate in lime soils, while *Alphaproteobacteria* was influenced more strongly by digestate in the no lime soil. In no lime treatments, digestate had a significant effect on more operational taxonomic units (146) compared to lime (127), indicating that lime application increased soil microbial community's stability. Liming and fertilizer had a significant effect on 16S rRNA gene copy numbers with the highest values observed in lime plus digestate treatments. Soil pH had a significant on AOA, *nosZ* and *nirS* gene copy numbers with the highest values observed in lime treatments. In the lime treatments digestate application had a positive impact on AOB gene copy numbers but this was not the case for soils without liming treatments. These results indicate that soil pH and fertilizer type should be taken into consideration for the management of functional gene abundance in agricultural soils.

The second incubation experiment was setup to analyze the effect of soil types (Kamishihoro and Arakida) and fertilizers (digestate, solid digestate, liquid digestate, chemical and control) on soil and plant (Japanese mustard spinach) nutrients, gene abundance and nitrous oxide (N<sub>2</sub>O)

emissions. Based on the results, soil type significantly influenced the above ground plant biomass and Nitrogen (N) content with higher values observed in Kamishihoro compared to Arakida. Kamishihoro had significantly higher soil inorganic N ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ ) content compared to Arakida soil. Fertilizer had a significant effect on soil  $\text{NO}_3^-$  (solid-Kamishihoro, digestate-Arakida) and  $\text{NH}_4^+$  (liquid-Kamishihoro, chemical-Arakida) content. The results indicate that digestate and its derived fertilizers (solid digestate, liquid digestate) can be used to influence soil inorganic N content in agricultural soils. Digestate derived fertilizers influenced soil pH with highest values observed in liquid in both Kamishihoro and Arakida soils. Arakida soil had significantly higher 16S rRNA and ITS gene copy numbers compared to Kamishihoro. Regarding soil  $\text{N}_2\text{O}$  emissions, Kamishihoro soil had significantly higher emissions compared to Arakida. Additionally, digestate and solid fertilizer had significantly higher  $\text{N}_2\text{O}$  emissions compared to liquid in both Kamishihoro and Arakida soils. The results indicate the importance of soil type and digestate derived fertilizers in modulating soil gene abundance and  $\text{N}_2\text{O}$  emissions in agricultural soils. Furthermore, the separation of digestate into solid and liquid fractions can be considered a reliable method to regulate aspects of soil nutrient dynamics in agricultural soils.

## **Acknowledgements**

I would like to thank the Almighty God for his grace and mercies without which I would not have been able to complete this thesis.

Special thanks goes to my supervisor Associate Professor Yoshitaka Uchida for his support and guidance throughout the study. I am also grateful to Prof. Ishiguro and Prof. Toma for the role he played towards completion of my thesis.

I thank Sekiko Kurazono for the significant technical support towards my research. Thanks to the Uchida lab members (Akira Nagata, Miyuki Oka, Johnson Chidozie Oraegbunam, Akari Kimura, Takamitsu Ohigashi, Anna Saito, Gen Takahashi, Ruka Kiyama, Ikabongo Mukumbuta, Chikae Tatsumi, Lin Jin-Feng, Terumasa Yamamoto, Nao Ishige and Hiroki Nishimine) for their assistance and support during my research.

Deepest gratitude to Yufita Dwichinta for her counsel on the scientific process.

My heartfelt appreciation goes to my brothers Mike, Leslie, Lwarim and sisters Valary, Lindah for their encouragement which played a vital role in completion of my thesis. Special thanks to my mother Catherine Madegwa who is a constant source of strength and motivation during the study.

Finally, I would like to express my sincere gratitude to MEXT Scholarship (grant number 180782) for their financial support without which the study would not have been possible.

## Table of content

<b>Abstract</b> .....	i
<b>Acknowledgements</b> .....	iv
<b>Table of content</b> .....	v
<b>Chapter 1 Introduction</b> .....	1
1.1 Background .....	1
1.2 Research Objectives .....	2
<b>Chapter 2 Literature Review</b> .....	4
2.1 Role of soil microorganisms in agriculture .....	4
2.2 Soil microorganisms and the nitrogen cycle .....	6
2.3 Soil microbial diversity and stability .....	8
2.3.1 Diversity.....	8
2.3.2 Stability .....	10
2.4 Factors that affect soil microbial properties.....	10
2.4.1 Land use .....	10
2.4.2 Agricultural conventional tillage practices .....	11
2.4.3 Effect of agricultural management practices on microbial diversity and stability .....	12
2.4.4 Seasonality .....	13
2.4.5 Fertilizers .....	14
2.4.6 Soil pH .....	20
References .....	24
<b>Chapter 3 Land use and season drive changes in soil microbial communities and related functions in agricultural soils</b> .....	40
3.1 Abstract .....	40
3.2 Introduction.....	41
3.3 Materials and methods .....	44
3.3.1 Study site and treatments .....	44
3.3.2 Soil sampling .....	45

3.3.3 Measurement of soil physical and chemical properties .....	46
3.3.4 Soil DNA extraction .....	46
3.3.5 16S rRNA Gene Amplicon Library Preparation and Sequencing .....	46
3.3.6 Sequence processing and data analysis.....	47
3.3.7 Soil microbial and chemical analysis.....	48
3.4 Results.....	50
3.4.1 Evenness, richness and microbial diversity .....	50
3.4.2 Distribution of OTUs under imposed treatments.....	53
3.4.3 Effect of imposed treatments on soil functions.....	55
3.4.4 Relationship between microbial community structure and soil properties .....	56
3.5 Discussion.....	57
3.5.1 Effect of land use on the soil microbial community .....	57
3.5.2 Effect of season on soil microbial community .....	60
3.5.3 Effect of season on shared phyla and predicted functions in cropland.....	61
3.5.4 Effect of season on shared phyla and predicted functions in grassland.....	63
3.6 Conclusion .....	64
References.....	65
Supplementary .....	78
<b>Chapter 4 Liming improves the stability of soil microbial community structures against the application of anaerobic digestate made from dairy wastes .....</b>	<b>84</b>
4.1 Abstract.....	84
4.2 Introduction.....	85
4.3 Materials and methods .....	87
4.3.1 Soil sampling and treatments .....	87
4.3.2 Incubation setup.....	88
4.3.3 Soil sampling during the incubation experiments, chemical and microbial analysis ..	88
4.3.4 Soil DNA extraction .....	88
4.3.5 16S rRNA gene amplicon library preparation and sequencing .....	89
4.3.6 Sequence processing and data analysis.....	89
4.3.7 Quantitative Polymerase Chain Reaction (qPCR).....	90
4.3.8 Analysis of bacterial community composition .....	92



4.3.9 Statistical analysis .....	92
4.4 Results .....	93
4.4.1 Beta Diversity .....	93
4.4.2 Changes in OTU distribution and stability .....	93
4.4.3 Class level microbes that were influenced by fertilizer application based on their abundance .....	95
4.4.4 Alpha Diversity .....	96
4.4.5 Gene Abundances .....	99
4.5 Discussion .....	102
4.5.1 Influence of soil pH on microbial response to digestate application .....	102
4.5.2 Influence of soil pH on the effect of digestate on OTU stability .....	103
4.5.3 Influence of soil pH on OTU response to digestate application .....	103
4.5.4 Influence of soil pH and fertilizers on microbial diversity .....	104
4.5.5 Effect of soil pH and fertilizer application on the abundances of bacteria and their functional genes .....	105
4.6 Conclusion .....	108
References .....	109
Supplementary .....	121
<b>Chapter 5 The impact of separating anaerobic digestate on soil nutrients, yields and microbial functions in agricultural soils .....</b>	<b>126</b>
5.1 Abstract .....	126
5.2 Introduction .....	127
5.3 Materials and methods .....	128
5.3.1 Study site and soil characteristics .....	128
5.3.2 Digestate separation to solid and liquid phase .....	129
5.3.3 Incubation setup .....	129
5.3.4 Soil and plant sampling and analysis .....	130
5.3.5 Nitrogen mass balance analysis .....	130
5.3.6 N <sub>2</sub> O gas sampling and analysis .....	130
5.3.7 Soil DNA extraction and microbial abundance .....	131
5.3.8 Statistical analysis .....	131
5.4 Results .....	132

5.4.1 Plant Analysis .....	132
5.4.2 Soil Analysis .....	133
5.4.3 Leachate Analysis .....	136
5.4.4 Nitrogen mass balance .....	138
5.4.5 Gene Abundances .....	139
5.4.6 Nitrous oxide emissions .....	139
5.5 Discussion .....	140
5.5.1 The effect if separated digestate on soil inorganic nitrogen content .....	140
5.5.2 Influence of soil types .....	142
5.5.3 Leachate analysis .....	142
5.5.4 Nitrogen mass balance .....	143
5.5.5 16S rRNA gene abundance .....	143
5.5.6 N <sub>2</sub> O emissions .....	143
5.5.7 Plant Analysis .....	144
5.6 Conclusion .....	145
References .....	146
Supplementary .....	151
<b>Chapter 6 Synthesis and Recommendations for further research .....</b>	<b>155</b>
6.1 Overall summary and future research recommendations .....	155
6.1.1 Land use and season drive changes in soil microbial communities and related functions in agricultural soils .....	155
6.1.2 Liming improves the stability of soil microbial community structures against the application of digestate made from dairy wastes .....	156
6.1.3 The effect of solid-liquid separation of digestate on soil nutrient dynamics and Japanese mustard spinach ( <i>Brassica rapa</i> var. <i>perviridis</i> ) yields in different soils .....	157
6.2 General Comments .....	158

## **List of Tables**

Table 2.1: Role of soil microbes in provisioning and regulating services provided by soil ecosystems (adapted from Dominati et al., 2010).....	5
Table 4.1: Primer sequences and qPCR conditions used for the study.....	91

## List of Figures

Figure 2.1: Partial soil nitrogen cycling processes with emphasis on nitrification, denitrification and related biomarker genes .....	8
Figure 2.2: Distribution of the principal constituents after solid–liquid separation (data based on own investigations and various references. Abbreviations; TN-Total Nitrogen, P-Phosphorous, K-Potassium, C-Carbon (Adapted form Bauer et al., 2009).....	18
Figure 2.3: Principle of flocculation (Adapted form Fechter and Kraume, 2016) .....	19
Figure 2.4: Soil biochemical processes and their relations to soil pH (Adopted from Neina, 2019) .....	21
Figure 3.1: Kamishihoro soil sampling sites.....	45
Figure 3.2: Box plots illustrating pielou evenness at OTU level based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points. p values calculated based on Kruskal-Wallis test. ....	50
Figure 3.3: Box plots illustrating observed OTU numbers based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points. p values calculated based on Kruskal-Wallis test. ....	51
Figure 3.4: Box plots illustrating Shannon diversity index at OTU level based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points. p values calculated based on Kruskal-Wallis test.....	52
Figure 3.5: Principal coordinate analysis (PCoA) by weighted unifrac based on land use and season in cropland and grassland. Axes represent the two dimensions explaining greatest proportion of variances in microbial communities. p values calculated based on permutational multivariate analysis of variance. ....	53
Figure 3.6: Venn diagrams showing the number of shared and unique OTUs based on land use and season. Percentage values represent relative abundance of OTUs in each section.....	55
Figure 3.7: The effect of season on predicted soil functions in cropland and grassland. Error bars show statistical differences of selected 16S rRNA gene-predicted functional profiles. Abbreviations ; nifH-nitrogenase, nirK-nitrite reductase, amoA-ammonia monooxygenase. ....	56
Figure 3.8: Nonmetric dimensional scaling plot based on bray curtis dissimilarity of microbial community structure in soils among seasonal treatments in cropland and grassland. Arrows show direction of significant ( $p < .05$ ) environmental parameters obtained by fitting environmental factors in the sample ordination space. ....	57
Figure 4.1: Principal coordinate analysis (PCoA) by unweighted unifrac based on soil pH and fertilizer treatments. Axis represent the two dimensions explaining greatest proportion of	

variances in microbial communities. p values calculated based on permutational multivariate analysis of variance.....	93
Figure 4.2: Volcano plots showing the distribution of OTUs abundance according to the adjusted p value ( $-\log_{10}$ scale) and the log twofold change comparing mix versus digestate. Black nodes in the right side represent OTUs significantly more abundant in the digestate treatments while black one in the left side represent to OTUs more abundant in mix treatments. ....	95
Figure 4.3: Ternary plots showing the relative occurrence of OTUs belonging to the top 10 class level microbes. Axis represent fertilizers (digestate, urea, control) treatments in lime and no lime treatments. Points closer to corners indicate that a great proportion of the class total relative abundance found in the fertilizer treatment. ....	96
Figure 4.4: Line graphs illustrating alpha diversity based on Pielou evenness, observed OTU numbers and Shannon diversity based on soil pH, fertilizer (F) and sampling day (D). Error bars show mean standard deviation mean $\pm$ SD (n=3). Significant levels are displayed as * $p < 0.05$ . ....	99
Figure 4.5: The temporal change of 16S rRNA, AOA, AOB, nosZ and nirS gene copy numbers in soil pH and fertilizer treatments. Error bars show mean standard deviation mean $\pm$ SD (n=3). Significant levels are displayed as * $p < 0.05$ , ** $p < 0.01$ , *** $p < 0.001$ . ....	102
Figure 5.1: The effect of fertilizer on dry weight of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation (n=3). Abbreviations; S-Soil, F-Fertilizer. ....	132
Figure 5.2: : Effect of fertilizer on temporal variations in $\text{NO}_3^-$ -N concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer.....	133
Figure 5.3: : Effect of fertilizer on temporal variations in $\text{NH}_4^+$ -N concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer.....	134
Figure 5.4: Effect of fertilizer on temporal variations in $\text{P}_2\text{O}_5$ concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer. ....	135
Figure 5.5: Effect of fertilizer on temporal variations of soil pH in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer. ....	136
Figure 5.6: Effect of fertilizer on temporal variations of leachate $\text{NO}_3^-$ content in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer. ....	137
Figure 5.7: Effect of fertilizer on temporal variations of $\text{NH}_4^+$ leachate content in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer. ....	138

Figure 5.8: Effect of fertilizer on nitrogen mass balance in percentage in the Kamishihoro and Arakida soil. .... 139

Figure 5.9: Temporal changes in soil N<sub>2</sub>O emissions in Kamishihoro and Arakida fertilizer treatments. Error bars show mean standard deviation mean  $\pm$  SD (n=3). .... 140

## Supplementary Figures

Figure S3.1: Cladogram showing significantly abundant taxa of shared OTUs based on land use, season and fertilizer. Taxa with significantly different abundance are represented by red and green dots. From the center outwards each ring represents kingdom, phylum, class, order, family and genus levels. The colored shadows represent trends of significant ( $p < .05$ ) taxa based on Kruskal–Wallis test.....	81
Figure S4.1: The relative abundance of phylum level OTUs that significantly increased in digestate treatments in L and NL treatments based on volcano plots.....	121
Figure S4.2:Volcano plots showing the distribution of OTUs abundance according to the adjusted p value ( $-\log_{10}$ scale) and the $\log_2$ fold change between lime and no lime treatments. Each point represents an individual OTUs. Black points indicate OTUs that were not significant ( $p > 0.05$ ), blue dots indicate OTUs that were significant ( $p < 0.05$ ), green dots indicate OTUs that had a $\log_2$ old change that was greater or less than 2 and red dots indicate OTUs that were significant ( $p < 0.05$ ) and had a $\log_2$ fold change that was greater or less than 2.....	122
Figure S4.3:Relative abundance of the most abundant microbes at phylum level in lime and no lime treatments.....	123
Figure S4.4:Relative abundance of the most abundant microbes at phylum level in digestate..	123
Figure S4.5:The temporal change of soil pH in control, digestate and urea treatments. Error bars show mean standard deviation $\text{mean} \pm \text{SD}$ ( $n=3$ ). Significant levels are displayed as * $p < 0.05$ , ** $p < 0.01$ , *** $p < 0.001$ .....	124
Figure S4.6:Relative abundance of microbes at phylum level in digestate.....	125
Figure S5.1:The effect of fertilizer on carbon content of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; S-Soil, F-Fertilizer. ....	153
Figure S5.2:The effect of fertilizer on nitrogen content of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; S-Soil, F-Fertilizer. ....	154

## Supplementary Tables

Table S3.1: The significant effects of land use on soil physical and chemical properties. Asterisks indicate a significant effect of each treatment (***) 0.001, ** 0.01, * 0.05) and “ns” shows “no significance” based on ANOVA. ....	78
Table S3.2: The significant effects of season on soil chemical properties in cropland and grassland. Asterisks indicate a significant effect of each treatment (***) 0.001, ** 0.01, * 0.05) and “ns” shows “no significance” based on ANOVA. ....	79
Table S3.3: Correlation analysis (spearman) among soil properties (pH, bulk density-BD, nitrogen-N, potassium-K, magnesium-Mg, cation exchange capacity-CEC, calcium-Ca and humus) and selected phyla based on land use (cropland and grassland) and season treatments. Selected microbes showed significant increase in each treatment. Red color shows significant ( $p < .05$ ) interactions. ....	80
Table S4.1: Digestate and Kamishihoro soil chemical properties .....	121
Table S5.1: Physical and chemical properties of Kamishihoro and Arakida soils (means $\pm$ standard deviation, $n=3$ ). ....	151
Table S5.2: Chemical properties of anaerobic digestate and separated solid and liquid fractions. ....	152



# **The use of anaerobic digestate from animal wastes and its impacts on soil nutrients and microbiome**

## **Chapter 1 Introduction**

### **1.1 Background**

Soil microbes are one of the largest pools of biodiversity on Earth (Bhattacharyya, 2012). Soil microbes are important towards the maintenance of soil health, which is defined as the “continuous capacity of soil to function as a vital living system” (Sathya et al., 2016). These microbes maintain soil health through the provision of ecosystem services (Singh et al., 2014; Yuliar et al., 2015). Some of these services include (Davison, 1998; Marque et al., 2014) 1) nutrient cycling such as carbon, nitrogen 2) production of plant growth hormones 3) reduced plant pathogen activity 4) enhanced soil structure 5) mineralization of pollutants. The maintenance of soil health has been identified as a central part in the establishment and maintenance of sustainable agricultural practices (Sathya et al., 2016). To ensure sustainable establishment of ecosystem services soil microbes should be diverse.

Microbial diversity is defined as the different types of microbes that are present in a specific habitat (Hendrick et al., 2000). Microbial diversity measurements include three main aspects; richness (microbes present in each habitat), evenness (relative abundance of microbes in each habitat) and community composition (the identity of the microbes in each habitat) (Hooper et al., 2005). High microbial diversity has been associated with sustainable provision of ecosystem services, increased soil health and microbial stability (Bell et al., 2005; Loreau, 2010). In contrast, low microbial diversity has been associated with the loss of soil health, microbial stability and related ecosystem services (Bell et al., 2005; Trivedi et al., 2016). Essentially for the sustainable creating and maintenance of soil health and related ecosystem service, high microbial diversity should be established. Despite the significance effect of microbial diversity to the provision soil health and ecosystem services, there is lack of adequate research and information on the effect of agricultural management practices on different aspects of the soil microbial diversity.

To address this gap in research, the current research sought to understand the effect of different agricultural management practices (land use, seasonality, fertilizers, soil pH) on soil microbial properties (abundance, composition, diversity), functions and stability.

## 1.2 Research Objectives

The objectives of this research were to

Determine the effect of land use, seasonality and fertilizer application on soil microbial communities and related functions in agricultural soils; Chapter 3.

Determine the effect of soil pH on microbial stability against the application of anaerobic digestate from dairy wastes; Chapter 4.

Determine the impact of separating digestate on soil and plant nutrients, nitrogen use efficiency, Japanese spinach yields, N<sub>2</sub>O emissions and microbial abundance in agricultural soils; Chapter 5.

## Thesis Structure

The thesis is divided into 6 chapters.

Chapter 1 This chapter introduces the importance of soil microbes in the agricultural environment. The chapter also highlights the current gap in research that will be addressed by the current research.

Chapter 2 This chapter is a literature review that describe the current available information on soil microbes and their importance in agriculture with emphasis on the nitrogen cycle. The chapter also highlights the importance of microbial diversity and factors that may influence microbial diversity within the agricultural environment.

- Chapter 3 This chapter reports on the results of soil sampling that was conducted in farmers' fields in Kamishihoro, Hokkaido, Japan. Sampling was conducted to determine the effect of land use, seasonality and fertilizer application on soil microbial communities and related functions in agricultural soils.
- Chapter 4 This chapter reports on the results of an incubation experiment that was designed to determine the effect of soil pH on microbial stability against the application of anaerobic digestate from dairy wastes. Two levels of soil pH were established during the experiment i.e., 6.5 (with lime) and 5.5 (without lime).
- Chapter 5 This chapter reports on the results of an incubation experiment that was designed to determine the impact of separating anaerobic digestate on soil nutrients, crop yields and microbial abundance in agricultural soils. Anaerobic digestate was separated into solid and liquid portions using a flocculant.
- Chapter 6 This chapter summarizes major findings from the research and recommendations for future research.

## Chapter 2 Literature Review

### 2.1 Role of soil microorganisms in agriculture

The agricultural sector is one of the largest engineered ecosystems, consisting of a third of the global land area (Zhang et al., 2007). Agriculture is an essential component for food, shelter and fiber production on Earth (Muhamad et al., 2020). There is an expected increase in demand of agricultural products due to the rapid increase in human population, with projections showing that the world population is likely to reach 9.7 billion people by 2050 (Roser, 2014). Soils are the most essential part of the agricultural production system, as the main source of plant nutrients (Tahat et al., 2020). Establishment and maintenance of soil health is significant towards provision of soil nutrients to plants. Soil health is defined as “the capacity of a soil to function as a vital living system within ecosystem and land use boundaries to sustain plant and animal production, maintain or enhance water and air quality, and promote plant and animal wellbeing” (Doran and Zeiss, 2000). Various researchers have identified that soil microorganisms play vital roles in the establishment and maintenance of soil health (Muller et al., 2016). Soil microorganisms directly influence soil health through provision of ecosystem services that offer many beneficial services within the agricultural environment (Singh et al., 2014).

Ecosystem services have been defined as “the conditions and processes through which natural ecosystems, and the species that make them up, sustain and fulfill human life” (Daily, 1977). Within the agricultural environment, ecosystem services are determined by the soil health, which is established and maintained by soil microorganisms (Kibblewhite et al., 2007). Ecosystem services provided by soil microorganisms include provisioning and regulatory services (Table 2.1).

Table 2.1: Role of soil microbes in provisioning and regulating services provided by soil ecosystems (adapted from Dominati et al., 2010).

Soil service	Role of soil microbes
Provisioning services – products obtained from the agricultural ecosystems	
Physical support	Organic matter decomposition, nutrient cycling, and improved soil aggregation.
Plant growth medium	Soil microbes mobilize nutrients from insoluble minerals to support plant growth
Regulating services – provide a stable and healthy environment	
Buffering water flows	Soil macropores are formed by soil biota which depend on microbes for food and fuel.
Detoxification and waste cycling	Achieved through microbial mineralization and immobilization of wastes and toxic substances. These microbes depend on soil nutrient availability, which rely on soil microbes
Filtering contaminants	Soil microbes influence soil properties such as hydrophobicity and wettability which effect ability of soil to filter contaminants
Biological control of pests, weeds and pathogens	Soil microbes such as bacteria increase nutrient availability to plants and outcompeting invading pathogens.
Carbon storage and regulation of greenhouse gas emissions	Methane producing and consuming microbes and denitrifying bacteria regulate soil nitrous oxide (N <sub>2</sub> O) and methane (CH <sub>4</sub> ) emissions.
Nutrient cycling	Soil microbes drive the Carbon and Nitrogen nutrient cycles.

## 2.2 Soil microorganisms and the nitrogen cycle

Nitrogen is an important element in agricultural production, constituting the main component of plants organic compounds (Stark and Richards, 2008). Although 78% of the atmosphere consists of N gas, plants cannot take up the nutrient in this form (Hirsch and Mauchline, 2015). A significant part of N occurs in organic forms and it must be converted to inorganic forms i.e.  $\text{NH}_4^+$  ( $\text{NH}_4^+$ ) and nitrate ( $\text{NO}_3^-$ ) for plant uptake (Boyle et al., 2008). Through the N cycle organic N is converted to inorganic forms, availing N for plants and removing excess amounts from the system (Dixon and Kahn, 2004). Soil microorganisms play important roles in all processes of the N cycle thus regulating plant N availability (Yao et al., 2011). The N cycle involves different process which include  $\text{N}_2$  fixation ( $\text{N}_2$  to  $\text{NH}_4^+$ ), nitrification ( $\text{NH}_4^+$  to  $\text{NO}_3^-$ ), denitrification ( $\text{NO}_3^-$  to  $\text{N}_2$ ) and ammonification (Figure 2.1) (Dueri et al., 2007).

### *N Fixation*

Nitrogen fixation involves the conversion of  $\text{N}_2$  gas from the atmosphere to ammonia ( $\text{NH}_3$ ) Fixation is carried out by free living and symbiotic bacteria and archaea that contain the nitrogenase enzyme (Franché et al., 2009). Some of the bacteria involved in the process are *Azotobacter*, *Bacillus Pseudomonas* among others (Kennedy et al., 1997; Li et al., 1992; Rozycki et al., 1999).

### *Nitrification*

Nitrification is the process through which  $\text{NH}_3$  is then converted to  $\text{NH}_4^+$  and then  $\text{NO}_3^-$  in a twostep process (Thomson et al., 2012) (Figure 2.1). The first step of the process, which involves conversion of  $\text{NH}_4^+$  to nitrites ( $\text{NO}_2^-$ ) is considered the most significant and rate limiting step. The process is driven by  $\text{NH}_3$ -oxidizing microorganisms which include  $\text{NH}_3$ -oxidizing bacteria (AOB) and  $\text{NH}_3$ -oxidizing archaea (AOA) that contain  $\text{NH}_3$  monooxygenase, an enzyme that converts  $\text{NH}_4^+$  to  $\text{NO}_2^-$  (Prosser and Nicol, 2008; 2012). The AOB microbes include *Nitrosomonas*, *Proteobacteria*, *Nitrosospira* among others (Norton, 2014). The AOA microbes include *Euryarchaeota* and *Crenarchaeota* among others (Zhou et al., 2015). The second step of the nitrification process (nitration) involves the conversion of  $\text{NO}_2^-$  to  $\text{NO}_3^-$ , which is driven by nitrobacter bacteria (Moreira and Siqueira, 2006). The  $\text{NO}_3^-$  form of N is readily taken up by plants and is very mobile in the soil and easily lost through leaching. This process of the N cycle

is responsible for  $\text{NO}_3^-$  water pollution and related consequences such as reduced crop yields, agricultural overapplication of N fertilizers, soil acidification and eutrophication (Davidson and Hackler, 1994).

### *Denitrification*

Denitrification is the stepwise conversion of  $\text{NO}_3^-$  or  $\text{NO}_2^-$  through nitric oxide (NO) to  $\text{N}_2\text{O}$  or  $\text{N}_2$  gas (Figure 2.1). The process is driven by different types of microbes including bacteria, archaea and eukaryotes (Shoun et al., 2012). The first step involves the conversion of  $\text{NO}_3^-$  to  $\text{NO}_2^-$  which is controlled by bacteria and fungi that contain the membrane-bound  $\text{NO}_3^-$  reductase (*Nar*) or the periplasmic  $\text{NO}_3^-$  reductase (*Nap*) (González et al., 2006). This is followed by conversion of  $\text{NO}_2^-$  to  $\text{NO}^-$  driven by bacteria that encode for the nitrite reductase (*Nir*) enzyme. More specifically, there are two different types of the *Nir* enzyme have been found among denitrifiers. The copper containing (Cu-Nir) *nirK* gene and heme *c* and heme *d*<sub>1</sub> (*cd*<sub>1</sub>-Nir) *nirS* gene. The two are not functionally different and do not occur in the same cell (Priemé and Tiedje, 2002). Microbes involved in the process include *Proteobacteria*, *Firmicutes*, *Bacteroidetes*, among others (Heylen et al., 2006). The produced nitric oxide (NO) is a very reactive gas and therefore soil accumulation should be prevented. The process by which NO is converted to  $\text{N}_2\text{O}$  gas (during incomplete denitrification) is facilitated by prokaryotes which encode for nitric oxide reductase (*Nor*) (Figure 2.1) (Braker and Tiedje, 2006). Three different types of the *Nor* gene have been characterized; *cNor*, *qNor* and *qCuANor* (Vries S and Pouvreau, 2007; Spanning 2011). Some microbes involved in the process include *Paracoccus halodenitrificans*, *Pseudomonas*, *Alcaligenes faecalis* among others (Sakurai and Sakurai, 1998; Zumft et al., 1994). This step in the denitrification process has received significant attention as most of the  $\text{N}_2\text{O}$  gas produced from agricultural soil has been attributed to denitrification (Signor and Cerri, 2013).  $\text{N}_2\text{O}$  is a greenhouse gas with a global warming potential of 298 (Signor and Cerri, 2013). Complete denitrification will convert NO to  $\text{N}_2$  gas, resultantly removing N from the soil without the environmentally harmful  $\text{N}_2\text{O}$  gas emission (Butterbach-Bahl et al. 2013). This is a process that is driven by bacteria and archaea which encode for the nitrous oxide reductase (*nosZ*) enzyme (Burger and Matiasek 2009). Some of the microbes that encode for the gene are *alpha*, *beta* and *gamma* *proteobacteria* among others (Orellana et al., 2014).




Figure 2.1: Partial soil nitrogen cycling processes with emphasis on nitrification, denitrification and related biomarker genes

## 2.3 Soil microbial diversity and stability

### 2.3.1 Diversity

Microbial diversity is defined as the number of different microbes in a given habitat (Hendrick et al., 2000). Components of microbial diversity include richness (number of microbes present in each habitat), evenness (relative abundance of microbes in a given habitat), community composition (identity of microbes present in a given habitat) (Hooper et al., 2005). Microbial diversity analysis includes the specific number of different microbes present and how even they are distributed. For example, a community with a high number of different microbial species is more diverse. However, a community with high evenness is considered more diverse than one that is less even with the same number of microbial species (Hendrick et al., 2000).



### 2.3.1.1 Components of microbial diversity

#### 2.3.1.1.1 Richness

High microbial species richness has a positive effect on overall ecosystem functioning. Positive species interactions have been observed in habitats that have high species richness (Chapin et al., 1994). High richness leads to establishment of microbes with different ecological niches and feeding preferences, which causes efficient use of resources and higher productivity (Chapin et al., 1994). Jiang (2007) and Langenherder (2010) reported that changes in microbial richness could have a direct effect on ecosystem services. Bell et al. (2005) conducted research on the effect of bacterial richness on ecosystem services and their results showed that increasing bacterial richness enhanced rates of respiration. These results demonstrate the importance of microbial richness within the agricultural environment

#### 2.3.1.1.2 Evenness

Microbial evenness has a significant impact on provision of agricultural ecosystem services. Research by Wittebole et al. (2009) showed that evenness played a significant role in establishing and maintaining functional stability in denitrifier microbes. Hillebrand et al. (2008) similarly found that microbial evenness and stability were directly related. High evenness enhanced microbial stability and related ecosystem services compared to low evenness. Specifically, low evenness values indicate that a specific habitat is dominated by few microbes, which reduces microbial stability. The resistance to external disturbances in such conditions would be achieved only if the dominant species are tolerant to disturbance (Hillebrand et al., 2008).

#### 2.3.1.1.3 Microbial community composition

Microbial community composition has a direct effect on provision of agricultural related ecosystem services. A report by Strickland et al. (2009) found that biogeochemical processes and microbial stability were influenced by microbial community composition. Specifically, Bell et al. (2005) and Salles et al (2012) reported that if changes in microbial community composition caused the loss or introduction of functional groups such as denitrifiers, related ecosystem services would be affected. Similarly, Hallin et al. (2009) found a positive correlation between the soil's bacterial community composition and the N cycle related soil functions such as nitrification. For sustainable provision of ecosystem services, management practices that reduce negative changes in microbial community composition should be established.

In summary high microbial diversity ensures the continued provision of agricultural ecosystem services. Additionally, high diversity increases the systems tolerance to disturbances such as land use, seasonal changes, pH and fertilizer application (Griffiths and Philippot, 2013; Loreau, 2010). Decreased diversity would reduce the systems stability and cause a loss of related ecosystem services (Trivedi et al., 2016).

### 2.3.2 Stability

Stability is defined as soil microbe's response to disturbances. Stability is usually measured based on two main concepts: resistance and resilience. Resistance is the degree to which microbes withstand a disturbance and resilience is the rate at which the community returns to its original state after a disturbance (Allison and Martiny, 2008; Shade et al., 2012). High resistance and resilience is accompanied with increased microbial stability and vice versa (Griffiths et al., 2001). Bearing in mind the significant role microbes play in providing agricultural ecosystem services, the effects of disturbances on their stability an associated function should be considered.

Research show that microbial diversity modulates stability. For instance, higher diversity values have been found to increase microbial stability and vice versa (Wertz et al., 2007). Girvan et al. (2005) conducted research on the effect of different toxins (copper sulphate and benzene) on the stability of soils with different natural diversity levels. Based on their results there was a positive relationship between microbial diversity and stability. On the contrary research by Wertz et al. (2007) found that a decline in diversity did not influence the resistance and resilience of key soil microbial groups following a disturbance. Therefore, research needs to be carried out to clarify the relationship between microbial stability and diversity.

## 2.4 Factors that affect soil microbial properties

### 2.4.1 Land use

Agricultural lands make up one third of the global land area (Ellis and Ramankutty, 2008; Ramankutty et al., 2008). Some of the common agricultural land use types include croplands for yields and grassland for animal feed (Yu et al., 2017). Cropland and grassland are characterized by specific management practices which impact soil microbial diversity and stability (Ramankutty et al., 2018).

Cropland conventional tillage practices are usually focused on increasing crop yields at the cost of soil health (Tilston et al., 2010). Common practices associated with conventional tillage practices include inorganic chemical fertilizers use and continuous tillage and monoculture (Zingore et al., 2005). However, grassland management systems are characterized by very little if any inorganic fertilizer use, reduced tillage and high plant diversity (Bissett et al., 2014; Lauber et al., 2009).

#### 2.4.2 Agricultural conventional tillage practices

Research show that conventional tillage practices such as inorganic chemical fertilizer use, continuous tillage and monoculture have a negative effect on soil physical, chemical and microbial properties (Zingore et al., 2005).

Inorganic fertilizer use changes the soil nutrient content and decreases soil pH, subsequently affecting soil microbial abundance and diversity (Bünemann et al., 2006). There have been reported positive and negative effects of inorganic N fertilizers within the agroecosystem. Apart from increasing crop yields, inorganic N fertilizer use has been associated with increased abundance of some functional genes in the soil. Zhang et al. (2016) reported that inorganic N fertilizer increased the abundance of AOB. However, there is extensive research showing the negative effects of chemical fertilizers on soil health and functional gene abundance. Tiston et al. (2010) reported that inorganic N fertilizers were the main cause of soil acidification on agricultural soils. Research show that high levels of acidity negatively affect soil nutrient immobilization and mineralization (Fageria and Baligar, 2008). Acidification increases loss of microbial biomass, essential plant nutrients (Lucas et al., 2011), increased  $\text{NO}_3^-$  leaching (Otto et al., 2016) and agricultural  $\text{N}_2\text{O}$  emissions (Clough et al., 2003). Additionally, inorganic N fertilizers have a negative effect on some functional gene abundance in agricultural soils. Sterngen et al. (2015) and Yaying et al. (2013) reported that fertilizers reduced the abundance of  $\text{NH}_3$  oxidizing archaea and nitrous oxide reductase, which are related to the N cycle.

Soil tillage practices affect soil physical, chemical and biological properties, subsequently impacting soil health and productivity (Mathew et al., 2012). Cropland related conventional tillage practices disrupt soil aggregates and have a negative effect on soil physiochemical properties (Alam et al., 2016). Various scientists have reported that conventional tillage practices have a negative effect on soil bulk density, soil aggregate distribution and water holding capacity (Alam

et al., 2016; Trajan and Linden et al., 1998; Wander et al., 1998). These changes create shifts in soil microbial habitats and community structures (Alguacil et al., 2008; Feng et al., 2003; Kandeler et al., 1999).

Continuous monocropping is a common conventional agricultural management, practiced to ensure maximum crop yield production. However, research shows that the practice has a negative effect on soil physiochemical properties and microbial habitat (Zhou et al., 2012). Fu et al. (2016) conducted research on the effect of continuous monoculture on soil microbial properties and enzyme activities. Based on their results monocropping decreased most quality indicators which included substrate richness, Shannon diversity, soil urease, available N and potassium. Gu et al. (2012) similarly reported that continuous monocropping reduced soil microbial diversity.

#### 2.4.3 Effect of agricultural management practices on microbial diversity and stability

Conventional tillage practices such as inorganic N fertilizer use, frequent tillage and monoculture have been linked to decreased microbial diversity and stability (Zingore et al., 2005).

Inorganic N fertilizer use has been associated with reduced soil diversity compared to use of organic fertilizers such as anaerobic digestate. Wu et al. (2020) conducted research on the effect of organic fertilizer application and reduced chemical N fertilizers on soil properties and bacterial community of an agricultural soil. Their results showed that organic fertilizer treatments significantly increased bacterial diversity compared to chemical N fertilizer treatments. Similar results were reported by Sun et al. (2015) and Yuan et al. (2018).

Various researchers (Ceja-Navaro et al., 2010; Ovreas and Torsvik, 1998) have reported that increased tillage intensity in croplands reduced microbial stability and diversity compared to reduced or no tillage that was practiced in grasslands. In agreement, Quadros et al. (2012) conducted research on the effect of tillage on soil microbial composition and diversity. Based on their results, frequent tillage had significantly lower Shannon diversity index values, compared to treatments with reduced tillage.

The practice of monoculture has been found to have negative effects on soil microbial stability and diversity. Zhao et al. (2018) conducted research on the long term effect of monoculture on soil and microbial properties. Based on their results, continuous cropping decreased soil bacterial alpha

diversity. In agreement, Liu et al. (2014) conducted research on microbial community diversities and taxa abundances in soils with potato monoculture. They found that increased monocropping reduced microbial diversity. In addition, Tiemann et al. (2015) found that increased above ground diversity, commonly observed in grasslands stimulated microbial functionality in the soil. These results show that although conventional tillage practices may be effective in increasing crop yields, they may not be sustainable for maintenance of soil health and microbial diversity.

#### 2.4.4 Seasonality

Seasonal changes are usually accompanied with shifts in rainfall, temperature and soil nutrient content which affect soil microbial properties (Kim et al., 2016; Luo et al., 2019). Additionally, seasonal changes are related to farmer management practices such as planting, fertilizer application schedules and harvesting that affect the soil physical, chemical and microbial properties (Bevivino et al., 2014). These seasonal changes have been identified as some of the main factors that drive changes in soil microbial structure and diversity (Shigyo et al., 2019).

Previous reports have identified the effect of seasonal changes on the soil microbial community within the agricultural environment (Bevivino et al., 2014; Lacerda-Junior et al., 2019; Li et al., 2019). However due to differences in soil physical, chemical and microbial properties, there have been contrasting results on the effects of seasonal treatments on specific soil microbes. Li et al. (2019) conducted research on the effect of seasonal changes i.e., summer, autumn and spring in microbial diversity and its relationship with soil chemical properties. Their results showed that there was a significant increase in microbial properties in summer due to the high rainfall and temperatures experienced during the season. Bevivino et al. (2014) conducted research in the soil bacterial community response to differences in agricultural management and seasonal changes in the Mediterranean region. Their results showed that seasonal variations affected soil chemical and structural fractions of the soil organic matter. Additionally, soil bacterial communities were affected by seasonal changes. For example, *Bacilli* and *Actinobacteria* were identified as the dominant microbes in spring while *Actinobacteria* and *Proteobacteria* were the dominant microbes in summer. Lacerda-Junior et al. (2019) conducted research on the effect of land use and seasonal effects on the soil microbiome. Their results showed a significant increase in

*Actinobacteria* and *Cyanobacteria* during the dry season, while *Proteobacteria*, *Acidobacteria* and *Bacteroidetes* significantly increased during the rainy season.

Therefore, research needs to be conducted to clarify the effect of seasonal variations on specific soil microbes within the agricultural environment, to develop sustainable management practices that are specific to each season (Shigyo et al., 2019).

#### 2.4.5 Fertilizers

Fertilizers are organic or inorganic based compounds that add nutrients to the soil (Maximum Yield Inc. 2019). Agricultural fertilizer use has been credited with increased soil fertility and crop yields (Vitousek et al., 2002). The ‘Green Revolution’ is credited for the significant increase in inorganic fertilizers in agriculture (Erisman et al., 2008). The term ‘Green Revolution’ was developed by William S. Gaud in 1968 and related to the introduction of various technologies and policies in developing countries with assistance from developed nations to increase agricultural food production (Conway, 1997). One of the main methods to achieve this objective was the use of chemical fertilizers to improve soil nutrient levels and subsequently crop yields (Parayil, 1992). The use of inorganic fertilizer involves animal excreta and plant residues. Farmers process animal excreta and use it in different forms within the agricultural environment. Some of the common processes and forms used include composting (the anaerobic fermentation of animal manure), anaerobic digestate (by product of the biogas production process) (Bernal et al. 2009; Yabe 2013) among others.

Agricultural fertilizer use is an important factor in ensuring global food security (Roberts, 1999). According to Roser (2014) the world global population is expected to reach 9.7 billion by 2050. As expected, this will increase global food demand by 70%, increasing dependency on fertilizers (Dawson and Hilton, 2011; FAO, 2012).

##### 2.4.5.1 Inorganic nitrogen fertilizers

Inorganic N fertilizers were first generated during the 20<sup>th</sup> century using the Harber-Bosch process developed by Fritz Haber (Erisman et al., 2008). Considering the rise in human population, the current society would not exist with the use of inorganic N fertilizers, to ensure food security (Erisman et al., 2008). The N is one of the nutrients that are essential for plant growth and development (Sharma and Bali, 2017). The nutrient is involved critical plant processes such as leaf

expansion, plant growth and biomass production. More specifically, plant molecules such as chlorophyll, amino acids and nucleic acids contain N and are necessary for biological processes such as photosynthesis, carbon metabolism, among others (Crawford et al., 2002; Frink et al., 1999). In addition, N application improves root growth and length, plant nutrient uptake and dry mass production (Diaz et al., 2006; Stitt and Krapp, 1999). Therefore, N is considered one of the main limiting nutrients for plant development in agriculture (Rütting et al., 2018).

The quantity of N fertilizer in agricultural soils has been increasing over the last 50 years (Zahoor et al., 2014). This practice as led to excessive inorganic N fertilizer application, which had led to various environmental and ecological problems (Ahmad et al., 2017). Some of these include eutrophication, greenhouse gas emissions (N<sub>2</sub>O), soil acidification among others (Bouwman et al., 2005; Guo et al., 2010).

#### 2.4.5.2 Organic anaerobic digestate fertilizer

Due to the negative effects of inorganic N fertilizer use on the environment and agricultural production sustainability, farmers have been turning to organic fertilizers. The biogas production process is a technology that is gaining popularity where organic matter is converted to biogas and the nutrient rich byproduct anaerobic digestate (Risberg et al., 2016). More specifically, biogas production is a biological process involving organic matter decomposition and stabilization under anaerobic conditions. One of the main benefits of anaerobic digestate is that various organic materials can be used as feed stock for the process. Some of the organic materials used as feedstock include animal wastes, energy crops, plant biomass, food industry and agricultural wastes (Appels et al., 2011; Yan et al., 2008). The microbial based process produces biogas (methane and carbon dioxide) and anaerobic digestate. Biogas is used as a source of energy for producing electricity, heat and vehicle fuel (Sun et al., 2016). Anaerobic digestate is considered a fertilizer with the potential of reducing the use of fossil fuel dependent inorganic N fertilizers (Holm-Nielsen et al., 2009; Walsh et al., 2012).

##### 2.5.5.2.1 Biogas production process

The biogas production process consists of four phases: enzymatic hydrolysis, acidogenesis and methanogenesis (Odwuge et al., 2020).

During the enzymatic hydrolysis process, large polymers are broken down by facultative and obligate anaerobic bacteria (*Streptococcus* and *Enterococcus*). Polysaccharides are degraded to

oligosaccharides and monosaccharides, proteins are degraded to peptides and amino acids, lipids are degraded to glycerol and fatty acids (Paritosh et al., 2017).

During acidogenesis, hydrolysis products are fermented to volatile fatty acids by facultative anaerobic bacteria (*Ruminococcus*, *Paenibacillus*, *Clostridium*). The volatile fatty acids from the process include acetate, valerate, butyrate, isobutyrate and propionate. Furthermore, NH<sub>3</sub>, hydrogen and carbon dioxide gas are produced along with fatty acids (Paritosh et al., 2017).

In the methanogenesis phase, methane is produced through reduction of carbon dioxide and fermentation of acetic acid. Methane is produced from acetic acid by acetolactic methanogens archaea. Additionally, methane is produced through the reduction of carbon dioxide by hydrogen by hydrogenotrophic methanogens. About 70% of the methane is produced from acetic acid fermentation while 30% is produced from carbon dioxide reduction (Karakashev et al., 2005). Anaerobic digestate is produced as a byproduct of the process.

#### 2.4.5.2.2 Anaerobic digestate characteristics and utilization

Anaerobic digestate is composed of a mixture of water, organic, inorganic substrates and various nutrients (Möller et al., 2015). Anaerobic digestate fertilizer contains higher mineralized nutrients than untreated feedstocks (Arthurson, 2009). During the biogas production process, nutrient transformations of the organic substrate occurs. Easily decomposable carbon compounds are removed, leaving behind compounds that are more difficult to decompose (Glowacka et al., 2020). The N in the substrate is transformed to NH<sub>4</sub><sup>+</sup>. Gemmeke et al. (2009) reported that the anaerobic digestate NH<sub>4</sub><sup>+</sup> content could increase by a factor of three depending on substrates used. Essentially the NH<sub>4</sub><sup>+</sup> content of anaerobic digestate is generally higher than the substrates, allowing the nutrient to be immediately available for plant uptake (Arthurson, 2009). Additionally, pathogenetic microbes such as parasites, viruses and bacteria are destroyed. Anaerobic digestate has an increase in vitamin and amino acid content, decrease in C/N ratio and no changes in macro and micronutrients (Glowacka et al., 2020).

Various researchers have reported increased soil nutrient and yield content due to soil anaerobic digestate application. Tampio et al. (2015) reported that anaerobic digestate fertilizer had a high NH<sub>4</sub><sup>+</sup> content (4.07 mg N g<sup>-1</sup>). Šimon et al. (2015) similarly reported that anaerobic digestate application resulted in higher wheat yields due to increased soil NH<sub>4</sub><sup>+</sup> content. Risberg et al. (2019) conducted research on characterization of anaerobic digestate and its effects on soil microbial



activity. Based on their results, anaerobic digestate increased the soil organic carbon and N content. Other researchers have found that anaerobic digestate soil application has been linked with increased soil pH, phosphorous, potassium, magnesium, soil carbon sequestration and increased water holding capacity (Pivato et al., 2016; Pranagal et al., 2019; Smith et al., 2014). Therefore, the application of anaerobic digestate is considered a practical substitute for reducing use of inorganic N fertilizers in the agro ecosystem (Risberg et al., 2016; Weiland, 2009).

#### 2.4.5.2.3 Solid Liquid Separation of Anaerobic digestate

The use of anaerobic digestate as an organic fertilizer has been increasing in recent years. However there have been some challenges associated with anaerobic digestate use as a fertilizer. One of the main challenges that has been observed with its use as a fertilizer is the high costs related to its transport and storage, compared to chemical fertilizers (Drosg et al., 2015). The high costs are caused by anaerobic digestates' low nutrient and large volume compared to chemical fertilizers. Bojesen et al. (2014) reported that transportation of substrates and anaerobic digestate are responsible for approximately 33% of the biogas production process cost. Another challenge is that due to anaerobic digestates' high moisture content, there is a risk of temporary flooding associated with its use during the rainy season especially in humid environments. In addition, farmers are interested in specific nutrient concentrations of anaerobic digestate so that they can apply the required amounts of nutrients based on soil requirements (Drosg et al., 2015). To address these challenges, anaerobic digestate can be separated to remove the solid and liquid fractions before field application (Heviánková et al., 2015). Research shows that separated fractions have substantially lower transportation costs due to reduced water contents, especially for the solid fraction. Additionally, the solid-liquid separation results in changes in nutrient distribution between solid and liquid fractions allowing for more precise nutrient applications based on soil requirements (Figure 2.2).




Figure 2.2: Distribution of the principal constituents after solid–liquid separation (data based on own investigations and various references. Abbreviations; TN-Total Nitrogen, P-Phosphorous, K-Potassium, C-Carbon (Adapted form Bauer et al., 2009)

#### 2.4.5.2.4 Anaerobic digestate Flocculation

Flocculation is a chemical treatment used to separate anaerobic digestate into solid and liquid fractions (Nowostawska et al., 2005). Anaerobic digestate particles have negative charges that tend to repel each other and therefore do not sediment. This characteristic makes the process of anaerobic digestate solid liquid fraction a bit more complicated (Fuchs et al., 2010). To enhance aggregation of the solid fraction, multivalent cations that cause coagulation are added in addition to polymers that enhance flocculation ( Hjorth et al., 2010).

Anaerobic digestate flocculation involves two steps. In the first step, the flocculant, a water soluble metal salt such as iron-III-chloride, is mixed with anaerobic digestate. Cations formed from the metal salt react with the negatively charged anaerobic digestate particles to form bigger particles (Figure 2.3). A flocculant aid polymer is usually added to increase the size of the solid particles for separation. The flocculant aid polymer is a long negatively charged chain where metal cations from the polymer connect to form a high particle for more efficient solid-liquid separation (Fechter and Kraume, 2016).




Figure 2.3: Principle of flocculation (Adapted from Fechter and Kraume, 2016)

#### 2.4.5.3 Effect of fertilizers on microbial abundance and stability

The use of organic and inorganic N fertilizers has a direct and indirect effect on soil physical and chemical properties (Dong et al., 2014; González-Chávez et al., 2010). Changes in soil physical and chemical properties have been identified as the main drivers in changes in microbial abundance and stability in agricultural soils (Martin et al., 2015). Thus, the use of fertilizers influences soil microbial abundance and stability (Bell et al., 2015).

The use of inorganic N fertilizers has been associated with reduced microbial abundance and stability. One of the main reasons is the effect of inorganic N fertilizers on soil pH. Inorganic N fertilizer use has been associated with agricultural soil acidification, which has a negative effect on microbial abundance and diversity (Guo et al., 2010; Sun et al., 2015). Brady and Weil (1996) and Wang et al. (2012) identified pH as one of the main factors affecting the diversity and microbial community composition. Soil pH influences other soil factors such as soil nutrient availability and cations solubility which influence the soil microbial community. Fierer and Jackson (2006) conducted research on factors that influence the diversity and biogeography of soil bacterial communities. Based on their results, soil pH was one of the main factors influencing the microbial abundance and diversity. Diversity was highest in soils with near neutral pH and lowest in acidic soils.

Organic fertilizers have a high impact on the soil microbial community than chemical fertilizers (Wei et al., 2017). This can be attributed to organic fertilizers improving different aspects of soil

fertility, organic carbon content, soil aggregate stability and microbial biomass which cause shifts in soil microbial community (Darwish et al., 1995; Lazcano, 2013; Marschner, 2003). Sapp et al. (2015), Hartman et al. (2014) and Coelho et al. (2020) conducted research on the effect of anaerobic digestate as one of the treatments on the soil microbial community. Based on their results it can be concluded that anaerobic digestate fertilizers were a major driver of microbial diversity. Specifically, anaerobic digestate increased various aspects of diversity including community composition, richness and evenness. Although there are few studies that have analyzed the effect of organic fertilizers on microbial stability, research shows that high microbial diversity is associated with enhanced microbial stability (McCann, 2002). Therefore, the increased diversity with organic fertilizer use is expected to result in enhanced microbial stability.

#### 2.4.6 Soil pH

Soil pH is defined as the negative logarithm to the base of 10 (p) of the soil's hydrogen ion ( $H^+$ ) concentration i.e.,  $pH = -\log_{10}[H^+]$  (Shukla et al., 2014; Zhou, 2008). Therefore, a change in pH denotes a tenfold change in  $H^+$  concentration. The pH scale, which ranges from 0 to 14, is used to show the level of acidity ( $<7$ ) or alkalinity ( $>7$ ) (Blackson, 1989). Soil pH is defined as a master soil variable due to the significant effect it has on the soil physical, chemical and biological properties (Figure 2.4) (Minasny et al., 2016; Neina et al., 2019).




Figure 2.4: Soil biochemical processes and their relations to soil pH (Adopted from Neina, 2019)

#### 2.4.6.1 Factors that affect agricultural soil pH




##### 2.4.6.1.1 Inorganic nitrogen fertilizers

Inorganic N fertilizer use is considered one of the most efficient methods to increase the soil N content and subsequent crop yields (Chen et al., 2017). However, the increased use of inorganic N fertilizers has caused the acidification of agricultural soils, which is negatively affecting the soil physical, chemical and microbial properties (Schroder et al., 2011). About 50% of the global agricultural lands have been negatively affected by soil acidification due to the use of inorganic N fertilizers (Dai et al.2017). Soil acidification has been identified as one of the most significant yield limiting factors in agricultural soils (Sumner and Noble, 2003). The N fertilizers themselves are not acidic, but their additions to soils are acid forming (Schroder et al., 2011). The N fertilizers cause soil acidification through the nitrification process through which  $\text{NH}_4^+$  to  $\text{NO}_3^-$ , producing  $\text{H}^+$  which acidify the soil (Kariuki et al., 2007; Tang et al., 2002).

Schroder et al. (2011) conducted research on the effect of long term use of N fertilizers on soil acidification. Their results showed that the long term use of inorganic N fertilizers significantly decreases soil pH levels, enhancing soil acidity. These results agree with various researchers who have reported that the continuous use of inorganic N fertilizers caused soil acidification in agricultural soils (Bouman et al., 1995; Chien et al., 2008).

#### 2.4.6.1.2 Liming

Liming is the most common and effective method to reduce increasing soil acidification in agricultural soils (Mkhonza et al., 2020). Most of the liming materials contain Calcium which are effective in reducing soil acidity (Rasnake et al., 2002). Adding lime to the soil reduces soil acidity by neutralizing H<sup>+</sup> in the soil (Equation 1) (Bolan et al., 2003; Buni, 2014).



It has previously been reported that liming agricultural soils reduces soil acidity (Alemayehu and Taffesse, 1999). Buni (2014) conducted research on the effect of liming acidic soils on soil properties and yields of Haricot bean. Based on their results, soil pH increased with increasing lime application. In their experiment, soil pH values increased from 5.03 (soils without lime) to 6.72 (soils with lime-3750 kg CaCO<sub>3</sub> ha<sup>-1</sup>). Additionally, Temesgen, et al. (2016) conducted research on the effect of lime and phosphorous fertilizer on acidic soils in the central highlands of Ethiopia. They reported that liming at the rate of 0.55, 1.1, 1.65 and 2.2 t ha<sup>-1</sup> increased soil pH by 0.48, 0.71, 0.85 and 1.1 units, respectively.

#### 2.4.6.2 Effect of soil pH on soil microbial community

Soil pH is one of the most significant factors affecting the soil microbial community in agricultural soils (Singh, 2018). Soil pH values that are near neutral to basic tend to increase soil microbial abundance compared to acidic pH values. Msimbira and Smith (2020) reported that the ideal pH for most soil microbes in the agricultural environment was from 5.5 to 6.5. Shen et al. (2010) conducted research on the effect of long-term fertilizer application on the abundance and composition of soil bacterial communities. Their results showed that less acidic soil conditions had higher bacterial populations compared to acidic conditions. Rousk et al. (2009) conducted

research on the effect of soil pH on fungal and bacterial growth. Their findings showed that decreasing soil pH levels from 8.3 to 4.0 caused a five-fold reduction in soil bacterial growth.

Soil pH levels significantly influence the soil N cycles nitrifier and denitrifier community (Zhang et al., 2016). Various studies have showed that soil pH levels influence the niche separation of AOA and AOB. More specifically, AOA prefer acidic environments of pH values of <5.5 while AOB prefer basic soil pH (He et al., 2012; Zhang et al., 2016). Additionally, soil pH has been found to influence the denitrification *nosZ* and *nirS* abundance. Čuhel et al. (2010) and Herold et al. (2018) reported that increasing the soil pH values through liming, enhanced the abundance of *nosZ* and *nirS* genes. Considering the significance of soil pH on different aspects of the microbial community and related soil functions, consideration should be taken on the effect of farmer management practices on soil pH values.

## References

- Ahmed, M., Rauf, M., Mukhtar, Z., & Saeed, N. A. (2017). Excessive use of nitrogenous fertilizers: an unawareness causing serious threats to environment and human health. *Environmental Science and Pollution Research*, 24(35), 26983–26987. <https://doi.org/10.1007/s11356-017-0589-7>
- Alam, M. K., Islam, M. M., Salahin, N., & Hasanuzzaman, M. (2014). Effect of Tillage Practices on Soil Properties and Crop Productivity in Wheat-Mungbean-Rice Cropping System under Subtropical Climatic Conditions. *The Scientific World Journal*, 2014, 1–15. <https://doi.org/10.1155/2014/437283>
- Alguacil, M. M., Lumini, E., Roldán, A., Salinas-García, J. R., Bonfante, P., & Bianciotto, V. (2008). The impact of tillage practices on arbuscular mycorrhizal fungal diversity in subtropical crops. *Ecological Applications*, 18(2), 527–536. <https://doi.org/10.1890/07-0521.1>
- Allison, S. D., & Martiny, J. B. H. (Eds.). (2008). Resistance, resilience, and redundancy in microbial communities (Vol. 105). United States of America: Proceedings of the National Academy of Sciences. <https://doi.org/10.1073/pnas.0801925105>
- Allison, V. J., Condon, L. M., Peltzer, D. A., Richardson, S. J., & Turner, B. L. (2007). Changes in enzyme activities and soil microbial community composition along carbon and nutrient gradients at the Franz Josef chronosequence, New Zealand. *Soil Biology and Biochemistry*, 39(7), 1770–1781. <https://doi.org/10.1016/j.soilbio.2007.02.006>
- Andersson, S., Nilsson, S. I., & Saetre, P. (2000). Leaching of dissolved organic carbon (DOC) and dissolved organic nitrogen (DON) in mor humus as affected by temperature and pH. *Soil Biology and Biochemistry*, 32(1), 1–10. [https://doi.org/10.1016/s0038-0717\(99\)00103-0](https://doi.org/10.1016/s0038-0717(99)00103-0)
- Arthurson, V. (2009). Closing the Global Energy and Nutrient Cycles through Application of Biogas Residue to Agricultural Land – Potential Benefits and Drawback. *Energies*, 2(2), 226–242. <https://doi.org/10.3390/en20200226>
- Bauer, A., Mayr, H., Hopfner-Sixt, K., & Amon, T. (2009). Detailed monitoring of two biogas plants and mechanical solid–liquid separation of fermentation residues. *Journal of Biotechnology*, 142(1), 56–63. <https://doi.org/10.1016/j.jbiotec.2009.01.016>



- Bell, C. W., Asao, S., Calderon, F., Wolk, B., & Wallenstein, M. D. (2015). Plant nitrogen uptake drives rhizosphere bacterial community assembly during plant growth. *Soil Biology and Biochemistry*, 85, 170–182. <https://doi.org/10.1016/j.soilbio.2015.03.006>
- Bell, T., Newman, J. A., Silverman, B. W., Turner, S. L., & Lilley, A. K. (2005a). The contribution of species richness and composition to bacterial services. *Nature*, 436(7054), 1157–1160. <https://doi.org/10.1038/nature03891>
- Bell, T., Newman, J. A., Silverman, B. W., Turner, S. L., & Lilley, A. K. (2005b). The contribution of species richness and composition to bacterial services. *Nature*, 436(7054), 1157–1160. <https://doi.org/10.1038/nature03891>
- Bernal, M. P., Albuquerque, J. A., & Moral, R. (2009). Composting of animal manures and chemical criteria for compost maturity assessment. A review. *Bioresource Technology*, 100(22), 5444–5453. <https://doi.org/10.1016/j.biortech.2008.11.027>
- Blackstock, J. C. (1989). The physical chemistry of aqueous systems. *Guide to Biochemistry*, 11–19. <https://doi.org/10.1016/b978-0-7236-1151-6.50008-x>
- Bhattacharyya, P. N. (2012). Diversity of Microorganisms in the Surface and Subsurface Soil of the Jia Bharali River Catchment Area of Brahmaputra Plains. PhD Thesis. Guwahati: Gauhati University
- Bojesen, M., Boerboom, L., & Skov-Petersen, H. (2015). Towards a sustainable capacity expansion of the Danish biogas sector. *Land Use Policy*, 42, 264–277. <https://doi.org/10.1016/j.landusepol.2014.07.022>
- Bouman, O. T., Curtin, D., Campbell, C. A., Biederbeck, V. O., & Ukrainetz, H. (1995). Soil Acidification from Long-Term Use of Anhydrous Ammonia and Urea. *Soil Science Society of America Journal*, 59(5), 1488–1494. <https://doi.org/10.2136/sssaj1995.03615995005900050039x>
- Bouwman, A. F., Van Drecht, G., Knoop, J. M., Beusen, A. H. W., & Meinardi, C. R. (2005). Exploring changes in river nitrogen export to the world's oceans. *Global Biogeochemical Cycles*, 19(1). <https://doi.org/10.1029/2004gb002314>
- Braker, G., & Tiedje, J. M. (2003). Nitric Oxide Reductase (norB) Genes from Pure Cultures and Environmental Samples. *Applied and Environmental Microbiology*, 69(6), 3476–3483. <https://doi.org/10.1128/aem.69.6.3476-3483.2003>

- Bünemann, E. K., Schwenke, G. D., & Van Zwieten, L. (2006). Impact of agricultural inputs on soil organisms—a review. *Soil Research*, 44(4), 379. <https://doi.org/10.1071/sr05125>
- Burger, M. & Matiassek, M. (2009). New approaches to distinguish between nitrification and denitrification as sources of nitrous oxide. In: Sheldon, A.L. & Barnhart, E.P (eds) *Nitrous oxide emissions research progress*. Nova Science Publishers, New York (USA), pp 167–178
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., & Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), 20130122. <https://doi.org/10.1098/rstb.2013.0122>
- Ceja-Navarro, J. A., Rivera, F. N., Patiño-Zúñiga, L., Govaerts, B., Marsch, R., Vila-Sanjurjo, A., & Dendooven, L. (2009). Molecular characterization of soil bacterial communities in contrasting zero tillage systems. *Plant and Soil*, 329(1–2), 127–137. <https://doi.org/10.1007/s11104-009-0140-9>
- Chapin, F. S., Walker, L. R., Fastie, C. L., & Sharman, L. C. (1994). Mechanisms of Primary Succession Following Deglaciation at Glacier Bay, Alaska. *Ecological Monographs*, 64(2), 149–175. <https://doi.org/10.2307/2937039>
- Chien, S. H., Gearhart, M. M., & Collamer, D. J. (2008). The effect of different ammonical nitrogen sources on soil acidification. *Soil Science*, 173(8), 544–551. <https://doi.org/10.1097/ss.0b013e31817d9d17>
- Chuka-ogwude, D., Ogbonna, J., & Moheimani, N. R. (2020). A review on microalgal culture to treat anaerobic digestate food waste effluent. *Algal Research*, 47, 101841. <https://doi.org/10.1016/j.algal.2020.101841>
- Coelho, J. J., Hennessy, A., Casey, I., Bragança, C. R. S., Woodcock, T., & Kennedy, N. (2020). Biofertilisation with anaerobic digestates: A field study of effects on soil microbial abundance and diversity. *Applied Soil Ecology*, 147, 103403. <https://doi.org/10.1016/j.apsoil.2019.103403>
- Collins, S. L., Sinsabaugh, R. L., Crenshaw, C., Green, L., Porras-Alfaro, A., Stursova, M., & Zeglin, L. H. (2008). Pulse dynamics and microbial processes in aridland ecosystems. *Journal of Ecology*, 96(3), 413–420. <https://doi.org/10.1111/j.1365-2745.2008.01362.x>

- Conway, G. (1997). *The doubly green revolution: food for all in the 21st century*. London, United Kingdom: Penguin Books.
- Crawford, N. M., & Forde, B. G. (2002). Molecular and Developmental Biology of Inorganic Nitrogen Nutrition. *The Arabidopsis Book*, 1, e0011. <https://doi.org/10.1199/tab.0011>
- Curtin, D., Peterson, M. E., & Anderson, C. R. (2016). pH-dependence of organic matter solubility: Base type effects on dissolved organic C, N, P, and S in soils with contrasting mineralogy. *Geoderma*, 271, 161–172. <https://doi.org/10.1016/j.geoderma.2016.02.009>
- Dai, Z., Zhang, X., Tang, C., Muhammad, N., Wu, J., Brookes, P.C., Xu, J. (2017). Potential role of biochars in decreasing soil acidification-a critical review. *Sci Total Environ* 581–582:601–611
- Davidson, E. A., & Hackler, J. L. (1994). Soil heterogeneity can mask the effects of ammonium availability on nitrification. *Soil Biology and Biochemistry*, 26(10), 1449–1453. [https://doi.org/10.1016/0038-0717\(94\)90232-1](https://doi.org/10.1016/0038-0717(94)90232-1)
- Davison, J. (1988). Plant Beneficial Bacteria. *Nature Biotechnology*, 6(3), 282–286. <https://doi.org/10.1038/nbt0388-282>
- Desalegn, T., Alemu, G., Adella, A., Debele, T., & Gonzalo J., J. (2016). Effect of lime and phosphorus fertilizer on acid soils and barley (*hordeum vulgare* L.) Performance in the central highlands of Ethiopia. *Experimental Agriculture*, 53(3), 432–444. <https://doi.org/10.1017/s0014479716000491>
- Diaz, C., Saliba-Colombani, V., Loudet, O., Belluomo, P., Moreau, L., Daniel-Vedele, F., . . . Masclaux-Daubresse, C. (2006). Leaf Yellowing and Anthocyanin Accumulation are Two Genetically Independent Strategies in Response to Nitrogen Limitation in *Arabidopsis thaliana*. *Plant and Cell Physiology*, 47(1), 74–83. <https://doi.org/10.1093/pcp/pci225>
- Dominati, E., Patterson, M., & Mackay, A. (2010). A framework for classifying and quantifying the natural capital and ecosystem services of soils. *Ecological Economics*, 69(9), 1858–1868. <https://doi.org/10.1016/j.ecolecon.2010.05.002>
- Dong, W.-Y., Zhang, X.-Y., Dai, X.-Q., Fu, X.-L., Yang, F.-T., Liu, X.-Y., . . . Schaeffer, S. (2014). Changes in soil microbial community composition in response to fertilization of paddy soils in subtropical China. *Applied Soil Ecology*, 84, 140–147. <https://doi.org/10.1016/j.apsoil.2014.06.007>

- Doran, J. W., & Zeiss, M. R. (2000). Soil health and sustainability: managing the biotic component of soil quality. *Applied Soil Ecology*, 15(1), 3–11.  
[https://doi.org/10.1016/s0929-1393\(00\)00067-6](https://doi.org/10.1016/s0929-1393(00)00067-6)
- Drosg, B., Baxter, D., IEA Bioenergy Programme, & IEA Bioenergy Programme Staff. (2015). Nutrient Recovery by Biogas Digestate Processing. IEA Bioenergy.
- Dueri, S., Calanca, P. L., & Fuhrer, J. (2007). Climate change affects farm nitrogen loss – A Swiss case study with a dynamic farm model. *Agricultural Systems*, 93(1–3), 191–214.  
<https://doi.org/10.1016/j.agry.2006.05.005>
- Edmondson, W.T.(1995). Eutrophication. *Encyclopedia of Environmental Biology*, vol. 1. Academic Press, New York, pp.697–703.
- Effects of Liming Acidic Soils on Improving Soil Properties and Yield of Haricot Bean. (2014). *Journal of Environmental & Analytical Toxicology*, 05(01). <https://doi.org/10.4172/2161-0525.1000248>
- Ellis, E. C., & Ramankutty, N. (2008). Putting people in the map: anthropogenic biomes of the world. *Frontiers in Ecology and the Environment*, 6(8), 439–447.  
<https://doi.org/10.1890/070062>
- Erisman, J. W., Sutton, M. A., Galloway, J., Klimont, Z., & Winiwarter, W. (2008a). How a century of ammonia synthesis changed the world. *Nature Geoscience*, 1(10), 636–639.  
<https://doi.org/10.1038/ngeo325>
- Erisman, J. W., Sutton, M. A., Galloway, J., Klimont, Z., & Winiwarter, W. (2008b). How a century of ammonia synthesis changed the world. *Nature Geoscience*, 1(10), 636–639.  
<https://doi.org/10.1038/ngeo325>
- Feng, Y., Motta, A. C., Reeves, D. W., Burmester, C. H., van Santen, E., & Osborne, J. A. (2003). Soil microbial communities under conventional-till and no-till continuous cotton systems. *Soil Biology and Biochemistry*, 35(12), 1693–1703.  
<https://doi.org/10.1016/j.soilbio.2003.08.016>
- Fechter, M & Kraume M. (2016). Digestate treatment techniques. Faculty of Process Sciences, Technical University of Berlin.
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences*, 103(3), 626–631.  
<https://doi.org/10.1073/pnas.0507535103>

- Förstner, U.(1995). “Land contamination by metals—global scope and magnitude of problem,” in *Metal Speciation and Contamination of Soil*, H. E. Allen, C. P. Huang, G. W. Bailey, and A. R. Bowers, Eds., pp. 1–33, CRC Press, Inc., Boca Raton, FL, USA, 1995.
- Franche, C., Lindström, K., & Elmerich, C. (2008). Nitrogen-fixing bacteria associated with leguminous and non-leguminous plants. *Plant and Soil*, 321(1–2), 35–59.  
<https://doi.org/10.1007/s11104-008-9833-8>
- Frink, C. R., Waggoner, P. E., & Ausubel, J. H. (1999). Nitrogen fertilizer: Retrospect and prospect. *Proceedings of the National Academy of Sciences*, 96(4), 1175–1180.  
<https://doi.org/10.1073/pnas.96.4.1175>
- Fu, H., Zhang, G., Zhang, F., Sun, Z., Geng, G., & Li, T. (2017). Effects of Continuous Tomato Monoculture on Soil Microbial Properties and Enzyme Activities in a Solar Greenhouse. *Sustainability*, 9(2), 317. <https://doi.org/10.3390/su9020317>
- Garvin, D.F. & Carver, B.F.(2003). Role of the genotype in tolerance to acidity and aluminum toxicity. p. 387–406. In Z. Rengel (ed.) *Handbook of soil acidity*. Marcel Dekker, New York.
- Gemmeke, B., Rieger, C., Weiland, P .(2009). *Biogas-Messprogramm II, 61 Biogasanlagen im Vergleich*. FNR, Gülzow
- Girvan, M. S., Campbell, C. D., Killham, K., Prosser, J. I., & Glover, L. A. (2005). Bacterial diversity promotes community stability and functional resilience after perturbation. *Environmental Microbiology*, 7(3), 301–313. <https://doi.org/10.1111/j.1462-2920.2005.00695.x>
- Głowacka, A., Szostak, B., & Klebaniuk, R. (2020). Effect of Biogas Digestate and Mineral Fertilisation on the Soil Properties and Yield and Nutritional Value of Switchgrass Forage. *Agronomy*, 10(4), 490. <https://doi.org/10.3390/agronomy10040490>
- González, P. J., Correia, C., Moura, I., Brondino, C. D., & Moura, J. J. G. (2006). Bacterial nitrate reductases: Molecular and biological aspects of nitrate reduction. *Journal of Inorganic Biochemistry*, 100(5–6), 1015–1023.  
<https://doi.org/10.1016/j.jinorgbio.2005.11.024>
- González-Chávez, M. C. A., Aitkenhead-Peterson, J. A., Gentry, T. J., Zuberer, D., Hons, F., & Loeppert, R. (2010). Soil microbial community, C, N, and P responses to long-term

- tillage and crop rotation. *Soil and Tillage Research*, 106(2), 285–293.  
<https://doi.org/10.1016/j.still.2009.11.008>
- Griffiths, B. S., & Philippot, L. (2013). Insights into the resistance and resilience of the soil microbial community. *FEMS Microbiology Reviews*, 37(2), 112–129.  
<https://doi.org/10.1111/j.1574-6976.2012.00343.x>
- Guo, J. H., Liu, X. J., Zhang, Y., Shen, J. L., Han, W. X., Zhang, W. F., . . . Zhang, F. S. (2010). Significant Acidification in Major Chinese Croplands. *Science*, 327(5968), 1008–1010.  
<https://doi.org/10.1126/science.1182570>
- Hallin, S., Jones, C. M., Schloter, M., & Philippot, L. (2009). Relationship between N-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *The ISME Journal*, 3(5), 597–605. <https://doi.org/10.1038/ismej.2008.128>
- Hartmann, M., Frey, B., Mayer, J., Mäder, P., & Widmer, F. (2014). Distinct soil microbial diversity under long-term organic and conventional farming. *The ISME Journal*, 9(5), 1177–1194. <https://doi.org/10.1038/ismej.2014.210>
- Hedrick, D. B., Peacock, A., Stephen, J. R., Macnaughton, S. J., Brüggemann, J., & White, D. C. (2000). Measuring soil microbial community diversity using polar lipid fatty acid and denaturing gradient gel electrophoresis data. *Journal of Microbiological Methods*, 41(3), 235–248. [https://doi.org/10.1016/s0167-7012\(00\)00157-3](https://doi.org/10.1016/s0167-7012(00)00157-3)
- Heylen, K., Gevers, D., Vanparys, B., Wittebolle, L., Geets, J., Boon, N., & De Vos, P. (2006). The incidence of nirS and nirK and their genetic heterogeneity in cultivated denitrifiers. *Environmental Microbiology*, 8(11), 2012–2021. <https://doi.org/10.1111/j.1462-2920.2006.01081.x>
- Hillebrand, H., Bennett, D. M., & Cadotte, M. W. (2008). Consequences of Dominance: A Review Of Evenness Effects on Local And Regional Ecosystem Processes. *Ecology*, 89(6), 1510–1520. <https://doi.org/10.1890/07-1053.1>
- Hirsch, P. R., & Mauchline, T. H. (2015). The Importance of the Microbial N Cycle in Soil for Crop Plant Nutrition. *Advances in Applied Microbiology*, 45–71.  
<https://doi.org/10.1016/bs.aambs.2015.09.001>
- Holm-Nielsen, J. B., Al Seadi, T., & Oleskowicz-Popiel, P. (2009). The future of anaerobic digestion and biogas utilization. *Bioresource Technology*, 100(22), 5478–5484.  
<https://doi.org/10.1016/j.biortech.2008.12.046>

- Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., . . . Wardle, D. A. (2005). Effects of Biodiversity on Ecosystem Functioning: A Consensus Of Current Knowledge. *Ecological Monographs*, 75(1), 3–35. <https://doi.org/10.1890/04-0922>
- Jiang, L. (2007). Negative Selection Effects Suppress Relationships Between Bacterial Diversity and Ecosystem Functioning. *Ecology*, 88(5), 1075–1085. <https://doi.org/10.1890/06-1556>
- Kandeler, E., Tschirko, D., & Spiegel, H. (1999). Long-term monitoring of microbial biomass, N mineralisation and enzyme activities of a Chernozem under different tillage management. *Biology and Fertility of Soils*, 28(4), 343–351. <https://doi.org/10.1007/s003740050502>
- Karakashev, D., Batstone, D. J., & Angelidaki, I. (2005). Influence of Environmental Conditions on Methanogenic Compositions in Anaerobic Biogas Reactors. *Applied and Environmental Microbiology*, 71(1), 331–338. <https://doi.org/10.1128/aem.71.1.331-338.2005>
- Kariuki, S. K., Zhang, H., Schroder, J. L., Edwards, J., Payton, M., Carver, B. F., . . . Krenzer, E. G. (2007). Hard Red Winter Wheat Cultivar Responses to a pH and Aluminum Concentration Gradient. *Agronomy Journal*, 99(1), 88–98. <https://doi.org/10.2134/agronj2006.0128>
- Kennedy, I. R., Pereg-Gerk, L. L., Wood, C., Deaker, R., Gilchrist, K., & Katupitiya, S. (1997). Biological nitrogen fixation in non-leguminous field crops: Facilitating the evolution of an effective association between *Azospirillum* and wheat. *Plant and Soil*, 194(1/2), 65–79. <https://doi.org/10.1023/a:1004260222528>
- Kibblewhite, M. G., Ritz, K., & Swift, M. J. (2007). Soil health in agricultural systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1492), 685–701. <https://doi.org/10.1098/rstb.2007.2178>
- Langenheder, S., Bulling, M. T., Solan, M., & Prosser, J. I. (2010). Bacterial Biodiversity-Ecosystem Functioning Relations Are Modified by Environmental Complexity. *PLoS ONE*, 5(5), e10834. <https://doi.org/10.1371/journal.pone.0010834>
- Lazcano, C., Gómez-Brandón, M., Revilla, P., & Domínguez, J. (2012). Short-term effects of organic and inorganic fertilizers on soil microbial community structure and function. *Biology and Fertility of Soils*, 49(6), 723–733. <https://doi.org/10.1007/s00374-012-0761-7>

- Lichtfouse, E., Navarrete, M., Debaeke, P., Souchère, V., Alberola, C., & Ménassieu, J. (2009). Agronomy for sustainable agriculture. A review. *Agronomy for Sustainable Development*, 29(1), 1–6. <https://doi.org/10.1051/agro:2008054>
- Loreau, M. (2010). Linking biodiversity and ecosystems: towards a unifying ecological theory. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1537), 49–60. <https://doi.org/10.1098/rstb.2009.0155>
- M. Tahat, M., M. Alananbeh, K., A. Othman, Y., & I. Leskovar, D. (2020). Soil Health and Sustainable Agriculture. *Sustainability*, 12(12), 4859. <https://doi.org/10.3390/su12124859>
- Marschner, P. (2003). Structure and function of the soil microbial community in a long-term fertilizer experiment. *Soil Biology and Biochemistry*, 35(3), 453–461. [https://doi.org/10.1016/s0038-0717\(02\)00297-3](https://doi.org/10.1016/s0038-0717(02)00297-3)
- Mathew, R. P., Feng, Y., Githinji, L., Ankumah, R., & Balkcom, K. S. (2012). Impact of No-Tillage and Conventional Tillage Systems on Soil Microbial Communities. *Applied and Environmental Soil Science*, 2012, 1–10. <https://doi.org/10.1155/2012/548620>
- Minasny, B., Hong, S. Y., Hartemink, A. E., Kim, Y. H., & Kang, S. S. (2016a). Soil pH increase under paddy in South Korea between 2000 and 2012. *Agriculture, Ecosystems & Environment*, 221, 205–213. <https://doi.org/10.1016/j.agee.2016.01.042>
- Minasny, B., Hong, S. Y., Hartemink, A. E., Kim, Y. H., & Kang, S. S. (2016b). Soil pH increase under paddy in South Korea between 2000 and 2012. *Agriculture, Ecosystems & Environment*, 221, 205–213. <https://doi.org/10.1016/j.agee.2016.01.042>
- Möller, K. (2015). Effects of anaerobic digestion on soil carbon and nitrogen turnover, N emissions, and soil biological activity. A review. *Agronomy for Sustainable Development*, 35(3), 1021–1041. <https://doi.org/10.1007/s13593-015-0284-3>
- Moreira, F. M. S & Siqueira, J. O. (2006) soil microbiology and biochemistry. 2. ed. Lavras: UFLa.
- Muhammad, D. (2020). Land Degradation and its Management: A Review. *International Journal of Environmental Sciences & Natural Resources*, 25(1). <https://doi.org/10.19080/ijesnr.2020.25.556157>
- Müller, D. B., Vogel, C., Bai, Y., & Vorholt, J. A. (2016). The Plant Microbiota: Systems-Level Insights and Perspectives. *Annual Review of Genetics*, 50(1), 211–234. <https://doi.org/10.1146/annurev-genet-120215-034952>



- Neina, D. (2019). The Role of Soil pH in Plant Nutrition and Soil Remediation. *Applied and Environmental Soil Science*, 2019, 1–9. <https://doi.org/10.1155/2019/5794869>
- Nortcliff, S. (2004). *Handbook of Soil Acidity*. Edited by Z. Rengel. New York: Marcel Dekker (2003), pp. 496, US\$195.00. ISBN 0–8247-0890-3. *Experimental Agriculture*, 40(3), 393–394. <https://doi.org/10.1017/s0014479704332059>
- Norton, J. M. (2014). Diversity and Environmental Distribution of Ammonia-Oxidizing Bacteria. *Nitrification*, 39–55. <https://doi.org/10.1128/9781555817145.ch3>
- Orellana, L. H., Rodriguez-R, L. M., Higgins, S., Chee-Sanford, J. C., Sanford, R. A., Ritalahti, K. M., . . . Konstantinidis, K. T. (2014). Detecting Nitrous Oxide Reductase (nosZ) Genes in Soil Metagenomes: Method Development and Implications for the Nitrogen Cycle. *MBio*, 5(3). <https://doi.org/10.1128/mbio.01193-14>
- Parayil, G. (1992). The Green Revolution in India: A Case Study of Technological Change. *Technology and Culture*, 33(4), 737. <https://doi.org/10.2307/3106588>
- Pivato, A., Vanin, S., Raga, R., Lavagnolo, M. C., Barausse, A., Rieple, A., . . . Cossu, R. (2016). Use of digestate from a decentralized on-farm biogas plant as fertilizer in soils: An ecotoxicological study for future indicators in risk and life cycle assessment. *Waste Management*, 49, 378–389. <https://doi.org/10.1016/j.wasman.2015.12.009>
- Priemé, A., Braker, G., & Tiedje, J. M. (2002). Diversity of Nitrite Reductase (nirK and nirS) Gene Fragments in Forested Upland and Wetland Soils. *Applied and Environmental Microbiology*, 68(4), 1893–1900. <https://doi.org/10.1128/aem.68.4.1893-1900.2002>
- Prosser, J. I., & Nicol, G. W. (2008). Relative contributions of archaea and bacteria to aerobic ammonia oxidation in the environment. *Environmental Microbiology*, 10(11), 2931–2941. <https://doi.org/10.1111/j.1462-2920.2008.01775.x>
- Prosser, J. I., & Nicol, G. W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. *Trends in Microbiology*, 20(11), 523–531. <https://doi.org/10.1016/j.tim.2012.08.001>
- Ramankutty, N., Evan, A. T., Monfreda, C., & Foley, J. A. (2008). Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles*, 22(1), n/a. <https://doi.org/10.1029/2007gb002952>
- Ramankutty, N., Mehrabi, Z., Waha, K., Jarvis, L., Kremen, C., Herrero, M., & Rieseberg, L. H. (2018). *Trends in Global Agricultural Land Use: Implications for Environmental Health*

- and Food Security. *Annual Review of Plant Biology*, 69(1), 789–815.  
<https://doi.org/10.1146/annurev-arplant-042817-040256>
- Rasnake, M., Murdock, L. Schwab, G. & Thom, B. (2002). Determining the Quality of Aglime: Using Relative Neutralizing Values. AGR-106, Cooperative Extension Service, University of Kentucky, Lexington.
- Roser, M. (2014). Future population growth. Retrieved July 7, 2020, from <https://OurWorldInData.org>
- Rousk, J., Brookes, P. C., & Bååth, E. (2009). Contrasting Soil pH Effects on Fungal and Bacterial Growth Suggest Functional Redundancy in Carbon Mineralization. *Applied and Environmental Microbiology*, 75(6), 1589–1596. <https://doi.org/10.1128/aem.02775-08>
- Rütting, T., Aronsson, H., & Delin, S. (2018). Efficient use of nitrogen in agriculture. *Nutrient Cycling in Agroecosystems*, 110(1), 1–5. <https://doi.org/10.1007/s10705-017-9900-8>
- Sakurai, N., & Sakurai, T. (1998). Genomic DNA Cloning of the Region Encoding Nitric Oxide Reductase in *Paracoccus halo denitrificans* and a Structure Model Relevant to Cytochrome Oxidase. *Biochemical and Biophysical Research Communications*, 243(2), 400–406. <https://doi.org/10.1006/bbrc.1998.8106>
- Salles, J. F., Le Roux, X., & Poly, F. (2012). Relating Phylogenetic and Functional Diversity among Denitrifiers and Quantifying their Capacity to Predict Community Functioning. *Frontiers in Microbiology*, 3, <https://doi.org/10.3389/fmicb.2012.00209>
- Sapp, M., Harrison, M., Hany, U., Charlton, A., & Thwaites, R. (2015). Comparing the effect of digestate and chemical fertiliser on soil bacteria. *Applied Soil Ecology*, 86, 1–9. <https://doi.org/10.1016/j.apsoil.2014.10.004>
- Sathya, A., Vijayabharathi, R., & Gopalakrishnan, S. (2016). Soil Microbes: The Invisible Managers of Soil Fertility. *Microbial Inoculants in Sustainable Agricultural Productivity*, 1–16. [https://doi.org/10.1007/978-81-322-2644-4\\_1](https://doi.org/10.1007/978-81-322-2644-4_1)
- Schroder, J. L., Zhang, H., Girma, K., Raun, W. R., Penn, C. J., & Payton, M. E. (2011). Soil Acidification from Long-Term Use of Nitrogen Fertilizers on Winter Wheat. *Soil Science Society of America Journal*, 75(3), 957–964. <https://doi.org/10.2136/sssaj2010.0187>
- Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., . . . Handelsman, J. (2012). Fundamentals of Microbial Community Resistance and Resilience. *Frontiers in Microbiology*, 3. <https://doi.org/10.3389/fmicb.2012.00417>

- Sharma, L., & Bali, S. (2017). A Review of Methods to Improve Nitrogen Use Efficiency in Agriculture. *Sustainability*, 10(2), 51. <https://doi.org/10.3390/su10010051>
- Signor, D., & Cerri, C. E. P. (2013). Nitrous oxide emissions in agricultural soils: a review. *Pesquisa Agropecuária Tropical*, 43(3), 322–338. <https://doi.org/10.1590/s1983-40632013000300014>
- Šimon, T., Kunzová, E., & Friedlová, M. (2016). The effect of digestate, cattle slurry and mineral fertilization on the winter wheat yield and soil quality parameters. *Plant, Soil and Environment*, 61(No. 11), 522–527. <https://doi.org/10.17221/530/2015-pse>
- Singh, B. (2018). Are Nitrogen Fertilizers Deleterious to Soil Health? *Agronomy*, 8(4), 48. <https://doi.org/10.3390/agronomy8040048>
- Singh, B. K., Quince, C., Macdonald, C. A., Khachane, A., Thomas, N., Al-Soud, W. A., . . . Campbell, C. D. (2014). Loss of microbial diversity in soils is coincident with reductions in some specialized functions. *Environmental Microbiology*, 16(8), 2408–2420. <https://doi.org/10.1111/1462-2920.12353>
- Smith, J., Abegaz, A., Matthews, R. B., Subedi, M., Orskov, E. R., Tumwesige, V., & Smith, P. (2014). What is the potential for biogas digesters to improve soil carbon sequestration in Sub-Saharan Africa? Comparison with other uses of organic residues. *Biomass and Bioenergy*, 70, 73–86. <https://doi.org/10.1016/j.biombioe.2014.01.056>
- Spanning, R.J. (2011). Structure, function, regulation and evolution of the nitrite and nitrous oxide reductases: denitrification enzymes with aβ-propeller fold, in *Nitrogen Cycling in Bacteria*, J. Moir, ed., Caister Academic Press, Norfolk, 2011, 135–161
- Stark, C.H. & Richards, K.G. (2008) The Continuing Challenge of Agricultural Nitrogen Loss to the Environment in the Context of Global Change and Advancing Research. *Dynamic Soil, Dynamic Plant*, 2, 1-12.
- Stitt, M., & Krapp, A. (1999). The interaction between elevated carbon dioxide and nitrogen nutrition: the physiological and molecular background. *Plant, Cell and Environment*, 22(6), 583–621. <https://doi.org/10.1046/j.1365-3040.1999.00386.x>
- Strickland, M. S., Lauber, C., Fierer, N., & Bradford, M. A. (2009). Testing the functional significance of microbial community composition. *Ecology*, 90(2), 441–451. <https://doi.org/10.1890/08-0296.1>

- Sun, C., Cao, W., Banks, C. J., Heaven, S., & Liu, R. (2016). Biogas production from undiluted chicken manure and maize silage: A study of ammonia inhibition in high solids anaerobic digestion. *Bioresource Technology*, 218, 1215–1223.  
<https://doi.org/10.1016/j.biortech.2016.07.082>
- Sun, P., Zhuge, Y., Zhang, J., & Cai, Z. (2012). Soil pH was the main controlling factor of the denitrification rates and N<sub>2</sub>/N<sub>2</sub>O emission ratios in forest and grassland soils along the Northeast China Transect (NECT). *Soil Science and Plant Nutrition*, 58(4), 517–525.  
<https://doi.org/10.1080/00380768.2012.703609>
- Sun, R., Zhang, X. X., Guo, X., Wang, D., & Chu, H. (2015). Bacterial diversity in soils subjected to long-term chemical fertilization can be more stably maintained with the addition of livestock manure than wheat straw. *Soil Biology and Biochemistry*, 88, 9–18.  
<https://doi.org/10.1016/j.soilbio.2015.05.007>
- Sumner, M. & Noble, A. (2003). Handbook of soil acidity. In: Rengel, Z. (ed) Marcel Dekker, New York, pp 1–28
- Taffesse, A. (1999). Effects of liming on maize yield and soil reaction. Proceedings of the 4th Conference of the Ethiopian Society of Soil Science. Addis Ababa, Ethiopia 272
- Tampio, E., Ervasti, S., & Rintala, J. (2015). Characteristics and agronomic usability of digestates from laboratory digesters treating food waste and autoclaved food waste. *Journal of Cleaner Production*, 94, 86–92. <https://doi.org/10.1016/j.jclepro.2015.01.086>
- Tang, Y., Garvin, D. F., Kochian, L. V., Sorrells, M. E., & Carver, B. F. (2002). Physiological Genetics of Aluminum Tolerance in the Wheat Cultivar Atlas 66. *Crop Science*, 42(5), 1541–1546. <https://doi.org/10.2135/cropsci2002.1541>
- Tiemann, L. K., Grandy, A. S., Atkinson, E. E., Marin-Spiotta, E., & McDaniel, M. D. (2015). Crop rotational diversity enhances belowground communities and functions in an agroecosystem. *Ecology Letters*, 18(8), 761–771. <https://doi.org/10.1111/ele.12453>
- Tilston, E. L., Sizmur, T., Dixon, G. R., Otten, W., & Harris, J. A. (2010). The Impact of Land-Use Practices on Soil Microbes. *Soil Microbiology and Sustainable Crop Production*, 273–295. [https://doi.org/10.1007/978-90-481-9479-7\\_7](https://doi.org/10.1007/978-90-481-9479-7_7)
- Trivedi, P., Delgado-Baquerizo, M., Anderson, I. C., & Singh, B. K. (2016). Response of Soil Properties and Microbial Communities to Agriculture: Implications for Primary

- Productivity and Soil Health Indicators. *Frontiers in Plant Science*, 7.  
<https://doi.org/10.3389/fpls.2016.00990>
- Van der Meer, J. J. M. (2000). Book review. *The Nature and Properties of Soils*. By N. C. Brady and R. R. Weil. 12th edn. Prentice Hall, Upper Saddle River, NJ, 1999; 881 pp.; ISBN 0 13 852444 0. *Sedimentology*, 47(6), 1230. <https://doi.org/10.1046/j.1365-3091.2000.0376c.x>
- Vitousek, P. M., Naylor, R., Crews, T., David, M. B., Drinkwater, L. E., Holland, E., . . . Zhang, F. S. (2009). Nutrient Imbalances in Agricultural Development. *Science*, 324(5934), 1519–1520. <https://doi.org/10.1126/science.1170261>
- Vries, S. & Pouvreau L.A.M (2007). Nitric oxide reductase: structural variations and catalytic mechanism, in *Biology of the Nitrogen Cycle*, Bothe, H., Ferguson, S.J., & Newton, W.E eds., Elsevier, Amsterdam, 2007, 57–66.
- Walsh, J. J., Jones, D. L., Edwards-Jones, G., & Williams, A. P. (2012). Replacing inorganic fertilizer with anaerobic digestate may maintain agricultural productivity at less environmental cost. *Journal of Plant Nutrition and Soil Science*, 175(6), 840–845. <https://doi.org/10.1002/jpln.201200214>
- Wei, M., Hu, G., Wang, H., Bai, E., Lou, Y., Zhang, A., & Zhuge, Y. (2017). 35 years of manure and chemical fertilizer application alters soil microbial community composition in a Fluvo-aquic soil in Northern China. *European Journal of Soil Biology*, 82, 27–34. <https://doi.org/10.1016/j.ejsobi.2017.08.002>
- Wertz, S., Degrange, V., Prosser, J. I., Poly, F., Commeaux, C., Guillaumaud, N., & Le Roux, X. (2007). Decline of soil microbial diversity does not influence the resistance and resilience of key soil microbial functional groups following a model disturbance. *Environmental Microbiology*, 9(9), 2211–2219. <https://doi.org/10.1111/j.1462-2920.2007.01335.x>
- White, R.E. (2006). *Principles and Practice of Soil Science: The Soil as a Natural Resource*, Blackwell Publishing, Oxford, UK.
- Wittebolle, L., Marzorati, M., Clement, L., Balloi, A., Daffonchio, D., Heylen, K., . . . Boon, N. (2009). Initial community evenness favours functionality under selective stress. *Nature*, 458(7238), 623–626. <https://doi.org/10.1038/nature07840>
- Wu, L., Jiang, Y., Zhao, F., He, X., Liu, H., & Yu, K. (2020). Increased organic fertilizer application and reduced chemical fertilizer application affect the soil properties and

- bacterial communities of grape rhizosphere soil. *Scientific Reports*, 10(1).  
<https://doi.org/10.1038/s41598-020-66648-9>
- Yabe, N. (2013). Environmental and economic evaluations of centralized biogas plants running on cow manure in Hokkaido, Japan. *Biomass and Bioenergy*, 49, 143–151.  
<https://doi.org/10.1016/j.biombioe.2012.12.001>
- Yan, Y., Zhang, L., Feng, L., Sun, D., & Dang, Y. (2018). Comparison of varying operating parameters on heavy metals ecological risk during anaerobic co-digestion of chicken manure and corn stover. *Bioresource Technology*, 247, 660–668.  
<https://doi.org/10.1016/j.biortech.2017.09.146>
- Yao, H., Bowman, D., & Shi, W. (2011). Seasonal variations of soil microbial biomass and activity in warm- and cool-season turfgrass systems. *Soil Biology and Biochemistry*, 43(7), 1536–1543. <https://doi.org/10.1016/j.soilbio.2011.03.031>
- Yu, P., Liu, S., Han, K., Guan, S., & Zhou, D. (2017). Conversion of cropland to forage land and grassland increases soil labile carbon and enzyme activities in northeastern China. *Agriculture, Ecosystems & Environment*, 245, 83–91.  
<https://doi.org/10.1016/j.agee.2017.05.013>
- Yuan, J., Yuan, Y., Zhu, Y., & Cao, L. (2018). Effects of different fertilizers on methane emissions and methanogenic community structures in paddy rhizosphere soil. *Science of The Total Environment*, 627, 770–781. <https://doi.org/10.1016/j.scitotenv.2018.01.233>
- Zhang, W., Ricketts, T. H., Kremen, C., Carney, K., & Swinton, S. M. (2007). Ecosystem services and dis-services to agriculture. *Ecological Economics*, 64(2), 253–260.  
<https://doi.org/10.1016/j.ecolecon.2007.02.024>
- Zhou, D. D. (2008). Microelectrodes for in-vivo determination of pH. *Electrochemical Sensors, Biosensors and Their Biomedical Applications*, 261–305. <https://doi.org/10.1016/b978-012373738-0.50012-x>
- Zhou, L., Wang, S., Zou, Y., Xia, C., & Zhu, G. (2015). Species, Abundance and Function of Ammonia-oxidizing Archaea in Inland Waters across China. *Scientific Reports*, 5(1).  
<https://doi.org/10.1038/srep15969>
- Zhou, X., & Wu, F. (2012). Dynamics of the diversity of fungal and *Fusarium* communities during continuous cropping of cucumber in the greenhouse. *FEMS Microbiology Ecology*, 80(2), 469–478. <https://doi.org/10.1111/j.1574-6941.2012.01312.x>

- Zhou, X., Yu, G., & Wu, F. (2012). Soil phenolics in a continuously mono-cropped cucumber (*Cucumis sativus* L.) system and their effects on cucumber seedling growth and soil microbial communities. *European Journal of Soil Science*, 63(3), 332–340.  
<https://doi.org/10.1111/j.1365-2389.2012.01442.x>
- Zingore, S., Manyame, C., Nyamugafata, P., & Giller, K. E. (2005). Long-term changes in organic matter of woodland soils cleared for arable cropping in Zimbabwe. *European Journal of Soil Science*, 0(0), 050912034650038-??? <https://doi.org/10.1111/j.1365-2389.2005.00707.x>
- Zumft, W.G. & Heinz, K. (2007). Chapter 5 - Nitrous oxide reductase. *Biology of the Nitrogen cycle*, 67-81.
- Zumft, W. G. (1997). Cell biology and molecular basis of denitrification. *Microbiology and Molecular Biology Reviews : MMBR*, 61(4), 533–616. <https://doi.org/10.1128/61.4.533-616.1997>
- Zumft, W. G., Braun, C., & Cuypers, H. (1994). Nitric oxide reductase from *Pseudomonas stutzeri*. Primary structure and gene organization of a novel bacterial cytochrome bc complex. *European Journal of Biochemistry*, 219(1–2), 481–490.  
<https://doi.org/10.1111/j.1432-1033.1994.tb19962.x>

## Chapter 3 Land use and season drive changes in soil microbial communities and related functions in agricultural soils

### 3.1 Abstract

Soil microorganisms are important for maintenance of soil health and related functions. Agricultural management practices such as land use, season and fertilizer affect soil microbial community structures. However, the effect of these management practices on soil microorganisms and related functions, influenced by regionally different soil types, is still not clear. Hence, the study was conducted in an Andosol (volcanic soil) dominated agricultural region in cool temperate climate to determine the effect of land use (cropland, grassland), season (spring, summer) and fertilizer (anaerobic digestate) on soil microorganisms and related functions. Soils were sampled from farmers' fields, DNA extracted and sequenced targeting 16S rRNA region. In result, land use had a significant effect on beta diversity and evenness with higher values recorded in cropland than grassland. However, grassland had a higher number of unique operational taxonomic units (OTUs) (10303) compared to cropland (5112). In cropland, season had a significant effect on beta diversity, evenness, OTU numbers and Shannon index with higher values recorded in summer compared to spring. Based on predicted soil functions, nitrogenase (*nifH*) had significantly higher values in cropland-summer while nitrite reductase (*nirK*) and ammonia monooxygenase (*amoA*) were significantly higher in cropland-spring. In grassland, season had a significant effect on beta diversity only. These results indicate that grassland microorganisms were stable and more resistant to seasonal changes than cropland, suggesting that conventional tillage practices have a negative effect on soil microbial stability. Additionally, grassland-spring (7059) had a higher number of unique OTUs than grassland-summer (2597). Based on predicted soil functions, *nifH* was significantly higher in grassland-spring while *nirK* and *amoA* were significantly higher in grassland-summer. These results indicate that the impact of seasons on soil microorganisms' distribution and abundance in cropland and grassland may directly affect soil functions.



### 3.2 Introduction

Soil microorganisms are significant to the maintenance of soil health by providing ecosystem services (Chaparro et al., 2012; Singh et al., 2014), such as decomposition of organic matter and nutrient cycling (Marques et al., 2014; Yuliar et al., 2015). Microbial diversity is an essential attribute of the soil microbial community (Bender et al., 2016). High microbial diversity enhances soil ecosystem services and tolerance to disturbances such as land use, seasonal changes and fertilizer application (Griffiths and Philippot, 2013; Loreau, 2010). On the contrary, a decrease in diversity would lead to potential loss of ecosystem services reducing the systems stability (Trivedi et al., 2016). Thus, soil microorganisms should be diverse to ensure sustainable provision of ecosystem services.

Agricultural soil disturbances are defined as factors that cause changes in the composition and structure of the soil (Thompson et al., 2002). Soil disturbances can be caused by farmer management, such as physical and chemical soil disturbances. Physical soil disturbances are usually caused by tillage practices, which are determined by the land use management system (Chen et al., 2011; Liu et al., 2019). Chemical soil disturbances are usually related to agricultural fertilizer application (Ogundijo et al., 2014). Additionally, soil disturbances can also occur naturally by shifts in rainfall and temperature, caused by seasonal changes, which induce soil disturbances (Luo et al., 2019). Although there are many other factors that can cause soil disturbance within the agricultural environment, site visits identified land use, fertilizer application and seasonality as some of the main factors that may have caused soil disturbance in the study site. Nonetheless, the influences of these factors, i.e., land use, seasonal changes and fertilizer application on soil microbial diversity and related soil functions in the study region are still not fully explored.

Although there are considerable research efforts made towards examining the effect of disturbances on soil microbial community composition on a global scale (Crowther et al., 2014; Johnson and Curtis, 2001), there is a lack of research on the same at a regional scale, focusing on the disturbance factors as stated above (i.e. land use, seasonality and fertilizer inputs). In addition, specific analyses of field-based observation of soil microbial diversity at a regional scale, on the same soil type, are essential because basic soil characteristics determine soil microorganisms' structures and related functions (Pan et al., 2014; Wang et al., 2019). Therefore, research needs to

be conducted to clarify the interaction of land use, seasonality and fertilizer application on soil microorganisms and related functions at a regional scale in agricultural soils.

Land use is one of the factors that drives soil microorganisms and functions (Degens et al., 2000; Pabst et al., 2013). Various studies have documented that land use types affect soil properties (soil pH, organic carbon, inorganic nitrogen, soil water content), which are intimately connected to microbial community structure, diversity and functions (Pabst et al., 2013; Sengupta et al., 2020; Bissett et al., 2014). Common land use types in agriculture involve grassland for animal feed and cropland for yields (Yu et al., 2017). Previous research shows that conventional cropland management practices such as intensive tillage, monoculture and inorganic fertilizer use, have been associated with a decrease in soil microbial diversity and related soil functions (Wardle, 1992; Zingore et al., 2005). However, due to differences in experimental locations, setups and soil properties among various studies, there have been contrasting effects of cropland and grassland management on specific microorganisms. For instance, Szoboszlay et al. (2017) reported that abundance of *Verrucomicrobiota* increased in cropland and decreased in grassland in European soils. On the other hand, Sengupta et al. (2020) reported that the abundance of *Acidobacteria* and *Gemmatimonadetes* increased in cropland and decreased in grassland soils.

In addition to the effects of land use, seasons drive changes in soil microbial diversity and functions (Bowles et al., 2014; Castro et al., 2009). Seasonal changes are usually accompanied with variations in rainfall, temperature and soil nutrient content (Smith et al., 2010). Seasons with high soil moisture and temperature like summer stimulate soil microbial activity and nutrient cycling compared to seasons with low soil moisture and temperature like spring (Galazka et al., 2017; Zifčáková., 2015). Considering cultivated agricultural soils, Li et al. (2019) observed that the relative abundance of *Acidobacteria* significantly increased in summer while *Actinobacteria* were significantly reduced in both spring and summer. However, research by Li et al. (2020) found that *Chloroflexi* and *Actinobacteria* were significantly higher in summer while *Acidobacteria* were significantly higher in spring. Therefore, research needs to be conducted to clarify the effects of season on specific soil microorganisms in agricultural soils.

Another factor controlling soil microbial diversity is fertilizer types, including animal waste based organic materials that have been found to increase microbial species richness in *Alphaproteobacteria*, *Betaproteobacteria*, and *Firmicutes* (Faisal et al., 2018). These microbes are

involved in decomposition of complex organic compounds (Buresova et al., 2019). As the price of chemical fertilizer soars (Ricker-Gilbert et al., 2014), not only livestock farmers but also crop farmers may utilize animal waste-based fertilizers in future, at the community level. To understand the effect of anaerobic digestate application on soil microbes, community level studies must be carried out. In some cases, animal wastes are collected from multiple farms and centralized to improve the efficiency of the effluent management. Community level field studies are critically important because interactions among soil types, the ratio between crop farmers and livestock farmers, sources of animal wastes and many other factors are unique to each community.

To evaluate the soils' microbial structure and potential changes in diversity, next generation sequencing (NGS) platforms will be used. Traditionally, researchers relied on culture dependent methods, which involved growing microbes in the laboratory. However, the main limitation of these methods is underestimation of diversity as some microorganisms do not grow well under laboratory culture conditions or as monocultures (Jo et al., 2016). To address these challenges, culture independent methods such as NGS platforms were developed. Through NGS platforms, soil microbial research has become cheaper, faster and convenient (Doolittle and Zhaxybayeva, 2010; Ghazanfar et al., 2009). To determine bacterial taxonomic classification and phylogeny, 16S rRNA gene is the most common genetic marker used (Janda and Abbott, 2007). This is because its present in almost all bacteria, its functions have not changed over time and the gene is large enough for informatics (Janda and Abbott, 2007).

Thus, this study aimed to address the following key questions; 1) In a regional scale, which of the imposed treatments (land use, season or fertilizer) is the biggest factor controlling soil microbes? and 2). Is it possible to identify the key microbes and related functions impacted by each imposed treatment? We therefore conducted research on soils from cropland and grassland applied with biogas slurry, mainly made of dairy wastes, during summer and spring. We hypothesized that land use would be the biggest factor controlling soil microbiome in Kamishihoro. More specifically, grassland-summer with anaerobic digestate application would enhance microbial diversity and related functions compared to other treatments.

### 3.3 Materials and methods

#### 3.3.1 Study site and treatments

Soils were sampled in 2018 from Kamishihoro (43°14'N, 143°18'E), located in Tokachi town, Hokkaido prefecture, Japan. Soils in the region are classified as Andosol (FAO/UNESCO). Treatments included two land use regimes; cropland and grassland (Figure 3.1), two seasons; spring (April to June) and summer (August to October) and fertilizer inputs; anaerobic digestate (AD), ammonium sulphate (AS) and control

Regarding land use, common management practices for cropland included crop rotation for 4 years involving potatoes, sorghum, wheat and beets. Beets (the focus of this study) were planted in April and August and harvested in July and November, respectively. The soil was tilled 4 times a year before seeding at a depth of 30 cm, using tractors. Fertilizers (Nitrogen-N, Phosphorous-P and Potassium-K) were applied at the rate of 40 kg N ha<sup>-1</sup>, 340 kg P ha<sup>-1</sup> and 158 kg K ha<sup>-1</sup> at planting. For grasslands farmers were practicing no till system and anaerobic digestate fertilizer was applied at the rate of 60 kg N ha<sup>-1</sup>. Grasslands and croplands were fertilized at the same time.

During the experimental period spring (April to June) had average temperature of 10.6°C and cumulative rainfall of 252 mm, while summer (July to October) had average temperature of 14°C and cumulative rainfall of 474 mm (Japan Meteorological Agency).




Figure 3.1: Kamishihoro soil sampling sites

### 3.3.2 Soil sampling

To avoid any contamination, all equipment and sampling procedures followed recommendations provided by Taberlet et al. (2012). For soil physical (bulk density-BD) analysis, soils were sampled from each fertilizer treatment using cores (100 cm<sup>3</sup>). For soil microbial and chemical analysis (pH, N, K, magnesium-Mg, cation exchange capacity-CEC and humus), 500 g of soil samples were sampled from each plot and fertilizer treatment using a shovel. For microbial analysis, samples were stored at -30°C before DNA extraction.

We took a total of 72 soils samples at a depth of 0-15 cm on 30<sup>th</sup> April 2018 (spring) and 30<sup>th</sup> August 2018 (summer). To benefit from periods of optimum plant growth and enhanced microbial activity, soils were sampled during seasons of high rainfall and temperature (spring and summer). Cropland samples were from a beet (*Beta vulgaris*) farm (36 samples) and two fertilizer treatments; 30 t ha<sup>-1</sup> AD (12 samples), 30 t ha<sup>-1</sup> AD + 100 kg ha<sup>-1</sup> AS (ADAS-12 samples) and control (12 samples). Grasslands were sampled from 4 plots with timothy (*Phleum pretense*) and alfalfa (*Medicago sativa*) grass species. Grassland fertilizer treatments were AD (18 samples) applied at

30 t ha<sup>-1</sup> and control (18 samples). To reduce risks and prevent biased results, each treatment was sampled at least three times during the experimental period.

### 3.3.3 Measurement of soil physical and chemical properties

Analysis for soil physical (pH, BD), and chemical (N, K, Mg, CEC and humus) properties from cropland and grassland was conducted by Tokachi Federation of Agricultural Cooperatives and data means provided as sample values (Table S3.1 and S3.2) using methods as described below. Soil pH was measured with a glass electrode pH meter. Three phase distribution was calculated according to the methods by (Dexter, 2004) with a sand plate and a pressure plate extractor, then BD was calculated. Total N was measured with CN coder (N.C-Analyzer, Sumika Chemical Analysis Service, Ltd). The K, Mg, CEC were measured according to the method by Scollenberger and Simon (1945) with an atomic absorption spectrophotometer. Soil humus was visually estimated with a soil color book based on the Mansell System.

### 3.3.4 Soil DNA extraction

Soil DNA was extracted by phenol-chloroform extraction method (Sagova-Mareckova et al., 2008). A 0.5 g of soil sample was mixed with 0.2 g of skim milk to reduce DNA attachment to Andosols (Hoshino and Matsumoto, 2005). Afterwards the sodium phosphate extraction buffer (600 µl) was added, and the mixture shaken for 10 minutes. Phenol-chloroform-isoamyl alcohol (300 µl) and chloroform-isoamyl alcohol (250 µl) were then added to separate the soil solution into an organic phase (lipids and cellular debris) and an aqueous phase (Deoxyribonucleic acid-DNA). Isopropanol (300 µl) and ethanol (1000 µl) were used to precipitate DNA from solution. Nuclease free water (30 µl) was then added to store DNA in solution. An extraction control with all the chemical reagents (without soil) was prepared to check for possible contamination during extraction. The DNA was quantified with Qubit ds DNA HS Assay Kit (Invitrogen, Carlsbad, CA, USA) based on manufacturer's protocol.

### 3.3.5 16S rRNA Gene Amplicon Library Preparation and Sequencing

A metabarcoding approach was used for library preparation and sequencing. Polymerase Chain Reaction (PCR) was conducted targeting V4 region of 16S bacterial gene using 515F (5'-

GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3') primer sets. Each PCR sample contained 10 µl of Amplitaq Gold® 360 Master Mix DNA polymerase, 0.4 µl of forward primer-F515, 0.4 µl of reverse primer-R806 (Caporaso et al., 2010), 8.2 µl of nuclease free water and 1 µl of DNA template, making up a final volume of 20 µl. Positive and negative controls were used during the amplification. The positive control was used to confirm that primers had attached to the DNA while the negative control was used to check for possible contamination during amplification. The controls contained the same mixture of polymerase, primers and water as the samples. However instead of 1 µl of DNA, the positive control contained 1 µl of *Escherichia coli* (Funakoshi Frontiers in Life Sciences, Bunkyo-ku, Tokyo, Japan) while the negative control contained 1 µl of nuclease free water.

PCR was conducted under the following conditions: 95°C for 600 sec and thereafter 30 cycles of 95°C for 30 sec, 57°C for 30 sec and 72°C for 60 sec with a final extension of 72°C for 60 sec. Second PCR was conducted to attach Ion P1 adaptor (Ion Torrent; Life Technologies), specific to each sample. Second PCR samples each contained 10 µl of Amplitaq Gold® 360 Master Mix DNA polymerase, 7.2 µl of nuclease free water, 0.4 µl of reverse primer (P1-R806), 0.4 µl of forward primer (Ion A-Barcode) and 2 µl of DNA template, making up a final volume of 20 µl. PCR was conducted under the following conditions: 95°C for 600 sec and thereafter 5 cycles of 95°C for 30 sec, 57°C for 30 sec and 72°C for 60 sec with a final extension of 72°C for 60 sec. Bioanalyzer High Sensitivity DNA Kit (Agilent Technologies, Palo Alto, CA, USA) was used to determine final library concentration and length, based on manufacturer's protocol. Libraries were then diluted to 50 pM using Low TE (Tris-EDTA; 19 mM Tris base, 0.1 mM EDTA, Ion Torrent; Life Technologies). Sequencing was done with Next Generation Sequencer (Ion PGM, Life Technologies). Ion Chef was used to load libraries to Ion 318 chip (Ion Torrent; Life Technologies) and thereafter sequencing performed with the Ion PGM™ Sequencer (Thermo Fisher Scientific K.K., Japan) using Ion PGM™ Hi-Q™ View Sequencing Kit.

### 3.3.6 Sequence processing and data analysis

Raw sequenced data was processed using QIIME 2 software package version 2019.7 (Boyle et al., 2009). DADA2 was used to cut primer sequences, denoise and select representative sequences (Callahan et al., 2016). Classify-sklearn classifier, which uses naïve bayes classifier to set

taxonomy, was used to classify representative sequences. Closed reference OTU picking was based on Greengenes 13\_8 database and clustered at 99% identity. The Greengenes database is used to classify archaea and bacteria. The classification is based on *denovo* automatic tree construction (McDonald et al., 2011). The Greengenes database has been used successfully in various research projects identify taxa accurately (Boers et al., 2019; McDonald et al., 2011). However, the database is incomplete and contains some unidentified and poorly interpreted sequences (Boers et al., 2019). To avoid bias due to different sampling depth, samples were rarefied to an even sampling depth, based on the sample with the lowest reads.

### 3.3.7 Soil microbial and chemical analysis

Alpha diversity, based on Pielou evenness, OTU counts and Shannon index, were generated in QIIME 2. Pielou evenness, which includes species abundance measurements, is considered a reliable measure of microbial community diversity analysis (Ehsani et al., 2019). As most alpha diversity analysis metrics are based on OTU data we analyzed microbial richness based on OTU counts (Chao, 1984; Chiu et al., 2014). The Shannon diversity index, which considers microbial richness and evenness, was used for alpha diversity analysis. Due to its reliability, Shannon index is one of the most reliable and widely used indices for microbial alpha diversity analysis (Strong, 2016). Weighted unifrac using Principal coordinate analysis (PCoA) was used for beta diversity analysis. Unlike most beta diversity measures, the Weighted unifrac measure takes into account species phylogeny and species relative abundance (Lozupone, 2011; Lozupone and Knight, 2005). To identify community composition variation between treatments, permutational multivariate analysis of variance (PERMANOVA) multivariate analysis using 999 permutations was performed in QIIME 2. Venn diagrams were constructed using phyloseq package (McMurdie and Holmes, 2013) in R software version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria) to visualize shared and unique OTUs between treatments. To determine the amount of variation in community composition explained by soil properties, Non-metric multidimensional scaling (NMDS) was conducted using vegan package (Dickson, 2003) in R software.

Linear Discriminant Analysis (LDA) with LEfSe was used to characterize microbial communities under imposed treatments (Segata et al., 2011). Significant ( $p < .05$ ) alphas were based on Kruskal–Wallis test.



Tax4Fun (Aßhauer et al., 2015) package was used to predict functional profiles from 16S rRNA database. Genes encoding key enzymes were identified using their KEGG orthologs. However, as some of the soil microbes are uncultured or unknown, not all sequenced microbes can be identified to KEGG organisms. Nonetheless, Tax4Fun is considered a reliable bioinformatics tool for predicting microbial genes, due to its high correlation with metagenomic sequencing of functional profiles (AßHauer et al., 2015). Tax4Fun provides a more practical procedure to predict functions compared to shotgun and quantitative polymerase chain reaction (qPCR), which are significantly expensive and prone to experimental variability, respectively (Smith and Osborn, 2009; Kaiser et al., 2016; AßHauer et al., 2015).

All statistical analyses for soil physical and chemical properties were performed using R software. In all analysis,  $p < 0.05$  were considered significant. Two-way and three-way analysis of variance (ANOVA) was performed to determine the effect of land use, season and fertilizer. For significant interactions, Tukey's HSD test was performed.

### 3.4 Results

#### 3.4.1 Evenness, richness and microbial diversity

Land use had a significant ( $p < .05$ ) effect on soil microbial evenness with higher values observed in cropland than grassland (Figure 3.2 A). In cropland, season had a significant ( $p < .05$ ) effect on microbial evenness indices (Figure 3.2 B), with higher values in cropland-summer than cropland-spring. Fertilizer treatments did not have a significant effect in both cropland and grassland (data not shown).




Figure 3.2: Box plots illustrating Pielou evenness at OTU level based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points. p values calculated based on Kruskal-Wallis test.

Land use did not have a significant effect on OTU numbers (Figure 3.3 A). In cropland, season had a significant ( $p < .05$ ) effect on observed OTU numbers (Figure 3.3 B), with more OTUs observed in cropland-summer than cropland-spring. In grassland although season was not significant (Figure 3.3 C), grassland-spring had more OTUs than grassland-summer. Fertilizer treatments did not have a significant effect in both cropland and grassland (data not shown).




Figure 3.3: Box plots illustrating observed OTU numbers based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points.  $p$  values calculated based on Kruskal-Wallis test.

Regarding Shannon Indices, land use did not have significant differences between cropland and grassland (Figure 3.4 A). In cropland, season had a significant ( $p < .05$ ) effect on diversity, with cropland-summer microorganisms being more diverse than cropland-spring (Figure 3.4 B). In grassland although season was not significant, grassland-spring microorganisms were more diverse than grassland-summer (Figure 3.4 C). Fertilizer treatments did not have a significant effect in both cropland and grassland (data not shown).




Figure 3.4: Box plots illustrating Shannon diversity index at OTU level based on land use and season in cropland and grassland. The boxes denote interquartile range with median as a black line and whiskers extending to the most extreme points.  $p$  values calculated based on Kruskal-Wallis test.

Regarding the soil beta diversity (weighted unfrac), samples from the same land use shared a high degree of similarity with significant ( $p < .001$ ) distinct clustering observed in cropland and grassland treatments (Figure 3.5 A). In addition, seasonal treatments shared a significant ( $p < .05$ ) high degree of similarity in both cropland (Figure 3.5 B) and grassland (Figure 3.5 C) treatments.




Figure 3.5: Principal coordinate analysis (PCoA) by weighted unifrac based on land use and season in cropland and grassland. Axes represent the two dimensions explaining greatest proportion of variances in microbial communities. p values calculated based on permutational multivariate analysis of variance.

### 3.4.2 Distribution of OTUs under imposed treatments

#### 3.4.2.1 Unique OTUs

Grassland had a higher number of unique OTUs than cropland (Figure 3.6 A). At phylum level, unique OTUs in cropland were dominated by *Protobacteria* (23.6%, 1156 OTUs), *Planctomycetes* (17.1%, 830 OTUs), *Acidobacteria* (12.9%, 356 OTUs), *Actinobacteria* (12.6%, 343 OTUs) and

*Chloroflexi* (11.5%, 556 OTUs). Unique OTUs in grassland were dominated by *Planctomycetes* (23%, 2006 OTUs), *Proteobacteria* (22.2%, 1855 OTUs), *Acidobacteria* (19.3%, 934 OTUs), *Chloroflexi* (10.2%, 1222 OTUs) and *Nitrospirae* (4.9%, 127 OTUs).

The effect of season varied based on land use. Cropland-summer had a higher number of unique OTUs than cropland-spring (Figure 3.6 B) while grassland-spring had a higher number of unique OTUs than grassland-summer (Figure 3.6 C). At phylum level, the unique OTUs in cropland-summer were dominated by *Planctomycetes* (26.3%, 575 OTUs), *Proteobacteria* (23.8%, 744 OTUs), *Acidobacteria* (10%, 296 OTUs), *Chloroflexi* (8.6%, 371 OTUs) and *Verrucomicrobia* (8%, 165 OTUs). In cropland-spring unique OTUs were dominated by *Proteobacteria* (30.9%, 404 OTUs), *Acidobacteria* (13.1%, 117 OTUs), *Actinobacteria* (10.3%, 125 OTUs), *Planctomycetes* (9.8%, 195 OTUs) and *Chloroflexi* (7.4%, 180 OTUs). At phylum level, grassland-summer unique OTUs were dominated by *Planctomycetes* (22.2%, 485 OTUs), *Proteobacteria* (19.1%, 450 OTUs), *Acidobacteria* (14.4%, 212 OTUs), *Chloroflexi* (11.8%, 306 OTUs) and *Actinobacteria* (5.9%, 122 OTUs). Grassland-spring unique OTUs were dominated by *Proteobacteria* (25.8%, 1290 OTUs), *Planctomycetes* (24.1%, 1294 OTUs), *Acidobacteria* (16.3%, 604 OTUs), *Chloroflexi* (9.2%, 893 OTUs) and *Actinobacteria* (6.1%, 341 OTUs).

In land use treatments all unique phyla in cropland were present in grassland. However, grassland treatment had 22 unique OTUs at phylum level that were not present in cropland. These unique OTUs were dominated by *Spirochaetes* (49 OTUs), *NKB19* (17 OTUs), *WS5* (13 OTUs), *WS1* (10 OTUs), *BHI80-139* (11 OTUs) and *OP8* (12 OTUs). In cropland seasonal treatments *GN04* (1 OTU) and *WWE1* (1 OTU) were found exclusively in cropland-spring while *Thermi* (5 OTUs), *Spirochaetes* (3 OTUs), *MVP-21* (2 OTUs), *BHI80-139* (1 OTU) and *NKB19* (1 OTU) were found exclusively in cropland-summer. In grassland seasonal treatments all unique phyla in grassland-summer were present in grassland-spring. However, grassland-spring treatments had 19 phyla that were not present in grassland-summer. These unique OTUs were dominated by *NKB19* (16 OTUs), *WS5* (13 OTUs), *WS1* (12 OTUs) and *OP8* (10 OTUs).




Figure 3.6: Venn diagrams showing the number of shared and unique OTUs based on land use and season. Percentage values represent relative abundance of OTUs in each section.

#### 3.4.2.2 Shared OTUs

Shared OTUs at phylum level based on land use and seasonal treatments in cropland and grassland were dominated by *Protobacteria Planctomycetes*, *Chloroflexi*, *Acidobacteria* and *Actinobacteria*. Considering the shared microorganisms (2611 OTUs) in land use treatments (Figure 3.6 A), *Actinobacteria*, *Firmicutes* and *Gemmatimonadetes* were significantly ( $p < .05$ ) higher in cropland. In grassland *Euryarchaeota*, *Bacteroidetes*, *Nitrospirae* and *WS3* were significantly ( $p < .05$ ) higher (Figure S3.1A). Of the shared microorganisms (2727 OTUs) in cropland seasonal treatments (Figure 3.6 A), *Crenarchaeota* were significantly ( $p < .05$ ) higher in cropland-spring while *Actinobacteria* and *Cyanobacteria* were significantly higher in cropland-summer (Figure S3.1B). Of the shared microorganisms (3258 OTUs) in grassland seasonal treatments (Figure 3.6 C) *Bacteroidetes* were significantly ( $p < .05$ ) higher in grassland-spring (Figure S3.1C).

#### 3.4.3 Effect of imposed treatments on soil functions

In cropland seasonal treatments, cropland-spring was significantly ( $p < .05$ ) enriched with predicted functions related to *amoA* and *nirK* while cropland-summer had significantly ( $p < .05$ ) enriched with *nifH* (Figure 3.7 A). In grassland season treatments, grassland-spring was significantly ( $p < .05$ ) enriched with predicted functions related to *nifH* while grassland-summer was significantly ( $p < .05$ ) enriched with *amoA* and *nirK* (Figure 3.7 B).




Figure 3.7: The effect of season on predicted soil functions in cropland and grassland. Error bars show statistical differences of selected 16S rRNA gene-predicted functional profiles. Abbreviations ; nifH-nitrogenase, nirK-nitrite reductase, amoA-ammonia monooxygenase.

#### 3.4.4 Relationship between microbial community structure and soil properties

Soil variables were fitted to the non-metric multidimensional scaling (NMDS) plot and distinct clustering was observed in cropland seasonal treatments (Figure 3.8 A). Significant ( $p < .05$ ) correlations between cropland community composition and soil properties were found for pH, K and Mg (Figure 3.8 B). In grassland, significant ( $p < .05$ ) correlations between grassland community composition and soil properties were found for K and BD (Figure 3.8 B).






Figure 3.8: Nonmetric dimensional scaling plot based on bray curtis dissimilarity of microbial community structure in soils among seasonal treatments in cropland and grassland. Arrows show direction of significant ( $p < .05$ ) environmental parameters obtained by fitting environmental factors in the sample ordination space.

### 3.5 Discussion

#### 3.5.1 Effect of land use on the soil microbial community

The land use influenced soil alpha (pielou evenness) and beta (weighted unifrac) diversity. Regarding the alpha diversity, significantly ( $p < .05$ ) higher soil microbial evenness was observed in cropland compared to grassland (Figure 3.2 A). Tillage practices mix the microbial habitat, unifying the microbial community structures and enhancing microorganism's evenness (Lienhard et al., 2013). Thus, our result may have been due to increased tillage intensity in cropland compared to grassland. Considering beta diversity, there was strong clustering of cropland and grassland on PCoA plot based on weighted unifrac distance metrics (Figure 3.5 A). This may have also been caused by differences in management practices between the two land use regimes. Specific to our

study, grassland management involved anaerobic digestate application and minimal soil disturbance, while cropland was characterized by relatively larger frequency of tillage and the addition of relatively larger amounts of inorganic fertilizers. Our results agree with Bissett et al. (2014) and Lauber et al. (2009) who reported that microbial community differences in cropland and grassland could be related to management practices that influenced soil physical and chemical properties, initiating changes in soil microorganisms' structure. Our results also indicate that management practices related to agricultural land use are critical in shaping microbial community structures in agricultural soils, by their influence on alpha (evenness) and beta (weighted unifrac) diversity.

Higher number of unique OTUs were observed in grassland compared to cropland (Figure 3.6 A). This may be due to cropland having lower soil pH (5.23) compared to grassland (5.90). Low soil pH in cropland may have been caused by continuous use of ammonium sulphate, an inorganic nitrogen fertilizer. Inorganic nitrogen fertilizer use lowers soil pH by enhancing oxidation of ammonium ( $\text{NH}_4^+$ ) to nitrate ( $\text{NO}_3^-$ ) which produces hydrogen ions ( $\text{H}^+$ ) (Garvin and Carver, 2003). Additionally, Msimbira and Smith (2020) found that the ideal pH for most soil microorganisms in agricultural soils was from 5.5 to 6.5, therefore their growth would be enhanced in grassland (pH 5.90) compared to cropland (pH 5.23). Furthermore, cropland management systems in Kamishihoro, that involved relatively stronger disturbances to the soils compared to the grasslands could be related to the decreased number of OTUs and microbial stability. Ovreas and Torsvik (1998) conducted research on the effect of soil management practices on microbial diversity and community structure in agricultural soils. They found that increased tillage intensity in cropland reduced soil microbial stability and diversity, in line with our results. Furthermore, a previous study (Atlas, 1984) reported that reduced bacterial stability was characterized by fewer soil microorganisms and low microbial stability, which was observed in cropland in our study. Our results suggest that management practices that directly or indirectly influence soil pH (such as fertilizer additions) and tillage influence the stability of microbial communities in agricultural soils.

The relative abundance of shared and unique OTUs in cropland and grassland were dominated by similar phyla. A similar trend was observed in seasonal treatments in cropland and grassland. The OTUs were dominated by microorganisms belonging to phyla *Proteobacteria*, *Planctomycetes*,

*Chloroflexi*, *Actinobacteria* and *Acidobacteria*. This may be explained in part by the phyla being identified as the dominant soil bacteria taxa in soil 16S rRNA studies (Janssen, 2006; Miyashita, 2015). Phyla *Proteobacteria*, *Planctomycetes*, *Chloroflexi*, *Actinobacteria* and *Acidobacteria* dominating both shared unique OTUs in different treatments (land use and season) suggests that they may play important roles in shaping the soil microorganism's community structure in agricultural soils.

Considering shared microbes at phylum level based on land use (Figure 3.6 A), *Actinobacteria*, *Firmicutes* and *Gemmatimonadetes* were significantly ( $p < .05$ ) higher in cropland compared to grassland (Figure S3.1A). The significantly ( $p < .01$ ) higher N content of cropland (0.51%) compared to grassland soils (0.39%) may have caused the significant increase of *Actinobacteria*, *Firmicutes* and *Gemmatimonadetes* in cropland treatments (Table S3.1). In addition, their abundance had a positive correlation with N suggesting that their growth is closely associated with increased soil N content (Table S3). Other authors (Dai et al., 2018; Madigan et al., 2009) have previously reported increased relative abundance of *Actinobacteria*, *Firmicutes* and *Gemmatimonadetes* in N fertilized cropland soils. Our results may be due to *Actinobacteria*, *Firmicutes* and *Gemmatimonadetes* being classified as copiotrophic. Microorganisms classified as copiotrophic have relatively faster growth rates with research showing that they rapidly increase under N rich conditions in agricultural soils (Fierer et al., 2011; Ho et al., 2017). Of the shared microbes, the significantly ( $p < .05$ ) higher relative abundance of *Euryarchaeota*, *Nitrospirae* and *WS3* in grassland compared to cropland (Figure S3.1A) may be attributed to higher pH in grassland (5.90) compared to cropland (5.23) (Table S3.1). Soil pH influences soil microbial growth and abundance by changing pH homeostasis or regulating soil nutrient availability (Zhalnina et al., 2014). Based on previous studies, *Euryarchaeota*, *Nitrospirae* and *WS3* were positively correlated and relatively more abundant as soil pH increased, in agreement with our research findings (Lauber et al., 2009; Rousk et al., 2010; Zhang et al., 2017). Additionally, the significantly ( $p < .05$ ) higher relative abundance of *Bacteroidetes* in grassland may be attributed to the higher anaerobic digestate application of grasslands compared to croplands. In our study *Bacteroidetes* was identified as one of the main phyla in anaerobic digestate fertilizer (Figure S3.2). Jiang et al. (2019) and Walter et al. (2018) similarly identified that *Bacteroidetes* was one of the major phyla in sequenced anaerobic digestate fertilizer samples. Grassland management in the study site involved 4 times annual application of anaerobic digestate while croplands involved chemical fertilizer

application before seeding. We therefore infer that the high *Bacteroidetes* content of anaerobic digestate fertilizer may have increased the phylum abundance in grassland compared to cropland.

Our results showed that cropland management practices involving beet crop influenced the crop microbial community. Huang et al.(2019) and Cerecetto et al. (2021) conducted research on the effects of continuous tillage of beet crops on the soil microbial community. Their results showed that conventional tillage practices reduced some aspects of microbial diversity, in agreement with our findings. Nonetheless, a limitation of our study is that we considered only one crop (beet) although research (Lange et al., 2015; Yang et al., 2020; Ugrinovic et al., 2014) shows that different crops influence soil properties, microbial structure and diversity in distinct ways. Further studies are therefore needed to determine the influence of different crops on soil microbial diversity.

### 3.5.2 Effect of season on soil microbial community

Significantly ( $p < .05$ ) higher evenness (Figure 3.2 B), richness (Figure 3.3 B) and diversity indices (Fig. 3.4 B and 3.5 B) were recorded in cropland-summer compared to cropland-spring. The same trend was observed in Venn diagrams (Figure 3.6 B) with more unique OTUs recorded in cropland-summer, than cropland-spring. Our findings may be attributed to higher rainfall experienced during cropland-summer compared to cropland-spring, that increased soil moisture content and subsequently microbial diversity. Soil moisture influences microbial structure by driving changes in soil temperature, pH, nutrient distribution, microbes cell metabolism and bacterial movements (Wang et al., 2014). Luo et al. (2019) conducted research on the effect of seasonal changes (summer, spring and autumn) on soil microbial diversity in agricultural soils. Based on their results, due to higher rainfall summer recorded higher diversity compared to spring in agreement with our findings.

In grassland, season did not have a significant ( $p < .05$ ) effect on microbial diversity. These results suggest that grassland had a relatively stable microbial community that could withstand the effect of seasonal changes. Jangrid et al. (2008) conducted research on the impact of land use, management intensity and fertilization on soil microbial community structure in agricultural systems. Based on their results, the absence of seasonal changes on microbial diversity in grassland

could be due to increased bacterial stability which enabled microbes to withstand seasonal changes, in corroboration with our findings.

Our results indicate that cropland microbes have low stability, indicated by significant changes in diversity due to seasonal changes compared to grassland microbes. This implies that conventional tillage practices in Kamishihoro cropland (frequent tillage, chemical fertilizer use) may have negative effects on soil microbial stability.

### 3.5.3 Effect of season on shared phyla and predicted functions in cropland

Of the shared microorganisms in cropland seasonal treatments, *Crenarchaeota* were significantly ( $p < .05$ ) higher in cropland-spring while *Actinobacteria* and *Cyanobacteria* were significantly ( $p < .05$ ) higher in cropland-summer (Figure S3.2B). Higher relative abundance of *Crenarchaeota* in cropland-spring compared to cropland-summer may have been caused by differences in chemical properties between the two seasons. In our experiment cropland-spring had lower pH (5) and higher chemical properties (Mg-30.8, K-50) than cropland-summer (pH-5.46, Mg-22.85, K-36.8) (Table S3.2), which may have enhanced the relative abundance of *Crenarchaeota*. Additionally, based on the correlation analysis, *Crenarchaeota* were negatively correlated to soil pH and positively correlated to Mg and K (Table S3.3). Lehtovirta et al. (2009) conducted research on how soil pH (4.5 to 7.5) regulates abundance and diversity of *Crenarchaeota* in soils. Their results showed that *Crenarchaeota* abundance declined as pH increased from 4.5 to 7.5, in agreement with our findings. Furthermore, Hoshino et al. (2011) and Furtak and Galazka (2019) reported that soil chemical properties such as Mg and K were significant in driving changes in abundance of *Crenarchaeota* in agricultural soils, with increasing values having higher abundance of the phyla.

In our study, predicted functional analysis showed that *amoA* and *nirK* gene was significantly ( $p < .05$ ) higher in cropland-spring compared to cropland-summer (Figure 3.7 A). Nicol and Schleper (2006) reported that *Crenarchaeota* contained genes related to *amoA*. Additionally, Mardanov et al. (2012) found that cultured and uncultured strains of *Crenarchaeota* contained the *nirK* gene which is associated with denitrification process of the nitrogen cycle. We therefore suggest that

the significantly higher relative abundance of *Crenarchaeota* in cropland-spring compared to cropland-summer may have led to higher expression of *amoA* and *nirK* gene in cropland-spring.

The significantly ( $p < .05$ ) higher relative abundance of *Actinobacteria* in cropland-summer may have been caused by higher pH observed in cropland-summer (5.46) compared to cropland-spring (5) (Table S3.2). This was further supported by the positive correlation between *Actinobacteria* and soil pH (Table S3.3). Wang et al. (2019) conducted research on the effect of soil pH (4.5 to 8.5) on the distribution and functions of microorganisms in farmlands in northeastern china. They reported that the relative abundance of *Actinobacteria* was positively correlated to soil pH with higher relative abundance observed as pH increased, in agreement with our study. Additionally, the significantly ( $p < .05$ ) higher relative abundance of *Cyanobacteria* in cropland-summer may be attributed to higher rainfall experienced in cropland-summer compared to cropland-spring. Similar findings have been reported by Moreno-Espíndola et al. (2018) who found that *Cyanobacteria* are positively correlated to increased rainfall, which would be enhanced during summer in our study.

Predicted functional analysis of our study showed that *nifH* was significantly ( $p < .05$ ) higher in cropland-summer compared to cropland-spring (Figure 3.7 A). Reports by Gtari et al. (2011) and Calderoli et al. (2017) identified that *Actinobacteria* and *Cyanobacteria* play functions related to nitrogen fixation through *nifH* gene expression. We therefore suggest that the significantly higher presence of members of *Actinobacteria* and *Cyanobacteria* in cropland-summer lead to higher expression of the *nifH* gene and subsequent functions related to nitrogen fixation in cropland-summer treatments.

Soil pH, Mg and K were significantly ( $p < .05$ ) correlated to differences in soil microbial communities between seasons in cropland (Figure 3.7A). Our results corroborate those of Wessen et al. (2010) and Wang et al. (2019) who reported that pH was one of the primary factors determining soil nutrient availability and therefore the distribution of soil microorganisms in agricultural soils. Additionally, Nicolitch et al. (2019) reported that base cations such as Mg and K had a significant effect on soil microbial distribution. More specifically, Mg and K have been found to improve the soil base saturation and increase nutrient availability, affecting microbial abundance and diversity in agricultural soils (Kreutzer, 1995; Rosberg et al., 2006).

In our study the significant changes in the relative abundance of specific microbes in seasonal treatments was linked to predicted soil functions in cropland. This indicates the importance of cropland seasonal treatments and their associated changes in soil physical and chemical properties, in relation to driving changes in soil microbial diversity, relative abundance and related soil functions. A limitation of our study is it was conducted for only two seasons i.e summer and spring, although the influence of land use has been shown to change in all seasons in various long term research experiments (Li et al., 2019; Sengupta et al., 2020). Therefore, continuous studies to monitor and clarify the effect of land use on soil microbes in different seasons over time should be conducted in Kamishihoro.

#### 3.5.4 Effect of season on shared phyla and predicted functions in grassland

Overall, the season had relatively minor impact on soil microbes in grasslands, when compared to those in croplands, but phylum *Bacterioidetes* was significantly ( $p < .05$ ) higher in grassland-spring than grassland-summer (Figure S3.1C). This may have been due to lower BD during grassland-spring (0.66) compared to grassland-summer (0.77) (Table S3.2). Although zero tillage was conducted in grasslands, tractors were used to apply anaerobic digestate once during spring (April) and the application repeated during summer (August). The higher BD of soils in grassland-summer compared to grassland-spring can therefore be related to soil compaction due to the use of the tractor twice for summer samples compared to once for spring samples. Alaoui et al.(2018) reported that the increased use of farm machinery like tractors contributed to increased soil compaction and BD, in agreement with our findings. In addition, *Bacterioidetes* were negatively correlated to BD further supporting the findings (Table S3.3). Our results are in line with Guo et al. (2019) whose research on the linkage between soil nutrients and microbial characteristics found that the abundance of *Bacterioidetes* were negatively correlated to soil BD. *Bacterioidetes* contain the nitrogen fixing gene *nifH* and subsequently play important roles in agricultural nitrogen fixation (Inoue et al., 2015). In our study predicted functional analysis showed that *nifH* was significantly ( $p < .05$ ) higher in grassland-spring compared to grassland-summer (Figure 3.7). We therefore infer that the significantly higher abundance of *Bacterioidetes* in grassland-spring may have resulted in higher expression of *nifH* gene. In our study K and BD were significantly ( $p < .05$ ) correlated to microbial distribution in grassland season treatments (Figure 3.8 B). Pan et al. (2014) similarly found that K was one of the nutrients that had strong correlations with bacterial

abundance in grassland soils. Additionally, Lee et al. (1996) and Li et al. (2002) reported soil BD to be one of the factors that influenced soil microbial populations, with increase in BD accompanied with a decline in bacterial abundance in agricultural soils. Our results suggest that both soil physical (BD) and chemical (K) properties are significant in causing changes in grassland microorganisms community structure in Kamishihoro grasslands.

### 3.6 Conclusion

Land use (grasslands and croplands) and season (summer and spring) significantly affected the soil microbial community in Kamishihoro region in Japan. Grassland soil microorganisms were resistant to seasonal changes compared to cropland. This was demonstrated by the significant changes in alpha and beta diversity observed in the cropland due to seasonal fluctuations. Conventional tillage practices in cropland may reduce microbial stability. Additionally, the effect of season on diversity varied based on land use with spring having higher values in grassland and summer in cropland. The relative abundance of both shared and unique OTUs in land use and seasonal treatments were dominated by similar microorganisms belonging to *Proteobacteria*, *Planctomycetes*, *Actinobacteria* and *Acidobacteria*. This suggests that these phyla play important roles in shaping regional core microbial community in the studied agricultural soils. Of the shared microorganisms, significantly higher soil microorganisms in cropland and grassland seasonal treatments seemed to influence relative abundance of predicted functions relating to the nitrogen cycle (*nifH*, *nirK*, *amoA*). These results show the importance of seasonal changes on soil microorganisms and the resultant provision of related soil functions within the agricultural environment. The study should provide more insight on the effect of combined effects of land use and seasonal treatments on specific microorganisms within the agricultural system at a regional level. Additionally, the study should contribute towards development of sustainable best management practices within the agricultural environment, that will ensure the continued provision of microbial ecosystem services.



## References

- Abubaker, J., Risberg, K., Jönsson, E., Dahlin, A. S., Cederlund, H., & Pell, M. (2015). Short-Term Effects of Biogas Digestates and Pig Slurry Application on Soil Microbial Activity. *Applied and Environmental Soil Science*, 2015, 1–15. doi.org/10.1155/2015/658542
- Alburquerque, J. A., de la Fuente, C., & Bernal, M. P. (2012). Chemical properties of anaerobic digestates affecting C and N dynamics in amended soils. *Agriculture, Ecosystems & Environment*, 160, 15–22. doi.org/10.1016/j.agee.2011.03.007
- Aßhauer, K. P., Wemheuer, B., Daniel, R., & Meinicke, P. (2015). Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data. *Bioinformatics*, 31(17), 2882–2884. doi.org/10.1093/bioinformatics/btv287
- Atlas, R. M., (1984) Use of microbial diversity measurements to assess environmental stress. In: Klug MJ, Reddy DA (ed) *Current perspectives in microbial ecology*. American Society for Microbiology, Washington DC, pp 540–545
- Banerjee, S., Baah-Acheamfour, M., Carlyle, C. N., Bissett, A., Richardson, A. E., Siddique, T., ... Chang, S. X. (2015). Determinants of bacterial communities in Canadian agroforestry systems. *Environmental Microbiology*, 18(6), 1805–1816. doi.org/10.1111/1462-2920.12986
- Bender, S. F., Wagg, C., & van der Heijden, M. G. A. (2016). An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. *Trends in Ecology & Evolution*, 31(6), 440–452. doi.org/10.1016/j.tree.2016.02.016
- Bissett, A., Abell, G. C. J., Brown, M., Thrall, P. H., Bodrossy, L., Smith, M. C., ... Richardsson, A. E. (2014). Land-use and management practices affect soil ammonia oxidiser community structure, activity and connectedness. *Soil Biology and Biochemistry*, 78, 138–148. doi.org/10.1016/j.soilbio.2014.07.020
- Bowles, T. M., Acosta-Martínez, V., Calderón, F., & Jackson, L. E. (2014). Soil enzyme activities, microbial communities, and carbon and nitrogen availability in organic

- agroecosystems across an intensively-managed agricultural landscape. *Soil Biology and Biochemistry*, 68, 252–262. doi.org/10.1016/j.soilbio.2013.10.004
- Buresova, A., Kopecky, J., Hrdinkova, V., Kamenik, Z., Omelka, M., & Sagova-Mareckova, M. (2019). Succession of Microbial Decomposers Is Determined by Litter Type, but Site Conditions Drive Decomposition Rates. *Applied and Environmental Microbiology*, 85(24). doi.org/10.1128/aem.01760-19
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, 37(8), 852–857. doi.org/10.1038/s41587-019-0209-9
- Calderoli, P. A., Collavino, M. M., Behrends Kraemer, F., Morrás, H. J. M., & Aguilar, O. M. (2017). Analysis of nifH-RNA reveals phylotypes related to Geobacter and Cyanobacteria as important functional components of the N<sub>2</sub>-fixing community depending on depth and agricultural use of soil. *MicrobiologyOpen*, 6(5), e00502. doi.org/10.1002/mbo3.502
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., ... Knight, R. (2010). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences*, 108(Supplement\_1), 4516–4522. doi.org/10.1073/pnas.1000080107
- Chaparro, J. M., Sheflin, A. M., Manter, D. K., & Vivanco, J. M. (2012). Manipulating the soil microbiome to increase soil health and plant fertility. *Biology and Fertility of Soils*, 48(5), 489–499. doi.org/10.1007/s00374-012-0691-4
- Chen, R., Blagodatskaya, E., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., & Kuzyakov, Y. (2012). Decomposition of biogas residues in soil and their effects on microbial growth kinetics and enzyme activities. *Biomass and Bioenergy*, 45, 221–229. doi.org/10.1016/j.biombioe.2012.06.014

- Crowther, T. W., Maynard, D. S., Leff, J. W., Oldfield, E. E., McCulley, R. L., Fierer, N., & Bradford, M. A. (2014). Predicting the responsiveness of soil biodiversity to deforestation: a cross-biome study. *Global Change Biology*, 20(9), 2983–2994. doi.org/10.1111/gcb.12565
- Dai, Z., Su, W., Chen, H., Barberán, A., Zhao, H., Yu, M., ... Xu, J. (2018). Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Global Change Biology*, 24(8), 3452–3461. doi.org/10.1111/gcb.14163
- Degens, B. (2000). Decreases in organic C reserves in soils can reduce the catabolic diversity of soil microbial communities. *Soil Biology and Biochemistry*, 32(2), 189–196. doi.org/10.1016/s0038-0717(99)00141-8
- Díaz, A. I., Oulego, P., Collado, S., Laca, A., González, J. M., & Díaz, M. (2018). Impact of anaerobic digestion and centrifugation/decanting processes in bacterial communities' fractions. *Journal of Bioscience and Bioengineering*, 126(6), 742–749. doi.org/10.1016/j.jbiosc.2018.05.024
- Dixon, P. (2003). VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science*, 14(6), 927–930. doi.org/10.1111/j.1654-1103.2003.tb02228.x
- Faisal, S., Yusuf Hafeez, F., Zafar, Y., Majeed, S., Leng, X., Zhao, S., ... Li, X. (2018). A Review on Nanoparticles as Boon for Biogas Producers—Nano Fuels and Biosensing Monitoring. *Applied Sciences*, 9(1), 59. doi.org/10.3390/app9010059
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences*, 103(3), 626–631. doi.org/10.1073/pnas.0507535103
- Fierer, N., Lauber, C. L., Ramirez, K. S., Zaneveld, J., Bradford, M. A., & Knight, R. (2011). Comparative metagenomic, phylogenetic and physiological analyses of soil microbial

- communities across nitrogen gradients. *The ISME Journal*, 6(5), 1007–1017.  
[doi.org/10.1038/ismej.2011.159](https://doi.org/10.1038/ismej.2011.159)
- Figuerola, E. L. M., Guerrero, L. D., Türkowsky, D., Wall, L. G., & Erijman, L. (2014). Crop monoculture rather than agriculture reduces the spatial turnover of soil bacterial communities at a regional scale. *Environmental Microbiology*, 17(3), 678–688.  
[doi.org/10.1111/1462-2920.12497](https://doi.org/10.1111/1462-2920.12497)
- Furtak, K., & Gałazka, A. (2019). Edaphic factors and their influence on the microbiological biodiversity of the soil environment. *Postępy Mikrobiologii - Advancements of Microbiology*, 58(4), 375–384. [doi.org/10.21307/pm-2019.58.4.375](https://doi.org/10.21307/pm-2019.58.4.375)
- Gałazka, A., Gawyjolek, K., Perzyński, A., Gałazka, R. I., & Jerzy, K. ě. ż. (2017). Changes in Enzymatic Activities and Microbial Communities in Soil under Long-Term Maize Monoculture and Crop Rotation. *Polish Journal of Environmental Studies*, 26(1), 39–46.  
[doi.org/10.15244/pjoes/64745](https://doi.org/10.15244/pjoes/64745)
- Galvez, A., Sinicco, T., Cayuela, M. L., Mingorance, M. D., Fornasier, F., & Mondini, C. (2012). Short term effects of bioenergy by-products on soil C and N dynamics, nutrient availability and biochemical properties. *Agriculture, Ecosystems & Environment*, 160, 3–14. [doi.org/10.1016/j.agee.2011.06.015](https://doi.org/10.1016/j.agee.2011.06.015)
- Garvin, D.F., and Carver, B.F. (2003). Role of the genotype in tolerance to acidity and aluminum toxicity. p. 387–406. In Z. Rengel (ed.) *Handbook of soil acidity*. Marcel Dekker, New York
- Griffiths, B. S., & Philippot, L. (2013). Insights into the resistance and resilience of the soil microbial community. *FEMS Microbiology Reviews*, 37(2), 112–129.  
[doi.org/10.1111/j.1574-6976.2012.00343.x](https://doi.org/10.1111/j.1574-6976.2012.00343.x)
- Gtari, M., Ghodhbane-Gtari, F., Nouioui, I., Beauchemin, N., & Tisa, L. S. (2011). Phylogenetic perspectives of nitrogen-fixing actinobacteria. *Archives of Microbiology*, 194(1), 3–11.  
[doi.org/10.1007/s00203-011-0733-6](https://doi.org/10.1007/s00203-011-0733-6)

- Guo, A., Zhao, Z., Zhang, P., Yang, Q., Li, Y., & Wang, G. (2019). Linkage between soil nutrient and microbial characteristic in an opencast mine, China. *Science of The Total Environment*, 671, 905–913. doi.org/10.1016/j.scitotenv.2019.03.065
- Castro, H. F., Classen, A. T., Austin, E. E., Norby, R. J., & Schadt, C. W. (2009). Soil Microbial Community Responses to Multiple Experimental Climate Change Drivers. *Applied and Environmental Microbiology*, 76(4), 999–1007. doi.org/10.1128/aem.02874-09
- Ho, A., Lonardo, D. P. D., & Bodelier, P. L. E. (2017). Revisiting life strategy concepts in environmental microbial ecology. *FEMS Microbiology Ecology*, fix006. doi.org/10.1093/femsec/fix006
- Hoshino, T. Y., & Matsumoto, N. (2005). Skim Milk Drastically Improves the Efficacy of DNA Extraction from Andisol, a Volcanic Ash Soil. *Japan Agricultural Research Quarterly: JARQ*, 39(4), 247–252. doi.org/10.6090/jarq.39.247
- Takada Hoshino, Y., Morimoto, S., Hayatsu, M., Nagaoka, K., Suzuki, C., Karasawa, T., ... Akiyama, H. (2011). Effect of Soil Type and Fertilizer Management on Archaeal Community in Upland Field Soils. *Microorganisms and Environments*, 26(4), 307–316. doi.org/10.1264/jsme2.me11131
- Inoue, J., Oshima, K., Suda, W., Sakamoto, M., Iino, T., Noda, S., ... Ohkuma, M. (2015). Distribution and Evolution of Nitrogen Fixation Genes in the Phylum Bacteroidetes. *Microorganisms and Environments*, 30(1), 44–50. doi.org/10.1264/jsme2.me14142
- Insam, H., Gómez-Brandón, M., & Ascher, J. (2015). Manure-based biogas fermentation residues – Friend or foe of soil fertility? *Soil Biology and Biochemistry*, 84, 1–14. doi.org/10.1016/j.soilbio.2015.02.006
- Jangid, K., Williams, M. A., Franzluebbers, A. J., Sanderlin, J. S., Reeves, J. H., Jenkins, M. B., ... Whitman, W. B. (2008). Relative impacts of land-use, management intensity and

- fertilization upon soil microbial community structure in agricultural systems. *Soil Biology and Biochemistry*, 40(11), 2843–2853. doi.org/10.1016/j.soilbio.2008.07.030
- Janssen, P. H. (2006). Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and 16S rRNA Genes. *Applied and Environmental Microbiology*, 72(3), 1719–1728. doi.org/10.1128/aem.72.3.1719-1728.2006
- Johnson, D. W., & Curtis, P. S. (2001). Effects of forest management on soil C and N storage: meta-analysis. *Forest Ecology and Management*, 140(2–3), 227–238. doi.org/10.1016/s0378-1127(00)00282-6
- Kou, X., Li, J., Liu, H., Li, B., Yu, X., Cao, X., ... Wang, L. (2020). Characteristics of bacterial biodiversity and community structure in non-rhizosphere soils along zonal distribution of plants within littoral wetlands in inner Mongolia, China. *Global Ecology and Conservation*, 24, e01310. doi.org/10.1016/j.gecco.2020.e01310
- Kreutzer, K. (1995). Effects of forest liming on soil processes. *Plant and Soil*, 168–169(1), 447–470. doi.org/10.1007/bf00029358
- Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., ... Gleixner, G. (2015). Plant diversity increases soil microbial activity and soil carbon storage. *Nature Communications*, 6(1), . doi.org/10.1038/ncomms7707
- Lauber, C. L., Strickland, M. S., Bradford, M. A., & Fierer, N. (2008). The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry*, 40(9), 2407–2415. doi.org/10.1016/j.soilbio.2008.05.021
- Lauber, C. L., Hamady, M., Knight, R., & Fierer, N. (2009). Pyrosequencing-Based Assessment of Soil pH as a Predictor of Soil Bacterial Community Structure at the Continental Scale. *Applied and Environmental Microbiology*, 75(15), 5111–5120. doi.org/10.1128/aem.00335-09
- Lee, W. J., Wood, C. W., Reeves, D. W., Entry, J. A., & Raper, R. L. (1996). Interactive effects of wheel-traffic and tillage system on soil carbon and nitrogen. *Communications in Soil*

- Science and Plant Analysis, 27(18–20), 3027–3043.  
doi.org/10.1080/00103629609369759
- Lehtovirta, L. E., Prosser, J. I., & Nicol, G. W. (2009). Soil pH regulates the abundance and diversity of Group 1.1c Crenarchaeota. *FEMS Microbiology Ecology*, 70(3), 367–376.  
doi.org/10.1111/j.1574-6941.2009.00748.x
- Li, C. H., Ma, B. L., & Zhang, T. Q. (2002). Soil bulk density effects on soil microbial populations and enzyme activities during the growth of maize (*Zea mays* L.) planted in large pots under field exposure. *Canadian Journal of Soil Science*, 82(2), 147–154.  
doi.org/10.4141/s01-026
- Lienhard, P., Terrat, S., Prévost-Bouré, N. C., Nowak, V., Régnier, T., Sayphoummie, S., ... Ranjard, L. (2013). Pyrosequencing evidences the impact of cropping on soil bacterial and fungal diversity in Laos tropical grassland. *Agronomy for Sustainable Development*, 34(2), 525–533. doi.org/10.1007/s13593-013-0162-9
- Liu, T., Zhou, X., Li, Z., Wang, X., & Sun, J. (2019). Effects of liquid digestate pretreatment on biogas production for anaerobic digestion of wheat straw. *Bioresource Technology*, 280, 345–351. doi.org/10.1016/j.biortech.2019.01.147
- Loreau, M. (2010). Linking biodiversity and ecosystems: towards a unifying ecological theory. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1537), 49–60. doi.org/10.1098/rstb.2009.0155
- Luo, X., Wang, M. K., Hu, G., & Weng, B. (2019). Seasonal Change in Microbial Diversity and Its Relationship with Soil Chemical Properties in an Orchard. *PLOS ONE*, 14(12), e0215556. doi.org/10.1371/journal.pone.0215556
- Ma, W., Yang, Z., Hou, S., Ma, Q., Liang, L., Wang, G., ... Zhao, T. (2020 preprint). Effects of living cover on the soil microbial communities and ecosystem functions of hazelnut orchard. *BMC Microbiology*. doi.org/10.21203/rs.3.rs-84985/v1

- Madigan, M. T., Martinko, J. M., Dunlap, P. V., & Clark, D. P. (2009). *Brock biology of microorganisms*, 12th edn. San Francisco, CA: Pearson Benjamin Cummings.
- Mahajan G. R. D., Sharma R. (2008). Bio-fertilizers-A way to sustainable agriculture. *Agrobios Newsletter*, 6, 36-37.
- Mardanov, A. V., Gumerov, V. M., Slobodkina, G. B., Beletsky, A. V., Bonch-Osmolovskaya, E. A., Ravin, N. V., & Skryabin, K. G. (2012). Complete Genome Sequence of Strain 1860, a Crenarchaeon of the Genus *Pyrobaculum* Able To Grow with Various Electron Acceptors. *Journal of Bacteriology*, 194(3), 727–728. doi.org/10.1128/jb.06465-11
- Marques, J. M., da Silva, T. F., Vollu, R. E., Blank, A. F., Ding, G.-C., Seldin, L., & Smalla, K. (2014). Plant age and genotype affect the bacterial community composition in the tuber rhizosphere of field-grown sweet potato plants. *FEMS Microbiology Ecology*, 88(2), 424–435. doi.org/10.1111/1574-6941.12313
- McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS ONE*, 8(4), e61217. doi.org/10.1371/journal.pone.0061217
- Miyashita, N. T. (2015). Contrasting soil bacterial community structure between the phyla Acidobacteria and Proteobacteria in tropical Southeast Asian and temperate Japanese forests. *Genes & Genetic Systems*, 90(2), 61–77. doi.org/10.1266/ggs.90.61
- Moreno-Espíndola, I. P., Ferrara-Guerrero, M. J., Luna-Guido, M. L., Ramírez-Villanueva, D. A., De León-Lorenzana, A. S., Gómez-Acata, S., ... Dendooven, L. (2018). The Bacterial Community Structure and Microbial Activity in a Traditional Organic Milpa Farming System Under Different Soil Moisture Conditions. *Frontiers in Microbiology*, 9, doi.org/10.3389/fmicb.2018.02737
- Msimbira, L. A., & Smith, D. L. (2020). The Roles of Plant Growth Promoting Microorganisms in Enhancing Plant Tolerance to Acidity and Alkalinity Stresses. *Frontiers in Sustainable Food Systems*, 4, doi.org/10.3389/fsufs.2020.00106



- Nicol, G. W., & Schleper, C. (2006). Ammonia-oxidising Crenarchaeota: important players in the nitrogen cycle? *Trends in Microbiology*, 14(5), 207–212.  
[doi.org/10.1016/j.tim.2006.03.004](https://doi.org/10.1016/j.tim.2006.03.004)
- Nicolitch, O., Feucherolles, M., Churin, J.-L., Fauchery, L., Turpault, M.-P., & Uroz, S. (2019). A microcosm approach highlights the response of soil mineral weathering bacterial communities to an increase of K and Mg availability. *Scientific Reports*, 9(1),  
[doi.org/10.1038/s41598-019-50730-y](https://doi.org/10.1038/s41598-019-50730-y)
- Ovreas, L., & Torsvik, V. (1998). Microbial Diversity and Community Structure in Two Different Agricultural Soil Communities. *Microbial Ecology*, 36(3), 303–315.  
[doi.org/10.1007/s002489900117](https://doi.org/10.1007/s002489900117)
- Pabst, H., Kühnel, A., & Kuzyakov, Y. (2013). Effect of land-use and elevation on microbial biomass and water extractable carbon in soils of Mt. Kilimanjaro ecosystems. *Applied Soil Ecology*, 67, 10–19. [doi.org/10.1016/j.apsoil.2013.02.006](https://doi.org/10.1016/j.apsoil.2013.02.006)
- Pan, Y., Cassman, N., de Hollander, M., Mendes, L. W., Korevaar, H., Geerts, R. H. E. M., ... Kuramae, E. E. (2014). Impact of long-term N, P, K, and NPK fertilization on the composition and potential functions of the bacterial community in grassland soil. *FEMS Microbiology Ecology*, 90(1), 195–205. [doi.org/10.1111/1574-6941.12384](https://doi.org/10.1111/1574-6941.12384)
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590–D596. [doi.org/10.1093/nar/gks1219](https://doi.org/10.1093/nar/gks1219)
- Ricker-Gilbert, J., Jumbe, C., & Chamberlin, J. (2014). How does population density influence agricultural intensification and productivity? Evidence from Malawi. *Food Policy*, 48, 114–128. [doi.org/10.1016/j.foodpol.2014.02.006](https://doi.org/10.1016/j.foodpol.2014.02.006)
- Røsberg, I., Frank, J., & Stuanes, A. O. (2006). Effects of liming and fertilization on tree growth and nutrient cycling in a Scots pine ecosystem in Norway. *Forest Ecology and Management*, 237(1–3), 191–207. [doi.org/10.1016/j.foreco.2006.09.045](https://doi.org/10.1016/j.foreco.2006.09.045)

- Rousk, J., Bååth, E., Brookes, P. C., Lauber, C. L., Lozupone, C., Caporaso, J. G., ... Fierer, N. (2010). Soil bacterial and fungal communities across a pH gradient in an arable soil. *The ISME Journal*, 4(10), 1340–1351. doi.org/10.1038/ismej.2010.58
- Sagova-Mareckova, M., Cermak, L., Novotna, J., Plhackova, K., Forstova, J., & Kopecky, J. (2008). Innovative Methods for Soil DNA Purification Tested in Soils with Widely Differing Characteristics. *Applied and Environmental Microbiology*, 74(9), 2902–2907. doi.org/10.1128/aem.02161-07
- Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C. (2011). Metagenomic biomarker discovery and explanation. *Genome Biology*, 12(6), R60. doi.org/10.1186/gb-2011-12-6-r60
- Sengupta, A., Hariharan, J., Grewal, P. S., & Dick, W. A. (2020). Bacterial community dissimilarity in soils is driven by long-term land-use practices. *Agrosystems, Geosciences & Environment*, 3(1), na. doi.org/10.1002/agg2.20031
- Singh, B. K., Quince, C., Macdonald, C. A., Khachane, A., Thomas, N., Al-Soud, W. A., ... Campbell, C. D. (2014). Loss of microbial diversity in soils is coincident with reductions in some specialized functions. *Environmental Microbiology*, 16(8), 2408–2420. doi.org/10.1111/1462-2920.12353
- Smith, J., Wagner-Riddle, C., & Dunfield, K. (2010). Season and management related changes in the diversity of nitrifying and denitrifying bacteria over winter and spring. *Applied Soil Ecology*, 44(2), 138–146. doi.org/10.1016/j.apsoil.2009.11.004
- Solomon, WGO, RW Ndana, and Y Abdulrahim (2012). The Comparative study of the effect of organic manure cow dung and inorganic fertilizer NPK on the growth rate of maize (*Zea mays* L.). *International Research Journal of Agricultural Science and Soil Science*, 2, 16-519.

- Sorensen, P. O., Germino, M. J., & Feris, K. P. (2013). Microbial community responses to 17 years of altered precipitation are seasonally dependent and coupled to co-varying effects of water content on vegetation and soil C. *Soil Biology and Biochemistry*, 64, 155–163. doi.org/10.1016/j.soilbio.2013.04.014
- Szoboszlay, M., Dohrmann, A. B., Poeplau, C., Don, A., & Tebbe, C. C. (2017). Impact of land-use change and soil organic carbon quality on microbial diversity in soils across Europe. *FEMS Microbiology Ecology*, 93(12), doi.org/10.1093/femsec/fix146
- Trivedi, P., Delgado-Baquerizo, M., Anderson, I. C., & Singh, B. K. (2016). Response of Soil Properties and Microbial Communities to Agriculture: Implications for Primary Productivity and Soil Health Indicators. *Frontiers in Plant Science*, 7, na. doi.org/10.3389/fpls.2016.00990
- Trivedi, P., Rochester, I. J., Trivedi, C., Van Nostrand, J. D., Zhou, J., Karunaratne, S., ... Singh, B. K. (2015). Soil aggregate size mediates the impacts of cropping regimes on soil carbon and microbial communities. *Soil Biology and Biochemistry*, 91, 169–181. doi.org/10.1016/j.soilbio.2015.08.034
- van der Heijden, M. G. A., & Wagg, C. (2012). Soil microbial diversity and agro-ecosystem functioning. *Plant and Soil*, 363(1–2), 1–5. doi.org/10.1007/s11104-012-1545-4
- Wang, C.-, Zhou, X., Guo, D., Zhao, J.-, Yan, L., Feng, G.-, ... Zhao, L.-. (2019). Soil pH is the primary factor driving the distribution and function of microorganisms in farmland soils in northeastern China. *Annals of Microbiology*, 69(13), 1461–1473. doi.org/10.1007/s13213-019-01529-9
- Wang, N., Wang, M., Li, S., Sui, X., Han, S., & Feng, F. (2014). Effects of variation in precipitation on the distribution of soil bacterial diversity in the primitive Korean pine and broadleaved forests. *World Journal of Microbiology and Biotechnology*, 30(11), 2975–2984. doi.org/10.1007/s11274-014-1725-x

- Wardle, D. A. (1992). A comparative assessment of factors which influence microbial biomass carbon and nitrogen levels in soil. *Biological Reviews*, 67(3), 321–358.  
doi.org/10.1111/j.1469-185x.1992.tb00728.x
- Wessén, E., Hallin, S., & Philippot, L. (2010). Differential responses of bacterial and archaeal groups at high taxonomical ranks to soil management. *Soil Biology and Biochemistry*, 42(10), 1759–1765. doi.org/10.1016/j.soilbio.2010.06.013
- Yu, P., Liu, S., Han, K., Guan, S., & Zhou, D. (2017). Conversion of cropland to forage land and grassland increases soil labile carbon and enzyme activities in northeastern China. *Agriculture, Ecosystems & Environment*, 245, 83–91.  
doi.org/10.1016/j.agee.2017.05.013
- Yuliar, Nion, Y. A., & Toyota, K. (2015). Recent Trends in Control Methods for Bacterial Wilt Diseases Caused by *Ralstonia solanacearum*. *Microorganisms and Environments*, 30(1), 1–11. doi.org/10.1264/jsme2.me14144
- Zhalnina, K., Dias, R., de Quadros, P. D., Davis-Richardson, A., Camargo, F. A. O., Clark, I. M., ... Triplett, E. W. (2014). Soil pH Determines Microbial Diversity and Composition in the Park Grass Experiment. *Microbial Ecology*, 69(2), 395–406. doi.org/10.1007/s00248-014-0530-2
- Zhang, C., Liu, G., Xue, S., & Wang, G. (2016). Soil bacterial community dynamics reflect changes in plant community and soil properties during the secondary succession of abandoned farmland in the Loess Plateau. *Soil Biology and Biochemistry*, 97, 40–49.  
doi.org/10.1016/j.soilbio.2016.02.013
- Zhang, Y., Shen, H., He, X., Thomas, B. W., Lupwayi, N. Z., Hao, X., ... Shi, X. (2017). Fertilization Shapes Bacterial Community Structure by Alteration of Soil pH. *Frontiers in Microbiology*, 8, na. doi.org/10.3389/fmicb.2017.01325

Žifčáková, L., Větrovský, T. š., Howe, A., & Baldrian, P. (2015). Microbial activity in forest soil reflects the changes in ecosystem properties between summer and winter. *Environmental Microbiology*, 18(1), 288–301. doi.org/10.1111/1462-2920.13026

Zingore, S., Manyame, C., Nyamugafata, P., & Giller, K. E. (2005). Long-term changes in organic matter of woodland soils cleared for arable cropping in Zimbabwe. *European Journal of Soil Science*, 0(0), 050912034650038-??? doi.org/10.1111/j.1365-2389.2005.00707.x

## Supplementary

Table S3.1: The significant effects of land use on soil physical and chemical properties. Asterisks indicate a significant effect of each treatment (\*\*\* 0.001, \*\* 0.01, \* 0.05) and “ns” shows “no significance” based on ANOVA.

	Land use		<i>p</i>
	Cropland	Grassland	
pH	5.23±0.43	5.90±0.30	***
N (%)	0.51±0.04	0.39±0.02	***
K	43.71±8.74	9.23±1.37	***
Mg	26.83±5.98	28.14±4.62	ns
CEC	36.78±3.25	30.50±2.11	***
BD (g/cm <sup>2</sup> )	0.73±0.08	0.70±0.57	ns
Humus	12.94±1.76	9.42±0.58	***

Table S3.2: The significant effects of season on soil chemical properties in cropland and grassland. Asterisks indicate a significant effect of each treatment (\*\*\*) 0.001, \*\* 0.01, \* 0.05) and “ns” shows “no significance” based on ANOVA.

	Cropland			Grassland		
	Summer	Spring	<i>p</i>	Summer	Spring	<i>p</i>
pH	5.46±0.31	5.00±0.42	**	5.80±0.20	5.98±0.36	ns
N (%)	0.52±0.01	0.50±0.03	ns	0.40±0.02	0.39±0.02	ns
K	36.80±3.41	50.63±6.61	***	9.65±0.94	8.80±1.63	ns
Mg	22.83±3.38	30.83±5.34	***	28.35±1.16	27.98±4.58	ns
CEC	37.03±3.60	36.53±3.00	ns	31.70±2.41	29.30±0.61	*
BD (g/cm <sup>2</sup> )	0.73±0.09	0.73±0.07	ns	0.75±0.03	0.66±0.03	***
Humus	13.00±1.59	12.87±1.99	ns	9.79±0.40	9.05±0.49	***

Table S3.3: Correlation analysis (spearman) among soil properties (pH, bulk density-BD, nitrogen-N, potassium-K, magnesium-Mg, cation exchange capacity-CEC, calcium-Ca and humus) and selected phyla based on land use (cropland and grassland) and season treatments. Selected microbes showed significant increase in each treatment. Red color shows significant (  $p < .05$ ) interactions.

<b>Cropland</b>		pH	N	Mg	K	Hu	CEC	Ca	BD
<i>Actinobacteria</i>	rho	-0.330	0.363	-0.176	0.382	0.397	0.372	-0.305	0.085
	<i>p</i>	0.005	0.002	0.140	0.001	0.001	0.001	0.009	0.475
<i>Firmicutes</i>	rho	-0.159	0.107	-0.148	0.286	0.170	0.175	-0.194	0.096
	<i>p</i>	0.183	0.373	0.215	0.015	0.154	0.142	0.102	0.425
<i>Gemmatimonadetes</i>	rho	-0.391	0.376	-0.125	0.492	0.421	0.438	-0.297	0.142
	<i>p</i>	0.001	0.001	0.294	0.001	0.001	0.001	0.011	0.236
<b>Cropland season</b>		pH	N	Mg	K	Hu	CEC	Ca	BD
<i>Actinobacteria</i>	rho	0.118	0.118	0.118	0.118	0.118	0.118	0.118	-0.039
	<i>p</i>	0.641	0.641	0.641	0.641	0.641	0.641	0.641	0.877
<i>Crenarchaeota</i>	rho	-0.131	-0.066	0.131	0.131	-0.066	-0.066	0.066	-0.066
	<i>p</i>	0.604	0.796	0.604	0.604	0.796	0.796	0.796	0.796
<i>Cyanobacteria</i>	rho	-0.177	-0.177	-0.177	-0.177	-0.177	-0.177	-0.177	0.413
	<i>p</i>	0.482	0.482	0.482	0.482	0.482	0.482	0.482	0.088
<b>Grassland</b>		pH	N	Mg	K	Hu	CEC	Ca	BD
<i>Nitrospirae</i>	rho	0.505	-0.473	0.093	-0.598	-0.524	-0.465	0.305	-0.290
	<i>p</i>	0.001	0.001	0.437	0.001	0.001	0.001	0.009	0.014
WS3	rho	0.438	-0.372	0.184	-0.444	-0.438	-0.368	0.338	-0.287
	<i>p</i>	0.001	0.001	0.121	0.001	0.001	0.001	0.004	0.014
<i>Euryarchaeota</i>	rho	0.389	-0.341	0.199	-0.447	-0.396	-0.339	0.343	-0.195
	<i>p</i>	0.001	0.003	0.093	0.001	0.001	0.004	0.003	0.100
<i>Bacteroidetes</i>	rho	-0.098	0.199	-0.143	0.098	0.201	0.207	-0.131	-0.061
	<i>p</i>	0.412	0.094	0.229	0.413	0.091	0.081	0.273	0.613
<b>Grassland-season</b>		pH	N	Mg	K	Hu	CEC	Ca	BD
<i>Bacteroidetes</i>	rho	-0.178	0.210	-0.231	-0.122	0.182	0.173	-0.240	-0.235



<i>p</i>	0.298	0.219	0.176	0.477	0.287	0.313	0.159	0.168
----------	-------	-------	-------	-------	-------	-------	-------	-------




Figure S 3.0.1A: Cladogram showing significantly abundant taxa of shared OTUs. Taxa with significantly different abundance are represented by red and green dots. From the center outwards each ring represents kingdom, phylum, class, order, family and genus levels. The colored shadows represent trends of significant ( $p < .05$ ) taxa based on Kruskal–Wallis test.

## B. Effect of season in cropland




Figure S3.1B: Cladogram showing significantly abundant taxa of shared OTUs based on season treatment in cropland. Taxa with significantly different abundance are represented by red and green dots. From the center outwards each ring represents kingdom, phylum, class, order, family and genus levels. The colored shadows represent trends of significant ( $p < .05$ ) taxa based on Kruskal–Wallis test.

### C. Effect of season in grassland




Figure S3.1C: Cladogram showing significantly abundant taxa of shared OTUs based on season treatment in cropland. Taxa with significantly different abundance are represented by red and green dots. From the center outwards each ring represents kingdom, phylum, class, order, family and genus levels. The colored shadows represent trends of significant ( $p < .05$ ) taxa based on Kruskal–Wallis test.

## **Chapter 4 Liming improves the stability of soil microbial community structures against the application of anaerobic digestate made from dairy wastes**

### 4.1 Abstract

Lime is used to reduce soil acidification in agricultural soils. However, its effects on the soil microbial community are not well understood. Additionally, the soil microbial community is known to be influenced by organic fertilizers. However, the question still remains whether liming influences the magnitude of organic fertilizers impact on soil microbial communities. Therefore, an incubation experiment was performed to understand the effect of lime application (pH = 6.5 and 5.5 for the soils with and without lime, respectively) and fertilizer (digestate, urea and control) on the soil microbial community structures, stability and gene functions. Soils were sampled weekly after the application of fertilizers for a month. For microbial community analysis, DNA was extracted and sequenced targeting 16S rRNA region. For gene abundances i.e 16S rRNA, ammonia oxidizing archaea (AOA), ammonia oxidizing bacteria (AOB), nitrous oxide reductase (*nosZ*) and nitrite reductase (*nirS*) quantitative PCR was conducted. In result, the relative abundance of *Actinobacteria* was influenced more strongly by digestate in lime soils, while *Alphaproteobacteria* was influenced more strongly by digestate in the no lime soil. In no lime treatments, digestate had a significant effect on more operational taxonomic units (146) compared to lime (127), indicating that lime application increased soil microbial community's stability. Liming and fertilizer had a significant effect on 16S rRNA gene copy numbers with the highest values observed in lime plus digestate treatments. Soil pH had a significant on AOA, *nosZ* and *nirS* gene copy numbers with the highest values observed in lime treatments. In the lime treatments digestate application had a positive impact on AOB gene copy numbers but this was not the case for soils without liming treatments. These results indicate that soil pH and fertilizer type should be taken into consideration for the management of functional gene abundance in agricultural soils.

## 4.2 Introduction

About 50% of agricultural areas globally are affected by soil acidity (Von Uexküll and Mutert, 1995). Within the agricultural environment, soil acidification is a natural process that occurs constantly (Goulding, 2016). However, the process can be enhanced or reduced based on farmer management practices (Kunhikrishnan et al., 2016). Application of ammonium-based fertilizers such as urea is one of the management practices that significantly enhances soil acidification (Goulding, 2016; Yang et al., 2012). High levels of acidity negatively affect soil nutrient immobilization and mineralization and (Fageria and Baligar, 2008). Additionally, acidification directly and indirectly causes shifts in microbial community composition and related soil functions (Pang et al., 2019). To reduce the negative effects of soil acidification, there have been considerable efforts towards identifying effective soil management practices that can reduce acidity in agricultural soils.

Liming has been identified as one of the most effective agricultural management practices that can reduce the negative effects of soil acidity (Caires et al., 2010; Chatzistathis et al., 2015; Kunhikrishnan et al., 2016; Mkhonza et al., 2020). Liming directly reduces acidity by neutralizing excessive hydrogen ions in the soil (Bolan et al., 2003). Research shows that reduced soil acidification through liming increases the availability of essential soil nutrients such as nitrogen among others (Fageria and Baligar, 2008; Mkhonza et al., 2020; Daniel et al., 2013, Pang et al., 2019). Changes in soil pH and nutrient availability due to liming are usually accompanied by shifts in the soil microbial community structure (Dai et al., 2018; Zhalnina et al., 2014).

Through its influence on soil pH and nutrient availability, lime application directly or indirectly influences the soil microbial community structure (Xue et al., 2010). Studies researching the effect of lime application on soil microbes are vital because soil microorganisms are significant to the maintenance of soil health by providing ecosystem services such as nitrogen cycling (Singh et al., 2016). Research by Xun et al. (2010) and Narendrula-Kotha et al. (2017) on the response of soil microbial community to changes in soil pH determined that liming regulated the soil microbial community, by enhancing the growth of neutrophilic soil microbes. However, there are contrasting results on the effect of liming on specific soil microbes. For instance, Xun et al. (2010) reported that short term liming enhanced the growth of *Alphaproteobacteria* and *Bacteroidetes* while decreasing the growth of *Actinobacteria* in agricultural soils. However, Narendrula-Kotha et al.

(2017) reported that the relative abundance of *Actinobacteria* and *Proteobacteria* were enhanced with lime application, while *Chloroflexi* and *Firmicutes* were reduced. Nevertheless, there are few studies researching the effect of lime application on the soil microbial community. Most of the available research on lime is focused on the effects of its application on soil pH, nutrients and crop yields (Cifu et al., 2004; Hollad et al., 2019; Li et al., 2018). Therefore, research needs to be conducted to clarify the effect of lime application on the soil microbial community in agricultural soils.

Increased soil pH by liming directly and indirectly affects the soil nitrogen transformation processes (Guo et al., 2019; Zhang et al., 2016). Considering that nitrogen is one of the nutrients that limits agricultural crop production, the effect of pH on its transformation should be studied (Hofstra and Bouwman, 2005). Through its influence on the nitrifier (AOB and AOA) and denitrifier (*nosZ* and *nirS*) community, liming affects soil inorganic nitrogen content and atmospheric nitrogen losses through N<sub>2</sub>O gas emissions (Jha et al., 2020; Liu et al., 2010). The first step of nitrification is catalyzed by AOA and AOB (Nunes-Alves, 2015). Studies on the ecological niches of AOA and AOB have discovered that soil pH is one of the major factors that influences niche separation of the two genes (Schleper, 2010; Xu et al., 2017). In addition, lime influences denitrification (*nirS* and *nosZ*) gene abundance within the agricultural environment. Jha et al. (2020) reported that lime application increased the abundance of *nirS* gene copy numbers in agricultural soils. Furthermore, lime application has the potential to decrease soil N<sub>2</sub>O emissions, through its influence on the denitrification *nosZ* gene (Clough et al., 2003; Zaman et al., 2007). However, due to variable findings (Huang et al., 2011; Jia and Conrad, 2009; Sun et al., 2015) and experimental setup differences among various studies, our understanding of the effect of liming on nitrifier and denitrifier gene abundance is still limited.

Potential changes induced by lime on soil microbes and related functional genes are dependent on the microbial community stability. One of the main aspects of microbial stability is resistance, which is described as the ability of soil microbes to withstand a disturbance over a specific period (Fan et al., 2013; Shade et al., 2012; Ziegler et al., 2017). The soil microbial community composition is one of the factors that may affect stability within the agricultural environment. Microbial communities dominated by oligotrophs, which are characterized by low growth rates and high resource use activity, increase microbial community stability compared to copiotroph

dominated communities (Fierer et al., 2006). Another factor that may affect microbial stability is diversity. Specifically, high microbial diversity has been associated with increased ecosystems stability compared to low diversity (McCann, 2000). Additionally, the soil nutrient content affects microbial stability. Management practices that increase soil nutrient content have been found to enhance microbial stability (De Vries and Shade, 2013). Previous research shows that lime application causes changes in soil microbial community structure, diversity and soil nutrient content (Pang et al., 2019; Wang et al., 2019). Therefore, soil lime additions are expected to influence soil microbial stability. Despite this, there is limited research investigating the specific effects of lime application on soil microbial stability. More information is therefore required to understand the effect of liming on soil microbial stability and how that relates to soil microbe's response to disturbances such as fertilizer additions.

Thus, this study aimed to investigate 1) the interaction between the effects of liming and fertilizer (digestate or urea) application on soil microbial community structures, and 2) the impacts of the interaction on some of the microbial functions, including nitrification and denitrification in soils. Also, we hypothesized that adding lime to soils before digestate application would reduce shifts of the soil microbial community after digestate application.

### 4.3 Materials and methods

#### 4.3.1 Soil sampling and treatments

Soils used in this research were sampled from fields located in Obihiro, Hokkaido, Japan (42° 45'01.9"N, 143° 08'20.4"E). Soils in the region are classified as Andosols (FAO/UNESCO). The sampled soils were sieved using a 4 mm mesh before incubation. Characterization of the initial soil chemical properties; pH, Phosphorous oxide ( $P_2O_5$ ), Potassium oxide ( $K_2O$ ), Nitrogen (N) and Carbon (C) (Table S1) was conducted by Tokachi Federation of Agricultural Cooperatives.

The experimental design was a completely randomized design with two levels of soil pH 5.5 without lime (NL) and 6.5 with lime (L) and three types of fertilizer inputs (digestate, urea and control), with three replicates. For lime treated soils, calcium carbonate ( $CaCO_3$ ) was applied at the rate of 2 g  $CaCO_3$   $kg^{-1}$  dry soil. Two fertilizer inputs were applied as urea and digestate at the rate of 50 kg  $NH_4^+$   $ha^{-1}$  and no fertilizers as the control. Digestate was provided by Hokkaido

biogas plant (Tokachi, Japan). The digestate was produced from dairy cow manure and their bedding materials (wood straw, wood chip). Characterization of digestate chemical properties; pH, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, N, ammonium (NH<sub>4</sub><sup>+</sup>) and nitrates (NO<sub>3</sub><sup>-</sup>) (Table S1) was conducted by Tokachi Federation of Agricultural Cooperatives.

#### 4.3.2 Incubation setup

For the incubation, 50 g of air-dried soil was placed in each plastic cup (6 cm of diameter and 5 cm of depth). The water filled pore space was maintained at 60 % using milli Q and bulk density established at 1 g cm<sup>-3</sup>. The plastic cups were covered with lids which had 6 holes to prevent contamination and excess evaporation. For lime treatments, CaCO<sub>3</sub> was added, and soils incubated for three months to stabilize the pH at 6.5. Fertilizers were applied to the soils after stabilization. Each treatment was replicated 3 times, totalling 72 soil cores.

#### 4.3.3 Soil sampling during the incubation experiments, chemical and microbial analysis

Soils were destructively sampled per week for soil chemical (pH and inorganic nitrogen) and DNA analysis. To determine soil pH, 5 g of fresh soils were mixed with milli Q in 1:5 ration (5:10) and shaken for 30 minutes. The soil pH was then determined using a pH meter (AS800, ASONE Co., Osaka, Japan).

To measure concentration of soil inorganic nitrogen (NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>), 5 g of fresh soils were mixed with 25 ml of 2 M KCl, solution and shaken for 30 minutes. The solution was filtered with 1 µl filter paper (Toyo Roshi Kaisha No. 5C filter paper, Tokyo Roshi Kaisha, Ltd., Tokyo, Japan). Colorimetric measurements for inorganic nitrogen were determined using a flow injection analyzer using the filtrate (AQLA-700, Aqualab Co., Ltd., Tokyo, Japan).

#### 4.3.4 Soil DNA extraction

Soil DNA was extracted from 0.5 g soil using the NucleoSpin ® Soil (Takara Bio Inc, Shiga, Japan) according to the manufacturer's instructions. Skim milk (0.2 g) was used to reduce DNA attachment to Andosol (Hoshino and Matsumoto, 2005). The extracted DNA was purified using Agencourt AM Pure XP Kit (Beckman Coulter, Fullerton, CA, USA) according to the manufactures protocol. The purified DNA was stored at -30°C until further analysis.



#### 4.3.5 16S rRNA gene amplicon library preparation and sequencing

Extracted soil DNA samples were diluted 10 times before the Polymerase Chain Reaction (PCR) targeting 16S rRNA gene using the primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5' GGACTACHVGGGTWTCTAAT-3'). Each PCR sample contained 10 µl of Amplitaq Gold® 360 Master Mix (Applied Biosystems™, Carlsbad, USA) DNA polymerase, 0.4 µl of forward primer-F515, 0.4 µl of reverse primer-R806, 8.2 µl of nuclease free water and 1 µl of DNA template, making up a final volume of 20 µl. PCR was conducted under the following conditions: 95°C for 600 sec and thereafter 30 cycles of 95°C for 30 sec, 57°C for 30 sec and 72°C for 60 sec with a final extension of 72°C for 60 sec. Amplicon purification was conducted using Agencourt AM Pure XP Kit in line with manufacturers protocol. Second PCR was thereafter conducted to attach Ion P1 adaptor (Ion Torrent, Life Technologies), specific to each sample. Second PCR samples each contained 10 µl of Amplitaq Gold® 360 Master Mix DNA polymerase, 7.2 µl of nuclease free water, 0.4 µl of reverse primer (P1-R806), 0.4 µl of forward primer (Ion A-Barcode) and 2 µl of DNA template, making up a final volume of 20 µl. The PCR was conducted under the same conditions as above but with 5 cycles. Bioanalyzer high sensitivity DNA Kit (Agilent Technologies, Palo Alto, CA, USA) was used to determine final library concentration and length, based on manufacturer's protocol. Libraries were then diluted to 50 pM using Low TE (Tris-EDTA, 19 mM Tris base, 0.1 mM EDTA, Ion Torrent, Life Technologies). Sequencing was done with Next Generation Sequencer (Ion PGM, Life Technologies). Ion Chef was used to load libraries to Ion 318 chip (Ion Torrent, Life Technologies) and thereafter sequencing performed with the Ion PGM™ sequencer (Thermo Fisher Scientific K.K., Japan) using Ion PGM™ Hi-Q™ view sequencing kit.

#### 4.3.6 Sequence processing and data analysis

Raw sequenced data was processed using QIIME 2 software package version 2019.7 (Caporaso et al., 2010). DADA2 was used to cut primer sequences, denoise and select representative sequences. Classify-sklearn classifier, which uses naïve brayes classifier to set taxonomy, was used to set representative sequences. Clustering of operational taxonomic units (OTUs) at 99% identity was conducted based on Greengenes 13\_8 database.

#### 4 3.7 Quantitative Polymerase Chain Reaction (qPCR)

Quantitative polymerase chain reaction (qPCR) was performed using the CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Inc, Hercules, USA).

For 16S rRNA, *nosZ* and *nirS* each sample contained 10.4 µl of KAPA SYBR Fast qPCR kit (Agilent Technologies, Palo Alto, CA, USA), 0.8 µl of forward and reverse primers, 7.0 µl of nuclease free water and 1 µl of DNA template, making up a final volume of 20 µl. For AOA and AOB each sample contained 12.5 µl of KAPA SYBR Fast qPCR kit (Agilent Technologies, Palo Alto, CA, USA), 0.5 µl of forward and reverse primers, 9.5 µl of nuclease free water and 2 µl of DNA template, making up a final volume of 25 µl.

The qPCR was conducted with primers and cycling conditions as described in Table 4.1.

Table 4.1: Primer sequences and qPCR conditions used for the study

Target	Primer names and sequences	qPCR condition
16S rRNA (Caporaso et al., 2011)	515F (5'GTGCCAGCMGCCGCGGTAA3') 806R (5'GGACTACHVGGGTWTCTAAT-3')	95°C (600 sec), [30 cycles 95°C (30 sec), 57°C (30 sec),72°C (60 secs)]
<i>nosZ</i> (Kloos et al., 2001)	<i>nosZ</i> 2F (CGCRACGGCAASAAGGTSMSSGT) <i>nosZ</i> 2R (CAKRTGCAKSGCRTGGCAGAA)	95°C (600 sec), [40 cycles 95°C (15 sec), 60°C (30 sec), 72°C (60 secs)]
<i>nirS</i> (Braker et al., 1988)	<i>nirS</i> 1F 5'-CCTA(C/T)TGGCCGCC(A/G)CA(A/G)T <i>NirS</i> 3R GCCGCCGTC(A/G)TG(A/C/G)AGGAA	95°C (600 sec), [40 cycles 95°C (15 sec), 60°C (30 sec), 72°C (60 secs)]
AOA (Meinhardt et al., 2015)	<i>nosZ</i> 2R (CAKRTGCAKSGCRTGGCAGAA) GenAOAR (CCAAGCGGCCATCCAGCTGTATGTCC)	96°C (180 sec), [40 cycles 95°C (30 sec), 65°C (60 sec), 72°C (60 sec)]
AOB (Meinhardt et al., 2015)	1Fmod (CTGGGGTTTCTACTGGTGGTC) GenAOBR (GCAGTGATCATCCAGTTGC)	95°C (600 sec), [40 cycles 95°C (30 sec), 58°C (60 sec), 72°C (60 secs)]

#### 4.3.8 Analysis of bacterial community composition

Permutational multivariate analysis of variance (PERMANOVA), alpha (Pielou evenness, OTU counts, Shannon) and beta (unweighted unifrac) diversity analysis was conducted in QIIME2. The “DESeq” function in R software (R Foundation for Statistical Computing, Vienna, Austria) was used to determine significant changes in OTUs based on different fertilizer treatments. Enhanced volcano package was used to visualize the changes in OTUs using volcano plots (Blighe et al., 2020). To determine the effect of fertilizers on microbial relative occurrence, ternary plot analysis was conducted using ggtern package (Hamilton and Ferry, 2018).

#### 4.3.9 Statistical analysis

All statistical analyses were performed using R software. In all analysis,  $p < 0.05$  were considered significant. For soil pH, inorganic nitrogen and gene abundance analysis, two-way and three-way analysis of variance (ANOVA) was performed to determine the effect of pH, fertilizers and sampling day. For significant interactions, Tukey’s HSD test was performed.

## 4.4 Results

### 4.4.1 Beta Diversity

Principal Coordinate Analysis (PCoA) based on unweighted UniFrac distances showed that samples from the same pH and fertilizer treatments had significant ( $p < 0.001$ ) distinct clustering between treatments (Figure 4.1). For fertilizer, digestate had its own cluster while urea and control treatments clustered together in both L and NL treatments. Therefore, samples from the same pH and fertilizer (digestate) treatments shared a high degree of similarity. Variability between fertilizer treatments was higher in NL compared to L treatments.




Figure 4.1: Principal coordinate analysis (PCoA) by unweighted unifrac based on soil pH and fertilizer treatments. Axis represent the two dimensions explaining greatest proportion of variances in microbial communities.  $p$  values calculated based on permutational multivariate analysis of variance.

### 4.4.2 Changes in OTU distribution and stability

Because our PCoA (Figure 4.1) analyses showed clear separation between the digestate treatment and other two treatments (control and urea referred to as “mix” treatment), volcano plots were

made to identify the OTUs significantly influenced by the digestate treatment, when compared to mix treatments.

Digestate significantly increased more OTUs (146 OTUs) in NL compared to L treatments (127 OTUs) (Figure 4.2). Considering the 127 OTUs that significantly increased due to digestate application in L treatments, most OTUs belonged to families *Sphingomonadaceae* (17 OTUs), *Xanthomonadaceae* (8 OTUs) and *Chthoniobacteraceae* (6 OTUs) (Figure 4.2 A). At phylum level most OTUs that were significantly increased with digestate application in L treatments belonged to *Proteobacteria* (52 OTUs), *Actinobacteria* (19 OTUs) and *Bacteroidetes* (10 OTUs) (Figure S4.1A). Furthermore, majority of OTUs that were significantly decreased in L treatments due to digestate application belonged to phyla *Acidobacteria* (9 OTUs), *Actinobacteria* (3 OTUs) *Euryarchaeota* (3 OTUs) and *Proteobacteria* (3 OTUs).

Considering the 146 OTUs that significantly increased due to digestate application in NL treatments, most OTUs belonged to families *Sphingomonadaceae* (17 OTUs), *Caulobacteraceae* (9 OTUs) and *Chitinophagaceae* (9 OTUs) (Figure 4.2 B). At phylum level majority of the OTUs that were significantly increased with digestate application in NL treatments belonged to *Proteobacteria* (65 OTUs), *Bacteroidetes* (19 OTUs) and *Actinobacteria* (15 OTUs) (Figure S4.1B). Additionally, majority of OTUs that were significantly decreased in NL digestate treatments belonged to phyla *Acidobacteria* (14 OTUs), *Actinobacteria* (4 OTUs) and *Planctomycetes* (4 OTUs).

Furthermore, we considered the overall effect of soil pH on OTU stability during the experimental period. In NL treatments more (121) OTUs significantly increased compared to L (96) treatments (Figure S4.2). The 121 OTUs that significantly increased in NL treatments, most belonged to families *Gaiellaceae* (15 OTUs) and *Sphingomonadaceae* (12 OTUs). At phylum level most of these OTUs belonged to *Actinobacteria* (40 OTUs), *Proteobacteria* (36 OTUs) and *Gemmatimonadetes* (10 OTUs). The 96 OTUs that significantly increased in L treatments, most belonged to families *0319-6A21* (6 OTUs), *Pirellulaceae* (3 OTUs) and *Gemmataceae* (3 OTUs). At phylum level most of these OTUs belonged to *Acidobacteria* (32 OTUs), *Actinobacteria* (13 OTUs) and *Chloroflexi* (11 OTUs).




Figure 4.2: Volcano plots showing the distribution of OTUs abundance according to the adjusted  $p$  value ( $-\log_{10}$  scale) and the log twofold change comparing mix versus digestate. Black nodes in the right side represent OTUs significantly more abundant in the digestate treatments while black one in the left side represent to OTUs more abundant in mix treatments.

#### 4.4.3 Class level microbes that were influenced by fertilizer application based on their abundance

To identify shifts in microbial community abundance due to fertilizer application, ternary plots were prepared (Figure 4.3). Considering OTUs in the top 10 abundant taxa at class level, *Alphaproteobacteria* and *Actinobacteria* were the more abundant in digestate in both L and NL treatments. *Alphaproteobacteria* were more influenced by digestate in NL treatments while *Actinobacteria* were more influenced in L treatments. In NL treatments, the OTUs that were more abundant in urea belonged to *Planctomycetia* and *Gemmatimonadetes*.




Figure 4.3: Ternary plots showing the relative occurrence of OTUs belonging to the top 10 class level microbes. Axes represent fertilizers (digestate, urea, control) treatments in lime and no lime treatments. Points closer to corners indicate that a great proportion of the class total relative abundance found in the fertilizer treatment.

#### 4.4.4 Alpha Diversity

Diversity indices for the bacterial community structures were also influenced by treatments. Regarding Pielou evenness, overall soil pH and sampling day (D) had a significant ( $p < 0.05$ ) effect with the highest values observed in L and D14, respectively (Figure 4.4 A and B). Considering digestate treatments, the interaction of pH and sampling day had a significant ( $p < 0.05$ ) effect with the highest values observed in L treatments on D28. For urea treatments, sampling day had a significant ( $p < 0.05$ ) effect with the highest values observed on D21. On D7, fertilizer had a significant ( $p < 0.01$ ) effect with the highest values observed in L treatments. On D28 the



interaction of soil pH and fertilizer was significant (  $p < 0.05$ ) with the highest values observed in L digestate treatments.

Considering OTU numbers, soil pH, fertilizer and sampling day did not have a significant (  $p < 0.05$ ) effect in both L and NL treatments (Figure 4.4 C and D). However, there were more OTUs observed in NL compared to L treatments. Additionally, digestate and urea fertilizer had the most OTUs in L and NL treatments, respectively.

Overall, the interaction of soil pH, fertilizer and sampling day had a significant (  $p < 0.05$ ) effect on Shannon diversity, with the highest values observed in NL treatments with digestate fertilizer on D21 (Figure 4.4 E and F).









Figure 4.4: Line graphs illustrating alpha diversity based on Pielou evenness, observed OTU numbers and Shannon diversity based on soil pH, fertilizer (F) and sampling day (D). Error bars show mean standard deviation mean  $\pm$  SD (n=3). Significant levels are displayed as \*  $p < 0.05$ .

#### 4.4.5 Gene Abundances

Overall, the effect of soil pH, fertilizer and sampling day had a significant ( $p < 0.05$ ) effect on 16S rRNA gene copy numbers with the highest values observed in L, digestate fertilizer on D21. Considering the effect of pH on stability of 16S rRNA gene copy numbers, fertilizer had a significant ( $p < 0.05$ ) effect in L with the highest values observed in digestate (Figure 4.5 A). In NL, the effect of sampling day was significant ( $p < 0.05$ ) on 16S rRNA gene copy numbers with the highest values observed on D21 (Figure 4.5 B).









Overall, the interaction of pH and sampling day had a significant ( $p < 0.001$ ) effect on AOA gene copy numbers with the highest values observed in NL on D21. Although the effect of fertilizer was not significant, digestate application had a negative effect on AOA gene copy numbers in both L and NL treatments (Figure 4.5 C and D). In control and urea treatments the interaction of pH and sampling day had a significant ( $p < 0.05$ ) effect on AOA gene copy numbers with the highest values observed in NL on D21. On D7 the AOA gene copy numbers were significantly ( $p < 0.001$ ) affected by soil pH with the highest values observed in L. On D21 the AOA gene copy numbers were significantly ( $p < 0.05$ ) affected by fertilizer with the highest values recorded in urea treatments. Additionally, on D28 the interaction of soil pH and fertilizer had a significant ( $p < 0.05$ ) effect on AOA gene copy numbers with the highest values observed in L urea treatments. Considering the effect of pH on stability of AOA gene copy numbers, sampling day had a significant ( $p < 0.001$ ) effect in NL with highest copy numbers recorded on D21 (Figure 4.5 D).

Overall, the interaction of fertilizer and sampling day had a significant ( $p < 0.001$ ) effect on AOB gene copy numbers with the highest values observed in urea on D21. The interaction of pH and sampling day was significant effect on AOB gene copy numbers ( $p < 0.01$ ) with the highest values observed in NL treatments on D21. The effect of fertilizer had a significant effect on all sampling days with the highest values observed in digestate except on D21 where urea was highest. Soil pH had a significant effect on D7 with the highest values observed in L treatments. The effect of fertilizers on AOB gene copy numbers varied based on soil pH. Highest AOB gene copy

numbers in L and NL treatments were observed in digestate and urea, respectively (Figure 4.5 E and F). In both L and NL treatments, fertilizer and sampling day had a significant ( $p < 0.05$ ) effect on AOB gene copy numbers. In L highest AOB gene copy numbers were observed in digestate on D21 and in NL the highest values were in urea on D21 (Figure 5E and F).

Although pH was not significant to *nosZ* gene copy numbers, higher values were observed in L compared to NL (Figure 4.5 G and H). Day had a significant ( $p < 0.01$ ) effect on *nosZ* gene copy numbers with higher values observed in L compared to NL treatments. In control treatments the interaction of soil pH and sampling day had a significant ( $p < 0.05$ ) effect on *nosZ* gene copy numbers with the highest values observed in L treatments on D7. In NL treatments, sampling day had a significant ( $p < 0.001$ ) effect on *nosZ* gene copy numbers with the highest values observed on D21.

Soil pH had a significant ( $p < 0.05$ ) effect on *nirS* gene copy numbers with higher values observed in L treatments. Considering the effect of fertilizers, in digestate treatments, pH and sampling day had a significant ( $p < 0.05$ ) effect on *nirS* gene copy numbers with the highest values observed in L treatments and D21. In Urea, pH had a significant ( $p < 0.05$ ) effect on *nirS* gene copy number numbers with the highest values observed in L treatments. Considering the effect sampling day, on D7 the interaction of soil pH and fertilizer had a significant ( $p < 0.05$ ) effect on *nirS* gene copy numbers with highest numbers observed in L treatments with urea fertilizers. On D14, pH had a significant effect on *nirS* gene copy numbers with the highest values observed in L treatments (Figure 4.5 I and J).






Figure 4.5: The temporal change of 16S rRNA, AOA, AOB, *nosZ* and *nirS* gene copy numbers in soil pH and fertilizer treatments. Error bars show mean standard deviation mean  $\pm$  SD (n=3). Significant levels are displayed as \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .

## 4.5 Discussion

### 4.5.1 Influence of soil pH on microbial response to digestate application

The effect of digestate on the relative abundance of soil bacterial community compositions was different depending on the liming treatment (Figure 4.3). For example, the abundance of *Alphaproteobacteria* and *Actinobacteria* increased in both L and NL treatments due to digestate application. However, *Alphaproteobacteria* was influenced more strongly by the digestate in NL compared to L treatments. To support our results, Shen et al. (2013) conducted research on the effect of soil pH (3.5 to 6.5) on the spatial distribution of bacterial communities and showed that the relative abundance of *Alphaproteobacteria* increased with decrease in soil pH in line with our results. Contrastingly, the abundance of *Actinobacteria* was influenced more by digestate in L compared to NL treatments. Similarly, Wang et al. (2019) conducted research on the effect of pH range from 4.5 to 9.0 on different microbes within the agricultural environments and reported that the abundance of *Actinobacteria* increased in higher pH levels. Thus, the impact of digestate on soil microbiome could be partially manipulated by liming the soil prior to application of digestate, but further studies are needed because our study was not fully focusing on the functional aspects of the microbiome.

#### 4.5.2 Influence of soil pH on the effect of digestate on OTU stability

Lime application increased the stability of soil microbiome (i.e. relatively smaller shifts in soil microbial community structure after the application of digestate in the L treatments compared to NL, Figure 4.1 and 4.2). Our findings can be attributed to the increased abundance of oligotrophic microbes in L compared to NL treatments (Figure S4.3). In our study, 7 out of 10 of the most abundant top 10 microbes at phylum level had higher relative abundance in L compared to NL treatments (Figure S4.3). More specifically *Acidobacteria*, *Chloroflexi*, *Planctomycetes*, *Verrucomicrobia*, *Nitrospirae*, *AD3* and *Crenarchaeota*, all characterized as oligotrophic (Janssen et al., 2002; Kabore et al., 2020; Kits et al., 2017; Moreno-Espíndola et al., 2018; Ramirez-Villanueva et al., 2015; Ward et al., 2009) were higher in L compared to NL treatments (Figure S4.2), and the increase of these microbes may have enhanced the stability of soil microbial community structure in L. De Vries and Shade (2013) reported that there was a direct relationship between soil microbial community structure and their response to external disturbances. Microbial communities dominated by oligotrophs (such as in our study) which have low growth rates and high resource use efficiency (Fierer et al., 2006) are more stable to external disturbance compared to communities dominated by copiotrophs (Bapiri et al., 2010; Lennon et al., 2012) in agreement with our results.

#### 4.5.3 Influence of soil pH on OTU response to digestate application

The current study showed that, in the L treatment, digestate increased the abundance of OTUs belonging to *Xanthomonadaceae* and *Chthoniobacteraceae*. *Xanthomonadaceae* is involved in the decomposition of carbon, nitrogen and phosphorous (Lueders et al., 2006; Folman et al., 2008). *Chthoniobacteraceae* play a role in methane oxidation, nitrogen fixation and polysaccharide degradation (Dunfield et al., 2007; Khadem et al., 2010). Contrastingly, in NL treatments, digestate increased the abundance of OTUs belonging to *Caulobacteraceae* and *Chitinophagaceae*. *Caulobacteraceae* is involved in the decomposition of lignin and cellulose (Pepe-Ranney et al., 2016; Wilhelm et al., 2019). *Chitinophagaceae* has been associated with cellulose decomposition and increased activity of  $\beta$ -glucosidase (Bailey et al., 2013; De Vries et al., 2015). Thus, the impact of digestate application on some soil microbes and related functions can be different depending on soil lime addition prior to the application of digestate.

Some of the families that increased due to digestate application i.e *Xanthomonadaceae*, and *Caulobacteraceae* are categorized as *Proteobacteria* at phylum level (De Vries et al., 2019; Kersters et al., 2006). Their increase in digestate treatments can be related to the copiotrophic nature of *Proteobacteria*. Copiotrophs rapidly increase in nutrient rich conditions (Fierer et al., 2011), which were observed in digestate (Table S4.1) in our study. In addition, *Chitinophagaceae* which belong to phylum *Bacteroidetes* increased in digestate NL but not in L treatments. In our study, *Bacteroidetes* was one of the families that was dominant in digestate fertilizer (Figure S4.4). We therefore infer that the high *Bacteroidetes* content of digestate fertilizer may have increased the phylum abundance in digestate NL treatments. Our results are in agreement with those of Jiang et al. (2019) and Walter et al. (2018) who reported that *Bacteroidetes* was one of the major phyla in digestate fertilizer samples.

Further phylum level analysis showed that *Synergistetes* and *Armatimonadetes* significantly increased exclusively in digestate NL treatments. Because they are not part of the soil major phyla, these species are poorly characterized and their soil functions not well understood (Bandhari and Gupta, 2012; Wang et al., 2019). Although the number of OTUs that were exclusively increased in NL treatments were few, we believe that the result provides more information on the phyla that may be stabilized with the addition of L in agricultural soils. Long term research should therefore be conducted to clarify the mechanisms and influence of soil pH on the stability of these phyla and related soil functions.

#### 4.5.4 Influence of soil pH and fertilizers on microbial diversity

Soil microbes in the L treatment had relatively higher Pielou evenness across all treatments, when compared to the NL treatments (Figure 4.4 A and B). Our results contrast with those of Narendrula-Kotha and Nkongolo (2017) who conducted research on the effect of lime (as one of the treatments) on the soil microbial community structure. Their results showed that lime did not have a significant effect on microbial evenness. The contrasting results may have been due to differences in soil types. Narendrula-Kotha and Nkongolo (2017) conducted their research on sandy soils while our research was conducted on an acidic volcanic soil with high nutrient content. Surprisingly, for Shannon diversity, NL treatments had significantly higher values compared to L (Figure 4.4 E and F). Although we are not sure why the effect of L was limited to evenness, we believe that the



results show the significance of soil pH in influencing specific diversity indices within the agricultural environment. Additionally, different diversity indices should be considered to understand the effect of soil pH on soil microbial diversity in agricultural soils.

Regarding the effect fertilizers on beta diversity, digestate had its own separate microbes cluster while urea and control treatments clustered together in both L and NL treatments (Figure 4.1). This might have been caused by differences in soil chemical properties between digestate and urea treatments. Although we applied the same amount of  $\text{NH}_4^+$  from digestate and urea ( $50 \text{ kg NH}_4^+ \text{ ha}^{-1}$ ), digestate contained other nutrients such as  $\text{P}_2\text{O}_5$  (0.03%),  $\text{K}_2\text{O}$  (0.27%),  $\text{NO}_3^-$  ( $1.92 \text{ mg NO}_3^- \text{ -N kg}^{-1} \text{ soil}$ ) and high pH (7.85) (Table S4.1) which may have enhanced shifts in the soil microbial community. To support this, the results by Tayyab et al. (2018; 2019) reported that increases in soil chemical properties such as pH, nitrogen, phosphorous and potassium caused changes in soil microbial properties, in agreement with our study. Our results demonstrate that in the short term, organic fertilizers such as digestate have more impact on the soil microbial community structure compared to inorganic fertilizers such as urea.

#### 4.5.5 Effect of soil pH and fertilizer application on the abundances of bacteria and their functional genes

##### 4.5.5.1 16S rRNA

An increase in soil pH enhanced the 16S rRNA gene copy numbers, averaging across the fertilizer treatments (Figure 4.5 A and B). Similar results were found by Rousk et al. (2010) who reported that increased soil pH due to liming had a positive correlation to 16S rRNA gene copy numbers. Shen et al. (2010) indicated that soil pH was a major factor affecting bacterial abundance in agricultural soils, with higher bacterial population size observed in less acidic soils, in agreement with our findings.

Digestate had the highest numbers of 16S rRNA gene copy numbers compared to urea and control (Figure 4.5 A and B). This may have been due to digestate fertilizer increasing the soil pH,  $\text{P}_2\text{O}_5$ ,  $\text{K}_2\text{O}$ , and  $\text{NO}_3^-$  (Table S4.1) which enhanced abundance of the 16S rRNA gene. A similar trend was observed by Nölvak et al. (2016) and Truu et al. (2008) who reported that digestate enhanced 16S rRNA gene copy numbers compared to chemical fertilizers. Their result was attributed to

digestates' high nutrient content that increased 16S rRNA gene abundance, in agreement with our study.

Additionally, in L treatments, the effect of fertilizer was significant with the highest values observed in digestate, while in NL treatments, the effect of fertilizers was not significant (Figure 4.5 A and B). Jezille et al. (2009) conducted research on the effect of liming on microbial activity and mineralization in manure amended soils. In agreement with our study, their results showed that fertilizer addition (manure) increased microbial activity in lime treatments than in treatments without lime. They attributed this observation to lime creating favorable conditions for soil microbes to grow efficiently. In their study, lime treatments provided favorable pH, increased soil  $\text{Ca}^{2+}$  and nitrogen content, which enhanced efficient microbial growth with manure treatments, in agreement with our study. Our findings show that soil pH and fertilizer type are important factors to consider in the expression of 16S rRNA gene copy numbers. More specifically, near neutral to basic pH combined with organic fertilizer such as digestate should enhance 16S rRNA gene copy numbers in acidic agricultural soils.

#### 4.5.5.2 Nitrification: AOA, AOB

Soil pH also had a significant effect on AOA gene copy numbers but was not significant for AOB (Figure 4.5). Regarding AOA gene copy numbers, unexpectedly, higher values were observed in L compared to NL treatments (Figure 4.5 C and D). These results contrast with previous reports (He et al., 2012, Zhang et al., 2016) which found that AOA gene copy numbers usually dominate in acidic ( $\text{pH} < 5.5$ ) soils. The discrepancy between the two studies may be attributed to differences in the range of soil pH. The acidic pH range for He et al. (2012) and Zhang et al. (2016) were below 5.5, which increased AOA gene abundance. In our research however, although L treatments had higher soil pH compared to NL, after addition of fertilizers, the NL treatments had a pH that was greater than 5.5 (Figure S4.5) which may have reduced AOA gene copy abundance in NL acidic soil. Alternatively, soil pH caused very little variation in AOB gene copy numbers (Figure 4.5 E and F). Nicol et al. (2008) and Wessen et al. (2010) similarly reported that changes in soil pH did not have a significant effect in AOB gene copy numbers in agreement with our study. Wessen et al. (2010) attributed this result to AOB microbial community having a high stability and resistance to changes in soil pH, which may have been the case in our study.

Fertilizers did not have a significant effect on AOA gene copy numbers but had a significant impact for AOB (Figure 4.5). Additionally, for AOA, digestate application had a negative effect on gene copy numbers in both L and NL treatments (Figure 4.5 C and D). Saunders et al. (2012) conducted research on bacterial responses to digestate application on incubated soils. Based on his results, digestate application reduced AOA gene copy numbers in corroboration with our study. A possible reason for the low gene copy numbers in digestate treatments may be due to AOA preference for oligotrophic environments which have low nutrient content (Erguder et al., 2009; Sterngen et al., 2015). We therefore suggest that the nutrient rich digestate fertilizer treatments would have decreased AOA gene abundance. The positive impact of fertilizer application on AOB gene copy numbers could be due to the high  $\text{NH}_4^+$  content of digestate and urea treatments that stimulated the AOB gene abundance. Our results are in agreement with Wang et al. (2013) and Zhang et al. (2016) who found that increase in soil  $\text{NH}_4^+$  content stimulated the growth of AOB gene copy numbers in agricultural soils. Based on our findings we concluded that the AOB abundance was enhanced by organic (urea) and inorganic (digestate) fertilizer, while organic fertilizers (digestate) reduced the abundance of AOA.

#### 4.5.5.3 Denitrification: *nosZ*, *nirS*

Soil pH had a significant effect on *nosZ* and *nirS* gene copy numbers (Figure 4.5). For *nosZ*, the significant effect of pH was observed on D7 with the highest values recorded in lime (Figure 4.5 G and H). In agreement with our findings, Bergaust et al. (2010) and Čuhel et al. (2010) reported that increasing soil pH stimulated *nosZ* gene copy numbers in agricultural soils, subsequently increasing the conversion of  $\text{N}_2\text{O}$  to  $\text{N}_2$  gas. Regarding *nirS*, our results are in agreement with those of Herold et al. (2018) who conducted research on the effect of pH (4.2 to 6.6) on *nirS* gene copy numbers. Their results showed that *nirS* abundance increased with soil pH in line with our study.

Fertilizers did not have a significant effect on *nosZ* gene copy numbers but had a significant effect on *nirS* (Figure 4.5). Additionally, for *nosZ*, higher gene copy numbers were observed in control treatments compared to digestate and urea (Figure 4.5 G and H). This may have been caused by fertilizers increased  $\text{NO}_3^-$  content that reduced the *nosZ* gene abundance. Our results agree with Huang et al. (2013) who conducted research on the effect of fertilizers on *nosZ* gene. Based on their findings, fertilizers did not enhance *nosZ* gene copy numbers as the gene's abundance was decreased by the presence of  $\text{NO}_3^-$ . Our findings show that fertilizers may increase agricultural

N<sub>2</sub>O emissions by reducing the *nosZ* gene abundance, although more field level studies are needed to confirm this. For *nirS* the interaction of soil pH and fertilizer was significant on D14 with the highest values observed in urea treatments (Figure 4.5 I and J). Our results for *nirS* agree with Dong et al. (2015) who found that fertilizers had a significant effect *nirS* gene copy numbers due to their high NH<sub>4</sub><sup>+</sup> content that enhance the genes abundance.

Based on our findings the gene copy numbers of 16S rRNA and nitrification (AOA and AOB) were fluctuating lesser in L compared to NL treatments (Figure 4.5). Although more long term studies need to be conducted to verify the underlying mechanisms that may have increased 16S rRNA and nitrification gene stability in lime treatments, we believe that the results are important towards understanding the influence of soil pH on soil functions, especially on the nitrification process of the nitrogen cycle.

#### 4.6 Conclusion

Liming increased the stability of soil microbial community structures, when digestate was applied. This was indicated by the lower number of operational taxonomic units that significantly changed due to digestate application in lime treatments compared to no lime. Digestate also had a larger impact on microbial community structures compared to urea. This indicates that in agricultural soils, organic fertilizers may have a larger influence on microbial diversity compared to inorganic fertilizers in the short term. Regarding the functionality of soil microbiomes, lime increased the relative abundance of 16S rRNA, AOA, *nosZ* and *nirS* gene copy numbers. Additionally, lime additions enhanced the positive effect of digestate on 16S rRNA gene copy numbers. The results show the importance of soil pH in modulating changes in 16S rRNA gene abundance and its response to inorganic fertilizer application. Although fertilizers (digestate and urea) increased the abundance of AOB, digestate decreased the abundance of AOA. This indicates that fertilizer sources are important to consider in the management of nitrification gene abundance in agricultural soils. The study should provide more information on the significance of soil pH as an important variable influencing changes in microbial community structure, diversity, stability and functional gene abundance. The study should contribute towards development of sustainable management practices within the agricultural environment that are focused on soil pH management.

## References

- Bailey, V.L., Fansler, S.J., Stegen, J.C., McCue, L.A., 2013. Linking microbial community structure to  $\beta$ -glucosidic function in soil aggregates. *ISME J.* 7, 2044–2053. <https://doi.org/10.1038/ismej.2013.87>.
- Bapiri, A., Bååth, E., Rousk, J., 2010. Drying–Rewetting Cycles Affect Fungal and Bacterial Growth Differently in an Arable. *Soil. Microb Ecol.* 60, 419–428. <https://doi.org/10.1007/s00248-010-9723-5>.
- Bergaust, L., Mao, Y., Bakken, L. R., Frostegård, A., 2010. Denitrification Response Patterns during the Transition to Anoxic Respiration and Posttranscriptional Effects of Suboptimal pH on Nitrogen Oxide Reductase in *Paracoccus denitrificans*. *Appl Environ Microbiol.* 76, 6387–6396. <https://doi.org/10.1128/AEM.00608-10>.
- Bhandari, V., Gupta, R. S., 2012. Molecular signatures for the phylum Synergistetes and some of its subclades. *ALJMAO.* 102, 517–540. <https://doi.org/10.1007/s10482-012-9759-2>.
- Blighe, K., Rana, S., Lewis, M., 2020. Enhanced Volcano: Publication-ready volcano plots with enhanced colouring and labeling. R package version 1.8.0. <https://github.com/kevinblighe/EnhancedVolcano>.
- Bolan, N. S., Adriano, D. C., Curtin, D., 2003. Soil acidification and liming interactions with nutrient and heavy metal transformation and bioavailability. *Adv. Agron.* 215–272. <https://doi.org/10.1016/s0065-2113, 0278006-1>.
- Braker, G., Fesefeldt, A., Witzel, K.P., 1998. Development of PCR primer systems for amplification of nitrite reductase genes (*nirK* and *nirS*) to detect denitrifying bacteria in environmental samples. *Appl. Environ. Microbiol.* 64, 3769–3775. <https://doi.org/64:3769-3775>
- Caires, E. F., Joris, H.A.W., Churka, S., 2010. Long-term effects of lime and gypsum additions on no-till corn and soybean yield and soil chemical properties in southern Brazil. *Soil Use and Manage.* 27, 45–53. <https://doi.org/10.1111/j.1475-2743.2010.00310.x>.

- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat.* 7, 335–336. <https://doi.org/10.1038/nmeth.f.303>.
- Chatzistathis, T., Alifragis, D., Papaioannou, A. (2015). The influence of liming on soil chemical properties and on the alleviation of manganese and copper toxicity in *Juglans regia*, *Robinia pseudoacacia*, *Eucalyptus* sp. and *Populus* sp. plantations. *J. Environ. Manage.* 150, 149–156. <https://doi.org/10.1016/j.jenvman.2014.11.020>.
- Cifu, M., Xiaonan, L., Zhihong, C., Zhengyi, H., Wanzhu, M., 2004. Long-term effects of lime application on soil acidity and crop yields on a red soil in Central Zhejiang. *PLSOA2*. 265, 101–109. <https://doi.org/10.1007/s11104-005-8941-y>.
- Clough, T.J., Sherlock, R.R., Kelliher, F.M., 2003. Can liming mitigate N<sub>2</sub>O fluxes from a urine-amended soil? *Aust. J. Soil Res.* 41, 439 – 457. <https://doi.org/10.1071/sr02079>.
- Čuhel, J., Šimek, M., Laughlin, R.J., Bru, D., Chèneby, D., Watson, C.J., Philippot, L., 2010. Insights into the Effect of Soil pH on N<sub>2</sub>O and N<sub>2</sub> Emissions and Denitrifier Community Size and Activity. *Appl. Environ. Microbiol.* 76, 1870–1878. <https://doi.org/10.1128/aem.02484-09>.
- Dai, Z., Su, W., Chen, H., Barberán, A., Zhao, H., Yu, M., Xu, J., 2018. Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Glob. Change Biol.* 24, 3452–3461. <https://doi.org/10.1111/gcb.14163>.
- Daniel, J.L.P., Santos, M.C., Zopollatto, M., Huhtanen, P., Nussio, L.G., 2013. A data-analysis of lime addition on the nutritive value of sugarcane in Brazil. *Anim. Feed Sci. Technol.* 184, 17–23. <https://doi.org/10.1016/j.anifeedsci.2013.05.009>.
- De Vries, F. T., Shade, A., 2013. Controls on soil microbial community stability under climate change. *Front Microbiol.* 1-6. <https://doi.org/10.3389/fmicb.2013.00265>.

- De Vries, M., Schöler, A., Ertl, J., Xu, Z., Schloter, M., 2015. Metagenomic analyses reveal no differences in genes involved in cellulose degradation under different tillage treatments. *FEMS Microbiol. Ecol.* 91, fiv069. <https://doi.org/10.1093/femsec/fiv069>.
- De Vries, H.J., Beyer, F., Jarzembowska, M., Lipińska, J., van den Brink, P., Zwijnenburg, A., Plugge, C.M., 2019. Isolation and characterization of Sphingomonadaceae from fouled membranes. *NPJ Biofilms Microbiomes* 5, 6. <https://doi.org/10.1038/s41522-018-0074-1>.
- Dong, Z., Zhu, B., Hua, K., Jiang, Y., 2015. Linkage of N<sub>2</sub>O emissions to the abundance of soil ammonia oxidizers and denitrifiers in purple soil under long-term fertilization. *J. Soil Sci.* 61, 5, 799–807. <https://doi.org/10.1080/00380768.2015.1049930>.
- Dunfield, P.F., Yuryev, A., Senin, P., Smirnova, A.V., Stott, M.B., Hou, S., Alam, M., 2007. Methane oxidation by an extremely acidophilic bacterium of the phylum Verrucomicrobia. *Nature*, 450, 879–882. <https://doi.org/10.1038/nature06411>.
- Erguder, T.H., Boon, N., Wittebolle, L., Marzorati, M., Verstraete, W., 2009. Environmental factors shaping the ecological niches of ammonia-oxidizing archaea. *FEMS Microbiology Reviews.* 33, 5, 855–869. <https://doi.org/10.1111/j.1574-6976.2009.00179.x>.
- Fageria, N.K., Baligar, V.C., 2008. Chapter 7 Ameliorating Soil Acidity of Tropical Oxisols by Liming for Sustainable Crop Production. *Adv. Agron*, 345–399. <https://doi.org/10.1016/s0065-2113, 0800407-0>.
- Fan, L., Liu, M., Simister, R., Webster, N.S., Thomas, T., 2013. Marine microbial symbiosis heats up: the phylogenetic and functional response of a sponge holobiont to thermal stress. *The ISME Journal.* 7, 991–1002. <https://doi.org/10.1038/ismej.2012.165>.
- Fierer, N. Jackson, R.B., 2006. The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 626-631.

- Fierer, N., Lauber, C.L., Ramirez, K.S., Zaneveld, J., Bradford, M.A., Knight, R., 2011. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *The ISME Journal*, 6, 1007–1017. <https://doi.org/10.1038/ismej.2011.159>.
- Folman, L.B., Klein Gunnewiek, P.J.A., Boddy, L., De Boer, W., 2008. Impact of white-rot fungi on numbers and community composition of bacteria colonizing beech wood from forest soil. *FEMS Microbiol. Ecol.* 63, 181–191. <https://doi.org/10.1111/j.1574-6941.2007.00425.x>.
- Goulding, K.W.T., 2016. Soil acidification and the importance of liming agricultural soils with particular reference to the United Kingdom. *Soil Use and Manag.* 32, 390–399. <https://doi.org/10.1111/sum.12270>.
- Guo, A., Ding, L., Tang, Z., Zhao, Z., Duan, G., 2019. Microbial response to CaCO<sub>3</sub> application in an acid soil in southern China. *J Environ Sci.* 79, 321–329. <https://doi.org/10.1016/j.jes.2018.12.007>.
- Hamilton, N.E, Ferry, M., (2018). “ggtern: Ternary Diagrams Using ggplot2.” *J. Stat. Softw.* 87, 1–17. <https://doi.org/10.18637/jss.v087.c03>.
- He, J.-Z., Hu, H.-W., Zhang, L.-M., 2012. Current insights into the autotrophic thaumarchaeal ammonia oxidation in acidic soils. *Soil Biol. Biochem.* 55, 146–154. <https://doi.org/10.1016/j.soilbio.2012.06.006>.
- Herold, M.B., Giles, M.E., Alexander, C.J., Baggs, E.M., Daniell, T.J., 2018. Variable response of *nirK* and *nirS* containing denitrifier communities to long-term pH manipulation and cultivation. *FEMS Microbiol. Lett.* 365, fny035. <https://doi.org/10.1093/femsle/fny035>.
- Hofstra, N., Bouwman, A.F., 2005. Denitrification in Agricultural Soils: Summarizing Published Data and Estimating Global Annual Rates. *Nutr. Cycling Agroecosyst.* 72, 267–278. <https://doi.org/10.1007/s10705-005-3109-y>.
- Holland, J.E., White, P.J., Glendining, M.J., Goulding, K.W.T., McGrath, S.P., 2019. Yield responses of arable crops to liming – An evaluation of relationships between yields and



- soil pH from a long-term liming experiment. *Eur J Agron.* 105, 176–188.  
<https://doi.org/10.1016/j.eja.2019.02.016>.
- Hoshino, Y.T., Matsumoto, N., 2005. Skim milk drastically improves the efficacy of DNA extraction from andisol, a volcanic ash soil. *Jarq.* 39, 247–252.  
<https://doi.org/10.6090/jarq.39.247>.
- Huang, R., Wu, Y., Zhang, J., Zhong, W., Jia, Z., Cai, Z., 2011. Nitrification activity and putative ammonia-oxidizing archaea in acidic red soils. *J. Soils Sediments.* 12, 420–428.  
<https://doi.org/10.1007/s11368-011-0450-4>.
- Huang, Y., Li, Y., Yao, H., 2013. Nitrate enhances N<sub>2</sub>O emission more than ammonium in a highly acidic soil. *J. Soils Sediments.* 14, 146–154. <https://doi.org/10.1007/s11368-013-0785-0>.
- Janssen, P.H., Yates, P.S., Grinton, B.E., Taylor, P.M., Sait, M., 2002. Improved Culturability of Soil Bacteria and Isolation in Pure Culture of Novel Members of the Divisions Acidobacteria, Actinobacteria, Proteobacteria, and Verrucomicrobia. *Appl. Environ. Microbiol.* 68, 5, 2391–2396. <https://doi.org/10.1128/aem.68.5.2391-2396.2002>.
- Jezile, G.G., Westfall, D.G., Peterson, G., Child, D.R., Turner, D.P., Averbek, W.V., 2009. Effects of liming on microbial activity and N mineralization in broiler manure-amended soils from Bizana, Eastern Cape, South Africa. *S Afr J Plant Soil.* 26, 18–23.  
<https://doi.org/10.1080/02571862.2009.10639927>.
- Jha, N., Palmada, T., Berben, P., Sagggar, S., Luo, J., McMillan, A.M.S., 2020. Influence of liming-induced pH changes on nitrous oxide emission, *nirS*, *nirK* and *nosZ* gene abundance from applied cattle urine in allophanic and fluvial grazed pasture soils. *Biol. Fertil. Soils.* 56, 811–824. <https://doi.org/10.1007/s00374-020-01460-1>.
- Jia, Z., Conrad, R., 2009. Bacteria rather than Archaea dominate microbial ammonia oxidation in an agricultural soil. *Environ. Microbiol.* 11, 1658–1671. <https://doi.org/10.1111/j.1462-2920.2009.01891.x>.

- Jiang, Y., Dennehy, C., Lawlor, P.G., Hu, Z., McCabe, M., Cormican, P., Gardiner, G.E., 2019. Exploring the roles of and interactions among microbes in dry co-digestion of food waste and pig manure using high-throughput 16S rRNA gene amplicon sequencing. *Biotechnol. Biofuels.* 12, 1-16. <https://doi.org/10.1186/s13068-018-1344-0>.
- Kaboré, O.D., Godreuil, S., Drancourt, M., 2020. Planctomycetes as Host-Associated Bacteria: A Perspective That Holds Promise for Their Future Isolations, by Mimicking Their Native Environmental Niches in Clinical Microbiology Laboratories. *Front. Cell. Infect. Microbiol.* 10, 2235-2988. <https://doi.org/10.3389/fcimb.2020.519301>.
- Kersters, K., De Vos, P., Gillis, M., Swings, J., Vandamme, P., Stackebrandt, E., 2006. Introduction to the Proteobacteria. In: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.H., Stackebrandt E., eds. *The Prokaryotes*, third ed, vol. 5. Springer, New York.
- Khadem, A.F., Pol, A., Jetten, M.S.M., Op den Camp, H.J.M., 2010. Nitrogen fixation by the verrucomicrobial methanotroph ‘*Methylacidiphilum fumariolicum*’. *SolV. Microbiology.* 156, 1052–1059. <https://doi.org/10.1099/mic.0.036061-0>.
- Kits, K.D., Sedlacek, C.J., Lebedeva, E.V., Han, P., Bulaev, A., Pjevac, P., Wagner, M., 2017. Kinetic analysis of a complete nitrifier reveals an oligotrophic lifestyle. *Nature*, 549, 269–272. <https://doi.org/10.1038/nature23679>.
- Kloos, K., Mergel, A., Rösch, C., Bothe, H., 2001. Denitrification within the genus *Azospirillum* and other associative bacteria. *Funct. Plant Biol.* 28, 991-998. <https://doi.org/10.1071/pp01071>.
- Kunhikrishnan, A., Thangarajan, R., Bolan, N.S., Xu, Y., Mandal, S., Gleeson, D.B., Naidu, R., 2016. Functional Relationships of Soil Acidification, Liming, and Greenhouse Gas Flux. *Adv. Agron.* 139, 1-71. <https://doi.org/10.1016/bs.agron.2016.05.001>.
- Lennon, J.T., Aanderud, Z.T., Lehmkuhl, B.K., Schoolmaster, D.R., 2012. Mapping the niche space of soil microorganisms using taxonomy and traits. *Ecology.* 93, 1867–1879. <https://doi.org/10.1890/11-1745.1>.

- Li, Y., Cui, S., Chang, S.X., Zhang, Q., 2018. Liming effects on soil pH and crop yield depend on lime material type, application method and rate, and crop species: a global meta-analysis. *Soils Sediments*. 19, 1393–1406. <https://doi.org/10.1007/s11368-018-2120-2>.
- Liu, B., Mørkved, P.T., Frostegård, A., Bakken, L.R., 2010. Denitrification gene pools, transcription and kinetics of NO, N<sub>2</sub>O and N<sub>2</sub> production as affected by soil pH. *FEMS Microbiol. Ecol.* 72, 407–417. <https://doi.org/10.1111/j.1574-6941.2010.00856.x>.
- Lueders, T., Kindler, R., Miltner, A., Friedrich, M.W., Kaestner, M., 2006. Identification of Bacterial Micropredators Distinctively Active in a Soil Microbial Food Web. *Appl. Environ. Microbiol.* 72, 5342–5348. <https://doi.org/10.1128/aem.00400-06>.
- Mccann, K.S., 2000. The diversity–stability debate. *Nature*. 405, 228–233. <https://doi.org/10.1038/35012234>.
- Meinhardt, K.A., Bertagnolli, A., Pannu, M.W., Strand, S.E., Brown, S.L., Stahl, D.A., 2015. Evaluation of revised polymerase chain reaction primers for more inclusive quantification of ammonia-oxidizing archaea and bacteria. *Environ. Microbiol. Rep.* 7, 354–363. <https://doi.org/10.1111/1758-2229.12259>.
- Mkhonza, N.P., Buthelezi-Dube, N.N., Muchaonyerwa, P., 2020. Effects of lime application on nitrogen and phosphorus availability in humic soils. *Sci Rep* 10, 8634. [//doi.org/10.1038/s41598-020-65501-3](https://doi.org/10.1038/s41598-020-65501-3).
- Moreno-Espíndola, I.P., Ferrara-Guerrero, M.J., Luna-Guido, M.L., Ramírez-Villanueva, D. A., De León-Lorenzana, A.S., Gómez-Acata, S., Dendooven, L., 2018. The Bacterial Community Structure and Microbial Activity in a Traditional Organic Milpa Farming System Under Different Soil Moisture Conditions. *Front. microbiol.* 9, 2737. <https://doi.org/10.3389/fmicb.2018.02737>.
- Narendrula-Kotha, R., Nkongolo, K.K., 2017. Microbial Response to Soil Liming of Damaged Ecosystems Revealed by Pyrosequencing and Phospholipid Fatty Acid Analyses. *PLOS ONE*. 12, e0168497. <https://doi.org/10.1371/journal.pone.0168497>.

- Nicol, G.W., Leininger, S., Schleper, C., Prosser, J.I., 2008. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environ. Microbiol.* 10, 2966–2978. <https://doi.org/10.1111/j.1462-2920.2008.01701.x>.
- Nölvak, H., Truu, M., Kanger, K., Tampere, M., Espenberg, M., Loit, E., Truu, J., 2016. Inorganic and organic fertilizers impact the abundance and proportion of antibiotic resistance and integron-integrase genes in agricultural grassland soil. *Sci. Total Environ.* 562, 678–689. <https://doi.org/10.1016/j.scitotenv.2016.04.035>.
- Pang, Z., Tayyab, M., Kong, C., Hu, C., Zhu, Z., Wei, X., Yuan, Z., 2019. Liming Positively Modulates Microbial Community Composition and Function of Sugarcane Fields. *Agron. J.* 9, 808. <https://doi.org/10.3390/agronomy9120808>.
- Pepe-Ranney, C., Campbell, A.N., Koechli, C.N., Berthrong, S., Buckley, D.H., 2016. Unearthing the Ecology of Soil Microorganisms Using a High Resolution DNA-SIP Approach to Explore Cellulose and Xylose Metabolism in Soil. *Front. microbiol.* 7, 703. <https://doi.org/10.3389/fmicb.2016.00703>.
- Ramirez-Villanueva, D.A., Bello-López, J.M., Navarro-Noya, Y.E., Luna-Guido, M., Verhulst, N., Govaerts, B., Dendooven, L., 2015. Bacterial community structure in maize residue amended soil with contrasting management practices. *Appl. Soil Ecol.* 90, 49–59. <https://doi.org/10.1016/j.apsoil.2015.01.010>.
- Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Fierer, N., 2010. Soil bacterial and fungal communities across a pH gradient in an arable soil. *The ISME Journal.* 4, 1340–1351. <https://doi.org/10.1038/ismej.2010.58>.
- Saunders, O.E., Fortuna, A.M., Harrison, J.H., Cogger, C.G., Whitefield, E., Green, T., 2012. Gaseous Nitrogen and Bacterial Responses to Raw and Digested Dairy Manure Applications in Incubated Soil. *Environ. Sci. Technol.* 46, 11684–11692. <https://doi.org/10.1021/es301754s>.

- Schleper, C., 2010. Ammonia oxidation: different niches for bacteria and archaea? *The ISME Journal*. 4, 1092–1094. <https://doi.org/10.1038/ismej.2010.111>.
- Shade, A., Peter, H., Allison, S.D., Baho, D.L., Berga, M., Bürgmann, H., Handelsman, J., 2012. Fundamentals of Microbial Community Resistance and Resilience. *Front. Microbiol.* 3, 417. <https://doi.org/10.3389/fmicb.2012.00417>.
- Shen, C., Xiong, J., Zhang, H., Feng, Y., Lin, X., Li, X., Chu, H., 2013. Soil pH drives the spatial distribution of bacterial communities along elevation on Changbai Mountain. *Soil Biol. Biochem.* 57, 204–211. <https://doi.org/10.1016/j.soilbio.2012.07.013>.
- Shen, J.P., Zhang, L.M., Guo, J.F., Ray, J.L., He, J.Z., 2010. Impact of long-term fertilization practices on the abundance and composition of soil bacterial communities in Northeast China. *Appl. Soil Ecol.* 46, 119–124. <https://doi.org/10.1016/j.apsoil.2010.06.015>.
- Singh, J.S., Kumar, A., Rai, A.N., Singh, D.P., 2016. Cyanobacteria: A Precious Bio-resource in Agriculture, Ecosystem, and Environmental Sustainability. *Front. Microbiol.* 7, 529. <https://doi.org/10.3389/fmicb.2016.00529>.
- Sterngren, A.E., Hallin, S., Bengtson, P., 2015. Archaeal Ammonia Oxidizers Dominate in Numbers, but Bacteria Drive Gross Nitrification in N-amended Grassland Soil. *Front. Microbiol.* 6, 1350. <https://doi.org/10.3389/fmicb.2015.01350>.
- Sun, R., Guo, X., Wang, D., Chu, H., 2015. Effects of long-term application of chemical and organic fertilizers on the abundance of microbial communities involved in the nitrogen cycle. *Appl. Soil Ecol.* 95, 171–178. <https://doi.org/10.1016/j.apsoil.2015.06.010>.
- Tayyab, M., Islam, W., Arafat, Y., Pang, Z., Zhang, C., Lin, Y., Zhang, H., 2018. Effect of Sugarcane Straw and Goat Manure on Soil Nutrient Transformation and Bacterial Communities. *Sustainability.* 10, 2361. <https://doi.org/10.3390/su10072361>.
- Tayyab, M., Islam, W., Lee, C.G., Pang, Z., Khalil, F., Lin, S., Zhang, H., 2019. Short-Term Effects of Different Organic Amendments on Soil Fungal Composition. *Sustainability.* 11, 198. <https://doi.org/10.3390/su11010198>.

- Truu, M., Truu, J., Ivask, M., 2008. Soil microbiological and biochemical properties for assessing the effect of agricultural management practices in Estonian cultivated soils. *Eur. J. Soil Biol.* 44, 231–237. <https://doi.org/10.1016/j.ejsobi.2007.12.003>.
- von Uexküll, H.R., Mutert, E., 1995. Global extent, development and economic impact of acid soils. *Plant soil.* 171, 1–15. <https://doi.org/10.1007/bf00009558>.
- Walter, A., Probst, M., Franke-Whittle, I.H., Ebner, C., Podmirseg, S.M., Etemadi-Shalamzari, M., Insam, H., 2018. Microbiota in anaerobic digestion of sewage sludge with and without co-substrates. *Water Environ J.* 33, 214–222. <https://doi.org/10.1111/wej.12392>.
- Wang, C., Zhou, X., Guo, D., Zhao, J., Yan, L., Feng, G., Zhao, L., 2019. Soil pH is the primary factor driving the distribution and function of microorganisms in farmland soils in northeastern China. *Ann. Microbiol.* 69, 1461–1473. <https://doi.org/10.1007/s13213-019-01529-9>.
- Wang, Y., Zhu, G., Song, L., Wang, S., Yin, C., 2013. Manure fertilization alters the population of ammonia-oxidizing bacteria rather than ammonia-oxidizing archaea in a paddy soil. *J. Basic Microbiol.* 54, 190–197. <https://doi.org/10.1002/jobm.201200671>.
- Wang, C.Y., Zhou, X., Guo, D., Zhao, J.H., Yan, L., Feng, G.Z., Zhao, L.P., 2019. Soil pH is the primary factor driving the distribution and function of microorganisms in farmland soils in northeastern China. *Ann. Microbiol.* 69, 1461–1473. <https://doi.org/10.1007/s13213-019-01529-9>.
- Ward, N. L., Challacombe, J.F., Janssen, P.H., Henrissat, B., Coutinho, P.M., Wu, M., Kuske, C.R., 2009. Three Genomes from the Phylum Acidobacteria Provide Insight into the Lifestyles of These Microorganisms in Soils. *Appl. Environ. Microbiol.* 75, 2046–2056. <https://doi.org/10.1128/aem.02294-08>.
- Wessén, E., Nyberg, K., Jansson, J.K., Hallin, S., 2010. Responses of bacterial and archaeal ammonia oxidizers to soil organic and fertilizer amendments under long-term management. *Appl. Soil Ecol.* 45,93–200. <https://doi.org/10.1016/j.apsoil.2010.04.003>.

- Wilhelm, R.C., Singh, R., Eltis, L.D., Mohn, W.W., 2018. Bacterial contributions to delignification and lignocellulose degradation in forest soils with metagenomic and quantitative stable isotope probing. *The ISME Journal*. 13, 413–429. <https://doi.org/10.1038/s41396-018-0279-6>.
- Xu, X., Liu, X., Li, Y., Ran, Y., Liu, Y., Zhang, Q., Di, H., 2017. High temperatures inhibited the growth of soil bacteria and archaea but not that of fungi and altered nitrous oxide production mechanisms from different nitrogen sources in an acidic soil. *Soil Biol. Biochem.* 107, 168–179. <https://doi.org/10.1016/j.soilbio.2017.01.003>.
- Xue, D., Huang, X., Yao, H., Huang, C., 2010. Effect of lime application on microbial community in acidic tea orchard soils in comparison with those in wasteland and forest soils. *J Environ Sci.* 22, 1253–1260. <https://doi.org/10.1016/s1001-0742, 0960246-1>.
- Yang, Y., Ji, C., Ma, W., Wang, S., Wang, S., Han, W., Smith, P., 2012. Significant soil acidification across northern China's grasslands during 1980s-2000s. *Glob. Change Biol.* 18, 2292–2300. <https://doi.org/10.1111/j.1365-2486.2012.02694.x>.
- Zaman, M., Nguyen, M.L., Matheson, F., Blennerhassett, J.D., Quin, B.F., 2007. Can soil amendments , zeolite or lime shift the balance between nitrous oxide and dinitrogen emissions from pasture and wetland soils receiving urine or urea-N? *Soil Res.* 5, 7, 543. <https://doi.org/10.1071/sr07034>.
- Zhalnina, K., Dias, R., de Quadros, P.D., Davis-Richardson, A., Camargo, F.A.O., Clark, I. M., Triplett, E.W., 2014. Soil pH Determines Microbial Diversity and Composition in the Park Grass Experiment. *Microb.* 69, 395–406. <https://doi.org/10.1007/s00248-014-0530-2>.
- Zhang, K., Chen, L., Li, Y., Brookes, P.C., Xu, J., Luo, Y., 2016. The effects of combinations of biochar, lime, and organic fertilizer on nitrification and nitrifiers. *Biology and Fertility of Soils*, 53,77–87. <https://doi.org/10.1007/s00374-016-1154-0>.

Ziegler, M., Seneca, F.O., Yum, L.K., Palumbi, S.R., Voolstra, C.R., 2017. Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat.* 8, 14213.  
<https://doi.org/10.1038/ncomms14213>.



Supplementary

Table S4.1: Digestate and Kamishihoro soil chemical properties

Parameter	Soil	Digestate
Water content %	-	98.4
pH (H <sub>2</sub> O)	5.5	7.85
P <sub>2</sub> O <sub>5</sub> (mg kg <sup>-1</sup> )	86	0.03 (%)
K <sub>2</sub> O (mg kg <sup>-1</sup> )	250	0.27 (%)
Total N %	0.35	0.22
NO <sub>3</sub> <sup>-</sup> -N (mg N kg <sup>-1</sup> )	-	1.92
NH <sub>4</sub> <sup>+</sup> -N (mg N kg <sup>-1</sup> )	-	2125.49
Total C %	4	-




Figure S4.1: The relative abundance of phylum level OTUs that significantly increased in digestate treatments in L and NL treatments based on volcano plots.




Figure S4.2: Volcano plots showing the distribution of OTUs abundance according to the adjusted p value ( $-\log_{10}$  scale) and the  $\log_2$  fold change between lime and no lime treatments. Each point represents an individual OTUs. Black points indicate OTUs that were not significant ( $p > 0.05$ ), blue dots indicate OTUs that were significant ( $p < 0.05$ ), green dots indicate OTUs that had a  $\log_2$  fold change that was greater or less than 2 and red dots indicate OTUs that were significant ( $p < 0.05$ ) and had a  $\log_2$  fold change that was greater or less than 2.




Figure S4.3: Relative abundance of the most abundant microbes at phylum level in lime and no lime treatments.




Figure S 4.4: Relative abundance of the most abundant microbes at phylum level in digestate.




Figure S4.5: The temporal change of soil pH in control, digestate and urea treatments. Error bars show mean standard deviation mean  $\pm$  SD (n=3). Significant levels are displayed as \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .




Figure S 4.6: Relative abundance of microbes at phylum level in digestate.

## **Chapter 5 The impact of separating anaerobic digestate on soil nutrients, yields and microbial functions in agricultural soils**

### 5.1 Abstract

The use of digestate as an organic fertilizer has been gaining popularity in recent years. High transportation costs and soil flood risks are some of the main challenges that have been associated with digestate use as a fertilizer. Separation of digestate in solid and liquid fractions has been proposed as a potential solution to address these challenges. However, the question remains whether the use of separated digestate fractions influences soil nutrient dynamics and microbial properties in agricultural soils. Therefore, an incubation experiment was setup to analyze the effect of soil types (Kamishihoro and Arakida) and fertilizers (digestate, solid digestate, liquid digestate, chemical and control) on soil and plant (Japanese mustard spinach) nutrient dynamics, microbial abundance and crop yields. Based on the results, soil type significantly influenced the above ground plant biomass and Nitrogen (N) content with higher values observed in Kamishihoro compared to Arakida. Kamishihoro had significantly higher soil inorganic N ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ ) content compared to Arakida soil. Fertilizer had a significant effect on soil  $\text{NO}_3^-$  (solid-Kamishihoro, digestate-Arakida) and  $\text{NH}_4^+$  (liquid-Kamishihoro, chemical-Arakida) content. The results indicate that digestate and its derived fertilizers (solid digestate, liquid digestate) can be used to influence soil inorganic N content in agricultural soils. Digestate derived fertilizers influenced soil pH with highest values observed in liquid in both Kamishihoro and Arakida soils. Regarding gene abundance, Arakida soil had significantly higher 16S rRNA gene copy numbers compared to Kamishihoro. Regarding soil  $\text{N}_2\text{O}$  emissions, Kamishihoro soil had significantly higher emissions compared to Arakida. Additionally, digestate and solid fertilizer had significantly higher  $\text{N}_2\text{O}$  emissions compared to liquid in both Kamishihoro and Arakida soils. The results indicate the importance of soil type and digestate derived fertilizers in modulating soil gene abundance and  $\text{N}_2\text{O}$  emissions in agricultural soils. Furthermore, the separation of digestate into solid and liquid fractions can be considered a reliable method to regulate various aspects of soil nutrient dynamics in agricultural soils.

## 5.2 Introduction

Digestate is the semi liquid byproduct of the biogas production process (Weiland, 2009). The use of digestate as an organic fertilizer within the agricultural environment has been increasing, attributed to its high nutrient content (Athurson, 2009). More specifically, digestate application increases the soil organic carbon (OC), inorganic nitrogen (N), phosphorous (P), potassium (K) among other nutrients (Šimon et al., 2016; Smith et al., 2014). Due to digestates' high ammonium ( $\text{NH}_4^+$ ) content, various researchers consider the fertilizer as a practical substitute for reducing the use of inorganic N fertilizers in the agricultural environment (Risberg et al., 2017; Weiland, 2009). Additionally, with the increasing negative effects of inorganic chemical N fertilizers use on the environment such as eutrophication, soil acidification and increased nitrous oxide  $\text{N}_2\text{O}$  emissions, (Bouwman et al., 2005; Guo et al., 2010), the use of organic fertilizers such as digestate has been increasing globally (Weiland, 2009; Gielnik et al., 2019).

Although the use of digestate has been increasing in recent years, there have been observed challenges associated with its use. Some of the challenges include the high storage and transportation costs, increased soil flood risks and challenges in applying specific nutrient concentrations of digestate compared to chemical fertilizers. Digestate use as a fertilizer has associated high transportation and storage costs due to its low nutrient content and large volume compared to chemical fertilizers (Camilleri-Rumbau et al., 2021). The high soil moisture (97%) of digestate presents agricultural challenges to its use as a fertilizer (Greenberg et al., 2019). This problem is amplified in humid environments or during periods of high rainfall when the use of digestate has been found to increase temporary soil flooding risks (Vaneckhaute et al., 2013). Additionally, the high soil moisture and varying nutrient contents complicate precision nutrient application of digestate based on soil nutrient requirements (Drosg et al., 2015).

The separation of digestate into solid and liquid fractions before soil application has been proposed as a potential solution to address the identified challenges accompanied with its use (Heviánková et al., 2015). Various methods have been developed to separate digestate into solid and liquid portions, with the most common being the combined use of press pretreatments and filters (Dosch, 1996). However, to increase efficiency of the process, the use of chemical treatments such as flocculants (water soluble metal salts) has been gaining popularity (Nowostawska et al., 2005; Meixner et al., 2015). Holm-Nielsen et al.(2009) reported that the separation of digestate to solid

and liquid fractions was a simple and efficient method to reduce high transport costs and simplify soil application of digestate. The separated solid and liquid fractions of digestate contain different soil nutrients, which helps farmers to apply the fertilizer based on soil requirements (Bauer et al., 2009). This is especially important for different soil types, with research showing that the impact of digestate application on soil nutrients varies based on soil types (Möller et al., 2015). The liquid fraction is considered a suitable substitute to the use of inorganic N fertilizers due to its high  $\text{NH}_4^+$ -N content (Sigurnjak et al., 2017). On the other hand, the solid fraction usually has a high OC and P content (Egene et al., 2020). These observed differences in soil nutrient content of the solid and liquid fractions are expected to influence the effect of their application on agricultural soils. Despite this observation, there is limited research investigating the effect of separated fractions of digestate on soil nutrient dynamics and microbial abundance in agricultural soils. Most of the available research is focused on the effect of digestate on crop yields (Tampio et al., 2015; Šimon et al., 2015; Risberg et al., 2017). Research therefore needs to be conducted to clarify the effects of separated solid and liquid fractions of digestate on soil and plant nutrient dynamics and microbial abundance in agricultural soils.

Thus, this study aimed to investigate 1) the impact of solid liquid separation of digestate on soil and plant nutrient dynamics and Japanese spinach yields 2) the impacts of solid liquid separation of digestate on soil  $\text{N}_2\text{O}$  emissions 3) the impacts of solid liquid separation of digestate on microbial abundance. We hypothesized that the digestate liquid fraction would have the highest soil N content, crop yields,  $\text{N}_2\text{O}$  emissions and microbial abundance compared to the solid fraction.

### 5.3 Materials and methods

#### 5.3.1 Study site and soil characteristics

Two types of soils (Andosols and Alluvial) were used for the study. The soils were sampled from Kamishihoro and Kawagoe, Japan. Kamishihoro soils, classified as Andosols (FAO/UNESCO), were sampled from farmers' fields (0-15 cm) located in Obihiro, Hokkaido, Japan ( $42^\circ 45'01.9''\text{N}$ ,  $143^\circ 08'20.4''\text{E}$ ). Soils from Kawagoe (Arakida soils), classified as Alluvial (FAO/UNESCO), were commercially sourced and had been sampled from the bottom layer of a paddy field in Kawagoe city, Saitama, Japan. Both soils were sieved using a 4 mm diameter mesh before incubation. Characterization of initial soil chemical and physical properties; pH, Phosphorous



oxide ( $P_2O_5$ ), Potassium oxide ( $K_2O$ ), Magnesium oxide ( $MgO$ ), Calcium oxide ( $CaO$ ), cation exchange capacity (CEC) and bulk density (BD) are as described in Table S5.1 and were measured using the following methods. Soil pH was measured with a glass electrode pH meter. The  $P_2O_5$ ,  $K_2O$  and  $MgO$  and  $CaO$  were analyzed using Flow Injection Analysis (FIA). CEC was measured according to the method by Scollenberger and Simon (1945), with an atomic absorption spectrophotometer. The soil BD was determined by comparing the change in soil moisture content before and oven drying to a constant weight .

### 5.3.2 Digestate separation to solid and liquid phase

Digestate was obtained from a biogas plant located in the Field Science Center for Northern Biosphere in Hokkaido University, Japan (N43.0781050, E141.334344). The digestate was produced from dairy cow excreta. A polymer flocculant was used to separate digestate into solid and liquid fractions. Analyzed chemical properties (pH,  $P_2O_5$ , nitrates-  $NO_3^-$ , ammonium-  $NH_4^+$ ) of digestate and its derived fertilizers are as described in Table S5.2.

### 5.3.3 Incubation setup

The experimental setup was a randomized design with 2 soils (Kamishihoro and Arakida) and 5 fertilizer treatments; Digestate (106 kg inorganic nitrogen [N]  $ha^{-1}$  and 27 kg  $P_2O_5$   $ha^{-1}$ ), ADS (200 g), ADL (200 g), chemical fertilizer-CF (120 kg inorganic-N  $ha^{-1}$  and 100 kg  $P_2O_5$   $ha^{-1}$ ) and no fertilizer control, replicated 3 times. Fertilizer application rates were based on recommended N standards for Japanese mustard spinach (Department of Agriculture, Hokkaido Government, 2015).

For the incubation, 1500 g of air-dry soil were placed in Wagner pots. Water filled pore space was maintained at 70 % using milli Q and BD established at  $0.7$  g  $cm^{-3}$  for Kamishihoro soils and  $1$  g  $cm^{-3}$  for Arakida soils. Fertilizers were added to the pots (Day 0) and Japanese mustard spinach planted 10 days after fertilizer application. Each fertilizer treatment had 3 replicates making up a total of 30 soil cores.

#### 5.3.4 Soil and plant sampling and analysis

For soil chemical and leachate analysis soils were sampled on day 1, 7, 16, 23, 30 and 37. To determine soil pH and inorganic N ( $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ), 5 g of soil was extracted with 2 M potassium chloride and shaken for 1 hour. Immediately after extraction, soil pH was determined using a pH meter (AS800; AS ONE Co., Osaka, Japan). The solution was then filtered with 1  $\mu\text{l}$  filter paper (Toyo Roshi Kaisha No. 5C filter paper; Tokyo) and colorimetric measurements for inorganic N determined using FIA (AQLA-700; Aqualab Co., Ltd., Tokyo, Japan).

To determine soil  $\text{P}_2\text{O}_5$ , 1 g of soil was extracted with 100 g mixture of ammonium sulfate and 0.5 M sulfuric acid. After shaking for 30 min, the solution was filtered with 1  $\mu\text{l}$  filter paper (Toyo Roshi Kaisha No. 5C filter paper; Tokyo) and  $\text{P}_2\text{O}_5$  colorimetric measurements determined using FIA (AQLA-700; Aqualab Co., Ltd., Tokyo, Japan).

Above ground biomass was collected on day 38. After oven drying at 65°C for 48 hours, carbon (C) and N measurements were analyzed using CN coder (Perkin Elmer 2400).

#### 5.3.5 Nitrogen mass balance analysis

The N mass balance was determined by analyzing the distribution of inorganic N from the soil and chemical fertilizers during the incubation period. For plants, N uptake was determined by multiplying plant dry weight with analyzed N content. For leachates, inorganic N loss was determined by multiplying collected leachate (ml) with analyzed leachate  $\text{NO}_3^-$ -N and  $\text{NH}_4^+$ -N content.

#### 5.3.6 $\text{N}_2\text{O}$ gas sampling and analysis

Gas samples were collected on day 7, 16, 23, 30 and 37. Each soil core were placed in a sealed plastic container (25 L) with a septum. At 0 and 60 min after sealing the container, 30 mL of headspace gas sample was collected through the septum using a gas-tight syringe. During the preliminary experiment, we confirmed that the  $\text{N}_2\text{O}$  gas concentration linearly increased in the plastic container during the first 60 min of incubation. Sampled gases were stored in pre vacuumed 20 mL glass vials. The  $\text{N}_2\text{O}$  was measured with a gas chromatograph (GC-2014, Shimadzu Co.,

Kyoto, Japan) and analyzed within 24 hours of sampling. The N<sub>2</sub>O flux was calculated by dividing the increased N<sub>2</sub>O concentration (from 0 to 60 min) by the incubation time (60 min).

### 5.3.7 Soil DNA extraction and microbial abundance

For microbial analysis, soils were sampled on day 1 and 37. Using 0.5 g soil, DNA was extracted with the NucleoSpin ® Soil (Takara Bio Inc, Shiga, Japan) according to the manufacturer's instructions. To reduce DNA attachment to the soils, 0.2 g of skim milk was added during extraction (Hoshino and Matsumoto, 2005). Extracted DNA was purified using Agencourt AM Pure XP Kit (Beckman Coulter, Fullerton, CA, USA) according to the manufacturer's protocol and thereafter stored at -30°C until further analysis.

To determine microbial abundance, the qPCR (Quantitative Polymerase Chain Reaction) was performed using CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Inc, Hercules, USA). DNA samples were 50-fold diluted before qPCR targeting 16S rRNA. Each sample contained 10.4 µl of KAPA SYBR Fast qPCR kit (Agilent Technologies, Palo Alto, CA, USA), 0.8 µl of forward (515F-5'-GTGCCAGCMGCCGCGGTAA-3') and reverse (806R-5'-GGACTACHVGGGTWTCTAAT-3') primers, 7.0 µl of nuclease free water and 1 µl of DNA template, making up a final volume of 20 µl. The qPCR was conducted under the following conditions: 95°C for 30 sec and thereafter 30 cycles of 95°C for 30 sec, 57°C for 30 sec with a final extension of 72°C for 60 sec.

### 5.3.8 Statistical analysis

All statistical analysis was conducted in R software version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria). To obtain a normal distribution, N<sub>2</sub>O emissions data was log transformed (log 10 (flux+1)) before analysis. In all analysis,  $P < 0.05$  were considered significant. Two-way analysis of variance (ANOVA) with multiple pairwise comparisons were conducted to determine the effect of soil type and fertilizer on soil and plant chemical properties, N<sub>2</sub>O emissions and microbial abundance. For significant interactions, Tukey's HSD test was performed.

## 5.4 Results

### 5.4.1 Plant Analysis

Regarding above ground biomass, Kamishihoro soil had significantly ( $P < 0.001$ ) higher values compared to Arakida soil (Figure 5.1). Fertilizer had a significant ( $P < 0.001$ ) effect on above ground biomass with highest values observed in chemical fertilizer in both Kamishihoro and Arakida soil. Below ground biomass was not significantly affected by soil type and fertilizer (data not shown).




Figure 5.1: The effect of fertilizer on dry weight of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation (n=3). Abbreviations; S-Soil, F-Fertilizer.

For above ground plant biomass C, there was an interaction between fertilizer and soil type ( $P < 0.05$ ). The highest above ground plant biomass C content was recorded in Arakida control treatments (Figure S5.1). Below ground plant biomass C content was not significantly affected by soil type and fertilizer (data not shown).

For plant biomass N, the effect of soil type and fertilizer had a significant ( $P < 0.001$ ) effect with highest values recorded in Kamishihoro soil and chemical fertilizer, respectively (Figure S5.2). Considering Kamishihoro and Arakida soil separately, fertilizer had a significant ( $P < 0.01$ ) effect

on above ground N content with highest values recorded in chemical fertilizer in both soils. Soil type had a significant effect on below plant biomass N content with highest values observed in Kamishihoro soil (Data not shown).

#### 5.4.2 Soil Analysis

Regarding soil  $\text{NO}_3^-$  content Kamishihoro soil had significantly ( $P < 0.001$ ) higher values compared to Arakida soil (Figure 5.2). Considering Kamishihoro and Arakida soil separately, fertilizer had a significant ( $P < 0.05$ ) effect on soil  $\text{NO}_3^-$  content with highest values recorded in solid (Kamishihoro) and digestate (Arakida). Soil  $\text{NO}_3^-$  content was more responsive to fertilizer in Kamishihoro soil compared to Arakida soil (Figure 5.2).




Figure 5.2: : Effect of fertilizer on temporal variations in  $\text{NO}_3^-$ -N concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer.

Considering soil  $\text{NH}_4^+$  content, there was an interaction between soil types and fertilizer (Figure 5.3). The highest soil  $\text{NH}_4^+$  content was recorded in Kamishihoro soil in chemical fertilizer. Considering Kamishihoro and Arakida soil separately, fertilizer had a significant ( $P < 0.01$ ) effect on soil  $\text{NH}_4^+$  content. In Kamishihoro significantly higher values were observed on in liquid and chemical fertilizer while in Arakida significantly highest values were observed in chemical

fertilizer. Soil  $\text{NH}_4^+$  content was more responsive to fertilizer in Kamishihoro soil compared to Arakida soil (Figure 5.3).




Figure 5.3: : Effect of fertilizer on temporal variations in  $\text{NH}_4^+$ -N concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; D-Day, F-Fertilizer.

For soil  $\text{P}_2\text{O}_5$  content, there was an interaction between soil types and fertilizer ( $P < 0.05$ ). The highest soil  $\text{P}_2\text{O}_5$  content was recorded in Arakida soil in chemical fertilizer (Figure 5.4). Arakida soil had significantly ( $P < 0.05$ ) higher soil  $\text{P}_2\text{O}_5$  content compared to Kamishihoro soil. Considering Kamishihoro and Arakida soil separately, fertilizer had a significant ( $P < 0.05$ ) effect on soil  $\text{P}_2\text{O}_5$  content in both soils, with highest values recorded in chemical fertilizer. Soil  $\text{P}_2\text{O}_5$  content was more responsive to fertilizer in Arakida soil compared to Kamishihoro soil (Figure 5.4).




Figure 5.4: Effect of fertilizer on temporal variations in  $P_2O_5$  concentration in the Kamishihoro and Arakida soil. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; D-Day, F-Fertilizer.

For soil pH, there was an interaction between soil types and fertilizer ( $P < 0.05$ ). The highest soil pH content was recorded in Arakida soil in liquid fertilizer (Figure 5.5). Considering Kamishihoro and Arakida separately, liquid fertilizer recorded significantly ( $P < 0.05$ ) higher soil pH values in both soils, while chemical fertilizer had lowest soil pH values. Soil pH was more responsive to fertilizer in Arakida soil compared to Kamishihoro soil (Figure 5.5).




Figure 5.5: Effect of fertilizer on temporal variations of soil pH in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation (n=3). Abbreviations; D-Day, F-Fertilizer.

### 5.4.3 Leachate Analysis

Regarding leachate  $\text{NO}_3^-$  content, Kamishihoro soil had significantly ( $P < 0.05$ ) higher values compared to Arakida soil. Chemical fertilizer had significantly ( $P < 0.05$ ) higher soil leachate  $\text{NO}_3^-$  content in both Kamishihoro and Arakida soil (Figure 5.6).






Figure 5.6: Effect of fertilizer on temporal variations of leachate  $\text{NO}_3^-$  content in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; D-Day, F-Fertilizer.

**For** the leachate  $\text{NH}_4^+$  content, there was an interaction between soil types and fertilizer ( $P < 0.05$ ). The highest leachate  $\text{NH}_4^+$  content was recorded in Arakida soil in chemical fertilizers (Figure 5.7). Considering Kamishihoro soil separately, fertilizer did not have a significant effect on leachate  $\text{NH}_4^+$  content. However, in Arakida soil, chemical fertilizer had significantly higher leachate  $\text{NH}_4^+$  content compared to other fertilizers.




Figure 5.7: Effect of fertilizer on temporal variations of  $\text{NH}_4^+$  leachate content in Kamishihoro and Arakida soil. Vertical bars indicate standard deviation ( $n=3$ ). Abbreviations; D-Day, F-Fertilizer.

#### 5.4.4 Nitrogen mass balance

Overall, Kamishihoro soil had higher ( $234.32 \text{ mg N kg}^{-1}$ ) inorganic N content compared to (Arakida  $63.72 \text{ mg N kg}^{-1}$ ). Considering fertilizer treatments, digestate had highest inorganic N content ( $164.50 \text{ mg N kg}^{-1}$  Kamishihoro soil and  $147.50 \text{ mg N kg}^{-1}$  Arakida soil), followed by liquid ( $119.19 \text{ mg N kg}^{-1}$  Kamishihoro soil and  $106.87 \text{ mg N kg}^{-1}$  Arakida soil) and solid fertilizer ( $91.20 \text{ mg N kg}^{-1}$  Kamishihoro soil and  $81.78 \text{ mg N kg}^{-1}$  Arakida soil).

In Kamishihoro soil, solid and liquid fertilizer had higher above ground percentage distribution compared to digestate. In solid and digestate fertilizer, about 30 % of the added N remained in the soil while 25 % was lost in other ways other than leaching (Figure 5.8). In liquid fertilizer, 12 % of added N remained in the soil while 44 % was lost in other ways other than leaching. In Arakida soil, liquid fertilizer treatments had higher N accumulated in leachates and aboveground plant biomass compared to digestate and solid fertilizer (Figure 8). About 28 %, 10%, and 3% of N remained in the soil under the digestate, solid, and liquid fertilizer treatments, respectively. Additionally, more than 50 % of N applied to the solid and liquid treatments were lost in other forms other than leaching.




Figure 5.8: Effect of fertilizer on nitrogen mass balance in percentage in the Kamishihoro and Arakida soil.

#### 5.4.5 Gene Abundances

Arakida soil had significantly ( $P < 0.001$ ) higher 16S rRNA gene copy numbers compared to Kamishihoro soil. In Kamishihoro soil, fertilizer did not have a significant effect on 16S rRNA gene copy numbers. However, in Arakida soil, fertilizer had a significant ( $P < 0.001$ ) effect on 16S rRNA gene copy numbers with the highest values observed in solid fertilizer.

#### 5.4.6 Nitrous oxide emissions

Regarding soil  $N_2O$  emissions, there was an interaction between soil type and fertilizer ( $P < 0.05$ ). The highest soil  $N_2O$  emissions were recorded in Kamishihoro soil in solid treatments (Figure 5.9). Fertilizer had a significant ( $P < 0.05$ ) effect on soil  $N_2O$  emissions in Kamishihoro and Arakida soils. In both Kamishihoro and Arakida soils, digestate, solid and chemical fertilizer had significantly higher soil  $N_2O$  emissions compared to other fertilizers (Figure 5.9).




Figure 5.9: Temporal changes in soil  $N_2O$  emissions in Kamishihoro and Arakida fertilizer treatments. Error bars show mean standard deviation mean  $\pm$  SD (n=3).

## 5.5 Discussion

### 5.5.1 The effect of separated digestate on soil inorganic nitrogen content

Digestate and its derived fertilizers influenced the soil inorganic nitrogen content (Figure 5.2 and 5.3). However, the effect of application varied based on soil types. For  $NO_3^-$  in Kamishihoro solid had the higher values while in Arakida digestate had the highest values. Solid having the highest values of  $NO_3^-$  in Kamishihoro is expected as results from the separated digestate chemical analysis showed that solid had higher  $NO_3^-$  content compared to digestate and liquid (Table S5.2). Möller and Müller (2012) similarly reported that the solid fraction of digestate contained a high amount of available nitrogen. These results contrast with those of various researchers (Drosg et al., 2015; Tambone et al., 2017) who have reported that after separation of digestate, the liquid portion contains high inorganic N content. The differences in results may be attributed to methods used to separate the digestate. Barampouti et al. (2020) reported that the different methods of digestate separation such as mechanical technologies, polymers and flocculants had different effects on the nutrient distribution in the solid and liquid fractions.

Considering soil  $NH_4^+$  content, in Arakida chemical fertilizer had highest values (Figure 5.3). However, in Kamishihoro soils highest values were recorded in both liquid and chemical fertilizers (Figure 5.3). Chemical nitrogen fertilizers having high  $NH_4^+$  content is expected as they directly increase the soil  $NH_4^+$  content (Giday, 2019). Regarding liquid digestate  $NH_4^+$  content, Sigurnjak

(2017) conducted research on nitrogen release and mineralization potential of various nitrogen fertilizers. Based on their results, liquid fraction of digestate had a high  $\text{NH}_4^+$  content in agreement with our study. Subsequently it can be used as a substitute for inorganic nitrogen fertilizer (Sigurnjak, 2017). Tambone et al.(2017) and Möller and Müller (2012) similarly reported that due to the high  $\text{NH}_4^+$  content (about 45–80% of N in the liquid fraction of digestate) the liquid part of digestate was a better fertilizer quality than digestate and could be used as a fast-acting nitrogen fertilizer.

Considering the soil  $\text{P}_2\text{O}_5$  content, digestate derived fertilizers had an influence in Arakida soils (Figure 5.4). Although the values were not significantly different, solid had the highest values compared to digestate and liquid (Figure 5.4). Egene et al.(2020) and Möller and Müller (2012) reported that the solid fraction of digestate had a high phosphorous content and could subsequently be used as a phosphorous fertilizer. Bachman et al.(2015) conducted research on phosphorous distribution and availability in untreated and separated biogas digestate. Based on their results the solid fraction had higher phosphorous concentration than the liquid fraction in agreement with our study.

Digestate and derived fertilizers had a significant influence on soil pH values (Figure 5.5). In both Kamishihoro and Arakida liquid, solid and digestate had the highest soil pH. These results may have been due to the high soil pH of applied digestate and its derived fertilizers (Table S5.2). Previous research has found that application of digestate and its derived fertilizers increased soil pH content of agricultural soils in line with our findings (Galvez et al., 2012, Valentinuzzi et al., 2020). Chemical fertilizer had the lowest soil pH in both Kamishihoro and Arakida soils. Sun et al.(2020) and Gouling (2016) similarly reported that the use of chemical fertilizers reduced soil pH levels and was one of the main reasons for agricultural soil acidification. Chemical fertilizer use cause soil acidity in agricultural soils due to enhancing the oxidation of ammonium ( $\text{NH}_4^+$ ) to nitrate ( $\text{NO}_3^-$ ) which produces hydrogen ions ( $\text{H}^+$ ) (Garvin and Carver, 2003). Our results show that digestate and its derived fertilizers can be used to reduce soil acidity in agricultural soils.

### 5.5.2 Influence of soil types

Inorganic N content was responsive to fertilizer additions in Kamishihoro soils compared to Arakida soils (Figure 5.2 and 5.3). However, for soil P<sub>2</sub>O<sub>5</sub> content and soil pH, Arakida soils were more responsive to fertilizer additions. Ohshiro et al.(2016) conducted research on the effects of soil types and fertilizers on growth, yield, and quality of edible *Amaranthus tricolor* lines in Japan. Based on their results, nutrient availability varied in different soil types based on differences in soil physical, chemical and biological properties, which may have been the case in our study. Lapwayi et al.(2012) similarly reported that the response of soil properties to fertilizers was dependent on soil types. These results indicate that soil type should be taken into consideration when determining agricultural fertilizer application rates. For example, the high clay minerals (allophane, imogolite) present in Kamishihoro soils have a high phosphorous retention capacity which may have reduced the increase in phosphorous content due to fertilizer application (Nyanzo, 2002). Additionally, the low difference in soil pH in Kamishihoro soils due to fertilizer additions may have been caused by the high soil pH buffering capacity of the soils. Shoji and Takahashi (2002) found that Andosols have high organic matter and cation exchange capacity which increase the soil pH buffering capacity in agreement with our study.

### 5.5.3 Leachate analysis

Interestingly digestate and its derived fertilizers did not influence the inorganic N content of leachates (Figure 5.6 and 5.7) . This may have been due to the soil types used in the experiment. The Kamishihoro soils had a high organic carbon content while Arakida soils had high clay content, which both reduce leaching losses. Kanthle et al. (2016) similarly reported that soils with high organic carbon content reduced leaching losses. Cameron and Moir (2013) found that soils with high clay content had reduced leaching losses in agreement with our study. Chemical fertilizers had the highest inorganic N leachate losses compared to digestate and its derived fertilizers in Kamishihoro and Arakida soils. Wang et al.(2019) conducted research on the effect of fertilizer types on soil N leaching losses. Based on their results, chemical fertilizer treatments had the highest N leaching losses compared to organic fertilizers in agreement with our study. These results show that on soils with high clay and organic matter content, digestate and its derived fertilizers can be used without affecting inorganic N leaching losses.

#### 5.5.4 Nitrogen mass balance

Considering digestate and its derived fertilizers, liquid had the highest N loss compared to digestate and solid fertilizer treatments (Figure 5.8). This may have been due to increased N losses through  $\text{NH}_3$  volatilization in liquid fertilizers. To prevent these losses the liquid part of digestate is better suited as a fast acting fertilizer (Tambone et al., 2017, Nakamura et al., 2007). Nakamura et al.(2012) also reported that incorporating digestate into the soil immediately after application could reduce  $\text{NH}_3$  volatilization losses. The solid digestate fertilizer treatments stored more loaded nitrogen into the soil compared to liquid, showing that it would be appropriate as a slow acting fertilizer. Nakamura et al.(2007) similarly reported that solid digestate contained N in organic form and would be best suited as a slow release fertilizer.

#### 5.5.5 16S rRNA gene abundance

The effect of digestate and its derived fertilizers on 16S rRNA gene copy numbers varied based on soil type. In Kamishihoro the effect of digestate and its derived fertilizers was not significant. However, in Arakida, solid had significantly higher 16S rRNA gene copy numbers compared to digestate and liquid. This may have been due to the differences in soil pH and chemical properties of the digestate fertilizers after separation. Specifically, solid digestate had the highest soil pH (8.72),  $\text{NO}_3^-$  (178.59 mg N  $\text{kg}^{-1}$ ) and  $\text{P}_2\text{O}_5$  (1462.47 mg  $\text{kg}^{-1}$ ) that may have enhanced the abundance of the 16S rRNA gene. Zhou et al.(2015) conducted research on the influence of fertilizers on soil bacterial communities in agricultural soils. Their results showed that soil  $\text{NO}_3^-$  content was one of the most important factors in shaping soil bacterial communities with higher abundance observed in higher  $\text{NO}_3^-$  concentrations. Regarding  $\text{P}_2\text{O}_5$  our results contrast with those of Wang et al.(2018) who reported that soil phosphorous content did not have a significant effect on 16S rRNA gene copy numbers in agricultural soils. The variations in results may have been due to differences in soil types and experimental time periods. Wang et al.(2018) conducted their research on Ultisols for 35 years while our experiment was conducted on Andosol for 37 days.

#### 5.5.6 $\text{N}_2\text{O}$ emissions

Kamishihoro soils had higher  $\text{N}_2\text{O}$  emissions compared to Arakida soils (Figure 5.9). This can be related to Kamishihoro soils having higher  $\text{NH}_4^+$  content compared to Arakida soils, which

stimulated nitrification and N<sub>2</sub>O emissions. N<sub>2</sub>O emissions peaked during the first week of the experiment. This can be attributed to nitrification of the added NH<sub>4</sub><sup>+</sup> from fertilizer additions, which were highest during the first week of the experiment. Similar trends in N<sub>2</sub>O emissions were reported by De La Fuente et al.(2013) and Albuquerque et al.(2012). Digestate and its derived fertilizers influenced soil N<sub>2</sub>O emissions with digestate and solid having the highest soil emissions compared to liquid. Askri et al.(2016) conducted research on the influence of origin and post treatment in greenhouse gas emissions after digestate application to soils. Based on their results, digestate had the highest N<sub>2</sub>O emissions while liquid had the lowest emissions in line with our results. They related this to increased availability of decomposable organic matter in digestate and solid which stimulated denitrification, which may have been the case in our study. These results show that solid liquid separation of digestate may be used to reduce greenhouse gas emissions from agricultural soils. Specifically, the liquid digestate can be used to decrease N<sub>2</sub>O emissions from digestate in agricultural soils.

#### 5.5.7 Plant Analysis

Digestate and its derived fertilizers did not influence the Japanese mustard spinach yields (Figure 5.1). This result shows that digestate and its derived fertilizers can be used interchangeably on Japanese mustard spinach without affecting crop yields. Chemical fertilizer had the highest yields in both Kamishihoro and Arakida soils. This can be attributed to chemical fertilizer N P K application being suited to Japanese mustard spinach demand based on the Hokkaido government recommendation for the crop. However, for digestate and its derived fertilizers, only N was taken into account for this incubation. Specifically, P for digestate and its derived fertilizers was below the recommended amounts, which may have reduced Japanese mustard spinach yields. Ehmann et al.(2018) conducted research on the fertilizing potential of separated digestate on annual and perennial biomass production systems. Based on their results, increased maize yields were recorded in chemical fertilizers compared to digestate and its derived fertilizers. They attributed this to chemical fertilizers providing nutrients based on crop requirements, which may have been the case in our study. These results indicate that plant yields can be enhanced in digestate and its derived fertilizers by supplementing with chemical fertilizers, based on plant requirements.



Unexpectedly control treatments had significantly higher above ground biomass C compared to other fertilizers in both Kamishihoro and Arakida soils. This may have been due to the inorganic N in control treatments being below plan requirements. Hermans et al.(2006) and Hirai et al. (2004) similarly reported that soil N deficiency caused the accumulation of carbohydrates such as sugars and starch in leaves, increasing above ground C content.

## 5.6 Conclusion

Digestate and its derived fertilizers influenced soil inorganic nitrogen content. However, the influence varied based on soil type. In highest  $\text{NO}_3^-$  was recorded in solid-Kamishihoro and digestate-Arakida. Considering  $\text{NH}_4^+$  highest values were recorded in liquid-Kamishihoro and chemical-Arakida. This indicates that soil types should be taken into consideration for soil inorganic nitrogen content when applying digestate and its derived fertilizers. Additionally, digestate and its derived fertilizers increased soil pH in Kamishihoro and Arakida soils, while chemical fertilizers decreased soil pH. Digestate and its derived fertilizers can therefore be used to reduce soil acidification in agricultural soils. Solid digestate enhanced the 16S rRNA in Arakida soils but not in Kamishihoro. Solid digestate can therefore be used to increase microbial abundance in alluvial agricultural soils. Considering  $\text{N}_2\text{O}$  emissions, liquid had the lowest emissions compared to digestate and solid fertilizer in both Kamishihoro and Arakida soils. This indicates that digestate separation can be used to reduce agricultural greenhouse  $\text{N}_2\text{O}$  emissions. Digestate and its derived fertilizers did not influence Japanese mustard spinach above ground biomass. However, chemical fertilizers had the highest above ground biomass of Japanese mustard spinach. This indicates that chemical fertilizers can be used to supplement digestate and its derived fertilizers to increase Japanese mustard yields. Digestate separation can therefore be used to modulate soil inorganic nitrogen content and  $\text{N}_2\text{O}$  emissions in agricultural soils. The study should contribute towards the development of efficient digestate management practices that can enhance soil nutrient content while reducing greenhouse gas ( $\text{N}_2\text{O}$ ) emissions.

## References

Alburquerque, J.A., De La Fuente, C., Ferrer-Costa, A., Carrasco, L., Cegarra, J., Abad, M., Bernal, M.P.: Assessment of the fertilizer potential of digestates from farm and agro industrial residues. *Biomass Bioenergy* 40, 181–189 (2012)

Arthurson, V. (2009). Closing the Global Energy and Nutrient Cycles through Application of Biogas Residue to Agricultural Land – Potential Benefits and Drawback. *Energies*, 2(2), 226–242. <https://doi.org/10.3390/en20200226>

Barampouti, E., Mai, S., Malamis, D., Moustakas, K., & Loizidou, M. (2020). Exploring technological alternatives of nutrient recovery from digestate as a secondary resource. *Renewable and Sustainable Energy Reviews*, 134, 110379. <https://doi.org/10.1016/j.rser.2020.110379>

Bauer, A., Mayr, H., Hopfner-Sixt, K., & Amon, T. (2009). Detailed monitoring of two biogas plants and mechanical solid–liquid separation of fermentation residues. *Journal of Biotechnology*, 142(1), 56–63. <https://doi.org/10.1016/j.jbiotec.2009.01.016>

Bouwman, A. F., Van Drecht, G., Knoop, J. M., Beusen, A. H. W., & Meinardi, C. R. (2005). Exploring changes in river nitrogen export to the world’s oceans. *Global Biogeochemical Cycles*, 19(1). <https://doi.org/10.1029/2004gb002314>

Camilleri-Rumbau, M. S., Briceño, K., Fjerbæk Søtoft, L., Christensen, K. V., Roda-Serrat, M. C., Errico, M., & Norddahl, B. (2021). Treatment of Manure and Digestate Liquid Fractions Using Membranes: Opportunities and Challenges. *International Journal of Environmental Research and Public Health*, 18(6), 3107. <https://doi.org/10.3390/ijerph18063107>

De La Fuente, C., Alburquerque, J.A., Clemente, R., Bernal, M.P.: Soil C and N mineralisation and agricultural value of the products of an anaerobic digestion system. *Biol. Fertil. Soils* 49, 313–322 (2013)

Drosg, B., Fuchs, W., Al Seadi, T., Madsen, M., & Linke, B. (2015). *Nutrient recovery by biogas digestate processing* (Vol. 2015). IEA Bioenergy Dublin. [https://www.ieabioenergy.com/wp-content/uploads/2015/08/NUTRIENT\\_RECOVERY\\_RZ\\_web2.pdf](https://www.ieabioenergy.com/wp-content/uploads/2015/08/NUTRIENT_RECOVERY_RZ_web2.pdf)

Egene, C. E., Sigurnjak, I., Regelink, I. C., Schoumans, O. F., Adani, F., Michels, E., . . . Meers, E. (2020). Solid fraction of separated digestate as soil improver: implications for soil fertility and carbon sequestration. *Journal of Soils and Sediments*, 21(2), 678–688.

<https://doi.org/10.1007/s11368-020-02792-z>

Garvin, D.F., and Carver, B.F. (2003). Role of the genotype in tolerance to acidity and aluminum toxicity. p. 387–406. In Z. Rengel (ed.) *Handbook of soil acidity*. New York :Marcel Dekker

Gielnik, A., Pechaud, Y., Huguenot, D., Cébron, A., Esposito, G., & van Hullebusch, E. D. (2019). Bacterial seeding potential of digestate in bioremediation of diesel contaminated soil. *International Biodeterioration & Biodegradation*, 143, 104715.

<https://doi.org/10.1016/j.ibiod.2019.06.003>

Greenberg, I., Kaiser, M., Gunina, A., Ledesma, P., Polifka, S., Wiedner, K., . . . Ludwig, B. (2019). Substitution of mineral fertilizers with biogas digestate plus biochar increases physically stabilized soil carbon but not crop biomass in a field trial. *Science of The Total Environment*, 680, 181–189. <https://doi.org/10.1016/j.scitotenv.2019.05.051>

Guo, J. H., Liu, X. J., Zhang, Y., Shen, J. L., Han, W. X., Zhang, W. F., . . . Zhang, F. S. (2010). Significant Acidification in Major Chinese Croplands. *Science*, 327(5968), 1008–1010.

<https://doi.org/10.1126/science.1182570>

Hermans, C., Hammond, J. P., White, P. J., & Verbruggen, N. (2006). How do plants respond to nutrient shortage by biomass allocation? *Trends in Plant Science*, 11(12), 610–617.

<https://doi.org/10.1016/j.tplants.2006.10.007>

Hirai, M. Y., Yano, M., Goodenowe, D. B., Kanaya, S., Kimura, T., Awazuhara, M., . . . Saito, K. (2004). From The Cover: Integration of transcriptomics and metabolomics for understanding of global responses to nutritional stresses in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences*, 101(27), 10205–10210. <https://doi.org/10.1073/pnas.0403218101>

Holm-Nielsen, J., Al Seadi, T., & Oleskowicz-Popiel, P. (2009). The future of anaerobic digestion and biogas utilization. *Bioresource Technology*, 100(22), 5478–5484.

<https://doi.org/10.1016/j.biortech.2008.12.046>

- Lupwayi, N. Z., Lafond, G. P., Ziadi, N., & Grant, C. A. (2012). Soil microbial response to nitrogen fertilizer and tillage in barley and corn. *Soil and Tillage Research*, 118, 139–146. <https://doi.org/10.1016/j.still.2011.11.006>
- Meixner, K., Fuchs, W., Valkova, T., Svardal, K., Loderer, C., Neureiter, M., . . . Drogg, B. (2015). Effect of precipitating agents on centrifugation and ultrafiltration performance of thin stillage digestate. *Separation and Purification Technology*, 145, 154–160. <https://doi.org/10.1016/j.seppur.2015.03.003>
- Möller, K. (2015). Effects of anaerobic digestion on soil carbon and nitrogen turnover, N emissions, and soil biological activity. A review. *Agronomy for Sustainable Development*, 35(3), 1021–1041. <https://doi.org/10.1007/s13593-015-0284-3>
- Nakamura, M., Fujikawa, T., Yuyama, Y., Yamaoka, M., & Oritate, F. (2012). Influence of application method on emissions of ammonia and nitrous oxide following application of methane fermentation-digested slurry [in Japanese]. *Japanese Journal of Soil Science and Plant Nutrition*, 83(2), 139–146. [https://doi.org/10.20710/dojo.83.2\\_139](https://doi.org/10.20710/dojo.83.2_139)
- Nakamura, M., Yuyama, Y., Yamaoka, M., & Fujikawa, T. (2007). Material Balance in Biogas Plant and Fertilizer Characteristics of Digested Liquid and Digested Liquid Supernatant [in Japanese]. *Transactions of The Japanese Society of Irrigation, Drainage and Reclamation Engineering*, 75(3), 337–343. <https://doi.org/10.11408/jsidre1965.2007.337>
- Nowostawska, U., Sander, S. G., McGrath, K. M., & Hunter, K. A. (2005). Effect of trivalent metal sulfates on the coagulation and particle interactions of alumina colloids. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 266(1–3), 200–206. <https://doi.org/10.1016/j.colsurfa.2005.06.041>
- Nanzyo, M. (2002). Unique properties of volcanic ash soils. *Global Environmental Research*, 6(2), 99–112.
- Ohshiro, M., Hossain, M. A., Nakamura, I., Akamine, H., Tamaki, M., Bhowmik, P. C., & Nose, A. (2016). Effects of soil types and fertilizers on growth, yield, and quality of edible *Amaranthus tricolor* lines in Okinawa, Japan. *Plant Production Science*, 19(1), 61–72. <https://doi.org/10.1080/1343943x.2015.1128087>

- Risberg, K., Cederlund, H., Pell, M., Arthurson, V., & Schnürer, A. (2017). Comparative characterization of digestate versus pig slurry and cow manure – Chemical composition and effects on soil microbial activity. *Waste Management*, 61, 529–538.  
<https://doi.org/10.1016/j.wasman.2016.12.016>
- Sigurnjak, I., De Waele, J., Michels, E., Tack, F. M. G., Meers, E., & De Neve, S. (2017). Nitrogen release and mineralization potential of derivatives from nutrient recovery processes as substitutes for fossil fuel-based nitrogen fertilizers. *Soil Use and Management*, 33(3), 437–446.  
<https://doi.org/10.1111/sum.12366>
- Šimon, T., Kunzová, E., & Friedlová, M. (2016a). The effect of digestate, cattle slurry and mineral fertilization on the winter wheat yield and soil quality parameters. *Plant, Soil and Environment*, 61(No. 11), 522–527. <https://doi.org/10.17221/530/2015-pse>
- Šimon, T., Kunzová, E., & Friedlová, M. (2016b). The effect of digestate, cattle slurry and mineral fertilization on the winter wheat yield and soil quality parameters. *Plant, Soil and Environment*, 61(No. 11), 522–527. <https://doi.org/10.17221/530/2015-pse>
- Smith, J., Abegaz, A., Matthews, R. B., Subedi, M., Orskov, E. R., Tumwesige, V., & Smith, P. (2014). What is the potential for biogas digesters to improve soil carbon sequestration in Sub-Saharan Africa? Comparison with other uses of organic residues. *Biomass and Bioenergy*, 70, 73–86. <https://doi.org/10.1016/j.biombioe.2014.01.056>
- Tambone, F., Orzi, V., D'Imporzano, G., & Adani, F. (2017). Solid and liquid fractionation of digestate: Mass balance, chemical characterization, and agronomic and environmental value. *Bioresource Technology*, 243, 1251–1256. <https://doi.org/10.1016/j.biortech.2017.07.130>
- Tampio, E., Ervasti, S., & Rintala, J. (2015). Characteristics and agronomic usability of digestates from laboratory digesters treating food waste and autoclaved food waste. *Journal of Cleaner Production*, 94, 86–92. <https://doi.org/10.1016/j.jclepro.2015.01.086>
- Wang, Q., Wang, C., Yu, W., Turak, A., Chen, D., Huang, Y., . . . Huang, Z. (2018). Effects of Nitrogen and Phosphorus Inputs on Soil Bacterial Abundance, Diversity, and Community Composition in Chinese Fir Plantations. *Frontiers in Microbiology*, 9. Published.  
<https://doi.org/10.3389/fmicb.2018.01543>

Weiland, P. (2009). Biogas production: current state and perspectives. *Applied Microbiology and Biotechnology*, 85(4), 849–860. <https://doi.org/10.1007/s00253-009-2246-7>

Zhou, J., Guan, D., Zhou, B., Zhao, B., Ma, M., Qin, J., . . . Li, J. (2015). Influence of 34-years of fertilization on bacterial communities in an intensively cultivated black soil in northeast China. *Soil Biology and Biochemistry*, 90, 42–51. <https://doi.org/10.1016/j.soilbio.2015.07.005>

Supplementary

Table S5.1: Physical and chemical properties of Kamishihoro and Arakida soils (means  $\pm$  standard deviation, n=3).

Parameter	Kamishihoro	Arakida soil
pH (H <sub>2</sub> O)	5.5	6.8 $\pm$ 0.1
P <sub>2</sub> O <sub>5</sub> (mg kg <sup>-1</sup> )	135	178 $\pm$ 2
K <sub>2</sub> O (mg kg <sup>-1</sup> )	253	123 $\pm$ 1
MgO (mg kg <sup>-1</sup> )	511	824 $\pm$ 16
CaO (mg kg <sup>-1</sup> )	2523	2582 $\pm$ 31
Lime saturation (%)	33.7	59.3 $\pm$ 1.3
Base saturation (%)	46.3	87.3 $\pm$ 1.2
Phosphate absorption coefficient	1384	893 $\pm$ 10
Cation exchange capacity (me kg <sup>-1</sup> )	281	155 $\pm$ 2
Bulk density (g cm <sup>-3</sup> )	0.78	1.10 $\pm$ 0.01

Table S5.2: Chemical properties of anaerobic digestate and separated solid and liquid fractions.

Parameter	Digestate	Solid fraction	Liquid fraction
Water content (%)	95.08 ± 0.22	93.09 ± 1.92	-
pH	8.27	8.72	8.42
NO <sub>3</sub> <sup>-</sup> -N (mg N kg <sup>-1</sup> )	1.92	178.59	2.31
NH <sub>4</sub> <sup>+</sup> -N (mg N kg <sup>-1</sup> )	2125.49	1002.08	1085.72
P <sub>2</sub> O <sub>5</sub> (mg kg <sup>-1</sup> )	544.65	1462.47	103.51
Mass distribution (%)	-	39.57	60.43






Figure S5.1: The effect of fertilizer on carbon content of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation (n=3). Abbreviations; S-Soil, F-Fertilizer.




Figure S5..2: The effect of fertilizer on nitrogen content of aboveground plant biomass in Kamishihoro and Arakida. Vertical bars indicate standard deviation (n=3). Abbreviations; S-Soil, F-Fertilizer.

## Chapter 6 Synthesis and Recommendations for further research

This chapter synthesizes the research and identifies emerging opportunities for advancement of further research

### 6.1 Overall summary and future research recommendations

#### 6.1.1 Land use and season drive changes in soil microbial communities and related functions in agricultural soils

The study involved sampling of soils from farmers' fields in the Kamishihoro, Japan. The objective was to determine the effect of land use (cropland, grassland), season (spring, summer) and fertilizer (anaerobic digestate) on soil microorganisms and related functions. After sampling DNA extracted and sequenced targeting 16S rRNA region. Land use and seasons were identified as the main factors driving changes in soil microbial communities in the Kamishihoro region. Specifically, land use drove changes in alpha (evenness) and beta diversity (unifrac) indices with higher values observed in cropland compared to grassland. However, grasslands had a higher number of unique operational taxonomic units (OTUs) (10303) compared to cropland (5112). Based on these results, land use influenced different aspects of the soil microbial diversity indices. The influence of seasons on the soil microbial community varied based on land use. In cropland, season influenced microbial alpha (Shannon, evenness, OTU counts) and beta (unifrac) diversity. In grassland however season influenced beta (unifrac) diversity only. Thus, conventional tillage practices in cropland could have a negative effect on microbial stability in agricultural soils. Considering predicted soil functions, nitrogenase (*nifH*) had significantly higher values in cropland-summer while nitrite reductase (*nirK*) and ammonia monooxygenase (*amoA*) were significantly higher in cropland-spring. In grassland however, *nifH* was significantly higher in grassland-spring while *nirK* and *amoA* were significantly higher in grassland-summer. These results indicated that the impact of seasons on soil microorganisms' distribution and abundance in cropland and grassland may directly affected soil functions.

Our study was conducted on one crop (beet) although research shows that various crops influence the soil microbial community and related functions in different ways (Yang et al., 2020). Future research should therefore be conducted in Kamishihoro region involving different crops to clarify the influence of crop variations on the soil microbial community. Additionally, we conducted soil

sampling for 2 seasons (summer and spring), although research shows that all 4 seasons have an impact on the soil microbial community (Li et al., 2019; Sengupta et al., 2020). Therefore, future experiments should conduct this research on a continuous basis (with all seasons) for an extended time period to compare the long-term effects of seasonality on soil microbes in Kamishihoro region.

#### 6.1.2 Liming improves the stability of soil microbial community structures against the application of digestate made from dairy wastes

The incubation experiment was conducted for a month to determine the effect of lime application (pH = 6.5 and 5.5 for the soils with and without lime, respectively) and fertilizer (digestate, urea and control) on the soil microbial community structures, stability and gene functions. Soils were sampled for pH and microbial analysis every seven days after fertilizer application. Microbial analysis DNA was extracted and sequenced targeting the 16S rRNA region. To determine gene abundance for 16S rRNA, ammonia oxidizing archaea (AOA), ammonia oxidizing bacteria (AOB), nitrous oxide reductase (*nosZ*) and nitrite reductase (*nirS*), quantitative PCR was conducted. Based on the result, soil pH influenced the relative abundance of soil microbes. For example, *Actinobacteria* was influenced more strongly by digestate in lime soils, while *Alphaproteobacteria* was influenced more strongly by digestate in the no lime soil. Lime increased the stability of operational taxonomic units against the application of digestate. However, this was not the case for urea, where treatments without lime were more stable. The combination of lime and digestate fertilizer increased the 16S rRNA gene copy numbers in Kamishihoro soils. Considering nitrification and denitrification genes, higher values of AOA, *nosZ* and *nirS* gene copy numbers were observed in lime treatments compared to those without lime. Additionally, digestate increased AOB gene copy numbers but that was not the case for AOA. Based on the results soil pH and fertilizers play significant roles towards the management of microbial stability and functional gene abundance in agricultural soils.

Our study was conducted without plants for one month, although research shows that plants such as legumes increase soil acidity (Monaghan et al., 1998; Kiiya et al., 2010). Future research should include crops to clarify the effect of lime on stability of soil microbes in a cultivated agricultural environment. Additionally, we focused on the effect of soil pH on the nitrogen cycle. However,

soil pH may influence other nutrient cycles directly and indirectly. Malik et al.(2018) reported that soil pH influenced carbon accumulation in agricultural soils, decomposition rates among other processes. Therefore, future research should include analysis of different nutrient cycles and the potential effects of soil pH on their related functions.

### 6.1.3 The effect of solid-liquid separation of digestate on soil nutrient dynamics and Japanese mustard spinach (*Brassica rapa* var. *perviridis*) yields in different soils

The incubation experiment was conducted for 37 days to determine the effect of soil types (Kamishihoro and Arakida) and fertilizers (digestate, solid digestate, liquid digestate, chemical and control) on soil and plant (Japanese mustard spinach) nutrient dynamics, microbial abundance and crop yields. In the results, soil type influenced the above ground biomass and nitrogen content with higher values recorded in Kamishihoro compared to Arakida. A similar trend was observed in soil inorganic nitrogen analysis with higher values observed in Kamishihoro. This indicates that soil types should be taken into consideration when using digestate and its derived fertilizers within the agricultural environment. The effect of separated digestate on soil inorganic nitrogen varied based on soil types as well. Specifically, fertilizer had a significant effect on soil  $\text{NO}_3^-$  (solid-Kamishihoro, digestate-Arakida) and  $\text{NH}_4^+$  (liquid-Kamishihoro, chemical-Arakida) content. Therefore, digestate and its derived fertilizers can be used to regulate soil inorganic nitrogen content in agricultural soils. Soil pH was increased by digestate and its derived fertilizers showing their potential to reduce acidification in agricultural soils. Liquid had the lowest  $\text{N}_2\text{O}$  emissions compared to solid and digestate. This indicates that separating digestate can reduce agricultural emissions of the greenhouse  $\text{N}_2\text{O}$  gas. Therefore, the separation of digestate into solid and liquid fractions may be a reliable method to reduce agricultural soil acidification, regulate nutrient dynamics and reduce greenhouse ( $\text{N}_2\text{O}$ ) gas emissions from agricultural soils.

In our study we used a flocculant to separate the digestate. However, there are different methods that are used to separate digestate into solid and liquid fractions. Some of these include mechanical technologies and polymers, among others. Barampouti et al. (2020) reported that the method used to separate digestate effected nutrient distribution in the solid and liquid fractions. Future research should therefore include different methods of separation to determine the potential effect this would have on the use of digestate derived fertilizers on soil and plant nutrient dynamics in

agricultural soils. Additionally, based on our experimental results, digestate and its derived fertilizers had low above ground biomass production compared to chemical fertilizers. A possible solution to this problem would be using chemical fertilizers to supplement nutrient deficiencies in digestate and its derived fertilizers. Therefore, future research should be conducted to clarify whether this is a viable method to improve the yield production of digestate and its derived fertilizers.

## 6.2 General Comments

- In chapter 3 we considered bulk soil samples for our analysis. Research has shown that the rhizosphere environment is very different from the bulk soil which leads to the establishment of distinct microbial communities. Including the rhizosphere soils could have provided a different perspective on the effect of imposed treatments on soil microbes.
- In chapter 4 stabilizing the soil pH at 6.5 took about 3 months. It is therefore important to take this adjustment time into consideration when conducting pH adjustment experiments in andosols as they have a high pH buffering capacity. We also considered one soil type (andosol) for our experiment. The pH adjustment period is expected to change based on different soils buffering capacity and should be taken into consideration for lime based experiments.
- In chapter 5 we considered only the plant required nitrogen recommendations for digestate and its derived fertilizers, due to the challenge associated with the use of organic fertilizers for precision nutrient application in agricultural soils. Resultantly, chemical fertilizer, where we considered plant required nitrogen, phosphorous and potassium requirements, had the highest plant yields compared to digestate and its derived fertilizers. Therefore, to improve yields it would be more practical to try and supplement digestate and its derived fertilizers with the appropriate chemical fertilizers.