



Title	Relationships between mite allergen levels, mold concentrations, and sick building syndrome symptoms in newly built dwellings in Japan
Author(s)	Saijo, Y.; Kanazawa, A.; Araki, A.; Morimoto, K.; Nakayama, K.; Takigawa, T.; Tanaka, M.; Shibata, E.; Yoshimura, T.; Chikara, H.; Kishi, R.
Citation	Indoor Air, 21(3), 253-263 https://doi.org/10.1111/j.1600-0668.2010.00698.x
Issue Date	2011-06
Doc URL	http://hdl.handle.net/2115/87326
Rights	This is the pre-peer-reviewed version of the following article: Saijo, Y., Kanazawa, A., Araki, A., Morimoto, K., Nakayama, K., Takigawa, T., Tanaka, M., Shibata, E., Yoshimura, T., Chikara, H. and Kishi, R. (2011), Relationships between mite allergen levels, mold concentrations, and sick building syndrome symptoms in newly built dwellings in Japan. Indoor Air, 21: 253-263, which has been published in final form at https://doi.org/10.1111/j.1600-0668.2010.00698.x
Type	article (author version)
File Information	74_Indoor Air, 2011.pdf



[Instructions for use](#)

1 **Relationships between Mite Allergen Levels, Mold Concentrations, and**
2 **Sick Building Syndrome Symptoms in Newly Built Dwellings in Japan**

3 Yasuaki Saijo¹, Ayako Kanazawa², Atsuko Araki², Kanehisa Morimoto³, Kunio
4 Nakayama³, Tomoko Takigawa⁴, Masatoshi Tanaka⁵, Eiji Shibata⁶, Takesumi
5 Yoshimura⁷, Hisao Chikara⁷, and Reiko Kishi²

6

7 Affiliations

8 ¹Department of Health Science, Asahikawa Medical University, Midorigaoka E2-1-1-1,
9 Asahikawa, 078-8510, Japan

10 ² Department of Public Health, Hokkaido University Graduate School of Medicine, Kita
11 15, Nishi 7, Kita-ku, Sapporo, 060-8638, Japan

12 ³Department of Social and Environmental Medicine, Osaka University Graduate School
13 of Medicine, 2-2 Yamadaoka, Suita, 565-0871, Japan

14 ⁴Department of Public Health, Okayama University Graduate School of Medicine,
15 Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama, 700-
16 8558, Japan

17 ⁵Fukushima College, 1-1 Chigoike Miyashiro, Fukushima, 960-8505, Japan

18 ⁶Department of Health and Psychosocial Medicine, Aichi Medical University School of

1 Medicine, Nagakute-cho, Aichi Prefecture, 480-1195, Japan

2 ⁷Fukuoka Institute of Health and Environmental Sciences, 39 Mukouzano, Dazaifu,

3 818-0135, Japan

4

5 Running head: Dust mite allergens, airborne fungi, and subjective symptoms

6

7 Corresponding author: R. Kishi

8 Department of Public Health, Hokkaido University School of Medicine, North-15,

9 West-7, Kita-ku, Sapporo 060-8638, Japan

10 Tel: +81 11 706 5907 Fax: +81 11 706 7805 E-mail: rkishi@med.hokudai.ac.jp

11

12

1 **Abstract**

2 This study investigated the possible relationships between exposures to mite allergen
3 and airborne fungi with sick building syndrome (SBS) symptoms for residents living in
4 newly built dwellings. We randomly sampled 5709 newly built dwellings in 6
5 prefectures from northern to southern Japan. A total of 1479 residents in 425 households
6 participated in the study by completing questionnaire surveys and agreeing to
7 environmental monitoring for mite allergen (Der 1), airborne fungi, aldehydes, and
8 volatile organic compounds. Stepwise logistic regression analyses adjusted for
9 confounders were used to obtain odds ratios (OR) of mite allergen and fungi for SBS
10 symptoms. Der 1 had a significantly high OR for nose symptoms. *Rhodotorula* had a
11 significantly high OR for any symptoms, and *Aspergillus* had significantly high OR for
12 eye symptoms. However, the total colony-forming units had a significantly low OR for
13 throat and respiratory symptoms. *Eurotium* had a significantly low OR for skin
14 symptoms. In conclusion, dust mite allergen levels and indoor airborne *Rhodotorula* and
15 *Aspergillus* concentrations may result in SBS symptoms in newly built dwellings.

16

17 Key words: Sick building syndrome; Mite allergen; Der 1; Mold; Colony-forming unit

18

1 **Practical implications**

2 Various factors can cause sick building syndrome symptoms. This study focused on
3 biological factors such as dust mite allergen and airborne fungi in newly built dwellings
4 in Japan. Dust mite allergen levels were significantly associated with higher rates of
5 nose symptoms, airborne *Rhodotorula* concentrations were significantly associated with
6 higher rates of any symptoms, and *Aspergillus* concentrations were significantly
7 associated with higher rates of eye symptoms. Measures should be taken to reduce mite
8 allergen levels and fungal concentrations in these dwellings.

9

1 **Introduction**

2 Although sick building syndrome (SBS) is associated with mucosal, skin, and
3 general symptoms that develop primarily in office buildings (Burge, 2004), SBS can
4 also develop in domiciliary environments (Engvall, 2001). SBS in dwellings has been
5 recently highlighted in Japan; some Japanese living in newly built or renovated
6 dwellings began to experience mucosal, skin, and general symptoms in the 1990s. This
7 has been called “sick house syndrome” in Japan, and its cause has often been attributed
8 to chemicals emitted from building materials (Torii, 2002).

9 However, causes of SBS include various factors other than chemicals. Biological
10 factors, such as molds or mites, also induce SBS symptoms. We reported that dampness
11 was significantly associated with SBS symptoms in solitary houses that were recently
12 built in the Hokkaido Prefecture, a northern island of Japan (Saijo, 2004, Takeda, 2009).
13 In a preliminary survey for this study, we also reported such significant associations in
14 solitary houses that were recently built in many other regions (6 prefectures) of Japan
15 (Kishi, 2009).

16 Several mechanisms are assumed to cause the dampness effect. Higher humidity in
17 dwellings facilitates mold growth (Garrett, 1998b) and proliferation of dust mites (Bemt,
18 2006, Garrett, 1998a, Hirsch, 1998), both of which can affect health of residents.

1 Possible relationships between increased airborne fungal concentrations and SBS
2 symptoms and asthma have been reported (Cooley, 1998, Ross, 2000). However,
3 inverse relationships between airborne fungal concentrations and respiratory symptoms
4 have been reported in primary school children (Kim, 2007). Thus, the effects of airborne
5 fungal concentrations on SBS symptoms have not been fully elucidated, and there are
6 no reports of these effects in newly built dwellings.

7 Because mite exposure has sensitization and allergic effects, it can aggravate and
8 induce subjective symptoms in people of affected buildings (Bornehag, 2004). Dust
9 mite allergen can cause asthma, rhinitis, and dermatitis (Tupker, 1998) that are defined,
10 in a narrow sense, as specific building-related illnesses when they primarily occur in
11 buildings (Redlich, 1997). However, it would appear that residents who have not been
12 diagnosed with the symptoms are probably treated for SBS (Laumbach, 2005).
13 Therefore, mite allergen exposure should be measured to elucidate its effect on SBS
14 symptoms.

15 In this study, we used the symptoms query part of the Japanese version of
16 MM040EA, a validated SBS questionnaire (Andersson, 1998, Mizoue, 2001), and
17 measured dust mite allergen levels and airborne fungal concentrations to evaluate the
18 possible associations of mite and mold exposures with SBS symptoms in newly built

1 dwellings in Japan.

2

3 **Materials and Methods**

4

5 **Study population and selection of homes**

6 In our previous survey (Kishi, 2009), we randomly sampled 5709 households in
7 dwellings that were ≤ 6 years old in 6 prefectures from northern to southern Japan:
8 Hokkaido (1240 households), Fukushima (910), Aichi (1070), Osaka (885), Okayama
9 (906), and Fukuoka (698). The household list was obtained from building plan approval
10 applications, which are the official data available for inspections. Preliminary
11 questionnaires on indoor air quality were sent to the households between November
12 2003 and January 2004, and residents of 2298 households (response rate = 40.3%)
13 responded to the questionnaire. Of these respondents, 1522 residents of 444 households
14 agreed to participate in this study, which was conducted from September to November
15 2004. Of these participants, 1479 residents in 425 households were used for analyses
16 after eliminating those with missing data. The participation rate for environmental
17 monitoring was 18.5% from the respondents in the preliminary questionnaire survey.
18 Informed consent was obtained from all participants, and the study was approved by the

1 institutional ethical board for epidemiological studies at Hokkaido University Graduate
2 School of Medicine, the principal investigating institution, and the ethical boards of the
3 regional universities involved with this study.

4

5 **Questionnaire survey**

6 We used 2 types of questionnaires: one for all residents and another for the head of the
7 household or his/her partner. The questionnaire for all residents queried for information
8 on personal characteristics and lifestyle, such as age, gender, current smoking habit,
9 time spent in the dwelling, working hours, and stress levels. Furthermore, the
10 questionnaire contained questions about the history of previous treatments (by a
11 physician) for asthma or allergies. The questionnaire contained the symptoms query part
12 of the Japanese version of MM040EA, a validated questionnaire designed for
13 epidemiological assessments of SBS symptoms (Andersson, 1998, Mizoue, 2001).

14 Symptoms surveyed for the past 3 months included the following: general symptoms
15 (fatigue, feeling heavy-headed, headache, nausea/dizziness, or difficulty concentrating);
16 eye symptoms (itching, burning, or irritation of the eyes); nose symptoms (irritated,
17 stuffy, or runny nose); throat and respiratory symptoms (hoarseness, dry throat, or
18 cough); and skin symptoms (dry or flushed facial skin, scaling/itching of the scalp or

1 ears, dry, itching, or red-skinned hands). For each symptom, the following answers were
2 possible: “Yes, often (every week);” “Yes, sometimes;” and “No, never.” An additional
3 question that regarded attributing a symptom to the home environment was included in
4 the questionnaire. SBS symptoms were scored as positive if at least one subsymptom
5 was found to occur often (every week)/sometimes and was believed to be attributed to
6 the home environment. In this study, “general symptoms,” “eye symptoms,” “nose
7 symptoms,” “throat and respiratory symptoms,” and “skin symptoms” indicate the
8 abovementioned MM040EA SBS symptoms as positive. In addition, “any symptoms”
9 were defined as at least 1 SBS symptom positive. The questionnaire for children who
10 could not read and/or write was answered by their parent.

11 The questionnaire for the head of the household or his/her partner contained
12 questions regarding dampness (condensation on window panes or walls, mold growth,
13 moldy odor, slow drying of wet towels in the bathroom, water leakage during the
14 previous 5 years or for a duration the person has lived in the dwelling if 5 years had not
15 passed) and pets in the dwelling. After summing the presence or absence of the 5
16 dampness indicators, an overall dampness index (0–5) was calculated (Kishi, 2009).
17 Data on the structure and age of the dwellings were obtained from a previous survey
18 conducted in 2003.

1

2 **Assessment of indoor environmental factors**

3 Indoor airborne fungi were collected on dichloran 18% glycerol agar (DG-18: 20 mL on
4 a 90 mm Petri dish) as a culture medium using an SAS air sampler (AINEX BIO-SAS
5 International Pbi, Italy; catalog number 29842 for standard 90-mm Petri dishes). An air
6 volume of 100 L was sampled at a flow rate of 100 L/min and height of 150 cm above
7 the floor in the living room. The DG-18 medium was incubated at 27°C, fungal colonies
8 were counted, and species were identified morphologically at the Mitsubishi Chemical
9 Medicine Corporation (Tokyo, Japan). The fungal concentrations were expressed as
10 colony-forming units per cubic meter of air (CFU/m³) after the cultured numbers on the
11 medium were multiplied by 10. A positive hole correction was not applied to correct for
12 coincident spore impaction, since the total number of spore counts on the mediums
13 rarely exceeded 300 colonies (Overberger, 1995).

14 Dust samples on the floor of the living room were collected at 0.5 m²/min with a
15 hand vacuum cleaner (HC-V15, National, Japan) equipped with a paper filter. The
16 levels of Der p1 and Der f1 in the dust samples were quantified by ELISA at the LCD
17 Allergic Center (Osaka, Japan); the quantification limit for Der p1 and Der f1 was 0.1
18 µg/g fine dust. Der 1 levels were calculated by summing the quantified Der p1 and Der

1 fl levels. For calculating Der 1, values of Der p1 and Der fl less than the lower
2 quantification limit were considered to be half of the lower quantification limit.

3 Indoor air monitoring for aldehydes, acetone, and volatile organic compounds
4 (VOCs) was performed in the living room of each dwelling. Air samples were collected
5 at 100–150 cm above the floor for 24 h with a DSD-DNPH diffusion sampler (Supelco,
6 Japan) for aldehydes and acetone and a VOC-SD diffusion sampler (Supelco) for VOCs.
7 Thirteen aldehydes and acetone were quantified by HPLC, and 29 VOCs were
8 quantified by GC-MS using a previously described method (Takigawa, 2004). The
9 quantification limit for each chemical was 1.0 $\mu\text{g}/\text{m}^3$. Total VOC (TVOC) was
10 calculated as the sum of all VOCs. Because this study dealt with controlling factors for
11 mold and mite risk analyses, only formaldehyde and TVOC were used (Takigawa,
12 2004). Due to their skewed distributions, formaldehyde and TVOC values were
13 transformed to common logarithmic values. For logarithmic transformations, values less
14 than the lower quantification limit were considered to be half of the lower quantification
15 limit.

16 Formaldehyde is a major air contaminant in residential and industrial occupational
17 environments and can cause mucosal irritation symptoms (Rumchev, 2002). TVOC is
18 one of the indicators of indoor air chemical environment and is possibly related to

1 subjective symptoms (Norback, 1990). Therefore, formaldehyde and TVOC
2 concentrations were used as minimum possible confounders of indoor chemical factors
3 in the statistical analyses, though many chemical have been detected in the indoor
4 environment.

5

6 **Statistical analysis**

7 Statistical analysis was performed using multiple logistic regression. First, crude odds
8 ratios with 95% confidence intervals (OR, 95% CI) for personal and dwelling factors,
9 each fungal concentration, and Der 1 levels for SBS symptoms were calculated. Due to
10 their skewed distributions, fungal concentrations and dust mite allergen levels were
11 transformed to common logarithmic values. For logarithmic transformation, a value of 0
12 CFU/m³ for fungus was changed to 0.5 CFU/m³.

13 Next, to obtain adjusted OR for SBS symptoms, mite allergen levels, total CFU,
14 and fungal concentrations were introduced separately into the model. We controlled for
15 age (≤ 19 , 20–39, 40–59, ≥ 60 years), gender, region (2 northern cities, Sapporo and
16 Fukushima, or others), and other personal or dwelling factors significantly related to
17 each symptom. Because time spent in the dwelling had several missing values, the most
18 frequent answer, 12–20 h/day, was assigned to the missing values. The dampness index

1 was not used for adjustments because it contained fungal exposure factors. Furthermore,
2 smoking was not used because it was only negatively related to skin symptoms and
3 appeared not to have a cause–effect relationship.

4 For additional adjusted analyses, the abovementioned adjustment factors were
5 forced into the model, and total CFU, fungal concentrations, and mite allergen levels
6 with $P < 0.2$ in the above adjusted analyses were entered into the models in a backward
7 stepwise manner.

8 A significance level of 5% was applied to all statistical analyses. All analyses used
9 IBM SPSS statistics software for Windows version 18.0 (SPSS Inc., Chicago, USA).

10

1 **Results**

2 The average age of the analyzed participants was 33 years (range 0–90 years). Females
3 spent longer periods of time in the dwellings and had a lower smoking rate (Table 1).

4 The number of dwellings in the 6 prefectures was as follows: Hokkaido (104),
5 Fukushima (65), Aichi (57), Osaka (78), Okayama (71), and Fukuoka (50). About 80%
6 of the dwellings were wooden; most had been built using the timber framework method
7 (Japanese conventional method) or the two-by-four construction. The most frequent
8 dwelling age category was 1 year (Table 2). SBS symptoms were more frequent in
9 females, and the most frequent symptoms were those of the nose (Table 3).

10 The median Der 1 level in the living rooms was 1.26 µg/g dust (Table 4). Table 5
11 shows fungal exposures in the living rooms. *Cladosporium* was identified as the most
12 prevalent fungal genera. Table 6 shows formaldehyde and TVOC concentrations in the
13 living rooms. The median concentrations of formaldehyde and TVOC were 40.6 and
14 112.3 µg/m³, respectively.

15 Crude analyses (Table 7) showed that Der 1 had a significantly high OR for eye
16 and nose symptoms. Total CFU and *Cladosporium* had a significantly low OR for throat
17 and respiratory symptoms. *Penicillium* had a significantly low OR for skin symptoms.
18 *Aspergillus* had a significantly high OR for eye symptoms and *Rhodotorula* had a

1 significantly high OR for any or all symptoms. *Candida* and *Cryptococcus* had a
2 significantly high OR for nose symptoms.

3 Adjusted analyses (Table 8) showed that Der 1 had a significantly high OR for eye
4 and nose symptoms. Total CFU and *Cladosporium* had a significantly low OR for throat
5 and respiratory symptoms. *Penicillium* had a significantly high OR for eye symptoms.
6 *Aspergillus* had a significantly high OR for eye symptoms, and *Rhodotorula* had a
7 significantly high OR for any symptoms. *Eurotium* had significantly low OR for skin,
8 and throat and respiratory symptoms.

9 Stepwise analyses (Table 9) showed that Der 1 had a significantly high OR for
10 nose symptoms (OR = 1.47; 95% CI: 1.14–1.88). *Aspergillus* had a significantly high
11 OR for eye symptoms (OR = 2.38; 95% CI: 1.29–4.39), and *Rhodotorula* had a
12 significantly high OR for any symptoms (OR = 0.68; 95% CI: 1.09–2.58). However,
13 total CFU had a significantly low OR for throat and respiratory symptoms, and
14 *Eurotium* had a significantly low OR for skin symptoms.

15

1 **Discussion**

2 Our results indicated relationships between SBS symptoms and Der 1,
3 *Aspergillus*, and *Rhodotorula* for residents living in newly built houses in Japan, even
4 after accounting for adjustments. However, total CFU and *Eurotium* had protective
5 relationships for SBS symptoms in our study. To our knowledge, this is the first study
6 that has investigated the relationships of airborne fungi and mite allergen levels with
7 SBS symptoms among more than 1000 residents in newly built houses.

8 In a German study, median Der 1 levels were reported to be 0.184, 0.224, and 0.480
9 $\mu\text{g/g}$ dust at 3 different times in the living rooms (Lau, 2000). In an Italian study,
10 geometric means for Der p1 and Der f1 were 0.15 and 0.83 $\mu\text{g/g}$ dust, respectively
11 (Moscatto, 2000). Furthermore, in a Korean study, geometric means for Der p1 and Der
12 f1 were 0.11 and 7.5 $\mu\text{g/g}$ dust, respectively (Nam, 2008) and in a Thai study it was 2.43
13 $\mu\text{g/g}$ dust for Der 1 (Trakultivakorn, 2004). The mite allergen levels on the floor of the
14 living rooms in our study were somewhat low; however, they appeared comparable with
15 those of previous reports.

16 As previously described, mite allergen exposure can aggravate and induce
17 subjective symptoms in people of affected buildings (Bornehag, 2004). In this study, the
18 Der 1 concentrations were significantly associated with increased nose symptoms that

1 primarily occurred in the dwellings. Japan has a high prevalence of allergic rhinitis, and
2 cedar pollen and mites are the predominant allergens associated with allergic rhinitis
3 (Sakashita, 2009). Japanese cedar pollen-induced allergic rhinitis usually develops in
4 spring and early summer, and therefore, the present survey was conducted in the fall to
5 clearly estimate the effects of mite allergen on nose symptoms without interference
6 from cedar pollen.

7 Mite allergen had no significant effect on throat and respiratory symptoms in this
8 study. Mite allergen is a major allergen associated with bronchial asthma (Richardson,
9 2005), and people with asthmatic symptoms are probably highly concerned with dust
10 mite allergen avoidance. This may have attenuated the effects of mite allergen exposure
11 on throat and respiratory symptoms in this cross-sectional setting.

12 Two Japanese studies reported that the most prevalent mold genus in indoor
13 environments was *Cladosporium*, and the second most prevalent was *Penicillium* or
14 *Aspergillus* (Sakai, 2003, Takahashi, 1997). Furthermore, these studies reported the
15 geometric means for indoor total CFU in DG-18 medium as follows: 138/m³ in
16 Yokohama, 301 and 237/m³ in the summer in Nagoya, and 88 and 79/m³ in the winter in
17 Nagoya. Thus, the indoor air fungal concentrations of our study appeared comparable
18 with previous Japanese studies.

1 In our final model, the indoor airborne *Aspergillus* concentrations were
2 significantly associated with a higher rate of eye symptoms. Indoor airborne *Aspergillus*
3 is considered as one of the causative agents for SBS symptoms (Cooley, 1998), and its
4 primary mechanism is speculated to be allergic reactions (Schwab, 2004). Although we
5 found no significant relationship between *Aspergillus* and throat and respiratory
6 symptoms, the main types of *Aspergillus* allergy symptoms reported were respiratory
7 symptoms. However, because severity of respiratory symptoms appear to provide more
8 of an incentive to be reported (Mari, 2003, Agarwal, 2009), further studies are needed to
9 clarify *Aspergillus* allergy effects on various symptoms in circumstances of relatively
10 low level exposures. In our final model, indoor airborne *Rhodotorula* concentrations
11 were significantly associated with a higher rate of any symptoms. The possibility of its
12 inducing allergic reactions has been reported (Savolainen, 1998, Greenberger, 2004);
13 however, to our knowledge, a relationship between indoor airborne *Rhodotorula*
14 concentrations and subjective symptoms has not been reported.

15 In our study, indoor airborne total CFU and *Eurotium* were significantly associated
16 with reduced throat and respiratory symptoms. In a Swedish study, total CFU had a
17 significant negative association with asthmatic symptoms in schools; the reason
18 speculated was that higher total CFU indicated higher air ventilation rates, which had

1 protective effects against asthmatic symptoms. (Kim, 2007) Indeed, outdoor airborne
2 total CFU are higher than indoor airborne total CFU in Japan and other countries (Lee,
3 2006, Shelton, 2002, Sakai, 2003, Takahashi, 1997). The outdoor airborne
4 *Cladosporium* concentrations were higher than the indoor ones (Lee, 2006, Shelton,
5 2002, Sakai, 2003, Takahashi, 1997), and *Cladosporium* was a major component of the
6 total CFU. If higher indoor airborne *Cladosporium* concentrations and total CFU
7 represented higher home ventilation rates, which could have an effect on diluting
8 contaminants originating from indoor materials or activities, which may be a reason for
9 the protective association of *Cladosporium* and total CFU on SBS symptoms. However,
10 because significant protective effects were only seen for throat and respiratory
11 symptoms, chance alone may be a reason for the protective effects. In addition,
12 ventilation rates and habits were not evaluated in this study. Further studies are needed
13 to clarify the relationships between ventilation rates, airborne fungal exposure, and SBS
14 symptoms.

15 Another study has found a protective effect of airborne *Cladosporium* concentrations
16 on allergic sensitization of infants (Osborne, 2006). The authors speculated a Th2
17 inhibiting mechanism, such as endotoxin (Niven, 2003), but this mechanism may not be
18 applicable to children and details were unknown.

1 *Eurotium* is a xerophilic mold, which has higher indoor airborne concentrations than
2 the outdoor ones in Japan (Takahashi, 1997). The possibility of allergy induction by
3 *Eurotium* has been reported (Akiyama, 2001). Thus, the association of *Eurotium* with
4 reduced SBS symptoms in our study was a surprising result. Because *Eurotium* was
5 present in outdoor airborne samples, although at lower concentrations than in indoor
6 airborne samples, the influence of outdoor concentrations and higher ventilation rates
7 may be a reflection of chance. Furthermore, because *Eurotium* is xerophilic and more
8 conspicuous in conditions of low dampness, it may be preferable to prevent SBS
9 symptoms and typical mold growth.

10 The present study has several limitations. First, the participation rate was relatively
11 low. People who are willing to be involved in an indoor air quality study and open to
12 extensive inspection tend to be those with more severe symptoms (Bornehag, 2006).
13 Therefore, the actual prevalence of SBS may be lower than that shown by our results.

14 Second, because this was a cross-sectional study, any causal relationships between
15 variables could not be determined. In addition, previous health problems of participants
16 may have altered their lifestyle habits such as cleaning frequency and ventilation time,
17 which would dilute the effects of exposure. And, we only evaluated the living rooms of
18 the dwellings for contaminants. Japanese houses are not large, and most residents seem

1 to stay in the living rooms for many hours, except for sleep; therefore, we consider that
2 the exposure levels of residents in the living rooms represented the overall exposure
3 levels in the dwellings (Sakaguchi, 2003). Furthermore, we evaluated the airborne
4 fungi at a specific point in time during 1 min sampling period, and evaluated only
5 culturable airborne fungi. It has been reported that shorter sampling period has lower
6 representativeness (Pasanen, 2001). And, seasonal variations could not be evaluated in
7 this study. In particular, the seasonal variations of airborne fungal concentrations may
8 affect the results. Moreover, non-culturable fungi have allergic and toxic effects, and the
9 superiority of evaluating both has been reported (Osborne, 2006). However, air
10 sampling is one of the most common methods to evaluate fungal exposure level in
11 indoor environments, and as the possible health effects caused fungal exposure indoor
12 mainly are suspected to be respiratory symptoms, air sampling is believed to represent
13 the exposure (Holme, 2010). Thus, though airborne fungi measurements with longer
14 sampling time or frequent sampling will be required for further studies, we believe our
15 study is valuable because, to our knowledge, this is the first study of evaluate the
16 relationship of airborne fungi exposure on subjective symptoms in more than four
17 hundred relatively new dwellings.

18 Third, Sixth, socioeconomic status was not measured. However, because all

1 households owned newly built detached houses, it was considered that the participants
2 had socioeconomic statuses belonging to the middle class.

3 Forth, we were not able to clarify the sources of the airborne fungi in these relatively
4 new dwellings. The sources of indoor airborne fungi which have been reported are
5 outdoor air, pets, carpets, water damaged materials and damped materials, etc (Li, 2004,
6 Ren, 2001). The new materials are also possible sources (Li, 2004), and some dwellings
7 in the present study had condensation on window panes or walls, mold growth, moldy
8 odor, and/or water leakage.

9 Fifth, though it has been reported that fungi have pathogenic effects through allergic
10 reactions, microbial VOC (MVOC) emissions and their inflammatory components
11 (glucans) (Li, 2004), the pathogenic mechanisms of fungi were not able to be clarified
12 in the present study.

13 Finally, a BIO-SAS sampler was selected for this study because it is easy to use and
14 portable (battery availability). Furthermore, it has a high flow rate of 100 L/min and can
15 have a high impact efficiency (Whyte, 2007). However, a high flow rate may cause
16 more fungal injuries due to drying of viable spores (Verhoeff, 1990). The sampling
17 duration can also affect the sampling performance. Long sampling durations can reduce
18 the analytical detection limit compared with short sampling durations, although long

1 durations can also cause more fungal injuries due to drying of viable spores and
2 mechanical damage from shear and impaction forces (Saldanha, 2008). Longer nozzle-
3 to-agar distances increase the cutoff sizes (d_{50}) (Whyte, 2007) . However, the agar
4 amount in the Petri dishes used in this study, 20 mL, was that commonly used in studies
5 because the instruction manual did not suggest an ideal agar amount. Thus,
6 underestimations of fungal amounts may have occurred, although the method for
7 airborne fungi evaluation was standardized in all 6 regions. Thus, we believe that the
8 relationship between airborne fungi and SBS symptoms could be properly evaluated.

9 In conclusion, dust mite allergen levels and indoor airborne *Aspergillus* and
10 *Rhodotorula* concentrations were significantly related to SBS symptoms in newly built
11 dwellings. We should take measures to reduce the mite allergen levels and fungal
12 concentrations in dwellings. Further studies are needed to clarify seasonal variations for
13 the effects of biological contaminants, including nonviable fungi, and their effects in
14 various types of dwellings. Furthermore, it will be necessary to investigate whether
15 increased indoor airborne *Cladosporium* concentrations and total CFU actually have
16 protective relationships for symptoms in particular environments.

17

1 **Acknowledgments**

2 This study was supported by a Health Science Research Grant for Research on
3 Environmental Health from the Ministry of Health, Labor, and Welfare of Japan.

4

5 **References**

6 Agarwal, R., Aggarwal, A.N., Gupta, D. and Jindal, S.K. (2009) "Aspergillus
7 hypersensitivity and allergic bronchopulmonary aspergillosis in patients with
8 bronchial asthma: systematic review and meta-analysis", *Int J Tuberc Lung Dis*,
9 **13**, 936-944.

10 Akiyama, K. (2001) "Fungal allergy -clinical aspect-", *Nippon Ishinkin Gakkai Zasshi*,
11 **42**, 109-111 (in Japanese).

12 Andersson, K. (1998) "Epidemiological approach to indoor air problems.", *Indoor Air*, **8**
13 **(supple 4)**, 32-39.

14 Bemt, L., Vries, M.P., Knapen, L., Jansen, M., Goossens, M., Muris, J.W. and Schayck,
15 C.P. (2006) "Influence of mattress characteristics on house dust mite allergen
16 concentration", *Clin Exp Allergy*, **36**, 233-237.

17 Bornehag, C.G., Sundell, J., Bonini, S., Custovic, A., Malmberg, P., Skerfving, S.,
18 Sigsgaard, T. and Verhoeff, A. (2004) "Dampness in buildings as a risk factor for

- 1 health effects, EUROEXPO: a multidisciplinary review of the literature (1998-
2 2000) on dampness and mite exposure in buildings and health effects", *Indoor*
3 *Air*, **14**, 243-257.
- 4 Bornehag, C.G., Sundell, J., Sigsgaard, T. and Janson, S. (2006) "Potential self-selection
5 bias in a nested case-control study on indoor environmental factors and their
6 association with asthma and allergic symptoms among pre-school children",
7 *Scand J Public Health*, **34**, 534-543.
- 8 Burge, P.S. (2004) "Sick building syndrome", *Occup Environ Med*, **61**, 185-190.
- 9 Cooley, J.D., Wong, W.C., Jumper, C.A. and Straus, D.C. (1998) "Correlation between
10 the prevalence of certain fungi and sick building syndrome", *Occup Environ*
11 *Med*, **55**, 579-584.
- 12 Engvall, K., Norrby, C. and Norback, D. (2001) "Sick building syndrome in relation to
13 building dampness in multi-family residential buildings in Stockholm", *Int Arch*
14 *Occup Environ Health*, **74**, 270-278.
- 15 Garrett, M.H., Hooper, B.M. and Hooper, M.A. (1998a) "Indoor environmental factors
16 associated with house-dust-mite allergen (Der p 1) levels in south-eastern
17 Australian houses", *Allergy*, **53**, 1060-1065.
- 18 Garrett, M.H., Rayment, P.R., Hooper, M.A., Abramson, M.J. and Hooper, B.M.

- 1 (1998b) "Indoor airborne fungal spores, house dampness and associations with
2 environmental factors and respiratory health in children", *Clin Exp Allergy*, **28**,
3 459-467.
- 4 Greenberger, P.A. (2004) "Mold-induced hypersensitivity pneumonitis", *Allergy Asthma
5 Proc*, **25**, 219-223.
- 6 Hirsch, T., Range, U., Walther, K.U., Hederer, B., Lassig, S., Frey, G. and Leupold, W.
7 (1998) "Prevalence and determinants of house dust mite allergen in East German
8 homes", *Clin Exp Allergy*, **28**, 956-964.
- 9 Holme, J., Hagerhed-Engman, L., Mattsson, J., Sundell, J. and Bornehag, C.G. (2010)
10 "Culturable mold in indoor air and its association with moisture-related
11 problems and asthma and allergy among Swedish children", *Indoor Air*, **20**, 329-
12 340.
- 13 Kim, J.L., Elfman, L., Mi, Y., Wieslander, G., Smedje, G. and Norback, D. (2007)
14 "Indoor molds, bacteria, microbial volatile organic compounds and plasticizers
15 in schools--associations with asthma and respiratory symptoms in pupils",
16 *Indoor Air*, **17**, 153-163.
- 17 Kishi, R., Saijo, Y., Kanazawa, A., Tanaka, M., Yoshimura, T., Chikara, H., Takigawa, T.,
18 Morimoto, K., Nakayama, K. and Shibata, E. (2009) "Regional differences in

1 residential environments and the association of dwellings and residential factors
2 with the sick house syndrome: a nationwide cross-sectional questionnaire study
3 in Japan", *Indoor Air*.

4 Lau, S., Illi, S., Sommerfeld, C., Niggemann, B., Bergmann, R., Von Mutius, E. and
5 Wahn, U. (2000) "Early exposure to house-dust mite and cat allergens and
6 development of childhood asthma: a cohort study. Multicentre Allergy Study
7 Group", *Lancet*, **356**, 1392-1397.

8 Laumbach, R.J. and Kipen, H.M. (2005) "Bioaerosols and sick building syndrome:
9 particles, inflammation, and allergy", *Curr Opin Allergy Clin Immunol*, **5**, 135-
10 139.

11 Lee, T., Grinshpun, S.A., Martuzevicius, D., Adhikari, A., Crawford, C.M., Luo, J. and
12 Reponen, T. (2006) "Relationship between indoor and outdoor bio-aerosols
13 collected with a button inhalable aerosol sampler in urban homes", *Indoor Air*,
14 **16**, 37-47.

15 Li, D.W. and Yang, C.S. (2004) "Fungal contamination as a major contributor to sick
16 building syndrome", *Adv Appl Microbiol*, **55**, 31-112.

17 Mari, A., Schneider, P., Wally, V., Breitenbach, M. and Simon-Nobbe, B. (2003)
18 "Sensitization to fungi: epidemiology, comparative skin tests, and IgE reactivity

- 1 of fungal extracts", *Clin Exp Allergy*, **33**, 1429-1438.
- 2 Mizoue, T., Reijula, K. and Andersson, K. (2001) "Environmental tobacco smoke
3 exposure and overtime work as risk factors for sick building syndrome in Japan",
4 *Am J Epidemiol*, **154**, 803-808.
- 5 Moscato, G., Perfetti, L., Galdi, E., Pozzi, V. and Minoia, C. (2000) "Levels of house-
6 dust-mite allergen in homes of nonallergic people in Pavia, Italy", *Allergy*, **55**,
7 873-878.
- 8 Nam, H.S., Siebers, R., Lee, S.H., Park, J.S., Kim, Y.B., Choi, Y.J. and Crane, J. (2008)
9 "House dust mite allergens in domestic homes in Cheonan, Korea", *Korean J*
10 *Parasitol*, **46**, 187-189.
- 11 Niven, R. (2003) "The endotoxin paradigm: a note of caution", *Clin Exp Allergy*, **33**,
12 273-276.
- 13 Norback, D., Torgen, M. and Edling, C. (1990) "Volatile organic compounds, respirable
14 dust, and personal factors related to prevalence and incidence of sick building
15 syndrome in primary schools", *Br J Ind Med*, **47**, 733-741.
- 16 Osborne, M., Reponen, T., Adhikari, A., Cho, S.H., Grinshpun, S.A., Levin, L.,
17 Bernstein, D.I. and Lemasters, G. (2006) "Specific fungal exposures, allergic
18 sensitization, and rhinitis in infants", *Pediatr Allergy Immunol*, **17**, 450-457.

- 1 Overberger, P.A., Wadowsky, R.M. and Schaper, M.M. (1995) "Evaluation of airborne
2 particulates and fungi during hospital renovation", *Am Ind Hyg Assoc J*, **56**, 706-
3 712.
- 4 Pasanen, A.L. (2001) "A review: fungal exposure assessment in indoor environments",
5 *Indoor Air*, **11**, 87-98.
- 6 Redlich, C.A., Sparer, J. and Cullen, M.R. (1997) "Sick-building syndrome", *Lancet*,
7 **349**, 1013-1016.
- 8 Ren, P., Jankun, T.M., Belanger, K., Bracken, M.B. and Leaderer, B.P. (2001) "The
9 relation between fungal propagules in indoor air and home characteristics",
10 *Allergy*, **56**, 419-424.
- 11 Richardson, G., Eick, S. and Jones, R. (2005) "How is the indoor environment related to
12 asthma?: literature review", *J Adv Nurs*, **52**, 328-339.
- 13 Ross, M.A., Curtis, L., Scheff, P.A., Hryhorczuk, D.O., Ramakrishnan, V., Wadden, R.A.
14 and Persky, V.W. (2000) "Association of asthma symptoms and severity with
15 indoor bioaerosols", *Allergy*, **55**, 705-711.
- 16 Rumchev, K.B., Spickett, J.T., Bulsara, M.K., Phillips, M.R. and Stick, S.M. (2002)
17 "Domestic exposure to formaldehyde significantly increases the risk of asthma
18 in young children", *Eur Respir J*, **20**, 403-408.

- 1 Saijo, Y., Kishi, R., Sata, F., Katakura, Y., Urashima, Y., Hatakeyama, A., Kobayashi, S.,
2 Jin, K., Kurahashi, N., Kondo, T., Gong, Y.Y. and Umemura, T. (2004)
3 "Symptoms in relation to chemicals and dampness in newly built dwellings", *Int*
4 *Arch Occup Environ Health*, **77**, 461-470.
- 5 Sakaguchi, J. and Akabayashi, S. (2003) "Field survey of indoor air quality in detached
6 houses in Niigata Prefecture", *Indoor Air*, **13 Suppl 6**, 42-49.
- 7 Sakai, K., Tsubouchi, H. and Mitani, K. (2003) "Airborne concentrations of fungal and
8 indoor air pollutants in dwellings in Nagoya, Japan", *Nippon Koshu Eisei Zasshi*,
9 **50**, 1017-1029 (in Japanese).
- 10 Sakashita, M., Hirota, T., Harada, M., Nakamichi, R., Tsunoda, T., Osawa, Y., Kojima,
11 A., Okamoto, M., Suzuki, D., Kubo, S., Imoto, Y., Nakamura, Y., Tamari, M. and
12 Fujieda, S. (2009) "Prevalence of Allergic Rhinitis and Sensitization to Common
13 Aeroallergens in a Japanese Population", *Int Arch Allergy Immunol*, **151**, 255-
14 261.
- 15 Saldanha, R., Manno, M., Saleh, M., Ewaze, J.O. and Scott, J.A. (2008) "The influence
16 of sampling duration on recovery of culturable fungi using the Andersen N6 and
17 RCS bioaerosol samplers", *Indoor Air*, **18**, 464-472.
- 18 Savolainen, J., Kortekangas-Savolainen, O., Nermes, M., Viander, M., Koivikko, A.,

- 1 Kalimo, K. and Terho, E.O. (1998) "IgE, IgA, and IgG responses to common
2 yeasts in atopic patients", *Allergy*, **53**, 506-512.
- 3 Schwab, C.J. and Straus, D.C. (2004) "The roles of Penicillium and Aspergillus in sick
4 building syndrome", *Adv Appl Microbiol*, **55**, 215-238.
- 5 Shelton, B.G., Kirkland, K.H., Flanders, W.D. and Morris, G.K. (2002) "Profiles of
6 airborne fungi in buildings and outdoor environments in the United States", *Appl
7 Environ Microbiol*, **68**, 1743-1753.
- 8 Takahashi, T. (1997) "Airborne fungal colony-forming units in outdoor and indoor
9 environments in Yokohama, Japan", *Mycopathologia*, **139**, 23-33.
- 10 Takeda, M., Saijo, Y., Yuasa, M., Kanazawa, A., Araki, A. and Kishi, R. (2009)
11 "Relationship between sick building syndrome and indoor environmental factors
12 in newly built Japanese dwellings", *International Archives of Occupational and
13 Environmental Health*, **82**, 583-593.
- 14 Takigawa, T., Horike, T., Ohashi, Y., Kataoka, H., Wang, D.H. and Kira, S. (2004)
15 "Were volatile organic compounds the inducing factors for subjective symptoms
16 of employees working in newly constructed hospitals?", *Environ Toxicol*, **19**,
17 280-290.
- 18 Torii, S. (2002) "Sick house syndrome", *Nippon Rinsho*, **60 Suppl 1**, 621-627 (in

- 1 Japanese).
- 2 Trakultivakorn, M. and Krudtong, S. (2004) "House dust mite allergen levels in Chiang
3 Mai homes", *Asian Pac J Allergy Immunol*, **22**, 1-6.
- 4 Tupker, R.A., De Monchy, J.G. and Coenraads, P.J. (1998) "House-dust mite
5 hypersensitivity, eczema, and other nonpulmonary manifestations of allergy",
6 *Allergy*, **53**, 92-96.
- 7 Verhoeff, A.P., Van Wijnen, J.H., Boleij, J.S., Brunekreef, B., Van Reenen-Hoekstra, E.S.
8 and Samson, R.A. (1990) "Enumeration and identification of airborne viable
9 mould propagules in houses. A field comparison of selected techniques", *Allergy*,
10 **45**, 275-284.
- 11 Whyte, W., Green, G. and Albisu, A. (2007) "Collection efficiency and design of
12 microbial air samplers", *Journal of Aerosol Science*, **38**, 97-110.

13

14

1 Table 1 Characteristics of participants (n = 1479)

Characteristics	All (n = 1479)		Female (n = 770)		Male (n = 709)	
	n	(%)	n	(%)	n	(%)
Age (years)						
≤9	326	22.0	159	20.6	167	21.7
10–19	186	12.6	100	13.0	86	11.2
20–29	77	5.2	32	4.2	45	5.9
30–39	323	21.8	132	17.1	191	24.8
40–49	243	16.4	134	17.4	109	14.2
50–59	144	9.7	57	7.4	87	11.3
≥60	180	12.2	96	12.5	84	10.9
Current smoker	180	12.2	30	3.9	150	21.2
History of allergy or asthma	713	48.2	383	49.7	330	46.5
Time in the dwelling (h/day)						
<12	394	26.6	82	10.6	312	44.0
12–20	835	56.5	490	63.6	345	48.7
≥20	239	16.2	190	24.7	49	6.9
Unknown	11	0.7	8	1.0	3	0.4

2

3

1 Table 2 Numbers of dwellings and participants with regard to dwelling characteristics

	Number of dwellings (n = 425)		Number of participants (n = 1479)	
	n	(%)	n	(%)
City				
Sapporo	104	24.5	343	23.2
Fukushima	65	15.3	238	16.1
Nagoya	57	13.4	191	12.9
Osaka	78	18.4	283	19.1
Okayama	71	16.7	260	17.6
Kitakyushu	50	11.8	164	11.1
Age of dwelling (years)				
<1	7	1.6	23	1.6
1-<2	108	25.4	392	26.5
2-<3	92	21.6	310	21.0
3-<4	77	18.1	272	18.4
4-<5	59	13.9	203	13.7
5-<6	58	13.6	200	13.5
6-<7	22	5.2	75	5.1
7-<8	2	0.5	4	0.3
Size of household				
1-2	92	21.6	176	11.9
3-4	258	60.7	903	61.1
5-8	75	17.6	400	27.0
Number of rooms				
2-4	76	17.9	239	16.2
5-6	270	63.5	825	55.8
7-12	74	17.4	302	20.4
Unknown	4	0.9	13	0.9
Wooden house	339	79.8	1150	77.8
Pets at home	103	24.2	377	25.5
Dampness index				
0	76	17.9	232	15.7
1	90	21.2	294	19.9
2	160	37.6	581	39.3
3	74	17.4	280	18.9
4	21	4.9	80	5.4
5	4	0.9	12	0.8

2

3

1 Table 3 Prevalence of sick building syndrome symptoms in females and males

Type of symptoms	Females (n = 769)		Males (n = 710)		All (n = 1479)	
	n	(%)	n	(%)	n	(%)
	Skin symptoms	42	5.5	18	2.5	60
Eye symptoms	29	3.8	22	3.1	51	3.4
Nose symptoms	65	8.5	50	7.0	115	7.8
Throat and respiratory symptoms	62	8.1	40	5.6	102	6.9
General symptoms	20	2.6	9	1.3	29	2.0
Any symptoms	128	16.8	82	11.5	210	14.2

2
3

1 Table 4 Der 1 levels in living rooms ($\mu\text{g/g}$ dust) (n = 425)

	GM*	Median	Min.	25th percentile	75th percentile	Max.	Identification rate (%)
Der p1	0.20	<0.1	<0.1	<0.1	0.55	144.80	46.8
Der f1	0.98	0.84	<0.1	0.28	4.11	200.00	86.1
Der 1 (Der p1 + Der f1)	1.69	1.26	<0.1	0.44	6.73	200.05	89.4

2 *GM: geometric mean

1 Table 5 Total colony-forming units (CFU) and fungal genera CFU in living rooms (n =
 2 425)

	GM* (CFU/m ³)	Median (CFU/m ³)	Min. (CFU/m ³)	25th percentile (CFU/m ³)	75th percentile (CFU/m ³)	Max. (CFU/m ³)	Identification rate (%)
Total CFU	247.9	260	0	160	445	3370	98.6
Genera							
<i>Cladosporium</i>	91.9	120	0	60	260	2440	92.0
<i>Penicillium</i>	12	20	0	0	50	2490	74.8
<i>Aspergillus</i>	2.7	0	0	0	10	950	46.1
<i>Alternaria</i>	1.5	0	0	0	10	100	32.3
<i>Rhodotorula</i>	1.1	0	0	0	0	330	20.0
<i>Eurotium</i>	0.9	0	0	0	0	310	17.2
<i>Aureobasidium</i>	0.7	0	0	0	0	50	8.7
<i>Candida</i>	0.7	0	0	0	0	220	7.3
<i>Cryptococcus</i>	0.6	0	0	0	0	120	7.1

3 *GM: geometric mean

1 Table 6 Concentrations of formaldehyde and total volatile organic compounds (TVOC)
 2 in living rooms (n = 425)

	GM*	Median	Min.	25th percentile	75th percentile	Max.
Formaldehyde ($\mu\text{g}/\text{m}^3$)	31.2	40.6	<1.0	29.0	57.7	202.8
TVOC ($\mu\text{g}/\text{m}^3$)	121.5	112.3	16.0	67.8	203.6	1770.9

3 *GM: geometric mean

1 Table 7 Crude odds ratios for personal and dwelling factors, mites, and fungi for sick building syndrome symptoms

	Any	Skin	Eye	Nose	Throat and respiratory	General
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
	P value	P value	P value	P value	P value	P value
Age						
≤19	2.89 (1.50–5.55) 0.001	1.72 (0.65–4.58) 0.277	2.26 (0.78–6.58) 0.135	3.21 (1.35–7.61) 0.008	1.70 (0.74–3.92) 0.211	0.79 (0.24–2.59) 0.694
20–39	3.37 (1.74–6.53) <0.001	1.75 (0.64–4.75) 0.276	1.60 (0.52–4.92) 0.416	2.96 (1.22–7.14) 0.016	2.75 (1.21–6.26) 0.016	0.67 (0.19–2.40) 0.539
40–59	2.07 (1.05–4.10) 0.037	1.10 (0.39–3.23) 0.834	0.93 (0.28–3.13) 0.905	1.66 (0.66–4.20) 0.281	1.49 (0.62–3.55) 0.369	1.17 (0.36–3.77) 0.796
≥60	Reference	Reference	Reference	Reference	Reference	Reference
Female	1.53 (1.14–2.06) 0.005	2.22 (1.27–3.90) 0.005	1.23 (0.70–2.15) 0.480	1.22 (0.83–1.79) 0.312	1.47 (0.97–2.22) 0.067	2.08 (0.94–4.60) 0.070
Two northern cites	1.78 (1.33–2.39) <0.001	1.94 (1.16–3.27) 0.021	1.28 (0.73–2.25) 0.388	1.30 (0.89–1.91) 0.176	2.15 (1.43–3.23) <0.001	1.26 (0.60–2.64) 0.538
Current smoker	0.30 (0.48–1.25) 0.300	0.24 (0.06–0.99) 0.049	0.61 (0.22–1.70) 0.341	0.75 (0.39–1.42) 0.375	1.06 (0.58–1.94) 0.854	0.25 (0.03–1.88) 0.179
History of allergy or asthma	2.26 (1.67–3.06) <0.001	3.60 (1.48–5.56) 0.001	3.26 (1.72–6.17) <0.001	4.06 (2.59–6.37) <0.001	1.65 (1.10–2.49) 0.016	1.54 (0.73–3.27) 0.260
Time in the dwelling (h)						
<12 (n = 394)	Reference	Reference	Reference	Reference	Reference	Reference
12–20 (n = 835)	1.52 (1.04–2.21) 0.030	2.10 (1.00–4.37) 0.048	2.00 (0.91–4.34) 0.086	1.45 (0.91–2.46) 0.110	1.56 (0.91–2.68) 0.109	1.34 (0.53–3.44) 0.537
≥20 (n = 239)	1.94 (1.23–3.08) 0.005	2.26 (0.94–5.45) 0.069	2.11 (0.82–5.42) 0.122	1.98 (1.09–3.59) 0.025	2.55 (1.37–4.76) 0.003	1.67 (0.53–5.22) 0.382

Age of dwelling (year)						
0–1	Reference	Reference	Reference	Reference	Reference	Reference
2–3	1.13 (0.77–1.66) 0.532	0.75 (0.37–1.54) 0.439	2.00 (0.88–4.53) 0.098	0.94 (0.57–1.56) 0.806	1.12 (0.67–1.88) 0.667	0.95 (0.33–2.76) 0.825
4–7	1.68 (1.15–2.46) 0.007	1.71 (0.90–3.23) 0.100	2.32 (1.02–5.29) 0.046	1.60 (0.99–2.59) 0.056	1.34 (0.79–2.25) 0.278	2.10 (0.84–5.70) 0.108
Wooden house	0.80 (0.55–1.15) 0.230	1.40 (0.80–2.49) 0.249	0.74 (0.36–1.54) 0.423	0.81 (0.53–1.32) 0.404	0.58 (0.33–1.02) 0.061	0.91 (0.37–2.25) 0.839
Pets at home	1.11 (0.80–1.54) 0.553	1.37 (0.79–2.40) 0.264	0.71 (0.35–1.44) 0.329	1.03 (0.67–1.60) 0.878	1.44 (0.93–2.21) 0.101	1.32 (0.59–2.93) 0.490
Population density** (n = 1466)	1.03 (0.88–1.22) 0.703	1.02 (0.75–1.38) 0.923	1.03 (0.76–1.41) 0.833	0.93 (0.71–1.23) 0.620	1.11 (0.94–1.34) 0.195	1.04 (0.70–1.55) 0.854
Dampness index**	1.42 (1.24–1.61) <0.001	1.27 (1.01–1.60) 0.042	1.41 (1.10–1.81) 0.006	1.66 (1.39–1.97) <0.001	1.27 (1.06–1.51) 0.010	1.37 (0.99–1.90) 0.056
Formaldehyde*	1.85 (1.21–2.84) 0.005	1.17 (0.64–2.16) 0.609	1.93 (0.81–4.62) 0.139	1.48 (0.89–2.46) 0.130	1.64 (0.93–2.91) 0.087	1.77 (0.59–5.32) 0.312
TVOC*	1.17 (0.78–1.77) 0.441	0.68 (0.32–1.44) 0.315	1.19 (0.55–2.56) 0.656	1.07 (0.63–1.83) 1.795	1.99 (1.15–3.44) 0.013	1.43 (0.52–3.94) 0.486
Der 1*	1.12 (0.93–1.35) 0.224	1.10 (0.80–1.53) 0.555	1.45 (1.02–2.07) 0.039	1.39 (1.09–1.77) 0.007	1.04 (0.80–1.34) 0.792	1.20 (0.76–1.92) 0.438
Total CFU*	0.86 (0.61–1.21) 0.389	0.69 (0.39–1.23) 0.210	1.73 (0.84–3.54) 0.136	0.81 (0.52–1.25) 0.340	0.57 (0.37–0.88) 0.011	0.73 (0.32–1.67) 0.458
<i>Cladosporium</i> *	0.94 (0.74–1.20) 0.620	0.79 (0.52–1.20) 0.264	1.64 (0.99–2.74) 0.057	0.88 (0.64–1.21) 0.440	0.65 (0.47–0.90) 0.008	0.73 (0.41–1.32) 0.297
<i>Penicillium</i> *	0.94 (0.70–1.25) 0.664	0.57 (0.33–0.99) 0.047	1.60 (0.95–2.71) 0.078	0.96 (0.66–1.40) 0.824	0.95 (0.64–1.42) 0.816	0.74 (0.35–1.56) 0.423
<i>Aspergillus</i> *	1.15 (0.79–1.69)	0.78 (0.37–1.66)	2.34 (1.29–4.23)	1.45 (0.91–2.32)	0.89 (0.51–1.56)	2.01 (0.90–4.49)

	0.469	0.513	0.005	0.116	0.678	0.088
<i>Alternaria*</i>	0.89 (0.54–1.15)	1.01 (0.47–2.56)	0.69 (0.25–1.91)	1.13 (0.61–2.10)	1.11 (0.55–2.08)	0.70 (0.19–2.65)
	0.636	0.828	0.478	0.696	0.832	0.600
<i>Rhodotorula*</i>	1.97 (1.32–2.93)	1.92 (0.99–3.68)	1.79 (0.87–3.68)	1.17 (0.65–2.11)	1.52 (0.86–2.67)	1.91 (0.77–4.75)
	0.001	0.050	0.112	0.594	0.147	0.166
<i>Eurotium*</i>	0.80 (0.43–1.49)	0.21 (0.04–1.31)	0.87 (0.27–2.79)	0.70 (0.30–1.65)	0.71 (0.29–1.76)	1.50 (0.45–5.00)
	0.481	0.095	0.814	0.417	0.458	0.510
<i>Aureobasidium*</i>	0.60 (0.19–1.91)	0.27 (0.20–3.77)	0.21 (0.01–4.48)	1.28 (0.37–4.51)	0.62 (0.13–3.12)	1.54 (0.16–14.99)
	0.389	0.322	0.316	0.698	0.565	0.710
<i>Candida*</i>	1.40 (0.72–2.74)	0.56 (0.10–3.12)	1.60 (0.50–5.09)	2.12 (1.02–4.43)	1.58 (0.67–3.72)	2.15 (0.57–8.04)
	0.323	0.512	0.430	0.045	0.296	0.256
<i>Cryptococcus*</i>	1.91 (0.84–4.31)	1.76 (0.44–6.96)	2.21 (0.57–8.57)	2.56 (1.01–6.48)	1.96 (0.69–5.63)	2.34 (0.42–12.96)
	0.122	0.424	0.251	0.047	0.209	0.330

1 *OR was calculated using 1 unit change (after common log-transformation)

2 **OR of per 1 number increasing

3

1 Table 8 Adjusted odds ratios for mites and fungi for sick building syndrome symptoms

	Any ^a	Skin ^b	Eye ^c	Nose ^d	Throat and respiratory ^e	General ^f
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
	P value	P value	P value	P value	P value	P value
Der 1*	1.16 (0.96–1.41) 0.136	1.12 (0.80–1.57) 0.501	1.45 (1.01–2.10) 0.046	1.47 (1.14–1.88) 0.003	1.08 (0.83–1.41) 0.553	1.16 (0.73–1.85) 0.532
Total CFU*	0.81 (0.58–1.15) 0.239	0.67 (0.38–1.18) 0.160	1.60 (0.79–3.25) 0.192	0.75 (0.48–1.18) 0.213	0.51 (0.33–0.79) 0.003	0.74 (0.33–1.67) 0.467
<i>Cladosporium</i> *	0.90 (0.70–1.15) 0.380	0.78 (0.52–1.178) 0.229	1.54(0.93–2.58) 0.096	0.84 (0.61–1.15) 0.273	0.63 (0.46–0.87) 0.005	0.81 (0.45–1.45) 0.472
<i>Penicillium</i> *	0.94 (0.70–1.27) 0.687	0.59 (0.34–1.03) 0.062	1.69 (1.01–2.84) 0.047	1.01 (0.69–1.49) 0.950	0.90 (0.60–1.37) 0.614	0.74 (0.34–1.57) 0.426
<i>Aspergillus</i> *	1.22 (0.82–1.82) 0.322	0.83 (0.39–1.79) 0.637	2.38 (1.29–4.39) 0.006	1.49 (0.92–2.43) 0.105	0.97 (0.55–1.73) 0.92	2.06 (0.91–4.65) 0.081
<i>Alternaria</i> *	0.62 (0.36–1.05) 0.077	0.77 (0.32–1.86) 0.559	0.53 (0.18–1.55) 0.243	0.99 (0.51–1.92) 0.966	0.77 (0.38–1.53) 0.450	0.59 (0.15–2.37) 0.461
<i>Rhodotorula</i> *	1.66(1.08–2.55) 0.020	1.60 (0.80–3.19) 0.185	1.70 (0.80–3.64) 0.170	1.07 (0.58–1.99) 0.823	1.15 (0.63–2.08) 0.658	1.97 (0.75–5.15) 0.168
<i>Eurotium</i> *	0.49 (0.24–1.01) 0.054	0.08 (0.01–0.62) 0.015	0.73 (0.20–2.70) 0.636	0.50 (0.19–1.33) 0.165	0.35 (0.12–0.99) 0.048	1.36 (0.38–4.86) 0.637
<i>Aureobasidium</i> *	0.79 (0.24–2.54) 0.686	0.34 (0.03–4.66) 0.420	0.21 (0.01–4.34) 0.313	1.35 (0.38–4.73) 0.643	1.07 (0.21–5.39) 0.933	1.57 (0.15–16.07) 0.706
<i>Candida</i> *	0.93 (0.46–1.89) 0.831	0.35 (0.06–2.17) 0.261	1.17 (0.35–3.92) 0.801	1.74 (0.80–3.80) 0.165	1.09 (0.45–2.67) 0.844	2.07 (0.53–8.31) 0.292
<i>Cryptococcus</i> *	1.68 (0.70–4.04) 0.246	1.34 (0.32–5.59) 0.690	2.37 (0.57–9.78) 0.235	2.59 (0.95–7.07) 0.064	1.44 (0.47–4.42) 0.525	2.22 (0.38–12.97) 0.375

- 1 *transformed to common logarithmic values
- 2 ^aAdjusted for age, gender, regions, history of allergy, time spent in the dwelling, age of dwelling, log formaldehyde
- 3 ^bAdjusted for age, gender, regions, history of allergy, time spent in the dwelling
- 4 ^cAdjusted for age, gender, regions, history of allergy, age of dwelling
- 5 ^dAdjusted for age, gender, regions, history of allergy, time spent in the dwelling
- 6 ^eAdjusted for age, gender, regions, history of allergy, time spent in the dwelling, log TVOC
- 7 ^fAdjusted for age, gender, regions

1 Table 9 Adjusted odds ratios for mites and fungi for sick building syndrome symptoms in stepwise models

	Any ^a	Skin ^b	Eye ^c	Nose ^d	Throat and respiratory ^e	General ^f
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
	P value	P value	P value	P value	P value	P value
Der 1*	not selected		not selected	1.47 (1.14–1.88)		
				0.003		
Total CFU		not selected			0.55 (0.15–0.85)	
					0.008	
<i>Cladosporium</i> *			not selected			
<i>Penicillium</i> *		0.63 (0.37–1.08)	not selected			
		0.093				
<i>Aspergillus</i> *			2.38 (1.29–4.39)	not selected		2.06 (0.91–4.65)
			0.006			0.081
<i>Alternaria</i> *	not selected					
<i>Rhodotorula</i> *	1.68 (1.09–2.58)	not selected	not selected			not selected
	0.019					
<i>Eurotium</i> *	0.48 (0.23–1.00)	0.09 (0.01–0.66)		not selected	0.42 (0.14–1.20)	
	0.051	0.019			0.105	
<i>Aureobasidium</i> *						
<i>Candida</i> *				not selected		
<i>Cryptococcus</i> *				not selected		

- 1 *transformed to common logarithmic values
- 2 ^aAdjusted for age, gender, regions, history of allergy, time spent in the dwelling, age of dwelling, log formaldehyde
- 3 ^bAdjusted for age, gender, regions, history of allergy, time spent in the dwelling
- 4 ^cAdjusted for age, gender, regions, history of allergy, age of dwelling
- 5 ^dAdjusted for age, gender, regions, history of allergy, time spent in the dwelling
- 6 ^eAdjusted for age, gender, regions, history of allergy, time spent in the dwelling, log TVOC: because total CFU and *Cladosporium*
- 7 were highly correlated, Total CFU that had a higher P value was entered into the stepwise analyses
- 8 ^fAdjusted for age, gender, regions
- 9 Not selected: eliminated in backward stepwise analysis