



Title	The relationship between atopic dermatitis and indoor environmental factors : a cross-sectional study among Japanese elementary school children
Author(s)	Ukawa, Shigekazu; Araki, Atsuko; Kanazawa, Ayako; Yuasa, Motoyuki; Kishi, Reiko
Citation	International Archives of Occupational and Environmental Health, 86(7), 777-787 https://doi.org/10.1007/s00420-012-0814-0
Issue Date	2013-10-01
Doc URL	http://hdl.handle.net/2115/87328
Rights	The final publication is available at link.springer.com
Type	article (author version)
File Information	56_Int Arch Occup Environ Health, 2013.pdf



[Instructions for use](#)

1 **Shigekazu Ukawa • Atsuko Araki • Ayako Kanazawa • Motoyuki Yuasa • Reiko Kishi**

2

3 **The relationship between atopic dermatitis and indoor environmental factors: a cross-sectional**
4 **study among Japanese elementary school children**

5

6 **Shigekazu Ukawa • Ayako Kanazawa • Motoyuki Yuasa**

7 **Department of Public Health Sciences, Hokkaido University Graduate School of Medicine, N15 W7,**

8 **Kita-ku, Sapporo 060-8638, Japan**

9

10 **Atsuko Araki • Reiko Kishi (Corresponding author)**

11 **Center for Environmental and Health Sciences, Hokkaido University, N12 W7, Kita-ku, Sapporo**

12 **060-0812, Japan**

13 **Tel: +81-11-706-4746; Fax: +81-11-706-4725**

14 **E-mail: rkishi@med.hokudai.ac.jp**

15

1 **Introduction**

2 Atopic dermatitis, classified as an allergic disease (The ISAAC Steering Committee 1998), is a
3 common chronic inflammatory skin condition, that appears in early infancy and childhood (Leung et al.
4 2008). It is reported to have a negative effect on the quality of life of children and their parents (Al
5 Shobaili 2010; McKenna et al. 2005). The overall prevalence of atopic dermatitis in school-age children
6 is estimated to be between 15 and 20% (Baumer 2008) and is increasing in the industrialised world
7 (Torres-Borrego et al. 2008) . In Japan it is about 16.9% (Williams et al. 1999). While the prevalence is
8 similar among boys and girls (Saeki et al. 2005), while prevalence is similar among boys and girls (Saeki
9 et al. 2005), sensitization to allergens such as mite and pollen has been shown to be significantly higher in
10 boys (Govaere et al. 2007). A family history of atopic dermatitis as determined by genetic factors (Osawa
11 et al. 2011) and immunological factors (Wu et al. 2011) is reported to be a strong risk factor for atopic
12 dermatitis. Mutations in the gene encoding filaggrin (FLG), which plays a role in epidermal barrier
13 formation and hydration, have been identified in 30% of Japanese atopic dermatitis patients (Nomura et al.
14 2007). However, these factors do not sufficiently explain the aetiology of atopic dermatitis. Thus,
15 environmental factors are thought to be important aspects of its pathophysiology (Leung et al. 2004).

16 Health risks from indoor air pollution have become a major issue (Fisk et al. 2010). Although
17 the indoor environment of dwellings is important, few epidemiological studies showing an association
18 between home environment factors and atopic dermatitis have been reported. Some epidemiological
19 studies have demonstrated that dampness in buildings and mould have adverse health effects on

1 respiratory, nose, and skin systems in infancy (Bornehag et al. 2004; Bornehag et al. 2005; Fisk et al.
2 2010). However, these factors were not associated with atopic dermatitis in elementary school children in
3 a Taiwanese study in subtropical areas (Yang et al. 2000). In two German studies, NO₂ emission
4 (Eberlein-Konig et al. 1998) and keeping pets indoors (Schafer et al. 1999) were reported to be risk
5 factors for atopic dermatitis. However, similar results were not found in other studies (Yang et al. 2000);
6 (Ibargoyen-Roteta et al. 2007). Despite these important studies, environmental risk factors for atopic
7 dermatitis in elementary school children are still not well known.

8 The aim of this study was to ascertain (1) which home environmental factors were associated
9 with atopic dermatitis in Japanese elementary school children aged 6 through 12 years and (2) whether
10 susceptibility to the home environment differed by gender.

11

12 **Material and methods**

13

14 **Study population**

15 This cross-sectional study was conducted in Sapporo city on Hokkaido, the north island of
16 Japan. Due to financial restrictions, 35 of Sapporo's 202 public elementary schools were invited to
17 participate in the present study. To include schools from all ten wards of Sapporo city, we approached
18 schools with the assistance of Sapporo Board of Education and Sapporo City Head Teachers' Association.
19 A total of 12 schools in eight wards agreed to participate in the study. Questionnaires were distributed to

1 all 6,393 children attending these schools via the classroom teachers. Parents were asked to fill in the
2 questionnaires. Each classroom teacher collected the questionnaires from his or her students. This study
3 was conducted after obtaining informed consent from all participants, and the study protocol was
4 approved by the ethics board for epidemiological studies at Hokkaido University Graduate School of
5 Medicine and conformed to the principles outlined in the Declaration of Helsinki of 1975, as revised in
6 1983.

7

8 Definition of atopic dermatitis

9 To define atopic dermatitis, we used the International Study Asthma and Allergies in
10 Childhood (ISAAC) core questionnaire (The ISAAC Steering Committee 1998). Atopic dermatitis was
11 defined as: a) "Having an itchy rash that comes and goes for at least 6 months?"; b) "Having the
12 aforementioned itchy rash at any time during the last 12 months?"; and c) "Having the aforementioned
13 itchy rash affect one or several of the following areas: the folds of the elbows, behind the knees, in front
14 of the ankles, under the buttocks, or around the neck, ears, or eyes?"

15

16 Questionnaire for home environment and lifestyle

17 The questionnaire on home environment investigated the following: housing type (solitary or
18 other), housing architecture (wooden or other), age of the house, home renovation within 5 years (yes or
19 no), living near a crowded street (yes or no), mechanical ventilation in the living room and/or bedroom

1 (yes or no), wall-to-wall carpeting in the home (yes or no), heating system (electronic, other (gas,
2 kerosene, or wood stove) with a ventilation duct to the outside, or other (gas, kerosene, or wood stove)
3 without a ventilation duct to the outside), furry animals and/or birds in the home (yes or no), indoor
4 smoker at home (yes or no), visible mould (yes or no), perception of mouldy odour (yes or no),
5 condensation on windowpanes (yes or no) and episodes of water leakage within the past 5 years (yes or
6 no). Demographic information such as gender, school grade, number of siblings, being the firstborn child,
7 number of family members, and parental history of allergies was also collected. Parental history of
8 allergies was assessed by the question: Have you ever been diagnosed by a doctor as having allergic
9 diseases such as asthma, allergic rhinitis, pollen allergy, or atopic dermatitis? Answers were classified as:
10 neither parent, mother only, father only, or both parents. Queries about lifestyle included hours of sleep,
11 getting enough sleep (never, sometimes, almost every day, or every day), refreshing sleep (never,
12 sometimes, almost every day, or every day), and deep sleep (never, sometimes, almost every day, or
13 every day).

14

15 Statistical analysis

16 Continuous variables are presented as medians (25 percentile-75 percentile); categorical
17 variables are presented as numbers (percentages). Characteristics and home environment of the study
18 participants with or without atopic dermatitis were compared using the Mann-Whitney U test and χ^2 test.
19 Odds ratios (ORs) and 95% confidence intervals (95% CI) for atopic dermatitis were calculated using

1 logistic regression. To obtain multivariate-adjusted ORs for atopic dermatitis in relation to home
2 environment, we controlled for possible confounders, including gender, school grade, parental history of
3 allergies (neither, mother only, father only, or both parents), number of siblings, and being a firstborn
4 child (yes or no). Study participants were then divided into two groups according to gender, and a
5 stratified analysis was performed to obtain adjusted ORs for atopic dermatitis in relation to home
6 environment factors. An alpha level of 0.05 was considered to be statistically significant. All statistical
7 analyses were performed using JMP version 9.0.2 for Windows (SAS Institute Inc., Cary, NC, USA).

8

9 **Results**

10 Of the subjects we approached, 4,445 (69.5%) responded to the questionnaire. After
11 eliminating subjects with missing data for the outcome variable and gender, 4,254 participants (2,089
12 boys and 2,165 girls) were evaluated in the final analysis. The prevalence of atopic dermatitis was 16.7%
13 (boys, 16.8%; girls 16.6%).

14 Table 1 shows associations between atopic dermatitis and characteristics of study participants.
15 Variables such as school grade, being a firstborn child, and parental history of allergies were significantly
16 associated with atopic dermatitis. There prevalence of atopic dermatitis was higher in children in lower
17 school grades who were not the firstborn, and who had both parents with a history of allergies.

18 Table 2 shows the associations between atopic dermatitis and home environment factors. There
19 were significantly more reports of visible mould, mouldy odour, and condensation on windowpanes in the

1 houses of the atopic dermatitis group at 40.2%, 7.5%, and 58.1%, respectively.

2 Table 3 shows the multivariate analysis of atopic dermatitis and home environment factors.

3 Having visible mould (OR 1.25, 95% CI: 1.01-2.11), mouldy odour (OR 1.54, 95% CI: 1.10-2.14), and

4 condensation on windowpanes in the house (OR 1.25, 95% CI: 1.05-1.48) were found to be risk factors

5 for atopic dermatitis after adjusting for gender, school grade, parental history of allergies, number of

6 siblings, and being a firstborn child (yes or no). Compared with use of an electric heating system, using a

7 non-electric heating system without a ventilation duct to the outside, was shown to have no significant

8 influence in the crude analysis. After adjusting for gender and school grade, a significant difference

9 appeared (OR 1.47 95% CI: 1.01-2.11).

10 Table 4 and table 5 show the multivariate analysis of atopic dermatitis and home environment

11 factors for boys and girls, respectively. Having visible mould (OR 1.28, 95% CI: 1.00-1.64), and mouldy

12 odour in the house (OR 1.64, 95% CI: 1.00-1.64) were found to be risk factors for atopic dermatitis for

13 boys (table 4). All ORs remained statistically significant after adjusting for gender, school grade, parental

14 history of allergies, number of siblings, and being firstborn. Having condensation on windowpanes in the

15 house was found to be a risk factor in the crude analysis, but statistical significance disappeared after

16 adjusting for school grade and parental history of allergies.

17 For girls, living near a crowded street, visible mould, and condensation on windowpanes in the

18 house were shown to be risk factors in the crude analysis. However, statistical significance disappeared

19 after adjusting for school grade and parental history of allergies.

1

2 **Discussion**

3 In this study, we found a significant relationship between atopic dermatitis and using a heating
4 system other than an electronic one without a ventilation duct to the outside. In a Taiwanese study, no
5 association was reported between atopic dermatitis and indoor use of gas heating (Yang et al. 2000),
6 whereas using gas heating with a wall mounted exhaust pipe was positively associated with atopic
7 dermatitis in a German study with a skewed sample distribution (Schafer et al. 1999). Exposure to even
8 low concentrations of NO₂ emission (Eberlein-Konig et al. 1998) or particulate matter (PM) (Song et al.
9 2011) from gas heating is considered to influence skin physiology parameters, and indoor pollutants from
10 gas combustion are thought to increase the likelihood of initial sensitisation to house dust mite (Ponsonby
11 et al. 2001). When a gas, kerosene, or wood stove is used, adequate ventilation should be provided to
12 keep the skin from drying out. Indoor air quality in winter in Sapporo city may have more of an effect on
13 people's health than in summer because the average outside temperatures are 20.3°C in summer,
14 compared to -2.5°C in winter (World Meteorological Organization; Japan Meteorological Agency). In our
15 study, housing materials, housing type, age of the house, and home renovation within 5 years were not
16 risk factors for atopic dermatitis.

17 We found significant relationships between atopic dermatitis and visible mould, perception of
18 mouldy odour, and episodes of condensation on windowpanes in the house. We also found a significant
19 association between episodes of condensation on windowpanes and atopic dermatitis. One previous study

1 demonstrated that moisture provides suitable environmental conditions necessary for mould growth
2 (Eggleston 2003). Most fungal spores are known to contain allergens (Green et al. 2003).
3 Immunoglobulin E (IgE)-mediated sensitisation to fungi such as *Alternaria*, *Aspergillus*, *Cladosporium*,
4 and *Penicillium* species, the main indoor fungi in Sapporo city (Takeda et al. 2009), is a strong risk factor
5 for asthma (Chapman 2006) and exposure to mould significantly elevates the levels of IgE antibodies
6 (Savilahti et al. 2001). Mould seems to cause and maintain skin inflammation, but the precise
7 immunological pathway has not yet been elucidated (Roll et al. 2004). To clarify this point, further
8 experimental studies are needed.

9 In the stratified analysis by gender, visible mould and mouldy odour were significantly related
10 to atopic dermatitis among boys, and this relationship remained after adjusting for variables such as
11 school grade, parental history of allergies, number of siblings, and being a firstborn child. No significant
12 association was observed between home environment and atopic dermatitis among girls after adjusting for
13 school grade and parental history of allergies. Sex hormones such as estradiol (Yamatomo et al. 2001;
14 Narita et al. 2007), progesterone (Mitchell and Gershwin 2007), testosterone (Schroeder et al. 1997), and
15 dehydroepiandrosterone (Sudo et al. 2001; Tabata et al. 1997) have been shown to positively or
16 negatively influenced allergic sensitisation, although such influences are now controversial (Chen et al.
17 2008). However, since the average age of menarche in Japan is 12.4 years old (Fukuda et al. 2011), the
18 influence of sex hormones should not affect our results. To find whether boys are more susceptible to
19 moulds than girls in elementary school children, further experimental and epidemiological studies are

1 needed.

2 When one German study was conducted in spring (Schafer et al. 1996), there was a significant
3 association between atopic dermatitis and a short distance of home from busy roads, however, this
4 association was not found in the present study which was conducted in winter. As housing in Sapporo city,
5 especially in cold winter, seems to have greater airtightness than other regions in Japan (Kanazawa et al.
6 2010), indoor air quality could be affecting atopic dermatitis more than outside air quality. No
7 associations between atopic dermatitis and wall-to-wall carpeting, furry animals or birds, and indoor
8 smoker at home were found. These results are consistent with several reports that showed no association
9 between atopic dermatitis and carpets (Austin and Russell 1997; Yang et al. 2000), presence of cats or
10 dogs, and passive smoking (Ibargoyen-Roteta et al. 2007; Yang et al. 2000; Schafer et al. 1999). Several
11 studies have reported that atopic dermatitis causes sleep problems (Schmitt J et al. 2009; Bender BG et al.
12 2003, 2008). In this study, atopic dermatitis was associated with less refreshing sleep and less deep sleep.
13 Intermittent skin itching may cause lower quality sleep. Although atopic dermatitis was associated with
14 school grade, parental history of allergies, and being a firstborn child (yes or no), these variables were
15 independent of home environments in this population because little change was found in estimated ORs
16 after adjusting for these variables.

17 Statistical significance may have occurred by chance because we ran a number of analyses.
18 However, as mentioned previously, environmental factors in this study are consistent with previous
19 reports (Yang et al. 2000; Eberlein-Konig et al. 1998; Schafer et al. 1999; Ibargoyen-Roteta et al. 2007).

1 We conducted multiple regression analysis using all variables from table 2 as well as, gender, school
2 grade, parental history of allergies, number of siblings, and being a firstborn child. Significant
3 associations were not found with variables such as heating system (Using a non-electric heating system
4 with a ventilation duct to the outside vs. an electronic system, OR(95%CI): 1.24(0.88-1.79), p=0.23;
5 using non electric heating system without a ventilation duct to the outside vs. an electronic system,
6 OR(95%CI): 1.35(0.91-2.02), p=0.13); visible mould present (OR(95%CI): 1.13(0.91-1.41), p=0.28);
7 mouldy odour (OR(95%CI): 1.41(0.96-2.02), p=0.08); condensation on window panes (OR(95%CI):
8 1.23(0.91-1.40), p=0.28). Thus, each environmental variable was not independently associated with
9 atopic dermatitis, but these environmental factors are mutually related to each other. These variables
10 should be associated with atopic dermatitis because changes in the odds ratio were within 10% compared
11 to variables in Table 3. Furthermore, problems with multicollinearity may include the results of a multiple
12 regression model. This could lead to important factors being overlooked. Consequently, each
13 environmental variable was introduced into the model separately and adjusted for subjective
14 characteristics such as gender, school grade, parental history of allergies, number of siblings, and being a
15 firstborn child.

16 This study was conducted in public elementary schools located in eight out of ten wards in
17 Sapporo. We distributed questionnaires to all the children attending these schools, and approximately
18 70% replied. Demographic characteristics in this population are similar to those for Sapporo city.
19 Furthermore, the prevalence of atopic dermatitis in this study is consistent with previous studies (Baumer

1 2008; Williams et al. 1999). Therefore, our results could be generalised to elementary school children
2 living in Sapporo city.

3 The present study had several limitations. First, socio-economic status (Werner et al. 2002),
4 parental educational level (Shaw et al. 2011), and food allergens such as milk, eggs, and soybeans (Akdis
5 et al. 2006; Breuer et al. 2004) have been reported to be risk factors for atopic dermatitis. However, these
6 factors were not examined in the present study. Second, we did not measure allergen levels; determination
7 of home environment in this study relied on self-reporting and was therefore subjective. The association
8 between exposure to mould and asthma has been highlighted over the world (Ahluwalia and Matsui 2011;
9 Smit et al. 2011). Thus, if subjects with atopic dermatitis thought that mould, condensation on
10 windowpanes, and their heating system were also risk factors for atopic dermatitis, this could have led to
11 reporting bias (Larsson et al. 2011). Third, genetic factors were also not measured. Gene mutations for
12 Filaggrin, a key protein that facilitates the terminal differentiation of the epidermis and formation of the
13 skin barrier, has been reported as a risk factor for atopic dermatitis (Osawa et al. 2011; Nemoto-Hasebe et
14 al. 2009; Osawa et al. 2010). However, genetic factors should have been partly controlled as we
15 calculated odds ratio adjusting for parental history of allergies. Fourth, because we used the
16 well-validated ISAAC questionnaire (The ISAAC Steering Committee 1998), the definition of atopic
17 dermatitis should be valid. However, measuring immunological factors such as cytokine and allergen
18 specific immune globulin (Wu et al. 2011; Namkung et al. 2011; Yanagi et al. 2010; Flohr et al. 2004)
19 would make our results more robust. Fifth, we distributed questionnaires to all of the children in the

1 schools, and thus the same parent would fill out two or more questionnaires for brothers and/or sisters.
2 Therefore, our population was not entirely independent. Sixth, this study could not show causal
3 relationship since it was a cross sectional study. To resolve this point, a prospective birth cohort study is
4 needed.

5 In conclusion, although we know the limitations of cross-sectional studies, the results of this
6 study showed that using a heating system other than electric without a ventilation duct to the outside,
7 visible mould, perception of mouldy odour, and episodes of condensation on windowpanes in the house
8 are associated with self-reported prevalence of atopic dermatitis. Especially among boys, visible mould,
9 and perception of mouldy odour are associated with atopic dermatitis. Further study is needed to
10 corroborate these findings.

11

12 Conflict of Interest

13 The authors declare that we have no conflict of interest.

14

15 Acknowledgments

16 We thank all participants and the public elementary schools in Sapporo city for their assistance. This
17 study was conducted with support from the Ministry of Health, Labour and Welfare, Health and Labour
18 Sciences Research Grant, Japan (H20-Research on Community Health Crisis Management-Ippan-009)
19 and the Environment Research and Technology Development Fund (C-1151) from the Japan Ministry of

1 Environment.

1 **References**

- 2 Ahluwalia SK, Matsui EC (2011) The indoor environment and its effects on childhood asthma. *Curr Opin*
3 *Allergy Clin Immunol* 11 (2):137-143
- 4 Akdis CA, Akdis M, Bieber T, Bindslev-Jensen C, Boguniewicz M, Eigenmann P, Hamid Q, Kapp A,
5 Leung DY, Lipozencic J, Luger TA, Muraro A, Novak N, Platts-Mills TA, Rosenwasser L,
6 Scheynius A, Simons FE, Spergel J, Turjanmaa K, Wahn U, Weidinger S, Werfel T, Zuberbier T
7 (2006) Diagnosis and treatment of atopic dermatitis in children and adults: European Academy
8 of Allergology and Clinical Immunology/American Academy of Allergy, Asthma and
9 Immunology/PRACTALL Consensus Report. *Allergy* 61 (8):969-987
- 10 Al Shobaili HA (2010) The impact of childhood atopic dermatitis on the patients' family. *Pediatr*
11 *Dermatol* 27 (6):618-623
- 12 Austin JB, Russell G (1997) Wheeze, cough, atopy, and indoor environment in the Scottish Highlands.
13 *Arch Dis Child* 76 (1):22-26
- 14 Baumer JH (2008) Atopic eczema in children, NICE. *Arch Dis Child Educ Pract Ed* 93 (3):93-97
- 15 Bornehag CG, Sundell J, Bonini S, Custovic A, Malmberg P, Skerfving S, Sigsgaard T, Verhoeff A
16 (2004) Dampness in buildings as a risk factor for health effects, EUROEXPO: a
17 multidisciplinary review of the literature (1998-2000) on dampness and mite exposure in
18 buildings and health effects. *Indoor Air* 14 (4):243-257
- 19 Bornehag CG, Sundell J, Hagerhed-Engman L, Sigsgaard T, Janson S, Aberg N (2005) 'Dampness' at

- 1 home and its association with airway, nose, and skin symptoms among 10,851 preschool
2 children in Sweden: a cross-sectional study. *Indoor Air* 15 Suppl 10:48-55
- 3 Breuer K, Heratizadeh A, Wulf A, Baumann U, Constien A, Tetau D, Kapp A, Werfel T (2004) Late
4 eczematous reactions to food in children with atopic dermatitis. *Clin Exp Allergy* 34 (5):817-824
- 5 Chapman MD (2006) Challenges associated with indoor moulds: Health effects, immune response and
6 exposure assessment. *Med Mycol* 44:S29-S32
- 7 Chen W, Mempel M, Schober W, Behrendt H, Ring J (2008) Gender difference, sex hormones, and
8 immediate type hypersensitivity reactions. *Allergy* 63 (11):1418-1427
- 9 Eberlein-Konig B, Przybilla B, Kuhl P, Pechak J, Gebefugi I, Kleinschmidt J, Ring J (1998) Influence of
10 airborne nitrogen dioxide or formaldehyde on parameters of skin function and cellular activation
11 in patients with atopic eczema and control subjects. *J Allergy Clin Immunol* 101 (1 Pt
12 1):141-143
- 13 Eggleston PA (2003) Environmental control for fungal allergen exposure. *Curr Allergy Asthma Rep* 3
14 (5):424-429
- 15 Fisk WJ, Eliseeva EA, Mendell MJ (2010) Association of residential dampness and mold with respiratory
16 tract infections and bronchitis: a meta-analysis. *Environ Health* 9:72
- 17 Flohr C, Johansson SG, Wahlgren CF, Williams H (2004) How atopic is atopic dermatitis? *J Allergy Clin
18 Immunol* 114(1):150-8
- 19 Fukuda M, Fukuda K, Shimizu T, Nobunaga M, Andersen EW, Byskov AG, Andersen CY (2011)

1 Paternal smoking habits affect the reproductive life span of daughters. *Fertil Steril* 95
2 (8):2542-2544

3 Govaere E, Van Gysel D, Massa G, Verhamme KM, Doli E, De Baets F (2007) The influence of age and
4 gender on sensitization to aero-allergens. *Pediatr Allergy Immunol* 18 (8):671-678

5 Green BJ, Mitakakis TZ, Tovey ER (2003) Allergen detection from 11 fungal species before and after
6 germination. *J Allergy Clin Immunol* 111 (2):285-289

7 Ibarгойen-Roteta N, Aguinaga-Ontoso I, Fernandez-Benitez M, Marin-Fernandez B, Guillen-Grima F,
8 Serrano-Monzo I, Hermoso-de-mendoza J, Brun-Sandiumetge C, Ferrer-Nadal A, Irujo-Andueza
9 A (2007) Role of the home environment in rhinoconjunctivitis and eczema in schoolchildren in
10 Pamplona, Spain. *J Investig Allergol Clin Immunol* 17 (3):137-144

11 Kanazawa A, Saijo Y, Tanaka M, Yoshimura T, Chikara H, Takigawa T, Morimoto K, Nakayama K,
12 Shibata E, Kishi R (2010) Nationwide study of sick house syndrome: comparison of indoor
13 environment of newly built dwellings between Sapporo city and Southern areas including those
14 in Honshu and Kyushu. *Nihon Eiseigaku Zasshi* 65 (3):447-458

15 Larsson M, Hagerhed-Engman L, Moniruzzaman S, Janson S, Sundell J, Bornehag CG (2011) Can we
16 trust cross-sectional studies when studying the risk of moisture-related problems indoor for
17 asthma in children? *Int J Environ Health Res* 21 (4):237-247

18 Leung DY, Boguniewicz M, Howell MD, Nomura I, Hamid QA (2004) New insights into atopic
19 dermatitis. *J Clin Invest* 113 (5):651-657

- 1 Leung DYM, Eichenfield IF, Boguniewicz M (2008) Atopic Dermatitis. In: Wolff K, Goldsmith LA, Katz
2 SI, Gilchrist BA, Paller AS, J.Leffell D (eds) Fitzpatrick's Dermatology in General Medicine.
3 vol 1, 7 edn. The McGraw-Hill Companies, United States of America, pp 146-158
- 4 McKenna SP, Whalley D, Dewar AL, Erdman RA, Kohlmann T, Niero M, Baro E, Cook SA, Crickx B,
5 Frech F, van Assche D (2005) International development of the Parents' Index of Quality of Life
6 in Atopic Dermatitis (PIQoL-AD). *Qual Life Res* 14 (1):231-241
- 7 Mitchell VL, Gershwin LJ (2007) Progesterone and environmental tobacco smoke act synergistically to
8 exacerbate the development of allergic asthma in a mouse model. *Clin Exp Allergy* 37
9 (2):276-286
- 10 Namkung JH, Lee JE, Kim E, Park GT, Yang HS, Jang HY, Shin ES, Cho EY, Yang JM (2011) An
11 association between IL-9 and IL-9 receptor gene polymorphisms and atopic dermatitis in a
12 Korean population. *J Dermatol Sci* 62 (1):16-21
- 13 Narita S, Goldblum RM, Watson CS, Brooks EG, Estes DM, Curran EM, Midoro-Horiuti T (2007)
14 Environmental estrogens induce mast cell degranulation and enhance IgE-mediated release of
15 allergic mediators. *Environ Health Perspect* 115 (1):48-52
- 16 Nemoto-Hasebe I, Akiyama M, Nomura T, Sandilands A, McLean WH, Shimizu H (2009) FLG mutation
17 p.Lys4021X in the C-terminal imperfect filaggrin repeat in Japanese patients with atopic eczema.
18 *Br J Dermatol* 161 (6):1387-1390
- 19 Nomura T, Sandilands A, Akiyama M, Liao H, Evans AT, Sakai K, Ota M, Sugiura H, Yamamoto K,

- 1 Sato H, Palmer CN, Smith FJ, McLean WH, Shimizu H (2007) Unique mutations in the filaggrin
2 gene in Japanese patients with ichthyosis vulgaris and atopic dermatitis. *J Allergy Clin Immunol*
3 119 (2):434-440
- 4 Osawa R, Akiyama M, Shimizu H (2011) Filaggrin gene defects and the risk of developing allergic
5 disorders. *Allergol Int* 60 (1):1-9
- 6 Osawa R, Konno S, Akiyama M, Nemoto-Hasebe I, Nomura T, Nomura Y, Abe R, Sandilands A,
7 McLean WH, Hizawa N, Nishimura M, Shimizu H (2010) Japanese-specific filaggrin gene
8 mutations in Japanese patients suffering from atopic eczema and asthma. *J Invest Dermatol* 130
9 (12):2834-2836
- 10 Ponsonby AL, Dwyer T, Kemp A, Couper D, Cochrane J, Carmichael A (2001) A prospective study of
11 the association between home gas appliance use during infancy and subsequent dust mite
12 sensitization and lung function in childhood. *Clin Exp Allergy* 31 (10):1544-1552
- 13 Roll A, Cozzio A, Fischer B, Schmid-Grendelmeier P (2004) Microbial colonization and atopic dermatitis.
14 *Curr Opin Allergy Clin Immunol* 4(5):373-8
- 15 Saeki H, Iizuka H, Mori Y, Akasaka T, Takagi H, Kitajima Y, Tezuka T, Tanaka T, Hide M, Yamamoto
16 S, Hirose Y, Kodama H, Urabe K, Furue M, Kasagi F, Torii H, Nakamura K, Morita E, Tsunemi
17 Y, Tamaki K (2005) Prevalence of atopic dermatitis in Japanese elementary schoolchildren. *Br J*
18 *Dermatol* 152 (1):110-114
- 19 Savilahti R, Uitti J, Roto P, Laippala P, Husman T (2001) Increased prevalence of atopy among children

- 1 exposed to mold in a school building. *Allergy* 56 (2):175-179
- 2 Schafer T, Heinrich J, Wjst M, Krause C, Adam H, Ring J, Wichmann HE (1999) Indoor risk factors for
3 atopic eczema in school children from East Germany. *Environ Res* 81 (2):151-158
- 4 Schafer T, Vieluf D, Behrendt H, Kramer U, Ring J (1996) Atopic eczema and other manifestations of
5 atopy: results of a study in East and West Germany. *Allergy* 51 (8):532-539
- 6 Schroeder JT, MacGlashan DW, Jr., MacDonald SM, Kagey-Sobotka A, Lichtenstein LM (1997)
7 Regulation of IgE-dependent IL-4 generation by human basophils treated with glucocorticoids. *J*
8 *Immunol* 158 (11):5448-5454
- 9 Shaw TE, Currie GP, Koudelka CW, Simpson EL (2011) Eczema prevalence in the United States: data
10 from the 2003 National Survey of Children's Health. *J Invest Dermatol* 131 (1):67-73
- 11 Smit LA, Bouzigon E, Bousquet J, Le Moual N, Nadif R, Pin I, Lathrop M, Demenais F, Kauffmann F,
12 Siroux V (2011) Mold allergen sensitization in adult asthma according to integrin beta3
13 polymorphisms and Toll-like receptor 2/+596 genotype. *J Allergy Clin Immunol* 128
14 (1):185-191
- 15 Song S, Lee K, Lee YM, Lee JH, Lee SI, Yu SD, Paek D (2011) Acute health effects of urban fine and
16 ultrafine particles on children with atopic dermatitis. *Environ Res* 111 (3):394-399
- 17 Sudo N, Yu XN, Kubo C (2001) Dehydroepiandrosterone attenuates the spontaneous elevation of serum
18 IgE level in NC/Nga mice. *Immunol Lett* 79 (3):177-179
- 19 Tabata N, Tagami H, Terui T (1997) Dehydroepiandrosterone may be one of the regulators of cytokine

1 production in atopic dermatitis. Arch Dermatol Res 289 (7):410-414

2 Takeda M, Saijo Y, Yuasa M, Kanazawa A, Araki A, Kishi R (2009) Relationship between sick building
3 syndrome and indoor environmental factors in newly built Japanese dwellings. Int Arch Occup
4 Environ Health 82 (5):583-593

5 The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee (1998)
6 Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and
7 atopic eczema: ISAAC. Lancet 351 (9111):1225-1232

8 Torres-Borrego J, Molina-Teran AB, Montes-Mendoza C (2008) Prevalence and associated factors of
9 allergic rhinitis and atopic dermatitis in children. Allergol Immunopathol (Madr) 36 (2):90-100

10 Werner S, Buser K, Kapp A, Werfel T (2002) The incidence of atopic dermatitis in school entrants is
11 associated with individual life-style factors but not with local environmental factors in Hannover,
12 Germany. Br J Dermatol 147 (1):95-104

13 Williams H, Robertson C, Stewart A, Ait-Khaled N, Anabwani G, Anderson R, Asher I, Beasley R,
14 Bjorksten B, Burr M, Clayton T, Crane J, Ellwood P, Keil U, Lai C, Mallol J, Martinez F,
15 Mitchell E, Montefort S, Pearce N, Shah J, Sibbald B, Strachan D, von Mutius E, Weiland SK
16 (1999) Worldwide variations in the prevalence of symptoms of atopic eczema in the
17 International Study of Asthma and Allergies in Childhood. J Allergy Clin Immunol 103 (1 Pt
18 1):125-138

19 Wu KG, Li TH, Chen CJ, Cheng HI, Wang TY (2011) Correlations of serum Interleukin-16, total IgE,

1 eosinophil cationic protein and total eosinophil counts with disease activity in children with
2 atopic dermatitis. *Int J Immunopathol Pharmacol* 24 (1):15-23

3 Yamatomo T, Okano M, Ono T, Nakayama E, Yoshino T, Satoskar AR, Harn DA, Jr., Nishizaki K
4 (2001) Sex-related differences in the initiation of allergic rhinitis in mice. *Allergy* 56
5 (6):525-531

6 Yanagi T, Akiyama M, Nishihara H, Ishikawa J, Sakai K, Miyamura Y, Naoe A, Kitahara T, Tanaka S,
7 Shimizu H (2010) Self-improvement of keratinocyte differentiation defects during skin
8 maturation in ABCA12-deficient harlequin ichthyosis model mice. *Am J Pathol* 177 (1):106-118

9 Yang CY, Cheng MF, Hsieh YL (2000) Effects of indoor environmental factors on risk for atopic eczema
10 in a subtropical area. *J Toxicol Environ Health A* 61 (4):245-253

11

12

Table 1. Association between atopic dermatitis and subject characteristics (n=4254)

Variable	Code	Atopic dermatitis (n=710)	Without atopic dermatitis (n=3644)	P-value
Gender	Boys	350(49.3)	1739(49.1)	0.91
	Girls	360(50.7)	1805(50.9)	
School grade	1 st	128(18.1)	571(16.1)	0.03*
	2 nd	145(20.5)	576(16.3)	
	3 rd	118(16.7)	605(17.1)	
	4 th	118(16.7)	628(17.8)	
	5 th	101(14.3)	568(16.1)	
	6 th	98(13.8)	589(16.7)	
Number of siblings	Person	2(2-3)	2(2-3)	0.43
Firstborn child	Yes	343(48.7)	1945(55.5)	0.001*
	No	361(51.3)	1559(44.5)	
Number of family members	Person	4(4-5)	4(4-5)	0.72
Parental history of allergies	Neither parent	125(18.0)	1361(39.5)	<0.001*
	Mather only	199(28.8)	830(24.1)	
	Father only	92(13.3)	479(13.9)	
	Both parents	275(39.8)	777(22.5)	
Sleeping hours	Hours	9(9-9.8)	9(9-9.8)	0.67
Sleep enough	Never	93(13.1)	380(10.8)	0.07
	Sometimes	126(17.7)	561(15.9)	
	Almost every day	347(48.9)	1746(49.6)	
	Everyday	144(20.3)	835(23.7)	
Refreshing sleep	Never	133(18.7)	463(13.1)	0.004*
	Sometimes	146(20.6)	702(19.9)	
	Almost every day	322(45.3)	1705(48.3)	
	Everyday	109(15.4)	662(18.7)	
Deep sleep	Never	39(5.5)	87(2.5)	<0.001*
	Sometimes	70(9.9)	234(6.6)	
	Almost every day	334(47.1)	1453(41.1)	
	Everyday	266(37.5)	1758(49.8)	

Values are expressed as median (25th-75th) or number of children (percentage).

P values were calculated between atopic dermatitis and without atopic dermatitis by Mann-Whitney U test or χ^2 test.

P<0.05.

Table 2. Association between atopic dermatitis and home environment (n=4254)

Variables	Code	Atopic dermatitis (n=710)	Without atopic dermatitis (n=3644)	P-value
Housing type	Solitary	303(42.7)	1542(43.7)	0.64
	Other	406(57.3)	1989(56.3)	
Architecture	Wooden	356(50.5)	1816(51.7)	0.56
	Other	349(49.5)	1698(48.3)	
Age of house	Year	13(6-20)	13(6-20)	0.37
Home renovation within 5 years	No	591(85.9)	2957(86.7)	0.56
	YES	97(14.1)	453(13.3)	
Living near a crowded street	No	150(22.2)	838(23.9)	0.12
	YES	557(78.8)	2666(76.1)	
Mechanical ventilation in the living room or bed room	No	243(34.7)	1210(34.6)	0.98
	YES	458(65.3)	2285(65.4)	
Wall-to-wall carpeting in the home	No	303(42.9)	1507(42.8)	0.93
	Yes	403(57.1)	2018(57.2)	
Heating system	Electronic	49(7.1)	303(8.8)	0.14
	Using a without electric heating system with a ventilation duct to the outside	493(71.0)	2456(71.7)	
	Using a without electric heating system without a ventilation duct to the outside	152(21.9)	669(19.5)	
Furry animals or bird in the home	No	539(75.9)	2642(74.7)	0.50
	YES	171(24.1)	894(25.3)	
Indoor smoker at home	No	378(55.4)	1835(51.9)	0.47
	YES	330(46.6)	1699(48.1)	
Visible mould	No	423(59.8)	2330(66.0)	0.001*
	YES	285(40.2)	1201(34.0)	
Mouldy odour	No	654(92.5)	3360(95.1)	0.004*
	YES	53(7.5)	172(4.9)	
Water leakage	No	611(86.8)	3148(89.2)	0.06
	YES	93(13.2)	383(10.8)	
Condensation on windowpanes	No	297(41.9)	1729(49.0)	<0.001*
	YES	412(58.1)	1801(51.0)	

Values are expressed as median (25th-75th) or number of children (percentage).

P values were calculated between atopic dermatitis and without atopic dermatitis by Mann-Whitney U test or χ^2 test.

*P<0.05.

Table 3. Multivariate analysis of atopic dermatitis and home environment

Variables	Code	COR (95%CI)	AOR ^a (95%CI)	AOR ^b (95%CI)	AOR ^c (95%CI)
Housing type	Solitary	1.00	1.00	1.00	1.00
	Other	1.03(0.88-1.22)	1.03(0.87-1.21)	0.99(0.83-1.17)	0.97(0.81-1.15)
Architecture	Wooden	1.00	1.00	1.00	1.00
	Other	1.04(0.89-1.23)	1.04(0.89-1.23)	1.01(0.85-1.20)	1.01(0.85-1.19)
Age of house	Every 5 years	1.01(0.65-2.13)	1.01(0.97-1.05)	1.01(0.97-1.06)	1.01(0.97-1.05)
Home renovation within 5 years	No	1.00	1.00	1.00	1.00
	YES	1.07(0.84-1.35)	1.09(0.86-1.38)	1.09(0.85-1.39)	1.08(0.84-1.38)
Living near a crowded street	No	1.00	1.00	1.00	1.00
	YES	1.16(0.96-1.42)	1.16(0.96-1.42)	1.10(0.90-1.36)	1.10(0.90-1.36)
Mechanical ventilation in the living room or bed room	No	1.00	1.00	1.00	1.00
	YES	0.99(0.84-1.18)	0.99(0.83-1.17)	1.01(0.85-1.21)	1.04(0.87-1.25)
Wall-to-wall carpeting in the home	No	1.00	1.00	1.00	1.00
	YES	0.99(0.84-1.16)	1.00(0.85-1.18)	1.00(0.84-1.18)	0.99(0.84-1.18)
Heating system	Electronic	1.00	1.00	1.00	1.00
	Using a without electric heating system with a ventilation duct to the outside	1.24(0.91-1.72)	1.29(0.95-1.80)	1.33(0.96-1.83)	1.28(0.93-1.80)
	Using a without electric heating system without a ventilation duct to the outside	1.40(0.99-2.00)	1.47(1.03-2.11)*	1.50(1.05-2.18)*	1.45(1.01-2.11)*
Furry animals or bird in the home	No	1.00	1.00	1.00	1.00
	YES	0.93(0.77-1.12)	0.96(0.79-1.16)	0.98(0.80-1.19)	0.94(0.77-1.15)
Indoor smoker at home	No	1.00	1.00	1.00	1.00
	YES	0.94(0.80-1.10)	0.94(0.79-1.10)	0.99(0.84-1.17)	1.01(0.85-1.20)
Visible mould	No	1.00	1.00	1.00	1.00
	YES	1.30(1.10-1.54)*	1.33(1.12-1.57)*	1.28(1.08-1.52)*	1.25(1.05-1.49)*
Mouldy odour	No	1.00	1.00	1.00	1.00
	YES	1.58(1.14-2.16)*	1.57(1.13-2.15)*	1.55(1.10-2.14)*	1.54(1.10-2.14)*
Water leakage	No	1.00	1.00	1.00	1.00
	YES	1.25(0.97-1.58)	1.27(0.99-1.62)	1.17(0.91-1.51)	1.15(0.89-1.48)
Condensation on windowpanes	No	1.00	1.00	1.00	1.00
	YES	1.33(1.13-1.56)*	1.34(1.14-1.58)*	1.25(1.06-1.48)*	1.25(1.05-1.48)*

Each environmental variable was introduced into the model separately. COR, crude odds ratio; AOR, adjusted odds ratio; CI, confidence interval.*P<0.05

^a Adjusted for gender and school grade. ^b Adjusted for gender, school grade, and parental history of allergies.

^c Adjusted for gender, school grade, parental history of allergies, number of siblings, and firstborn child (yes or no).

Table 4. Multivariate analysis of atopic dermatitis and home environment (boys only)

Variables	Code	n	COR (95%CI)	AOR ^a (95%CI)	AOR ^b (95%CI)	AOR ^c (95%CI)
Housing type						
	Solitary	908	1.00	1.00	1.00	1.00
	Other	1171	1.09(0.87-1.38)	1.09(0.86-1.38)	1.10(0.87-1.14)	1.10(0.86-1.40)
Architecture						
	Wooden	1062	1.00	1.00	1.00	1.00
	Other	1008	1.13(0.89-1.42)	1.12(0.89-1.41)	1.08(0.85-1.37)	1.06(0.83-1.35)
Age of house	Every 5 years	2014	1.01(0.95-0.99)	1.01(0.96-1.07)	1.02(0.96-1.08)	1.01(0.96-1.07)
Home renovation within 5 years						
	No	1762	1.00	1.00	1.00	1.00
	YES	254	1.05(0.74-1.48)	1.10(0.76-1.54)	1.04(0.72-1.47)	1.05(0.72-1.49)
Living near a crowded street						
	No	473	1.00	1.00	1.00	1.00
	YES	1600	0.99(0.76-1.31)	0.99(0.76-1.31)	0.95(0.72-1.26)	0.93(0.70-1.25)
Mechanical ventilation in the living room or bed room						
	No	700	1.00	1.00	1.00	1.00
	YES	1356	0.89(0.70-1.14)	0.89(0.70-1.14)	0.91(0.71-1.17)	0.94(0.73-1.21)
Heating system						
	Electronic	170	1.00	1.00	1.00	1.00
	Using a without electric heating system with a ventilation duct to the outside	1442	1.05(0.69-1.65)	1.11(0.72-1.76)	1.14(0.74-1.83)	1.12(0.72-1.79)
	Using a without electric heating system without a ventilation duct to the outside	424	1.25(0.78-2.05)	1.31(0.82-2.17)	1.35(0.83-2.25)	1.34(0.83-2.24)
Wall-to-wall carpeting in the home						
	No	875	1.00	1.00	1.00	1.00
	YES	1202	0.91(0.72-1.15)	0.91(0.72-1.16)	0.89(0.70-1.13)	0.88(0.69-1.13)
Furry animals or bird in the home						
	No	1624	1.00	1.00	1.00	1.00
	YES	460	0.86(0.64-1.14)	0.88(0.66-1.17)	0.91(0.66-1.20)	0.88(0.65-1.18)
Indoor smoker at home						
	No	1096	1.00	1.00	1.00	1.00
	YES	987	0.90(0.71-1.13)	0.89(0.70-1.12)	0.95(0.74-1.20)	0.95(0.75-1.21)
Visible mould						
	No	1355	1.00	1.00	1.00	1.00
	YES	727	1.14(1.06-1.70)*	1.37(1.08-1.73)*	1.31(1.03-1.68)*	1.28(1.00-1.64)*
Mouldy odour						
	No	1965	1.00	1.00	1.00	1.00
	YES	114	1.67(1.06-2.57)*	1.68(1.06-1.98)*	1.61(1.00-2.51)*	1.64(1.00-1.64)*
Water leakage						
	No	1841	1.00	1.00	1.00	1.00
	YES	239	1.41(1.00-1.95)*	1.42(1.01-1.98)*	1.30(0.91-1.83)	1.30(0.91-1.82)
Condensation on windowpanes						
	No	988	1.00	1.00	1.00	1.00
	YES	1092	1.31(1.04-1.66)*	1.33(1.05-1.68)*	1.24(0.98-1.58)	1.23(0.96-1.57)

Each environmental variable was introduced into the model separately. COR, crude odds ratio; AOR, adjusted odds ratio; CI, confidence interval. *P<0.05.

^a Adjusted for school grade. ^b Adjusted for school grade and parental history of allergies

^c Adjusted for school grade, parental history of allergies, number of siblings, and firstborn child (yes or no).

Table 5. Multivariate analysis of atopic dermatitis and home environment (girls only)

Variables	Code	n	COR (95%CI)	AOR ^a (95%CI)	AOR ^b (95%CI)	AOR ^c (95%CI)
Housing type	Solitary	937	1.00	1.00	1.00	1.00
	Other	1224	0.99(0.79-1.24)	0.98(0.78-1.23)	0.93(0.73-1.17)	0.97(0.77-1.24)
Architecture	Wooden	1110	1.00	1.00	1.00	1.00
	Other	1039	0.97(0.78-1.23)	0.98(0.78-1.24)	0.96(0.76-1.21)	0.97(0.76-1.23)
Age of house	Every 5 years	2078	1.02(0.96-1.07)	1.02(0.96-1.08)	1.01(0.96-1.08)	1.01(0.95-1.07)
Home renovation within 5 years	No	1786	1.00	1.00	1.00	1.00
	YES	296	1.08(0.78-1.49)	1.11(0.80-1.52)	1.15(0.81-1.60)	1.11(0.78-1.55)
Living near a crowded street	No	515	1.00	1.00	1.00	1.00
	YES	1623	1.37(1.04-1.83) *	1.37(1.04-1.83) *	1.29(0.97-1.74)	1.32(0.99-1.79)
Mechanical ventilation in the living room or bed room	No	1387	1.00	1.00	1.00	1.00
	YES	753	1.11(0.88-1.42)	1.10(0.87-1.41)	1.16(0.90-1.49)	1.17(0.91-1.52)
Heating system	Electronic	182	1.00	1.00	1.00	1.00
	Using a without electric heating system with a ventilation duct to the outside	1507	1.47(0.94-2.41)	1.51(0.96-2.47)	1.55(0.98-2.58)	1.46(0.91-2.43)
	Using a without electric heating system without a ventilation duct to the outside	397	1.58(0.96-2.70)	1.63(0.99-2.78)	1.67(0.99-2.91)	1.56(0.92-2.72)
Wall-to-wall carpeting in the home	No	935	1.00	1.00	1.00	1.00
	YES	1219	1.08(0.86-1.36)	1.11(0.88-1.40)	1.13(0.89-1.43)	1.13(0.89-1.44)
Furry animals or bird in the home	No	1557	1.00	1.00	1.00	1.00
	YES	605	1.00(0.78-1.29)	1.04(0.80-1.34)	1.04(0.79-1.34)	0.99(0.76-1.30)
Indoor smoker at home	No	1117	1.00	1.00	1.00	1.00
	YES	1042	0.99(0.79-1.24)	1.00(0.80-1.25)	1.05(0.83-1.33)	1.08(0.85-1.38)
Visible mould	No	1398	1.00	1.00	1.00	1.00
	YES	759	1.28(1.01-1.61) *	1.31(1.03-1.65) *	1.26(0.99-1.60)	1.23(0.96-1.57)
Mouldy odour	No	2049	1.00	1.00	1.00	1.00
	YES	111	1.49(0.92-2.32)	1.48(0.92-2.32)	1.50(0.92-2.38)	1.46(0.88-2.33)
Water leakage	No	1918	1.00	1.00	1.00	1.00
	YES	237	1.10(0.76-1.83)	1.12(0.78-1.59)	1.04(0.71-1.49)	1.01(0.69-1.46)
Condensation on windowpanes	No	1038	1.00	1.00	1.00	1.00
	YES	1121	1.35(1.07-1.69) *	1.36(1.08-1.71) *	1.27(0.99-1.60)	1.27(0.99-1.61)

Each environmental variable was introduced into the model separately. COR, crude odds ratio; AOR, adjusted odds ratio; CI, confidence interval.*P<0.05

^a Adjusted for school grade. ^b Adjusted for school grade and parental history of allergies.

^c Adjusted for school grade, parental history of allergies, number of siblings, and firstborn child (yes or no).