

## HOKKAIDO UNIVERSITY

Title	Studies on the early physiologial responses governing heat stress-inducible gene expression in the red alga Neopyropia yezoensis [an abstract of dissertation and a summary of dissertation review]
Author(s)	HO, KHOA VIET
Citation	北海道大学. 博士(水産科学) 甲第15123号
Issue Date	2022-09-26
Doc URL	http://hdl.handle.net/2115/87553
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Туре	theses (doctoral - abstract and summary of review)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Ho_Viet_Khoa_abstract.pdf (論文内容の要旨)



## 学位論文内容の要旨

博士の専攻分野の名称:博士(水産科学)

氏名:Ho Viet Khoa

学位論文題目

Studies on the early physiological responses governing heat stress-inducible gene expression in

the red alga Neopyropia yezoensis

(スサビノリの高温ストレス誘導性遺伝子発現を司る初期生理応答の研究)

Heat stress responses are complex regulatory processes, including sensing, signal transduction, and gene expression. However, the exact mechanisms of these processes in seaweeds are not well known. We explored the relationship between membrane physical states and gene expression in the red alga *Neopyropia yezoensis*. To analyze heat stress-induced gene expression, we identified two homologs of the heat-inducible *high temperature response 2* (*HTR2*) gene in *Neopyropia seriata*, named *NyHTR2* and *NyHTR2L*. We found conservation of *HTR2* homologs only within the order Bangiales; their products contained a novel conserved cysteine repeat we designated the Bangiales cysteine-rich motif. A quantitative mRNA analysis showed that expression of *NyHTR2* and *NyHTR2L* was induced by heat stress. However, the membrane fluidizer benzyl alcohol (BA) did not induce expression of these genes, indicating that the effect of heat was not due to membrane fluidization. In contrast, expression of genes encoding multiprotein-bridging factor 1 (*NyMBF1*) and HSP70s (*NyHSP70-1* and *NyHSP70-2*) was induced by heat stress and by BA, indicating that it involved a membrane

fluidization-dependent pathway. In addition, dark treatment under heat stress promoted expression of NyHTR2, NyHTR2L, NyMBF1, and NyHSP70-2 but not NyHSP70-1; expression of NyHTR2 and NyHTR2L was membrane fluidization independent and that of other genes was membrane fluidization dependent. These findings indicate that the heat stress response in N. yezoensis involves membrane fluidization-dependent and -independent pathway.