



Title	Reproductive behavior and role in maintaining an aggregative form of the freshwater green alga Marimo, <i>Aegagropila linnaei</i> , in Lake Akan, Hokkaido, Japan
Author(s)	Umekawa, Taketo; Wakana, Isamu; Ohara, Masashi
Citation	Aquatic Botany, 168, 103309 https://doi.org/10.1016/j.aquabot.2020.103309
Issue Date	2021-01
Doc URL	http://hdl.handle.net/2115/87613
Rights	© 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Legends of supplemental data.pdf



[Instructions for use](#)

Table A1

Location and depth of each study site in Lake Akan.

Table A2

Morphological characteristics of cells of *Aegagropila linnaei* in each study site in Lake Akan. Statistics indicates mean \pm standard deviation (minimum and maximum number).

Fig. A1

Map showing study sites in Lake Akan.

Fig. A2

Landscape of Lake Akan (a) and habits of *Aegagropila linnaei* in each study site: aggregations in Churui (b) and Kinetanpe (c), free-floating filaments in Takiguchi (d), and epilithic filaments (arrows) in Jyagaiwa (e) and Shurikomabetsu (f).

Fig. A3

Habits of *Aegagropila linnaei* in Lake Akan. A group of free-floating type (a) and a filament of Takiguchi (b), and a group of epilithic type on a stone (c) and a filament having zoospores of Shurikomabetsu (arrowhead indicates a liberation pore) (d).