



Title	Effect of components of green tea extracts, caffeine and catechins on hepatic drug metabolizing enzyme activities and mutagenic transformation of carcinogens
Author(s)	NIKAIIDOU, Satoshi; ISHIZUKA, Mayumi; MAEDA, Yutaka; HARA, Yukihiko; KAZUSAKA, Akio; FUJITA, Shoichi
Citation	Japanese Journal of Veterinary Research, 52(4), 185-192
Issue Date	2005-02
DOI	10.14943/jjvr.52.4.185
Doc URL	http://hdl.handle.net/2115/877
Type	bulletin (article)
File Information	185-192.pdf



[Instructions for use](#)

Effect of components of green tea extracts, caffeine and catechins on hepatic drug metabolizing enzyme activities and mutagenic transformation of carcinogens

Satoshi NIKAIDOU^{§#}, Mayumi ISHIZUKA[§], Yutaka MAEDA[§], Yukihiro HARA[‡],
Akio KAZUSAKA[§] and Shoichi FUJITA^{§*}

(Accepted for publication : February 15, 2005)

Abstract

Green tea contains catechins and caffeine as major constituents. Treatment of rats with green tea (2.5% w/v) significantly increased 7-ethoxycoumarin O-deethylase (7-ECOD), caffeine N-1 demethylase (CN1D) and UDP-glucuronyltransferase (UGT) activities. Treatment with caffeine similarly activated CYP1A2 and related monooxygenases as well as UGT, while treatment with catechins induced UGT activity but not 7-ECOD or CN1D activity. Numbers of benzo[a]pyrene (BP)-induced revertant colonies in an Ames test (mutation assay) with *S. typhimurium* TA98 as the test strain were markedly larger when BP was preincubated with the liver S-9 (9000 ×g supernatant of liver homogenate) from green tea-treated rats than when preincubated with that from control rats. In a modified Ames assay system in which UGT was activated by the addition of UDP-glucuronic acid to the preincubation mixture, the numbers of revertant colonies in the assay using liver S-9 from green tea-treated rats decreased to a similar level to that in the assay using S-9 from controls. The acceleration of two enzymatic reactions may contribute to the rapid elimination of BP; the first step, the formation of a metabolic intermediate (which is mutagenic) by CYP1A2 and the second, the conjugation of active metabolic intermediates by UGT. We speculated that green tea can reduce the amount of time carcinogens reside in the body and the chance that body tis-

[§]: Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

[#]: Hokkaido Animal Research Center, 5-39 Shintoku-nishi, Shintoku-cho, Hokkaido, 081-0038, Japan

[‡]: Food Research Laboratories, Mitsui Norin Co. Ltd., Fujieda, Shizuoka 426, Japan

*Corresponding author : Prof. Shoichi Fujita

Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, N18W9, Sapporo 060-0818, Japan

Tel : 81-11-706-6948/Fax : 81-11-706-5105

E-mail : fujita@vetmed.hokudai.ac.jp

sues will be exposed to active metabolites of carcinogens thorough rapid elimination due to the simultaneous induction of CYP1A2 and UGT activities.

Key words : Cytochrome P450, Green tea, Mutagens, UDP-glucuronyltransferase

Introduction

Diet and beverages play an important role in determining the metabolism of xenobiotics and incidence of carcinogenesis²⁰. Green tea is one of the most popular beverages in Asian countries. It contains catechins, caffeine, vitamins and amino acids and is reported to possess anticarcinogenic and antimutagenic properties against changes induced by polycyclic aromatic hydrocarbons (PAH)¹³.

PAH carcinogens such as benzo[a]pyrene (BP) in the environment are metabolically activated by cytochrome P450 (CYP) to DNA-reactive and mutagenic forms¹⁰. In contrast, one of the phase II enzymes, UDP-glucuronyl transferase (UGT), catalyzes glucuronidation and contributes to the elimination of metabolically activated toxicants. Because of their strong antioxidative activity and inhibitory effects on the CYP-mediated activation of carcinogens, catechins have been believed to be the most important factor in cancer prevention due to ingestion of green tea¹⁸.

In contrast to a report that CYP-mediated reactions were inhibited by green tea *in vitro*²², a number of studies have found that treatment of rats with green tea increased CYP1A and UGT activities *in vivo*^{7,8,17,21}. The cytosolic aromatic hydrocarbon (Ah) receptor contributes to the induction of expression of the above enzymes, and Williams et al. have showed that tea extracts weakly activate Ah receptor-dependent transcription²³. Moreover, another study demonstrated that treatment of rats with caffeine significantly increased CYP1A2 activity in a

Ah receptor-independent manner⁴. Decaffeinated green tea extract was unlikely to alter CYP2D or CYP3A-dependent activity in drug-metabolism⁹. These observations raise doubt over, or even contradict, the hypothesis that the prevention of chemical carcinogenesis by green tea may be due, at least partly, to the inhibitory effect of catechins on the CYP-mediated activation of carcinogens.

A reduction in the activation of pre-mutagens by green tea extracts has been reported; however, the contribution of UGT is still unclear. In this study, we tested the effect of green tea, catechins and caffeine on hepatic enzyme activities and the mutagenic transformation of BP. The Ames test is a sensitive biological method for measuring the mutagenic potency of chemicals. We carried out the Ames test to see if the induction of CYP1A expression and the phase II conjugating enzyme UGT play any role in the prevention of the mutagenic transformation of carcinogens and elucidate the possible mechanism of cancer prevention by green tea.

Materials and methods

Chemicals.

Caffeine monohydrate, 7-ethoxycoumarin and *p*-nitrophenol (*p*-NP) were purchased from Wako Pure Chemical Co. (Tokyo, Japan). Glucose 6-phosphate, glucose 6-phosphate dehydrogenase and NADPH were obtained from Oriental Yeast Co. (Tokyo, Japan). UDP-glucuronic acid (UDPGA) was purchased from Sigma Chemical Co. (St. Louis, MO, USA). Goat anti-rat CYP1A antiserum was from Daiichi Pure Chemical Co. (Tokyo, Japan). Green tea was bought locally (a prod-

uct of Shizuoka, Japan). Catechin mixtures (catechins), (-)-epigallocatechin, (-)-epicatechin, (-)-epigallocatechin gallate and epicatechin gallate, were provided by the Food Research Laboratories, Mitsui Norin Co. (Shizuoka, Japan). The other reagents were of analytical grade.

Animals.

Male Wistar rats (4 and 7 weeks old) were purchased from Japan SLC Inc. (Shizuoka, Japan) and used for experiments after 1 week of acclimatization. Animals were housed at $24 \pm 1^\circ\text{C}$ with a 12 h light and 12 h dark cycle, and given laboratory feed and water *ad libitum*. Rats (5 weeks old) were treated with freshly prepared green tea (2.5% w/v) in drinking water for 4 weeks. All experiments using animals were performed under the supervision and with the permission of the University's Institutional Animal Care and Use Committee. Control animals were given water *ad lib*. Rats (8 weeks old) were injected intraperitoneally with caffeine (150 mg/kg/day) or catechins (800 mg/kg/day) in saline solution for three days in separate experiments. Animals were killed by decapitation 24 h after the last treatment.

Preparation of hepatic S-9, cytosol and microsomes.

Hepatic S-9, cytosol and microsomes were prepared as described by Omura and Sato¹⁹. All samples were frozen in liquid nitrogen and stored at -80°C until assayed. The protein concentration of hepatic samples was determined as described by Lowry *et al.*¹⁶. Microsomal cytochrome P 450 contents were determined as described by Omura and Sato¹⁹.

Preparation and analysis of tea components.

Aqueous extracts of green tea (2.5% w/v) were prepared as described by Sohn *et*

*al.*²¹. The quantitation of tea components was performed using high-pressure liquid chromatography (HPLC). Catechins or green tea extracts (2.5% w/v) dissolved in the mobile phase were subjected to HPLC. The HPLC was performed by the Food Research Laboratories, Mitsui Norin Co. (Shizuoka, Japan), using a Shimadzu LC-6A liquid chromatograph equipped with a data processor (Chromatopac C-R6A) and a SPD-6AV spectrophotometric detector. The column used was Inertsil ODS (4.6 mm i.d. \times 250 mm) (GL Science Inc., Tokyo, Japan). The mobile phase was a mixture of acetonitrile : ethyl acetate : 0.05% phosphoric acid (12:6:90, v/v), and the flow rate was 2.0 ml/min. The wavelength for absorbance was 280 nm.

Enzyme assays.

The assay of 7-ethoxycoumarin O-deethylase (7-ECOD) activity was described by Greenlee *et al.*¹¹. Caffeine N-1 demethylase (CN1D) activity was measured as described by Agundez *et al.*¹. UGT activity was measured as described by Bock *et al.*⁶.

CYP1A immunoblot analysis.

Immunoblot analysis for the determination of CYP1A protein was performed as described by Laemmli¹⁴ using antibody to rat CYP1A. The spectral configuration of the immunoblot was analyzed using NIH Image as described by Lennard¹⁵.

Ames assay.

The Ames assay was performed as described by Ames *et al.*³, with some minor modifications, using *S. typhimurium* TA98. Preincubation mixtures (0.7ml) consisted of an overnight culture of *S. typhimurium* TA98 and a hepatic S-9 and S-9 cofactor mix (Oriental East Co., Tokyo) containing a complete NADPH generating system and NADH. In ex-

periments to investigate the effect of the conjugation reaction (phase II) on the metabolic activation of BP by CYP (phase I), the substrate (5 µg BP/plate) was added to preincubation mixtures containing 15 mM UDP-glucuronic acid (UDPGA). Preincubation was initiated by addition of the substrate, and mixtures were preincubated at 37°C for 20 min. Immediately after the preincubation, 2 ml of soft agar containing 0.5 mM L-histidine and 0.5 mM biotin was added. The mixtures were poured onto a glucose agar plate. The numbers of revertant colonies were counted after 48 h of incubation at 37°C.

Analysis of results.

All results are expressed as the mean ± SD. Statistical significance was determined with an unpaired t-test and Turkey-test.

Results

The results of the quantitative analyses of components in the green tea and the catechin preparations used in the experiment are shown in Table 1. When lyophilized, 1 ml of green tea (2.5% w/v) gave 7.18 mg of tea solids. As shown in Table 1, the green tea prepa-

ration used contained 30% catechins and 7% caffeine.

Treatment of rats with green tea or caffeine produced a moderate, but significant, increase in 7-ECOD, CN1D and UGT activities. CN1D activity has been reported to be specific for CYP1A2-catalyzed enzyme activity (5). Treatment of rats with crude catechins also significantly increased UGT activity. 7-ECOD and CN1D activities were not affected by catechin treatment (Table 2).

Immunoblot analysis of hepatic micro-

Table 1. Quantitative analysis of components in tea preparations by HPLC.

Components	Green tea (2.5% w/v)		Crude catechins	
	mg/ml (percentage of total solids)			
Total solids	7.18	(100.0)	2.50	(100.0)
Total catechins	2.14	(29.8)	1.64	(65.6)
EGC	0.75	(10.4)	0.48	(19.2)
EC	0.41	(5.8)	0.25	(10.0)
EGCG	0.78	(10.9)	0.65	(26.0)
ECG	0.20	(2.8)	0.26	(10.4)
Caffeine	0.49	(6.9)	0.02	(0.8)

EGC : (-)-epigallocatechin, EC : (-)-epicatechin, EGCG : (-)-epigallocatechin gallate, ECG : (-)-epicatechin gallate.

Table 2. Comparison of P450 levels and phase I and phase II enzyme activities after pretreatment of rats with green tea, caffeine or catechins.

Pretreatment	P450.	CN1D	7-ECOD	UGT
	nmol/mg protein	pmol/min/mg protein	nmol/min/mg protein	
Control	0.60 ± 0.13	9.07 ± 1.97	0.63 ± 0.02	7.56 ± 0.93
Green tea	0.58 ± 0.08	16.30 ± 2.30**	0.95 ± 0.04**	9.73 ± 0.76**
Control	0.73 ± 0.05	10.60 ± 0.74	0.65 ± 0.15	6.87 ± 1.16
Caffeine	0.86 ± 0.13	17.67 ± 3.52*	0.97 ± 0.18*	9.57 ± 0.56*
Control	0.79 ± 0.02	6.48 ± 0.63	0.54 ± 0.10	9.03 ± 0.05
Catechins	0.83 ± 0.01	6.23 ± 0.92	0.45 ± 0.01	11.59 ± 0.66*

Rats were treated with green tea (2.5% w/v, 4 weeks) in drinking water, or injected with caffeine (150 mg/kg/day, 3 days) or catechins (800 mg/kg/day, 3 days) in saline in a separate experiment. Other experimental details are discussed in the text. Each value is expressed as the mean ± SD from 4 rats. Significantly different from control; * : P < 0.05. ** : P < 0.01. 7-ECOD: 7-ethoxycoumarin O-deethylase activity, CN1D: caffeine N-1 demethylase activity, UGT: UDP-glucuronyltransferase activity




Figure 1. Immunoblot analysis and intensity of CYP1A2 expression in hepatic microsomes from rats using anti-rat CYP1A antiserum. The amount of microsomal protein applied was 15 μ g. (A) The arrow indicates the position of CYP1A2 protein. (B) Relative expression levels of CYP1A2. Microsomes used were from control (untreated, 4 weeks) rats (lane 1), green tea (2.5% w/v, 4 weeks) -treated rats (lane 2), control (saline, 3 days) rats (lane 3), caffeine (150 mg/kg/day, 3 days) -treated rats (lane 4), control (saline, 3 days) rats (lane 5), crude catechins (800 mg/kg/day, 3 days) -treated rats (lane 6), 3-methylcholanthrene (40 mg/kg/day, 3 days) -treated rats (lane 7).

somes revealed a moderate increase in the level of CYP1A2 in hepatic microsomes from green tea- and caffeine-treated rats, but not in crude catechin-treated rats (Figure 1). The anti-CYP1A antibody that we used might cross-react with both isoforms of CYP1A1 and CYP1A2. Since CYP1A2 was the major CYP1A isoform in liver, we detected the constitutive expression of CYP1A2 in treated or untreated rat livers. In the normal condition, the expression level of CYP1A1 might be very low in the liver. Western blot experiments were repeated at least three times to confirm the results.

In the Ames assay using *S. typhimurium* TA98, preincubation of BP with hepatic S-9 from green tea-treated rats significantly increased the number of revertant colonies. However, the addition of UDPGA, a co-substrate of UGT, to preincubation mixtures decreased the numbers of revertant colonies to a level similar to that of the control (Figure 2).

Discussion

It has been reported that treatment of rats with green tea significantly increased




Figure 2. Effect of the addition of UDP-glucuronic acid on ability to transform BP into a mutagen using *S. typhimurium* TA98. The Ames assay was performed using hepatic S-9 from rats treated with green tea (2.5% w/v, 4 weeks) in drinking water. Control rats received water ad lib. Each value is expressed as the mean \pm SD from 3 or 4 rats. Significantly different from control; * :P<0.05.

CYP1A and UGT activities^{7,8,17,21}. In this study, we confirmed this finding and demonstrated that treatment with caffeine also produced a similar increase in CYP1A2 and UGT activities. On the other hand, treatment with catechins did not significantly affect CYP1A2. However, UGT activity was significantly increased in catechin-treated rats. Catechin preparations contained 0.8% caffeine as a contaminant, the effect of which we thought would be negligible in this study. The amount of caffeine-impurity was less than the level which could affect phase I and phase II enzymes. It has been reported that treatment with the ellagic acid of natural polyphenols such as catechins significantly increased UGT activity but not CYP1A-catalyzed enzyme activity². Williams et al. reported that treatment with a high concentration of green tea

extracts activated the promoter region of CYP1A1 in HepG2 cells²³. However, they also showed inhibiting effects of several catechins on the activation of the Ah receptor by TCDD. They suggested that green tea extracts contain both agonists and antagonists of the Ah receptor. Notably, (-)-epigallocatechin gallate was revealed to be a potent inhibitor of Ah receptor function, and did not induce CYP1A1 transcription. Taken together, these results indicate that caffeine in green tea is an inducer of CYP1A2 and UGT activities, while the catechins in green tea selectively induce UGT activity.

Cytochrome P450 enzymes have been shown to play an important role in the oxidative metabolic activation and detoxification of many mutagens and carcinogens¹². The metabolic activation of polycyclic aromatic hydrocarbon pro-carcinogens to DNA-reactive and mutagenic forms is often catalyzed by the CYP1A subfamily¹⁰. Treatment with green tea induced expression of CYP1A and, hence, an increase in active metabolites of BP in the Ames assay. However, it also induced UGT activity, which catalyzes the detoxification of such active metabolites. Since the conjugation of BP with UDPGA by UGT might be the detoxification route, the addition of UDPGA to the Ames test is associated with a reduction in BP mutagenicity. This reaction system is closer to the situation in vivo because UGT is operative. These results suggest that the elimination of BP from rats is accelerated by green tea, while the levels of active metabolic intermediates do not exceed the control levels. The acceleration of two enzymatic reactions may contribute to the rapid elimination of BP; the first step, the formation of a metabolic intermediate (which is mutagenic) by CYP1A2 and the second, a conjugation reaction of active metabolic intermediates involving UGT.

We showed the possible role of green tea in the prevention of chemical carcinogenesis through the induction of CYP1A and UGT activities using a modified Ames assay system in which UGT is operative. The induction of CYP1A2 and UGT activities by caffeine and the induction of UGT activity by catechins may contribute to the inactivation and accelerate the elimination of active metabolic intermediates of pro-carcinogens.

Acknowledgements :

This work was supported by a Grant-in-aid for Scientific Research (Project 15201009) from the Japanese Ministry of Education, Science, Sports, Culture and Technology.

References

- 1) Agundez, J.A., Luengo, A. and Benitez, J. 1992. Caffeine demethylase activity in human and dark Agouti rat liver microsomes. *Drug. Metab. Dispos.*, 20: 343-349.
- 2) Ahn, D., Putt, D., Kresty, L., Stoner, G.D., Fromm, D. and Hollenberg, P. H. 1996. The effect of dietary ellagic acid on rat hepatic and esophageal mucosal cytochromes P450 and phase II enzymes. *Carcinogenesis*, 17: 821-828.
- 3) Ames, B.N., McCann, J. and Yamasaki, E. 1975. Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test. *Mutat. Res.*, 31: 347-364.
- 4) Ayalogu, E.O., Snelling, J., Lewis, D.F., Talwar, S., Clifford, M.N. and Ioannides, C. 1995. Induction of hepatic CYP1A2 by the oral administration of caffeine to rats: lack of association with the Ah locus. *Biochim. Biophys. Acta.*, 1272: 89-94.
- 5) Berthou, F., Flinois, J.P., Ratanasavanh, D., Beaune, P., Riche, C. and Guillouzo, A. 1991. Evidence for the involvement of several cytochromes P - 450 in the first steps of caffeine metabolism by human liver microsomes. *Drug Metab. Dispos.*, 19: 561-567.
- 6) Bock, K.W., Burchell, B., Dutton, G.J., Hanninen, O., Mulder, G.J., Owens, I.S., Siest, G. and Tephly, T.R. 1983. UDP-glucuronosyltransferase activities. *Biochem. Pharmacol.*, 32: 953-955.
- 7) Bu-Abbas, A., Clifford, M.N., Walker, R. and Ioannides, C. 1994. Selective induction of rat hepatic CYP1 and CYP4 proteins and of peroxisomal proliferation by green tea. *Carcinogenesis*, 15: 2575-2579.
- 8) Bu-Abbas, A., Clifford, M.N., Walker, R., Ioannides, C. 1998. Contribution of caffeine and flavanols in the induction of hepatic Phase II activities by green tea. *Food. Chem. Toxicol.*, 36: 617-21.
- 9) Donovan, J.L., Chavin, K.D., Devane, C.L., Taylor, R.M., Wang, J.S., Ruan, Y., Markowitz, J.S. 2004. Green tea (Camellia sinensis) extract does not alter cytochrome p450 3A4 or 2D6 activity in healthy volunteers. *Drug Metab. Dispos.*, 32:906-8.
- 10) Gautier, J.C., Lecoœur, S., Cosme, J., Perret, A., Urban, P., Beaune, P. and Pompon, D. 1996. Contribution of human cytochrome P450 to benzo[a]pyrene and benzo[a]pyrene-7,8-dihydrodiol metabolism, as predicted from heterologous expression in yeast. *Pharmacogenetics*, 6: 489-499.
- 11) Greenlee, W.F. and Poland, A. 1978. An improved assay of 7-ethoxycoumarin O-deethylase activity : induction of hepatic enzyme activity in C57BL/6J and DBA/2 J mice by phenobarbital, 3-methylcholanthrene and 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J. Pharmacol. Exp. Ther.*, 205: 596-605.
- 12) Guengerich, F. P. 1988. Roles of cytochrome P - 450 enzymes in chemical car-

- cinogenesis and cancer chemotherapy. *Cancer Res.*, 48: 29-46.
- 13) Katiyar, S.K., Agarwal, R., Zaim, M.T. and Mukhtar, H. 1993. Protection against N-nitrosodiethylamine and benzo[a]pyrene-induced forestomach and lung tumorigenesis in A/J mice by green tea. *Carcinogenesis*, 14: 849-855.
- 14) Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227: 680-685.
- 15) Lennard, P.R. 1990. Image analysis for all. *Nature*, 347: 103-104.
- 16) Lowry, O.H., Rosenbrough, N.J., Far, A.L. and Randall, R.J. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- 17) Maliakal, P.P., Coville, P.F., Wanwimolruk, S. 2001. Tea consumption modulates hepatic drug metabolizing enzymes in Wistar rats. *J. Pharm. Pharmacol.*, 53: 569-77.
- 18) Muto, S., Fujita, K., Yamazaki, Y., Kamataki, T. 2001. Inhibition by green tea catechins of metabolic activation of procarcinogens by human cytochrome P450. *Mutat. Res.*, 479: 197-206.
- 19) Omura, T. and Sato, R. 1964. The carbon monoxide-binding pigment of liver microsomes. *J. Biol. Chem.*, 239: 2379-2385.
- 20) Rogers, A.E., Zeisel, S.H. and Groopman, J. 1993. Diet and carcinogenesis. *Carcinogenesis*, 14: 2205-2217
- 21) Sohn, O.S., Surace, A., Fiala, E.S., Richie, J.P. Jr., Colosimo, S., Zang, E. and Weisburger, J.H. 1994. Effects of green and black tea on hepatic xenobiotic metabolizing systems in the male F344 rat. *Xenobiotica*, 24: 119-127.
- 22) Wang, Z.Y., Das, M., Bickers, D.R. and Mukhtar, H. 1988. Interaction of epicatechins derived from green tea with rat hepatic cytochrome P-450. *Drug Metab. Dispos.*, 16: 98-103.
- 23) Williams, S.N., Shih, H., Guenette, D.K., Brackney, W., Denison, M.S., Pickwell, G. V. and Quattrochi, L.C. 2000. Comparative studies on the effects of green tea extracts and individual tea catechins on human CYP1A gene expression. *Chem. Biol. Interact.*, 128: 211-29.