



Title	Environmentally relevant lead (Pb) water concentration induce toxicity in zebrafish (Danio rerio) larvae
Author(s)	Kataba, Andrew; Botha, Tarryn L.; Nakayama, Shouta M. M.; Yohannes, Yared B.; Ikenaka, Yoshinori; Wepener, Victor; Ishizuka, Mayumi
Citation	Comparative biochemistry and physiology Part C: Toxicology & pharmacology, 252, 109215 https://doi.org/10.1016/j.cbpc.2021.109215
Issue Date	2022-02-01
Doc URL	http://hdl.handle.net/2115/87844
Rights	© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	http://creativecommons.org/licenses/by-nc-nd/4.0/
Type	article (author version)
File Information	Kataba et al_2nd Revision_20210916clear.pdf



[Instructions for use](#)

Environmentally relevant lead (Pb) water concentration induce toxicity in zebrafish (*Danio rerio*) larvae

Andrew Kataba^{a,b}, Tarryn L. Botha^{a,c,d}, Shouta M.M Nakayama^a, Yared B. Yohannes^{a,e}, Yoshinori Ikenaka^{a,c,f,g}, Victor Wepener^{a,c} and Mayumi Ishizuka^{a*}

^a Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Kita18, Nishi 9, Kita-ku, Sapporo, Hokkaido 060-0818, Japan

^b Department of Biomedical Sciences, School of Veterinary Medicine, The University of Zambia, Box 32379, Lusaka, Zambia

^c Water Research Group, Unit for Environmental Sciences and Management, North-West University, Private Bag X6001, Potchefstroom, South Africa

^d Institute for Nanotechnology and Water Sustainability, College of Science, Engineering and Technology, University of South Africa, Florida Science Campus, P/Bag X6, Roodepoort, 1709, South Africa.

^e Department of Chemistry, College of Natural and Computational Science, University of Gondar, P. O. Box 196, Gondar, Ethiopia

^f Translational Research Unit, Veterinary Teaching Hospital, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, 060-0818, Japan

^g One Health Research Center, Hokkaido University, Japan

24 * Corresponding author

25 Mayumi Ishizuka

26 Laboratory of Toxicology, Department of Environmental Veterinary Sciences, Faculty
27 of Veterinary Medicine, Hokkaido University, Kita 18 Nishi 9, Kita-ku, Sapporo 060-
28 0818, Japan

29 Email: ishizum@vetmed.hokudai.ac.jp

30 Tel: +81-11-706-6949

Abstract

Early developmental stages of aquatic organisms including fish are inherently vulnerable to lead (Pb) and other water metal contaminants. However, reports on the deleterious effects of environmentally relevant Pb levels are limited. To this end, we exposed 2.5 hours post fertilization (hpf) old zebrafish (*Danio rerio*) embryos to a range of Pb concentrations encompassing environmentally relevant levels (1, 10, 25, 50 and 100 µg/L Pb) until 96 hpf. Exposure negatively impacted the development and survival of zebrafish embryos by inducing embryo coagulation related mortalities in a concentration-dependent manner. At 24 hpf, the highest level of exposure (100 µg/L Pb) had impaired embryo activity characterized by reduced burst activity and the number of movements per minute made by embryos. At 72 hpf, newly hatched larvae exhibited adverse cardiovascular effects (100 µg/L Pb group) and neuromuscular effects (50 and 100 µg/L Pb groups). The antioxidant system dysregulation evidenced by downregulation of catalase, and upregulation of mRNA expression of glutathione S-transferase and cytochrome oxidase subunit I were observed. The pro-apoptotic tumor protein P53 (*TP53*) and the anti-apoptotic B cell lymphoma -2 (*Bcl-2*) mRNA expression levels were also affected. The former was downregulated across exposed groups and the latter was upregulated and downregulated in the groups with Pb concentrations less than 50 µg/L Pb and downregulated in 50 µg/L Pb, respectively. These findings suggest that Pb within environmentally relevant levels may be deleterious to developing zebrafish.

Keywords: zebrafish; lead; cardiovascular; apoptosis; twitching

1. Introduction

Lead (Pb) is a metal that naturally occurs in small amounts in the environment (Hailegnaw et al., 2015). High quantities of Pb in the environment are traceable to anthropogenic activities such as mining, Pb products production processes and their use (Komárek et al., 2008). An example of a heavily Pb polluted town with mining activities as a source is Kabwe, Zambia following 9 decades of unregulated waste management at the now closed lead-zinc mine (Yabe et al., 2018). The extensive Pb pollution of the Kabwe town has plunged the majority of its residents into chronic Pb poisoning especially children having blood Pb levels above the minimum Pb reference value of 5 µg/dL (Yabe et al., 2020, 2015). Domesticated animals, free-range chickens including free-roaming dogs around the region of the closed mine have been found with remarkable Pb levels in blood (Nakayama et al., 2011; Toyomaki et al., 2020; Yabe et al., 2013, 2011). Moreover, water sampled from natural water bodies and boreholes inside the region of the closed mine was found to have a Pb concentration range of 0.1- 94 µg/L against the country's regulatory limit of 50 µg/L Pb in water (Nachiyunde et al., 2013).

Lead exposure to developing fish embryos has been linked to undesirable effects including delayed hatching, premature hatching, and malformations of larvae, which leads to mortalities (Jezierska et al., 2009). In addition, Pb poisoning causes an imbalance of the antioxidants and eventual dysregulation of the antioxidant system (Kim and Kang, 2017). Studies have demonstrated that the antioxidant system dysregulation through increased generation of reactive oxygen species (ROS) is the major cause of oxidative-induced damage in fish exposed to Pb (Kim and Kang, 2017). A review paper citing Pb exposure studies revealed that early developmental stages of fish are more sensitive to

Pb-induced toxicity (Sfakianakis et al., 2015). However, the Pb exposure concentrations in most of the cited studies ranged from 100 to 10000 µg/L (Sfakianakis et al., 2015), which may not reflect the real prevailing environmental Pb levels.

Altered swimming behaviour accompanied by an increased oxidative stress response in larval zebrafish following acute exposure to water Pb concentrations in Kabwe, Zambia has been reported (Kataba et al., 2020). However, a dearth of data on the impacts of these Pb levels on the cardiovascular, neuromuscular and ROS-induced toxicity on the early developmental stages of fish exists. Moreover, to the best of our knowledge, the environmentally relevant Pb-induced neuromuscular toxicity in hatched embryos from the 50 and 100 µg/L Pb level of exposure has never been reported before. To bridge this information gap, we investigated the undesirable effects of the Kabwe water Pb levels on fish health by means of the zebrafish embryo toxicity testing (FET) protocol. Zebrafish (*Danio rerio*) has been known to be an ideal model for toxicological investigations owing to its morphological, biochemical and physiological data that is obtainable in early life stages such as embryos (Hill et al., 2005). Furthermore, the zebrafish lifecycle that can be managed in a laboratory environment, the embryo's ability to absorb compounds in water (Yin et al., 2017) and the shared sensitivity range with other fish species endemic to Africa (Botha et al., 2015) made the zebrafish embryos a choice model for the present study. The FET test was employed to investigate: 1) how safe are the water Pb levels reported from Kabwe, Zambia (range 0.1 - 94 µg/L Pb) on early developing stages of fish?; and 2) is the country's regulatory limit for Pb in water (50 µg/L) as reported by Nachiyunde et al., (2013) conducive to support early developmental stages of fish?

2. Materials and Methods

2.1. Fish husbandry and embryo collection

Wild-type zebrafish breeding stock kept at 26 – 28°C on a 14-hr light and 10-hr dark cycle in a ZebTec (Tecniplast, Italy) system at the North-West University's National Aquatic Bioassay Facility (NABF), South Africa were used to breed fertilized embryos. All the experiments conducted following and in strict adherence to research guidelines mandated by the North-West University AnimCare Ethics Committee (Approval number NWU-00269-16-A5). The breeding process and the collection of embryos were carried out as previously described by [Kataba et al. \(2020\)](#).

2.2. Lead (Pb) stock solution preparation and concentrations selection

A 10 mg/L Pb stock solution was prepared using pure grade (99.5%) lead acetate trihydrate in lead-free ultrapure water. From the Pb stock solution, five dilutions (1, 10, 25, 50, and 100 µg/L) were prepared using the embryo development medium (each litre contains 0.875 g NaCl, 0.038 g KCl, 0.120 g MgSO₄, 0.021 g KH₂PO₄, and 0.006 g Na₂HPO₄) constituted in deionized water (pH 8.2). The selections of the first four exposure concentrations (1-50 µg/L) were chosen to reflect the range reported in water samples from Kabwe, Zambia and the permissible water Pb level by the national authority body ([Nachiyunde et al., 2013](#)). The 100 µg/L Pb concentration was included in adherence with the Fish Embryo toxicity (FET) test protocol recommended by the Organisation for Economic Co-operation and Development (OECD) test guidelines (TG 236) for the testing of Chemicals ([Busquet et al., 2014](#)).

2.3 Fish Embryo Toxicity (FET) test and embryo activity analysis

Fertilized zebrafish embryos (2.5 hpf old) within an early developmental stage, following sorting under a Zeiss stemi microscope, were assigned to 5 concentrations of Pb at selected concentrations (0, 1, 10, 25, 50, and 100 µg/L) diluted with the embryo media in plastic 6-well plates (total volume 5000 µL/ well). A positive control with 3,4-dichloroaniline test solution was prepared. Six well plate replicates (n = 5 embryos per well plate) for the control with 30 total embryos, five replicates of total 25 embryos per treatment (1-100µg/L) were performed without renewal of the treatment solutions. The plates were covered with self-adhesive, oxygen-permeable sealing film (BRAND®, Sigma Alrich) to prevent evaporation of the test solution. The embryos were incubated at 28 °C for 96 h and their morphological condition was monitored every 24 h interval. Normal embryo morphology referencing was as described by [Kimmel et al., \(1995\)](#) and any dead or coagulated embryos were recorded and removed from the test plate. The six control replicates numbers (30 embryos) used were in line with the OECD guidelines (TG 236) for the testing of chemicals ([Busquet et al., 2014](#)).

2.4 Sub-lethal embryo activity, cardiology and twitching

A non-invasive video recording technique of assessing embryos within their chorions or hatched larvae were used. To assess embryo activity at 24 hpf, movements of embryos within test solutions were recorded for 1 min using a remote-controlled stereomicroscope (Zeiss, Germany) connected to a camera. Videos from 8 randomly selected individual embryos per replicates were assessed using DanioScope V1 software (Zeiss, Germany). The burst activity and the burst count/min were computed as a representation of embryo activity. The mean burst activity represents the percentage of time (from total

measurement duration) the embryo was moving, and the burst count/per minute represents the number of movements per minute.

At 72 hpf, blood flow and heart rate were assessed using 6 newly hatched larvae per group (n = 6) that were randomly selected using video recording followed by analysis. Individual zebrafish larvae were picked with a pipette and placed in a drop of the exposure media on a glass slide and videos were taken using a stereomicroscope (Zeiss, Germany) using a remote-controlled microscope camera. Heart rate videos were taken for 30 seconds with the heart in view while the larvae lay in lateral recumbence in the exposure media in a temperature-controlled room at the same used during the exposure **period**. The videos were imported in DanioScope and automatically the numbers of beats per minute (BPM) were calculated. The blood flow analysis video recordings were taken by focusing on the caudal artery caudal to the anal pore in view for 30 seconds. The blood flow was presented as an activity percentage.

The muscular activity of the larvae was assessed using video recordings which were later analyzed as an indicator of twitching movements. The full video was used, and movements were confirmed by watching the video as well as action potential output graphs. A peak of the action potential indicated a twitch count, and the action potential peak width represented the duration of the twitch over the 60 second time interval. The raw data was exported and used for time scale determinations (DanioScope V1 Software, Noldus Information Technology, Wageningen, Netherlands). In the present study, only action potentials with an amplitude of 5% and above were classified as muscular twitches (Fig. 3 C - H). Normal swimming or pectoral fin movement of larval zebrafish was not

considered as twitch activity. [Fig. SV1](#) shows the representative larvae video recordings for each Pb exposure level that were included in the current study.

2.5. RNA extraction and real-time PCR analysis

At the end of the exposure period (96 hpf), hatched embryos (larvae) from the different groups were sacrificed using ice-cold embryo media and 5-10 larvae were collected as a pooled sample and immediately preserved in RNA Later solution (SIGMA Life Science, St. Louis, MO, USA) at -80°C prior to their transportation to Hokkaido University, Japan. Following the sample transportation using a cold chain system on dry ice to the Laboratory of Toxicology, Faculty of Veterinary Medicine, Hokkaido University the samples were stored at -80°C until RNA extraction. The total RNA was extracted from 5-10 pooled zebrafish hatched embryos and the cDNA synthesis was done using the TOYOBO cDNA kit (TOYOBO Co., Ltd., Life Science Department, Osaka, Japan). The quantitative reverse transcription polymerase chain reaction conditions used were as previously described by [Kataba et al. \(2020\)](#). The primer sets used shown in [Table S1](#) and all primers underwent validation and were obtained as described by [Kataba et al. \(2020\)](#). The normalization and the mRNA expression levels calculations was done using the comparative ($2^{-\Delta\Delta C_t}$) method.

2.6. Verification of Pb in exposure concentrations

The nominal exposure concentrations that were prepared (0, 1, 10, 25, 50, and 100 µg/L) and used in the current study were verified using the inductively coupled plasma mass spectrometry (ICP-MS). Aliquots of freshly prepared Pb exposure concentrations in which zebrafish embryos and larvae were reared during the exposure period were used for verification by two independent analysts. The sample treatment and analysis were

done as previously described by [Kataba et al. \(2020\)](#). Recoveries from freshly prepared Pb solutions ranged from 97 to 110 % with actual concentrations of 0, 1.1, 9.7, 25.2, 49.6, and 100.7 µg/L Pb, respectively.

2.7. Data Analysis

The FET test data statistics were generated from the TOXRAT® software. GraphPad Prism software (Prism 7 for Windows; Version 5.02, California USA) was used to perform the rest of the data analysis including embryo activity, blood flow, heart rate, twitching and gene expression. The data were first tested for normality using Kolmogorov–Smirnov test. For normally distributed data, an analysis of variance (one-way ANOVA) and the differences among test groups were assessed using Tukey’s test. A non-parametric Kruskal–Wallis test followed by a Mann–Whitney test (all other comparisons) was applied for non-normally distributed data were used. The data were reported as mean and standard error of the mean (SEM). The hatching and overall survival rates proportions differences between the exposed and control groups were analysed by MedCalc® which uses the "N-1" Chi-squared test ([Campel, 2007](#)). The difference between groups was considered at two levels of significance and was marked at $p < 0.05$ (*) and at $p < 0.01$ (**). The graphical presentation of data was done using GraphPad Prism software (Prism 7 for Windows; Version 5.02, California, USA).

3. Results

3.1. Lead induced developmental toxicity in zebrafish embryo toxicity

The overall survival in the present study is defined as the total number of embryos that survived prior to hatching and those that hatched at 72 hpf. All the embryos that died during the exposure period were removed from testing plates. Hatching rates in the control and exposed groups are shown in Table 1. Lead exposure reduced the overall survival rate of exposed zebrafish embryos in a concentration-dependent manner; from 84% in the 1.0 µg/L to 52% in the 100 µg/L Pb. Lead-induced coagulation embryo mortalities that increased with the increase in Pb exposure concentration (Table 1) were recorded between 24 and 72 hpf. All surviving embryos hatched in all treatments by 72 hpf. No mortalities were observed in hatched larvae between 72 and 96 hpf periods.

Table 1. Overall survival and hatching rates of embryos


Treatment (µg/L)	Total introduced	Mortality 24 - 72 hpf	Mortality (%)	Hatched 72 hpf	Hatch 72 hpf (%)	Overall survival rate (%)
Control	30	2	6.7	28	93.3	93.3
1.0	25	4	16	21	84.0	84.0
10	25	13	52	12	48.0**	48.0
25	25	8	32	17	68.0*	68.0
50	25	8	32	17	68.0*	68.0
100	25	12	48	13	52.0**	52.0

(* $p < 0.05$ and ** $p < 0.01$ between exposure groups and the control group).

3.2. Lead exposure attenuated zebrafish embryo activity

Lead exposure affected the burst activity (Fig. 1A) and the burst count per minute (Fig. 1B) of zebrafish embryos at 24 hpf. Only the 100 µg/L Pb group ($0.10 \pm 0.05\%$) recorded significant ($p < 0.01$) lower burst activity compared to the control group (0.62

± 0.13 %). The burst count per minute was significantly lower ($p < 0.01$) in the 100 µg/L Pb group (0.24 ± 0.16 burst/min) relative to the control group (1.00 ± 0.19 burst/min). The embryo activity data retrieved for two concentrations namely; 25 and 50 µg/L Pb at 24 hpf from the video recordings were not sufficient for data analysis.



3.3. Lead exposure induced cardiovascular dysfunction

Lead exposure caused changes in cardiovascular responses of 72 hpf zebrafish for heart rate (Fig.2A) and blood flow (Fig.2B). The heart rate was significantly lower in the 100 µg/L Pb compared to the control ($p < 0.05$). No differences were observed in heart rate in exposed groups with Pb concentrations less than 100 µg/L. The blood flow at the 100 µg/L Pb exposure concentration was significantly elevated when compared to the control groups ($p < 0.05$).




Fig. 2 Effects of Pb on the cardiovascular system of larvae (72 hpf). (A) Heart rate in beats per minute (n = 6). (B). Blood flow (n = 6 for all groups). Values are presented as mean \pm SEM. The asterisk represents significant differences from the control using Mann-Whitney test (* p < 0.05).

3.4. Lead exposure induced muscular twitching

In the present study, muscular twitches were observed in zebrafish larvae in the 50 and 100 µg/L Pb exposed groups. The muscular twitching effects were absent in zebrafish larvae in exposed groups with less than 50 µg/L Pb concentration and the control group (Fig. 3). The twitching increased from 3.5 ± 1.4 twitches per min in the 50 µg/L exposure group to 17.0 ± 3.3 twitches per minute in the 100 µg/L Pb exposed group (Fig. 3A). The muscular twitch durations were 0.55 ± 0.16 and 0.51 ± 0.06 seconds in the 50 µg/L Pb and the 100 µg/L Pb groups, respectively (Fig. 3B).




Fig. 3. Lead induced involuntary muscular twitching (n = 6): **A.** Muscle twitching (number of twitches/min); **B.** Twitch durations (s) Values are presented as mean \pm SEM. The asterisk represents significant difference from the control using Mann-Whitney test (* p < 0.05; ** p < 0.01); **C, D, E, F, G and H.** Representative outputs files indicating twitch frequency and duration. The red line represents the 5% activity used as a cut-off point mark. Activity of less than 5% in a larva fish was considered a normal muscular activity and above 5% represented muscular twitching.

3.5. Lead exposure affected mRNA expression

Changes in the mRNA expression levels of antioxidant enzymes (CAT, GPX, SOD, GST, HO-1 and Nrf2) following exposure to different Pb concentrations are shown in Fig. 4. Lead induced significant down regulation of CAT mRNA expression in 50 µg/L Pb group (0.6-fold change) in relation to the control (Fig. 4A). The mRNA expression levels

of GPX, SOD and Nrf2 enzymes across the exposed groups remained unchanged when compared with the control (Fig. 4 B, C and F). The mRNA levels of GST were upregulated significantly in comparison with the control. The GST expression levels followed a concentration-dependent pattern, i.e. 1 µg/L Pb (2.5-fold change), 10 µg/L Pb (4.9-fold change), 25 µg/L Pb (8.0-fold change), 50 µg/L Pb (7.9-fold change) and 100 µg/L Pb (7.6-fold change), respectively (Fig. 4D). Furthermore, Pb exposure induced significant upregulation of HO-1 mRNA levels at 100 µg/L Pb with 2.2-fold change (Fig. 4E).




Fig. 4. Expression of mRNA in the pooled samples (n = 4). Values were normalized against Tubulin alpha-1A (used as house-keeping gene) and represent the mean mRNA expression value \pm SEM relative to those of the controls. The asterisk represents significant difference when compared with the controls (* p < 0.05 and ** p < 0.01; Mann-Whitney test)

3.6. Lead exposure affected mRNA expression of pro-apoptotic and anti-apoptotic enzymes

Mitochondrial related electron transport reactive oxygen species (ROS) associated oxidative stress response enzymes namely uncoupling protein-2 (*Ucp-2*) and cytochrome c oxidase subunit I (*CoxI*) mRNA levels were investigated. Lead exposure did not induce

Ucp-2 mRNA expression changes across the exposed groups (Fig. 5A). On the other hand, *CoxI* mRNA levels were significantly upregulated across all the exposed groups with 7.8-fold change (1 µg/L Pb), 5.9-fold change (10 µg/L Pb), 5.9-fold change (25 µg/L Pb), 5.4-fold change (50 µg/L Pb) and 7.3-fold change (100 µg/L Pb) compared to the control (Fig. 5B.)

The expression of the mRNA levels of the pro-apoptotic encoding protein tumour protein p53 (*TP53*) was significantly downregulated across all the exposure groups except in the 100 µg/L treatment when compared to the control (Fig. 5C). The B cell lymphoma-2 (*Bcl-2*) mRNA expression was significantly upregulated with 2-fold change (1 µg/L Pb), 1.8-fold change (10 µg/L Pb) and 1.7-fold change (25 µg/L Pb) when compared to the control. A significant downregulation of *TP53* mRNA expression with 0.8-fold at 50 µg/L Pb concentration was observed (Fig. 5D).




Fig. 5. Expression of mRNA of mitochondrial related enzymes (A, B, C and D) in the pooled samples (n = 4). Values normalized against Tubulin alpha-1A (used as house-keeping gene) and represent the mean mRNA expression value \pm SEM (n = 4 pooled samples) relative to those of the controls. The asterisk represents a statistically significant difference when compared with the controls (* $p < 0.05$; Mann-Whitney test).

4. Discussion

The current study sheds light on the negative effects of the environmental Pb water levels including the permissible value of 50 $\mu\text{g/L}$ that were reported by [Nachiyunde et al. \(2013\)](#) in Kabwe, Zambia on aquatic life. The permissible dissolved Pb in water Kabwe was comparable to the acute toxic criteria limit for dissolved Pb of 54.1 $\mu\text{g/L}$ at a hardness of 85 mg/L (as CaCO_3) as set by the United States Environmental Protection Agency (USEPA) as recently reported by [DeForest et al., \(2017\)](#). However, regulatory institutions around the globe seem not to have common criteria for dissolved Pb in water to ensure good water quality to support aquatic life ([Li et al., 2019](#)). For instance, the action level for dissolved Pb in water delivered to users for public consumption according to the Comprehensive Environmental Response Compensation and Liability Act (CERCLA), USA is 15 $\mu\text{g/L}$ ([DeForest et al., 2017](#)).

The present study showed that varying Pb concentrations induced toxicity in zebrafish embryos between 1-100 $\mu\text{g/L}$ Pb. The deleterious effects of Pb exposure in the development and survival of zebrafish embryos were observed within environmentally relevant and regulatory Pb concentrations. Neuromuscular, cardiovascular and antioxidant system effects due to Pb exposure with similar to what has been reported in studies which had higher Pb concentrations ([Chen et al., 2012](#); [Zhao et al., 2019](#)) are discussed.

In developmental toxicity studies involving zebrafish embryos, mortality and hatching of embryos are widely regarded as endpoints that influence the overall survival rate of zebrafish (Hallare et al., 2006). In this study, none of the surviving embryos failed to hatch at 72 hpf across all the exposed groups, pre-hatched embryo mortalities were observed with highest concentration having the highest percentage of the mortalities. All the mortalities were recorded between 24 and 72 hpf, a feature that was consistent with reports that indicated that the early embryonic stages after fertilization are more vulnerable to metal intoxication (Jeziarska et al., 2009). The high permeability of the embryo membrane to metal ions and rapid organogenesis accounts for the high sensitivity and vulnerability of embryos during the early stages (Frayse et al., 2006). The permeability of the embryo membrane or chorion imply that the concentration Pb in the exposure solution influences the level Pb ion that accumulates in the body of the embryo to cause residual toxicity. Although the mechanisms of accumulated Pb-induced toxicity in embryos may be complex, an indirect connection between survival rates and neuromuscular effects was observed in the present study suggesting Pb cumulative toxicity. For instance 50 and 100 µg/L Pb concentrations with larvae that had neuromuscular toxicity, the former that exhibited mild effects had 68% survival rate, and the latter with much pronounced effects had 52% survival rate.

Muscular twitching has been suggested as neuromuscular toxic effects of Pb (Van Den Avyle et al., 1989). In the present study, the 100 µg/L Pb group which had decreased embryo activity at 24 hpf had pronounced muscular twitching at 72 hpf suggestive of early onset of Pb-induced neuromuscular toxicity. Although altered spontaneous movements in zebrafish larvae have been reported at 1000 µg/L Pb (Chen et al., 2012), the neuromuscular toxicity being reported in our study at 50 µg/L Pb suggest that

zebrafish embryos may be vulnerable to Pb-induced toxicity even in low Pb concentrations. The lack of the mature functional blood-brain barrier and rapid growth that incorporates Pb into cellular processes due to the high affinity of Pb^{2+} for Ca^{2+} dependent processes could be among the reasons low Pb concentrations in the present study elicited similar effects with studies that employed high Pb concentrations.

Lead induced cardiovascular toxicity was observed at 100 $\mu\text{g/L}$ Pb level of exposure characterized by reduced heart rate and increased blood flow in the present study. These findings were in tandem with previously reported cardiovascular toxicity in zebrafish embryos exposed to much higher concentrations of Pb (3000, 6000 and 12 000 $\mu\text{g/L}$ Pb) than our study (Yin et al., 2017). The reduced heart rate may be linked to the antagonism between Pb and calcium and the ability of Pb^{2+} ions to block calcium channels causing impaired calcium availability for optimum heart function (Mattos et al., 2017). The increase in blood flow observed in the 100 $\mu\text{g/L}$ Pb exposed group corroborated evidence indicating that Pb may induce increased blood flow. Although the mechanisms that accounting for Pb-induced blood flow increase are complex, impaired nitric oxide system, inhibition of endothelial cell growth, oxidative stress and altered cellular Ca^{2+} tracking have been implicated in Pb-induced hypertension (ATSDR, 2019). Moreover, the Pb concentration at which the cardiovascular toxicity observed in the present study was close to the upper limit of the environmentally reported water Pb concentration (94 $\mu\text{g/L}$ Pb) in Kabwe, Zambia (Nachiyunde, 2013).

Lead induced oxidative stress is one of the mechanisms by which Pb exposure induces toxicity in animals including fish (Kim and Kang, 2017). Therefore, mRNA expression of the antioxidant system and related genes were analyzed in the present study. The major

antioxidant gene coding enzymes were dysregulated in the Pb exposed zebrafish larvae with the groups that showed cardiovascular and neuromuscular toxicity having enhanced dysregulation in some cases. For instance, catalase enzyme mRNA expression was downregulated in all exposed groups. Catalase enzyme gene downregulation is associated with the inhibition of catalase enzyme activity (Craig et al., 2007). Catalase offers cellular protection against oxidative damage due to its involvement in the detoxification and elimination of hydrogen peroxide generated from reactive oxygen species (Stancová et al., 2015). Hence the downregulation of the catalase transcripts observed in the current study could be a reflection of an overwhelmed antioxidant response due to excessive generation of non-radical hydrogen peroxide molecules. The other two key antioxidant encoding genes, heme oxygenase- 1 (*HO-1*) and glutathione-S-transferase (*GST*) mRNA expression levels were upregulated with the latter following a Pb concentration response. The *GST* enzyme is a vital metabolic and antioxidant enzyme whose gene expression pattern suggests an enhanced response in an attempt to decrease Pb toxicity through conjugation of Pb to glutathione to facilitate elimination (He et al., 2011).

It has been elucidated that ROS generation is linked to the activity of the mitochondria through the electron transport chain (Flora et al., 2012). Thus, we analyzed vital mitochondrial cytochrome c oxidase I (*CoxI*) and uncoupling protein-2 (*Ucp-2*) enzymes gene expression levels. *CoxI* is a terminal electron acceptor of the mitochondrial respiratory chain related to the generation of superoxide anion and *Ucp-2* mitigates against the impact of mitochondrial superoxide anion (Sohal et al., 2008). In the present study, *CoxI* was upregulated across all exposed groups without an accompanying upregulation of *Ucp-2* mRNA expression. This was in contrast to previous findings in larval zebrafish exposed to acute environmentally relevant Pb levels (Kataba et al., 2020).

The difference observed in the expression of the *CoxI* and *Ucp-2* mRNA expression in [Kataba et al. \(2020\)](#) and the current study may be due to the differences in the Pb exposure duration as well as the age of the zebrafish at the beginning of the exposure. The younger, the fish, the more susceptible it is to waterborne toxicants ([Jeziarska et al., 2009](#)). This upregulation of *CoxI* without the concurrent upregulation of *Ucp-2* may explain the Pb-induced toxic effects observed in zebrafish embryos as *Ucp-2* activity tend to neutralize the impact of *CoxI* ([Kataba et al., 2020](#)). Moreover, increased superoxide anion and disruption of the *Ucp-2* levels have been linked to neuronal oxidative damages and sustained neuronal oxidative damage causes neuronal apoptosis ([Wu et al., 2010](#)). The neuronal apoptosis may manifest as neurotoxicity a probable mechanism behind the neuromuscular toxicity observed in the present study.

In aquatic organisms, oxidative stress has been linked to enhanced apoptosis ([Livingstone, 2001](#)). In the current study, we used the B-cell lymphoma 2 (*Bcl-2*) and tumour protein p53 (*TP53*) genes as transcriptional markers related to the apoptosis signalling processes ([Jin et al., 2011](#)). The *Bcl-2* mRNA expression levels in Pb exposure concentrations less than 50 µg/L (1-25 µg/L Pb) were **upregulated** and downregulated in Pb exposure concentrations equal to or greater than 50 µg/L. The upregulation of the *Bcl-2* mRNA expression in Pb concentrations less than 50 µg/L Pb could be an indirect compensatory and protective response against Pb-induced apoptotic elements ([Bonneau et al., 2013](#)). Whereas, the downregulation of *Bcl-2* mRNA expression levels at 50 µg/L Pb could be a reflection of an exhausted *Bcl-2* protein compensatory mechanism due to the high presence of Pb ([Jin et al., 2010](#)). On the other hand, Pb exposure seemed to have triggered the downregulation of *TP53* mRNA expression across exposed groups. The *Bcl-2* gene encodes for protein that is a member of the *Bcl-2* family that regulates, suppresses

and prevents aberrant apoptosis (Jin et al., 2010). The prevention is achieved by neutralization of the pro-apoptotic proteins (p53) by Bcl-2 proteins (Pyati et al., 2007). Notwithstanding the striking non-Pb concentration-dependent mRNA expression of the antiapoptosis and proapoptosis genes, the mRNA expression pattern of these genes suggested that apoptosis could have been among the possible contributors to the neuromuscular twitching observed. Moreover, the significant downregulation of anti-apoptotic and pro-apoptotic gene transcripts at 50 µg/L Pb and non-significant downregulation at 100 µg/L Pb reflects a hormetic response, a feature that has been reported in larval zebrafish (Kataba et al., 2020). Overall, the study has demonstrated the deleterious effects of the water Pb levels in Kabwe on zebrafish embryos **and larvae** including cardiovascular and neuromuscular toxicity.

There are limitations to our study. The lack of accompanying enzymatic assays and non-enzymatic such as lipid peroxidase or protein carbonyl compound analyses and antioxidant tissue levels analyses are among the major limitations to support the gene expression **effects** thus observed (Mccarthy and Smyth, 2009). The non-retrieval of the embryo activity data at 24 hpf for 25 and 50 µg/L Pb poses another limitation on the comparison of the embryo activity and neuromuscular toxicity at these levels of exposure. Furthermore, in the light of anti-apoptotic and pro-apoptotic genes, mRNA dysregulation observed without accompanying apoptosis assays such as acridine orange staining limits the interpretation of our results. Notwithstanding, the present study has demonstrated that environmentally relevant Pb levels could affect the overall survival rates of zebrafish embryos through or accompanied by cardiovascular, neuromuscular, and antioxidant system aberrations.

5. Conclusions

Lead dissolved in water poses a threat to aquatic life even in lowest quantifiable amounts. Water Pb concentrations that are below and within the “permissible limit” (10 to 50 µg/L Pb) could be detrimental to zebrafish life especially at early developmental stages as evidenced by embryonic coagulation linked mortalities. Furthermore, our FET test concentrations provided additional insights on the Kabwe Pb water concentrations. Lead water concentrations of 50 to above 100 µg/L could even be more detrimental to developing fish embryos with a myriad Pb linked toxicities. Embryonic activity aberrations, cardiovascular toxicity (reduced heart rate, increased blood flow activity), oxidative stress system imbalance, antiapoptotic and proapoptotic balance and the neuromuscular toxicity (muscle twitching) are among the deleterious effects of environmentally relevant Pb levels. Further investigations on the impact of environmentally relevant water Pb concentrations and the permissible (regulatory) water Pb levels on reproduction, development and health of locally available fish species are needed.

6. Ethical statement

All experimental procedures were done with the due approval by the AnimCare animal research ethics committee (ethics approval number: NWU-00269-16-A5) at the North-West University. All animals were maintained, and procedures carried out in adherence with the code of ethics in research, training and testing of drugs in South Africa and complied with national legislation (NHREC reg. number AREC-130913-015).

7. Authors contribution

Andrew Kataba: Conceptualization; Data curation; Formal analysis; Methodology; Writing - original draft; Writing - review & editing. **Tarryn Lee Botha, Yared B. Yohannes:** Validation, Methodology, Software, Writing - review & editing. **Shouta M.M. Nakayama** and **Yoshinori Ikenaka:** Funding acquisition; Resources; Writing - review & editing. **Victor Wepener, Mayumi Ishizuka:** Funding acquisition Resources; Supervision; Writing - review & editing.

8. Declaration of Competing Interest

The authors declare that they have no conflict of interest relating to the work presented in this manuscript.

9. Acknowledgements

The chemical analyses were technically supported by Mr. Takahiro Ichise and Ms. Nagisa Hirano at Hokkaido University, Faculty of Veterinary Medicine and; Dr Hannes Erasmus at the North-West University, South Africa. This work was supported by the Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan awarded to M. Ishizuka (No. 16H0177906, 18K1984708, and 18KK028708), Y. Ikenaka (18H0413208), and S.M.M. Nakayama (No. 17KK0009, 20K20633). The work was also supported by the Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan awarded to M. Ishizuka (No. 16H0177906, 18K1984708, 18KK028708, 21H02351), Ikenaka (18H0413208), and S.M.M. Nakayama (No. 17KK0009 and 20K20633). This work was also supported by the foundation of JSPS Bilateral Open Partnership Joint Research Projects (JPJSBP120209902; SMMN). This work was also supported by JSPS CORE to CORE program (MI), The Japan Prize Foundation; and Hokkaido University's SOUSEI Support Program for Young Researchers in FY2020 (SMMN); Hokkaido University Specific Research Projects (MI). This research was also supported by JST/JICA, SATREPS (Science and Technology Research Partnership for Sustainable Development; No. JPMJSA1501) and aXis (Accelerating Social Implementation for SDGs Achievement; No. JPMJAS2001) funded by JST as well as Program for supporting introduction of the new sharing system (JPMXS0420100619). and the Hokkaido University Faculty of Veterinary Medicine Wise/Leading Travel and Subsistence grant.

511 Bioassays were conducted in the National Aquatic Bioassay Facility at North-West
512 University, South Africa (NRF Grant UID99024). This is publication number 555 of the
513 Water Research Group, North-West University.

514

References

- ATSDR, A. for T.S. and D.R., 2019. Toxicological Profile for Lead (Draft for Public Comment). ATSDR's Toxicol. Profiles. https://doi.org/10.1201/9781420061888_ch106
- Bonneau, B., Prudent, J., Popgeorgiev, N., Gillet, G., 2013. Non-apoptotic roles of Bcl-2 family: The calcium connection. *Biochim. Biophys. Acta - Mol. Cell Res.* 1833, 1755–1765. <https://doi.org/10.1016/j.bbamcr.2013.01.021>
- Bose-O'Reilly, S., Yabe, J., Makumba, J., Schutzmeier, P., Ericson, B., Caravanos, J., 2017. Lead intoxicated children in Kabwe, Zambia. *Environ. Res.* 0–1. <https://doi.org/10.1016/j.envres.2017.10.024>
- Botha, T.L., James, T.E., Wepener, V., 2015. Comparative Aquatic Toxicity of Gold Nanoparticles and Ionic Gold Using a Species Sensitivity Distribution Approach. *J. Nanomater.* 2015. <https://doi.org/10.1155/2015/986902>
- Busquet, F., Strecker, R., Rawlings, J.M., Belanger, S.E., Braunbeck, T., Carr, G.J., Cenijn, P., Fochtman, P., Gourmelon, A., Hübler, N., Kleensang, A., Knöbel, M., Kussatz, C., Legler, J., Lillicrap, A., Martínez-jerónimo, F., Polleichtner, C., Rzodeczko, H., Salinas, E., Schneider, K.E., Scholz, S., Brandhof, E. Van Den, Ven, L.T.M. Van Der, Walter-rohde, S., Weigt, S., Witters, H., Halder, M., 2014. OECD validation study to assess intra- and inter-laboratory reproducibility of the zebrafish embryo toxicity test for acute aquatic toxicity testing. *Regul. Toxicol. Pharmacol.* 69, 496–511. <https://doi.org/10.1016/j.yrtph.2014.05.018>
- Campel, I., 2007. Chi-squared and Fisher–Irwin tests of two-by-two tables with small sample recommendations. *Stat. Med.* 26, 3661–3675. <https://doi.org/10.1002/sim.2832>
- Chen, J., Chen, Y., Liu, W., Bai, C., Liu, X., Liu, K., Li, R., Zhu, J.H., Huang, C., 2012. Developmental lead acetate exposure induces embryonic toxicity and memory deficit in adult zebrafish. *Neurotoxicol. Teratol.* 34, 581–586. <https://doi.org/10.1016/j.ntt.2012.09.001>
- Craig, P.M., Wood, C.M., McClelland, G.B., 2007. Oxidative stress response and gene expression with acute copper exposure in zebrafish (*Danio rerio*). *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* 5, 1882–1892. <https://doi.org/10.1152/ajpregu.00383.2007>
- DeForest, D.K., Santore, R.C., Ryan, A.C., Church, B.G., Chowdry, J.M., Brix, Kevin, V., 2017. Development of biotic ligand model-based freshwater aquatic life criteria for lead following us environmental protection agency guidelines. *Environ. Toxicol. Chem.* 36, 2965–2973. <https://doi.org/10.1002/etc.3861>
- Flora, G., Gupta, D., Tiwari, A., 2012. Toxicity of lead: A review with recent updates. *Interdiscip. Toxicol.* 5, 47–58. <https://doi.org/10.2478/v10102-012-0009-2>
- Fraysse, B., Mons, R., Garric, J., 2006. Development of a zebrafish 4-day embryo-larval bioassay to assess toxicity of chemicals. *Ecotoxicol. Environ. Saf.* 63, 253–267.

556 <https://doi.org/10.1016/j.ecoenv.2004.10.015>

557 Hailegnaw, B., Kirmayer, S., Edri, E., Hodes, G., Cahen, D., 2015. Rain on
 558 Methylammonium Lead Iodide Based Perovskites: Possible Environmental Effects
 559 of Perovskite Solar Cells. *J. Phys. Chem.*
 560 <https://doi.org/10.1021/acs.jpcclett.5b00504>

561 Hallare, A., Nagel, K., Köhler, H.R., Triebkorn, R., 2006. Comparative embryotoxicity
 562 and proteotoxicity of three carrier solvents to zebrafish (*Danio rerio*) embryos.
 563 *Ecotoxicol. Environ. Saf.* 63, 378–388.
 564 <https://doi.org/10.1016/j.ecoenv.2005.07.006>

565 He, X., Nie, X., Wang, Z., Cheng, Z., Li, K., Li, G., Hung Wong, M., Liang, X., Tsui,
 566 M.T.K., 2011. Assessment of typical pollutants in waterborne by combining active
 567 biomonitoring and integrated biomarkers response. *Chemosphere* 84, 1422–1431.
 568 <https://doi.org/10.1016/j.chemosphere.2011.04.054>

569 Hill, A.J., Teraoka, H., Heideman, W., Peterson, R.E., 2005. Zebrafish as a model
 570 vertebrate for investigating chemical toxicity. *Toxicol. Sci.* 86, 6–19.
 571 <https://doi.org/10.1093/toxsci/kfi110>

572 Jezierska, B., Ługowska, K., Witeska, M., 2009. The effects of heavy metals on
 573 embryonic development of fish (a review). *Fish Physiol. Biochem.* 35, 625–640.
 574 <https://doi.org/10.1007/s10695-008-9284-4>

575 Jin, Y., Zhang, X., Shu, L., Chen, L., Sun, L., Qian, H., Liu, W., Fu, Z., 2010. Oxidative
 576 stress response and gene expression with atrazine exposure in adult female
 577 zebrafish (*Danio rerio*). *Chemosphere* 78, 846–852.
 578 <https://doi.org/10.1016/j.chemosphere.2009.11.044>

579 Jin, Y., Zheng, S., Pu, Y., Shu, L., Sun, L., Liu, W., Fu, Z., 2011. Cypermethrin has the
 580 potential to induce hepatic oxidative stress, DNA damage and apoptosis in adult
 581 zebrafish (*Danio rerio*). *Chemosphere* 82, 398–404.
 582 <https://doi.org/10.1016/j.chemosphere.2010.09.072>

583 Kataba, A., Botha, T.L., Nakayama, S.M.M., Yohannes, Y.B., Ikenaka, Y., Wepener,
 584 V., Ishizuka, M., 2020. Acute exposure to environmentally relevant lead levels
 585 induces oxidative stress and neurobehavioral alterations in larval zebrafish (*Danio*
 586 *rerio*). *Aquat. Toxicol.* 227, 105607.
 587 <https://doi.org/10.1016/j.aquatox.2020.105607>

588 Kim, J., Kang, J., 2017. Chemosphere Effects of sub-chronic exposure to lead (Pb) and
 589 ascorbic acid in juvenile rockfish: Antioxidant responses, MT gene expression,
 590 and neurotransmitters. *Chemosphere* 171, 520–527.
 591 <https://doi.org/10.1016/j.chemosphere.2016.12.094>

592 Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., Schilling, T.F., 1995. Stages
 593 of embryonic development of the zebrafish. *Dev. Dyn.* 203, 253–310.
 594 <https://doi.org/10.1002/aja.1002030302>

595 Komárek, M., Chrastný, V., Mihaljevi, M., 2008. Lead isotopes in environmental
 596 sciences: A review. *Environ. Int.* 34, 562–577.

597 <https://doi.org/10.1016/j.envint.2007.10.005>

598 Li, X., Kong, H., Ji, X., Gao, Y., Jin, M., 2019. Zebrafish behavioral phenomics applied
599 for phenotyping aquatic neurotoxicity induced by lead contaminants of
600 environmentally relevant level. *Chemosphere* 224, 445–454.
601 <https://doi.org/10.1016/j.chemosphere.2019.02.174>

602 Livingstone, D.R., 2001. Contaminant-stimulated reactive oxygen species production
603 and oxidative damage in aquatic organisms. *Mar. Pollut. Bull.* 42, 656–666.
604 [https://doi.org/10.1016/S0025-326X\(01\)00060-1](https://doi.org/10.1016/S0025-326X(01)00060-1)

605 Maiti, A.K., Saha, N.C., Paul, G., 2010. Effect of lead on oxidative stress, Na⁺+K⁺
606 +ATPase activity and mitochondrial electron transport chain activity of the brain of
607 *clarias batrachus* L. *Bull. Environ. Contam. Toxicol.* 84, 672–676.
608 <https://doi.org/10.1007/s00128-010-9997-9>

609 Mattos, G.F. De, Costa, C., Savio, F., Alonso, M., Nicolson, G.L., 2017. Lead
610 poisoning : acute exposure of the heart to lead ions promotes changes in cardiac
611 function and Cav1.2 ion channels 807–825. [https://doi.org/10.1007/s12551-017-](https://doi.org/10.1007/s12551-017-0303-5)
612 0303-5

613 McCarthy, D.J., Smyth, G.K., 2009. Testing significance relative to a fold-change
614 threshold is a TREAT. *Bioinformatics* 25, 765–771.
615 <https://doi.org/10.1093/bioinformatics/btp053>

616 Nachiyunde, K., Ikeda, H., Okuda, T., Nishijima, W., 2013. Assessment of Dissolved
617 Heavy Metal Pollution in Five Provinces of Zambia. *J. Environ. Prot. (Irvine, Calif.)* 04, 80–85. <https://doi.org/10.4236/jep.2013.41b015>

619 Nakata, H., Nakayama, S.M.M., Yabe, J., Liazambi, A., Mizukawa, H., Sobhy, W.,
620 Ikenaka, Y., Ishizuka, M., 2016. Reliability of stable Pb isotopes to identify Pb
621 sources and verifying biological fractionation of Pb isotopes in goats and chickens.
622 *Environ. Pollut.* 208, 395–403. <https://doi.org/10.1016/j.envpol.2015.10.006>

623 Nakayama, Shouta M M, Ikenaka, Y., Hamada, K., Muzandu, K., Choongo, K.,
624 Teraoka, H., Mizuno, N., Ishizuka, M., 2011. Metal and metalloid contamination in
625 roadside soil and wild rats around a Pb-Zn mine in Kabwe, Zambia. *Environ.*
626 *Pollut.* 159, 175–181. <https://doi.org/10.1016/j.envpol.2010.09.007>

627 Nakayama, Shouta M.M., Ikenaka, Y., Hamada, K., Muzandu, K., Choongo, K.,
628 Teraoka, H., Mizuno, N., Ishizuka, M., 2011. Metal and metalloid contamination in
629 roadside soil and wild rats around a Pb-Zn mine in Kabwe, Zambia. *Environ.*
630 *Pollut.* 159, 175–181. <https://doi.org/10.1016/j.envpol.2010.09.007>

631 Pyati, U.J., Look, A.T., Hammerschmidt, M., 2007. Zebrafish as a powerful vertebrate
632 model system for in vivo studies of cell death. *Semin. Cancer Biol.* 17, 154–165.
633 <https://doi.org/10.1016/j.semcancer.2006.11.007>

634 Sfakianakis, D.G., Renieri, E., Kentouri, M., Tsatsakis, A.M., 2015. Effect of heavy
635 metals on fish larvae deformities: A review. *Environ. Res.* 137, 246–255.
636 <https://doi.org/10.1016/j.envres.2014.12.014>

637 Sohal, R.S., Toroser, D., Brégère, C., Mockett, R.J., Orr, W.C., 2008. Age-related
638 decrease in expression of mitochondrial DNA encoded subunits of cytochrome c
639 oxidase in *Drosophila melanogaster*. *Mech. Ageing Dev.* 129, 558–561.
640 <https://doi.org/10.1016/j.mad.2008.04.006>

641 Stancová, V., Ziková, A., Svobodová, Z., Kloas, W., 2015. Effects of the non-steroidal
642 anti-inflammatory drug (NSAID) naproxen on gene expression of antioxidant
643 enzymes in zebrafish (*Danio rerio*). *Environ. Toxicol. Pharmacol.* 40, 343–348.
644 <https://doi.org/10.1016/j.etap.2015.07.009>

645 Toyomaki, H., Yabe, J., Nakayama, S.M.M., Yohannes, Y.B., Muzandu, K., Liazambi,
646 A., Ikenaka, Y., Kuritani, T., Nakagawa, M., Ishizuka, M., 2020. Factors
647 associated with lead (Pb) exposure on dogs around a Pb mining area, Kabwe,
648 Zambia. *Chemosphere* 247, 125884.
649 <https://doi.org/10.1016/j.chemosphere.2020.125884>

650 Van Den Avyle, M.J., Garvick, S.J., Blazer, V.S., Hamilton, S.J., Brumbaugh, W.G.,
651 1989. Skeletal deformities in smallmouth bass, *Micropterus dolomieu*, from
652 southern Appalachian reservoirs. *Arch. Environ. Contam. Toxicol.* 18, 688–696.
653 <https://doi.org/10.1007/BF01225007>

654 Wu, Z., Zhao, Y., Zhao, B., 2010. Superoxide anion, uncoupling proteins and
655 Alzheimer's disease. *J. Clin. Biochem. Nutr.* 46, 187–194.
656 <https://doi.org/10.3164/jcbn.09-104-2>

657 Yabe, J., Nakayama, S.M., Nakata, H., Toyomaki, H., Yohannes, Y.B., Muzandu, K.,
658 Kataba, A., Zyambo, G., Hiwatari, M., Narita, D., Yamada, D., Hangoma, P.,
659 Munyinda, N.S., Mufune, T., Ikenaka, Y., Choongo, K., Ishizuka, M., 2020.
660 Current trends of blood lead levels, distribution patterns and exposure variations
661 among household members in Kabwe, Zambia. *Chemosphere* 243, 125412.
662 <https://doi.org/10.1016/j.chemosphere.2019.125412>

663 Yabe, J., Nakayama, S.M.M., Ikenaka, Y., Muzandu, K., Choongo, K., Mainda, G.,
664 Kabeta, M., Ishizuka, M., Umemura, T., 2013. Metal distribution in tissues of free-
665 range chickens near a lead-zinc mine in Kabwe, Zambia. *Environ. Toxicol. Chem.*
666 32, 189–192. <https://doi.org/10.1002/etc.2029>

667 Yabe, J., Nakayama, S.M.M., Ikenaka, Y., Muzandu, K., Ishizuka, M., Umemura, T.,
668 2011. Uptake of lead, cadmium, and other metals in the liver and kidneys of cattle
669 near a lead-zinc mine in Kabwe, Zambia. *Environ. Toxicol. Chem.* 30, 1892–1897.
670 <https://doi.org/10.1002/etc.580>

671 Yabe, J., Nakayama, S.M.M., Ikenaka, Y., Yohannes, Y.B., Bortey-sam, N., Nketani,
672 A., Ntapisha, J., Mizukawa, H., Umemura, T., Ishizuka, M., 2018. *Chemosphere*
673 Lead and cadmium excretion in feces and urine of children from polluted
674 townships near a lead-zinc mine in Kabwe, Zambia. *Chemosphere* 202, 48–55.
675 <https://doi.org/10.1016/j.chemosphere.2018.03.079>

676 Yabe, J., Nakayama, S.M.M., Ikenaka, Y., Yohannes, Y.B., Bortey-Sam, N., Oroszlany,
677 B., Muzandu, K., Choongo, K., Kabalo, A.N., Ntapisha, J., Mweene, A.,
678 Umemura, T., Ishizuka, M., 2015. Lead poisoning in children from townships in

679 the vicinity of a lead-zinc mine in Kabwe, Zambia. *Chemosphere* 119, 941–947.
680 <https://doi.org/10.1016/j.chemosphere.2014.09.028>

681 Yin, J., Wang, A.P., Li, W.F., Shi, R., Jin, H.T., Wei, J.F., 2017. Sensitive biomarkers
682 identification for differentiating Cd and Pb induced toxicity on zebrafish embryos.
683 *Environ. Toxicol. Pharmacol.* 56, 340–349.
684 <https://doi.org/10.1016/j.etap.2017.10.010>

685 Zhao, J., Zhang, Q., Zhang, B., Xu, T., Yin, D., Gu, W., Bai, J., 2019. Developmental
686 exposure to lead at environmentally relevant concentrations impaired
687 neurobehavior and NMDAR-dependent BDNF signaling in zebrafish larvae.
688 *Environ. Pollut.* 113627. <https://doi.org/10.1016/j.envpol.2019.113627>

689