



Title	Planned seedling production in the distribution area of <i>Saccharina japonica</i> : Sorus formation through the induction of sporophyte maturation and the culture of the seedlings produced
Author(s)	Yotsukura, Norishige; Seki, Shohei; Sasaki, Sachiko; Yoshida, Masanori
Citation	Aquaculture research, 53(2), 676-683 https://doi.org/10.1111/are.15612
Issue Date	2021-10-04
Doc URL	http://hdl.handle.net/2115/87876
Rights	This is the peer reviewed version of the following article: Yotsukura, N., Seki, S., Sasaki, S., & Yoshida, M. (2022). Planned seedling production in the distribution area of <i>Saccharina japonica</i> : Sorus formation through the induction of sporophyte maturation and the culture of the seedlings produced. <i>Aquaculture Research</i> , 53, 676–683., which has been published in final form at https://doi.org/10.1111/are.15612 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.
Type	article (author version)
File Information	are_15612.pdf



[Instructions for use](#)

1 Planned seedling production in the distribution area of *Saccharina japonica*: sorus formation
2 through the induction of sporophyte maturation and the culture of the seedlings produced
3

4 Running title: Planned seedling production of *Saccharina*

5

6 Norishige Yotsukura¹, Shohei Seki², Sachiko Sasaki³, Masanori Yoshida³

7

8 ¹Field Science Center for Northern Biosphere, Hokkaido University, Sapporo 060-0809, Japan

9 ²Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan

10 ³Fisheries Institute, Aomori Prefectural Industrial Technology Research Center, Aomori 039-
11 3381, Japan

12

13 Correspondence

14 Norishige Yotsukura, Field Science Center for Northern Biosphere, Hokkaido University,

15 Sapporo 060-0809, Japan

16

17 Abstract

18 Inconsistency in maturation of saccharinan kelp due to changes in the marine
19 environment is a problem in terms of securing seedlings for aquaculture. In Hokkaido,
20 the main producing area of *Saccharina japonica*, it has become a challenge to stably
21 secure the parental algae of the species for aquaculture. When maturation was induced
22 by short-day conditions in naturally collected *S. japonica* sporophytes as well as in the
23 sporophytes collected from them and cultivated, sorus formation began in
24 approximately three weeks and spread to 0.4–16.8% on the upper side area and 23.2–

25 41.1% on the underside area after about five weeks, with the highest rates, 23.2–59.5%,
26 observed around the base part of the underside. Sorus formation began from the base
27 and middle parts in the top-intact thalli and from the marginal part in the top-cut thalli.
28 The released zoospore germlings grew steadily as aquaculture seedlings, and the blade
29 width and wet weight of the seedlings collected in August and subjected to mariculture
30 in November were significantly larger after approximately five and a half months
31 compared to those of thalli cultivated using a conventional method (i.e., thalli collected
32 just prior to the mariculture). Seedling production through maturation induction in
33 sporophytes would be a feasible option for seedling production facilities and would be
34 helpful for stably producing seedlings without the effect of the natural environment.

35

36 Keywords: aquaculture, maturation induction, *Saccharina japonica*, seedling
37 production, sorus formation

38

39 INTRODUCTION

40 *Saccharina japonica* (Areschoug) C.E. Lane, C. Mays, Druehl & G.W. Saunders grows along
41 the northern coasts of Japan and surrounding waters (Yotsukura et al., 2008); in Hokkaido, the
42 main producing area of the species, it has been harvested mainly as foodstuff for more than
43 1000 years (Oishi, 1987). However, full-scale aquaculture of *S. japonica* in Hokkaido only
44 began in 1966, and its yield, which has become increasingly stabilized thanks to advances in
45 technology, has compensated for the progressive decline in wild catches of the species.

46 Generally, this species is cultivated using a method in which zoospores released from the
47 sporophytes (parental algae) are attached to seedling threads for seedling production in indoor
48 tanks, and the seedling are then cultivated in coastal area (Kawashima, 1984; Su et al., 2017). In

49 Japan, to produce the seedlings, an important process for aquaculture, high-quality wild mature
50 *S. japonica* sporophytes are collected from the periphery of the aquaculture sites in autumn as
51 parental algae (Kawashima, 1984). However, in recent years, disruption in maturation of kelp
52 sporophytes due to changes in the marine environment has become serious. For example, in the
53 Oyasu district of Hakodate City, Hokkaido, the seedlings were formerly produced in mid-
54 September but were only produced on September 29 in 2019 and October 11 in 2020, indicating
55 the increasing difficulty in stable seedling production. This would be related to the slow decline
56 in water temperature during that season in recent years.

57 *Saccharina japonica* has also been cultivated in China, Korea, and Russia (Sohn, 1998;
58 Kawai et al., 2015; Shan et al., 2017; Hwang et al., 2019). Although not native to Chinese
59 waters, *S. japonica* aquaculture began in the mid-1950s using local populations originated from
60 Japan, and yield recently reached approximately 1.4 Mt (dry weight), well above Japan's yield
61 (4319 t in 2018), making China the largest producer (Hwang et al., 2019). In China, a "summer
62 sporeling technique" was developed in which seedling threads are made using farmed
63 sporophytes that mature in early summer, and now more than 90% of the *S. japonica* products
64 are produced using this technique (Tseng et al., 1955; Shan et al., 2011). This aquaculture
65 technique, which allows seedling production in summer, has various advantages over autumn
66 production techniques (Tseng et al., 1955; Shan et al., 2011) and is expected to increase the
67 quality of cultivated kelp products, which are generally lower in industrial value compared to
68 wild counterparts, since more time can be spent on the culture of seedlings, particularly in
69 indoor tanks and the sea. This technique is expected to also become widespread in Japan. In
70 China, the precocious sporophytes obtained from aquaculture are used as parental algae, which
71 allows stable production of seedlings (Su et al., 2017). However, along the Japanese coasts, it is
72 difficult to collect mature sporophytes of the species, which release zoospores most actively in

73 early summer (Kawashima, 2012). To obtain parental algae in this season, the zoospores need to
74 be obtained by artificially inducing maturation in the sporophytes.

75 It has been reported that maturation in sporophytes of *Saccharina* kelp can be induced by
76 removing the meristematic tissue followed by cultivation under favorable conditions (short-day
77 and dark conditions, water temperature around 10°C, and high nutrient salt concentration)
78 (Luning, 1988; Nimura and Mizuta, 2002a; Skriptsova and Titlyanov, 2003; Pang and Luning,
79 2004; Forbord et al., 2012; Boderskov et al., 2021). In *S. latissima*, a species widely distributed
80 around the world, sorus formation and zoospore release were confirmed after a six-week
81 cultivation of sporophytes, collected in August in Norway, under low-light and short-day
82 conditions. When zoospores derived from the sori formed during the tank-cultivation of fertile
83 sporophytes, obtained in December, under running seawater and low-light/short-day conditions
84 were pre-cultured in a hatchery for 42 days, germlings grew notably well in subsequent
85 deployment (Forbord et al., 2020). In regard to the sporophytes of *S. japonica*, a species found
86 in limited sea areas and is an important ingredient in Japanese food, it has found in a Chinese
87 cultivar and a Chinese hybrid cultivar that sorus formation was induced under short-day
88 conditions in sporophytes from which a blade part, not adjacent to the meristematic tissue, was
89 removed and that in the latter, the release of zoospores was also confirmed (Su et al., 2020). In
90 contrast, the only study conducted on maturation induction in sporophytes of Japanese *S.*
91 *japonica* using wild thalli as parental algae for aquaculture, reported that sori were formed in
92 young thalli (thalli not subjected to artificial removal of parts except for a marker bore) after
93 approximately one month of cultivation at 5–20°C under a short-day condition (light 9 h: dark
94 15 h), and at 15°C under a day-neutral condition, in a tank through which filtered seawater
95 flowed (Kiri-hara et al., 2003). However, the release of zoospores from sori was not confirmed.
96 To incorporate seedling production technique in which maturation is artificially induced into

97 existing aquaculture production systems in the future, it will be necessary, using candidate thalli,
98 to investigate their detailed maturation mechanisms and the growth of seedlings obtained from
99 them. Therefore, in this study, in order to examine the possibility of producing seedlings by
100 inducing maturation in the sporophytes and ultimately achieving stable seedling production and
101 high-quality products, maturation was induced in an indoor tank in wild thalli and thalli
102 cultivated from their zoospores. The formation process of sori was examined, and the germlings
103 were grown from the zoospores obtained, thereby investigating the possibility of producing
104 aquaculture seedlings through maturation induction.

105 **MATERIALS AND METHODS**

106 Materials used for this experiment are eight thalli of *Saccharina japonica*: seven sporophytes
107 originated from 2-year-old wild kelp collected the year before and cultured in the sea and one
108 wild sporophyte (Table 1). Within them, the four were collected in Hakodate City, on July, 2018
109 (Nos. 1 & 2, with sori formed on the top part; and Nos. 3 & 4, without sori) and immediately
110 transported to the Fisheries Institute, Aomori Prefectural Industrial Technology Research Center
111 in a refrigerated condition after seawater was wiped off their surfaces. The arrived sporophytes
112 were bored using a cork borer 15 cm above the blade base and cultivated in a 1400-l tank
113 (internal dimensions: 100 cm × 300 cm × 50 cm) inside a land facility, taking into consideration
114 the results of a preceding study (Kiriwara et al., 2003). The seawater used was pumped from the
115 nearby sea, filtered, adjusted to 15 °C, and flushed at 500 l/h. The temperature of the seawater in
116 the tank was recorded at 10-min intervals. The tank was covered with a blackout curtain, and a
117 fluorescent light was placed above the tank so that the photon flux density below at the water
118 surface was 117 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (short-day conditions of 9/15 h of light/dark). An ultra-miniature
119 data recorder (MDS-MkV, Alec Electronics Co., Ltd.) was used to measure the water
120 temperature and the light intensity. The cultivation was continued from July 25 to August 28,

121 2018, and blade length and the position of the marker hole were measured periodically during
122 the period. In addition, the presence or absence of sori was investigated for the blade and three
123 equal parts of the blade (i.e., top, middle, and base parts), and the percentage of sori area to
124 blade area was calculated using the image measurement support tool TouchDeMeasure
125 (Enomoto et al. 2017).

126 Two samples (Nos. 5 & 6) collected on June, 2019, in Hakodate City, with their top ends
127 being cut so that their blade length became 600 cm; one sample (No. 7, an immature sporophyte
128 collected from wild kelp the year before last and cultivated in the sea) collected on June, 2019,
129 in Rishirifuji Town; and one sample (No. 8) collected on July, 2019, in Rausu Town (Table 1),
130 were cultivated in the tank and measurements taken using the same method as in the previous
131 year. The cultivation period was June 26 to August 16, 2019, for Nos. 5 and 6; June 26 to
132 August 5, 2019, for No. 7; and July 8 to August 5, 2019, for No. 8. Six wild mature sporophytes
133 (Nos. 9–14) collected on December 16, 2019, in the Oniwaki district of Rishirifuji Town were
134 used for comparison. After the measurements were taken, the samples were turned upside-down,
135 and cultivation continued.

136 Among the cultivated samples, No. 8 was taken out of the tank at the end of the cultivation
137 period, and a piece (ca. 15 cm × ca. 30cm) was then cut from the mature portion. After being
138 washed with filtered seawater, the piece was wrapped in a paper towel and stored overnight in a
139 constant-temperature chamber at 10 °C. The next morning, the piece was placed gently in a 2-l
140 Petri dish filled with sterile filtered seawater (10 °C) to induce the release of zoospores. The
141 zoospores released in the dishes were then observed under a microscope (150×) after 15–20 min
142 following the placement of the piece, and the piece was removed from the dish after confirming
143 the release of more than 10 zoospores within the field of view. After being filtered using
144 bleached cotton, the liquid in the dish was poured into a 12-l tank, and the seedlings were

145 collected using a seedling collector wrapped with 50 m of Cremona thread. This tank was then
146 transferred into a constant-temperature chamber maintained at 10 °C, allowed to stand for 24 h
147 in darkness, and then subjected to aeration culture under medium day conditions (12/12 h of
148 light/dark) with a photon density of 25–65 $\mu\text{mol m}^{-2} \text{s}^{-1}$. The culture medium was PESI medium
149 (Tatewaki, 1966), which was changed every week. Cultivated seedlings were subjected to
150 mariculture at an aquaculture facility located at the shore where the original samples were
151 collected (44°12'28.60"N, 145°20' 28.09"E) from November 7, 2019, to June 29, 2020. The
152 shape, length, and width of the blade, as well as the wet weight, were measured for the 10
153 samples for comparison with control samples attached to seedling threads just prior to the
154 mariculture (i.e., samples cultivated using the conventional method), and the means of each trait
155 were compared by t-test after confirming equal variance by F-test. Between November 19,
156 2019, and May 2, 2020, the aquaculture rope was dropped to near the seafloor (to a water depth
157 of about 6 m in a 30-m depth area) to avoid flow-ice damage and the growth of sporophytes was
158 suppressed.

159 This work does not contain any experiments involving animals performed by any of the
160 authors. And so, ethical approval is not required.

161 **RESULTS**

162 No noticeable increases in blade length and width were observed in the top-intact thalli
163 during the cultivation period. The blades of Nos. 2 and 4 and No. 8 were shortened about one
164 week and about two weeks after the start of cultivation, respectively, via the loss of the top part.
165 However, growth (distance from the blade base to the marker hole) during the cultivation period
166 was longest in No. 6 at 4 cm; followed by No. 7 at 3.5 cm; No. 3 at 1.5 cm; Nos. 1, 5 and 8 at 1
167 cm; No. 2 at 0.5 cm; and No. 4 at 0 cm.

168 As shown Table 2, among the first-year cultivated thalli where a small number of sori were

169 confirmed on both sides of their blades at the start of the experiment, in No. 1, noticeable
170 expansion of sori occurred after the 19th day of the cultivation, especially in the base part on the
171 upper side and the entire area on the underside (Figure 1-A). In No. 2, sori were observed in the
172 whole area of both sides after the 13th day, and sori expansion afterward was remarkable on the
173 underside, especially near the base. The percentages of sori area to blade area after the 19th, 26th,
174 and 33rd days follow: for the upper side of No. 1, 12.2%, 14.3%, and 16.8%, respectively; and
175 correspondingly for the underside of No. 1, 22.8%, 39.4%, and 41.1%; for the upper side of No.
176 2, 1.2%, 4.1%, and 4.9%; and the underside of No. 2, 22.8%, 32.0%, and 29.6%.

177 However, among the first-year cultivated samples that were immature at the beginning of the
178 experiment, in No. 3, sori appeared clearly at the base part on the underside on the 19th day and
179 also appeared slightly at the top part on the underside and the base part on the upper side. By the
180 26th day, sori had spread to the entire areas on both sides, except for the top and middle parts on
181 the upper side. In No. 4, sori formation started on the underside by the 19th day and spread to
182 the upper side by the 26th day. The percentages of sori area to blade area after the 19th, 26th, and
183 33rd days follow: for the upper side of No. 3, 1.3%, 7.7%, and 8.3%, respectively; and
184 correspondingly for the underside of No. 3, 4.4%, 37.7%, and 39.0%; for the upper side of No.
185 4, 0%, 4.2%, and 4.0%; and for the underside of No. 4, 11.2%, 26.9%, and 40.0%.

186 Among the first-year top-cut cultivated samples that were immature at the beginning of the
187 experiment, in No. 5, on the 21st day, sori were confirmed near the cut site on the underside and
188 also on the upper side to a lesser extent (Figure 1-B), and by the 28th day, spread to both sides.
189 In No. 6, on the 21st day, sori were confirmed in the whole area on the underside and near the
190 cut site on the upper side and, by the 28th day, spread to the whole area on the upper side. The
191 percentages of sori area to blade area after the 21st, 28th, and 37th days follow: for the upper side
192 of No. 5, 0.7%, 1.6%, and 0.8%, respectively; and correspondingly for the underside of No. 5,

193 4.1%, 20.8%, and 25.1%; for the upper side of No. 6, 1.7%, 2.9%, and 3.1%; and for the
194 underside of No. 6, 7.7%, 16.1%, and 23.2%. In No. 7 (second-year cultivated thallus), sori, on
195 the 28th day, were confirmed in the middle and base parts on the underside and the top and
196 middle parts on the upper side and, by the 37th day, spread to the whole area on the underside
197 and the top and middle parts on the upper side (Figure 1-C). The percentages of sori area to
198 blade area after the 21st, 28th, and 37th days follow: for the upper side, 0%, 0.1%, and 0.4%,
199 respectively; and for the underside, 0%, 21.6%, and 28.8%, respectively. In No. 8 (second-year
200 wild thallus), sori were confirmed in the middle and base parts on the underside on the 16th day
201 and also in the middle and base parts on the upper side on the 25th day (Figure 1-D, E). The
202 percentages of sori area to blade area on the 16th and 25th days follow: for the upper side, 7.6%
203 and 11.7%, respectively; and for the underside, 24.6% and 37.2%, respectively. Partial damage
204 of blade, especially in the base part, occurred in most samples and a slight decrease in the
205 percentages of sori are to blade area was seen. For the control samples (wild thalli), the
206 percentages of sori area to blade area follow: for the upper side of Nos. 9–14, 0%, 0%, 0%,
207 3.6%, 6.6%, and 7.6%, respectively; and correspondingly for the underside, 50.0%, 50.7%,
208 77.7%, 31.3%, 31.7%, and 70.9%.

209 In the mariculture experiment, there were no significant differences in blade length between
210 the experimental and control samples ($p = 0.05$), but there were significant differences in blade
211 width and wet weight ($p = 0.002$ and $p = 0.003$, respectively) (Figure 2).

212 **DISCUSSION**

213 In this study, under the short-day conditions set based on the results of a preceding study
214 (Kirihara et al., 2003), in all experimental samples, sori were formed in approximately three
215 weeks after the start of maturation induction and spread to the whole area in approximately five
216 weeks, at which point the percentage of sori area to blade area did not significantly differ

217 between the experimental and wild thalli at the end of seedling production. A large number of
218 zoospores were obtained from the experimental thalli on the 29th day as well as from the wild
219 thalli, and their germlings grew healthily. Thus, the zoospores obtained using maturation
220 induction could be used for stably supplying seedlings for aquaculture as they were not
221 influenced by the natural environment. Note that in *S. japonica*, one study reported cultivation
222 of immature sporophytes of a Chinese hybrid cultivar with their blade lengths cut to a uniform
223 length of 1.5 m (using a recirculation system within a facility, which enabled the supply of
224 nutrient salts, but otherwise with conditions similar to those of the present study) (Su et al.,
225 2020). The blades were covered with sori after one month of cultivation and zoospores were
226 released from the sori 44 days later. However, the authors did not discuss what percentages of
227 the blade area were covered with sori one month after the start of cultivation or whether it was
228 possible to produce seedlings at this stage. We used wild thalli for our study, which have
229 maturation characteristics in the sea different to those of hybrid cultivars, and cultivated thalli
230 derived from wild thalli (e.g. Kawashima, 2012; Su et al., 2020). Although the percentages of
231 sori area to blade area varied among sample parts, the percentages around the base part of the
232 underside – the part with the highest formation rate – on days 26–37 of cultivation were higher
233 than about 50% in all thalli, showing that they were suitable as parental algae for cultivation. In
234 this study, between the seedlings produced using maturation induction and the seedlings using a
235 conventional method, significant differences were found in blade width and wet weight but not
236 in blade length. It is suggested that growth of the experimental seedlings was suppressed
237 because they were cultivated in the two-year aquaculture area, and the culture net was dropped
238 to a greater depth during winter. Since it is one-year aquaculture that would enjoy the benefits of
239 early marine input of seedlings, this technique needs to be verified in the future at southern sea
240 areas in Hokkaido where this type of aquaculture is practiced.

241 As mentioned above, in China, precocious sporophytes obtained from aquaculture are used as
242 parental algae, which allows the stable supply of seedlings (Su et al., 2017). However, as a
243 result of repeated aquaculture, a decrease in productivity and quality of products resulting from
244 the crossing of closely related seedlings may occur (Li et al., 2016; Zhao et al., 2016).
245 Therefore, in areas where *S. japonica* naturally occurs, it is desirable that wild thalli be used to
246 produce seedlings (in this case, the presence or absence of sori at the onset of maturation
247 induction is not an issue). However, in Hokkaido, the adherence of Hydrozoa (e.g., *Plumularia*),
248 Bryozoa (e.g., *Membranipora*), and Amphipoda (e.g., *Ceinina*) to kelp becomes more
249 pronounced from late summer (cf. Figure 2-D), and their effects on seedling production have
250 become serious (e.g., Akaike et al., 2002). We believe that systematically producing parental
251 algae for aquaculture in indoor facilities under suitable conditions will protect them from these
252 periphytons and would be very effective from the viewpoint of securing parental algae.
253 Normally, second-year thalli are used as parental algae for aquaculture in Hokkaido. In this
254 experiment, when the first-year and the second-year thalli were compared, sorus formation
255 began earlier in the former but the extent of the spread of sori at the height of sorus formation
256 was similar between the two. Since the second-year thalli are larger than the first-year thalli and
257 have a larger sori area, which makes it easier to judge their useful characteristics as marine
258 products, it would be more efficient to use the second-year thalli for maturation-induced
259 seedling production.

260 At the seedling production facilities of the Oyasu branch of the Toi Fisheries Cooperation,
261 Hakodate City, the main aquaculture area of this species in Hokkaido, the total length of
262 seedling threads used for one production event is 23,400 m (Mr. N. Azuma, Toi Fisheries
263 Cooperation, personal communication). Assuming 127 young thalli per centimeter of thread (set
264 based on the authors' preliminary observations [N = 10]) at the onset of marine aquaculture, the

265 total number of young thalli is approximately 300 million, which is usually obtained from 40–
266 50 parental algae. No more than 100 mature parental algae are needed to obtain 500 million
267 young *S. japonica* seedlings for aquaculture (Zhao et al., 2016), and despite any differences in
268 expectations toward aquaculture seedlings between the two countries, the seedlings needed
269 would basically be the same. In this study, the maturation of four samples could be induced in
270 one tank, which means that at the industrial level, one seedling production facility would need
271 around 10 tanks. This scale is feasible for existing facilities – e.g., the aforementioned Oyasu
272 facilities have 10 tanks.

273 In this study, in all experimental thalli and outdoor wild thalli, sorus formation started from
274 the underside of the blade and spread to their upper side. Wild *S. japonica* has a distinct phase
275 when sori are formed on one side of their blades during early maturation (Kawashima, 2012),
276 whereas few maturation-induced thalli showed such a phase. In this species, both sides of blade
277 pieces matured at the same time when both sides were cultivated under a uniform environment
278 (Nimura and Mizuta, 2002b), and we suggest that the results of our study were influenced by
279 the thalli being turned upside-down regularly.

280 In wild *S. japonica*, sorus formation starts from the base part (Kawashima, 2012). In this
281 study, sorus formation began near the cut side in the top-cut thalli, whereas it predominantly
282 occurred near the base and middle parts in the top-intact thalli. It has been speculated that in
283 saccharinan kelp, auxin, when transported from the base part to the marginal part, suppresses
284 sorus formation during the growth period (Buchholz and Lüning, 1999; Lüning et al., 2000;
285 Pang and Lüning, 2004; Kai et al., 2006). The fact that sori are formed near the cut site when a
286 part away from the base is removed supports this hypothesis (Su et al., 2020). However, there
287 have been no reports in intact *S. japonica* plants that compare the sorus formation process
288 between the top-cut and top-intact thalli. It is suggested that in this study, sorus formation

289 occurred as a result of the loss of the sorus formation inhibitor from the cut site, supporting the
290 said hypothesis. For Nos. 3 and 4, during tank cultivation, the loss of the top part was negligible
291 in the former but significant in the latter; however, there were no differences in sorus formation
292 sites between the two samples. In Hokkaido, when this species is cultivated, sori are formed in
293 the top parts of the blade during the growth period, and the margin of top parts are severely lost
294 in these algae. Movement of resources associated with growth from the marginal part to the
295 median and base parts has been suggested as well as the limitation of resources necessary for the
296 maturation of the marginal parts (Mizuta et al., 1999), but knowledge on the effect of the loss of
297 the marginal part on sorus formation is scarce. There is some indication that the localization of
298 an abscisic acid-like substance, a sori induction promotor, is involved in the maturation of *S.*
299 *japonica* (Nimura and Mizuta, 2002b, c). To be able to handle parental algae in an optimal
300 manner, detailed studies on the loss and artificial removal of the top part are required from the
301 viewpoints of both the inhibition and promotion of sorus formation.

302 **ACKNOWLEDGEMENTS**

303 The authors thank Dr. K. Noro for the kind advice and useful comments. The authors also
304 thank staffs of Toi Fisheries Cooperation, Mr. K. Tenjin (Rausu Fisheries Cooperation), and Mr.
305 R. Ueda (Rishiri Fisheries Cooperation) for collecting materials and cooperation with field
306 experiment. Prof. M. Toda (Niigata University) and Dr. K. Enomoto (The University of Shiga
307 Prefecture) are also thanked for consenting to use TouchDeMeasure. This research was
308 supported by a research project grant-in-aid for Scientific Research (18K05774) from the
309 Ministry of Education, Science, Sports, and Culture, Japan.

310

311 **CONFLICT OF INTEREST**

312 The authors declare no conflicts of interest associated with this manuscript.

313

314 **DATA AVAILABILITY STATEMENT**

315 The datasets generated and/or analyzed during the current study are available from the
316 corresponding author on reasonable request.

317

318 **REFERENCES**

- 319 Akaike, S., Takiya, A., Tsuda, F., Motoya, A., & Takahashi, K. (2002). Seasonal occurrence of a
320 kelp-boring amphipod, *Ceinia japonica* along the coasts of Hokkaido from 1997 to 2001.
321 Scientific Reports of Hokkaido Fisheries Experimental Station, 61, 25-18. (in Japanese)
- 322 Boderskov, T., Rasmussen, M. B., Bruhn, A. (2021). Obtaining spores for the production of
323 *Saccharina latissima*: seasonal limitations in nature, and induction of sporogenesis in darkness.
324 Journal of Applied Phycology, 33: 1035–1046. <https://doi.org/10.1007/s10811-020-02357-0>.
- 325 Buchholz, C., & Lüning, K. (1999). Isolated, distal frond discs of the brown alga *Laminaria*
326 *digitata* form sorus, but not discs near to the meristematic transition zone. Journal of Applied
327 Phycology, 11, 579– 584.
- 328 Enomoto, K., Toda, M., Kawasaki, T., & Shimizu, Y. (2017). Image measurement support
329 system "TouchDeMeasure" using touch operation. Image Laboratory, 28 (9), 1-7. (in Japanese)
- 330 Forboad, S., Skjermo, J., Arff, J., Handå, A., Reitan, K.I., Bjerregaard, R., & Lüning, K. (2012).
331 Development of *Saccharina latissima* (Phaeophyceae) kelp hatcheries with year-round
332 production of zoospores and juvenile sporophytes on culture ropes for kelp aquaculture. Journal
333 of Applied Phycology, 24, 393–399. DOI 10.1007/s10811-011-9784-y.
- 334 Forboad, S., Steinhovden, K.B., Solvang, T., Handå, A., & Skjermo, J. (2020). Effect of seeding
335 methods and hatchery periods on sea cultivation of *Saccharina latissima* (Phaeophyceae): a
336 Norwegian case study. Journal of Applied Phycology, 32, 2201-2212.

337 <https://doi.org/10.1007/s10811-019-01936-0>.

338 Hwang, E. K., Yotsukura, N., Pang, S. J., Su, L., & Shan, T. F. (2019). Seaweed breeding
339 programs and progress in the Eastern Asian countries. *Phycologia*, 58, 484-495.
340 <https://doi.org/10.1080/00318884.2019.1639436>.

341 Kai, T., Nimura, K., Yasui, H., and Mizuta, H. (2006). Regulation of sorus formation by auxin
342 in Laminariales sporophyte. *Journal of Applied Phycology*, 18, 95–101. DOI: 10.1007/s10811-
343 005-9020-8.

344 Kawai, T., Galanin, D., Krupnova, T., & Yotsukura, N. (2015). Harvest and cultivation of
345 *Saccharina japonica* in northern Hokkaido, Japan, and southern Sakhalin and Primorye, Russia:
346 A review. *Algal Resources*, 8, 155-163.

347 Kawashima, S. (1984). Kombu cultivation in Japan for human foodstuff. *The Japanese Journal*
348 *of Phycology*, 32, 379-394.

349 Kawashima, S. (2012). Morphology and taxonomy of the laminariaceous algae in cold water
350 area of Japan. Tokyo: Seibutsukenkyusha. (in Japanese)

351 Kirihara, S., Fujikawa, Y., & Notoya, M. (2003). Effect of temperature and day length on the
352 zoosporangial sorus formation and growth of sporophytes *Laminaria japonica* Areschoug
353 (Laminariales, Phaeophyceae) in tank culture. *SUISANZOUSHOKU*, 51, 385–390. (in
354 Japanese)

355 Li, X., Zhang, Z., Qu, S., Liang, G., Sun, J., Zhao, N., Cui, C., Cao, Z., Li, Y., Pan, J., Yu, S.,
356 Wang, Q., Li, X., Luo, S., Song, S., Guo, L., & Yang, G. (2016). Improving seedless kelp
357 (*Saccharina japonica*) during its domestication by hybridizing gametophytes and seedling-
358 raising from sporophytes. *Scientific Reports*, 6, 21 255. DOI 10.1038/srep21255.

359 Lüning, K. (1988). Photoperiodic control of sorus formation in the brown alga *Laminaria*
360 *saccharina*. *Marine Ecology Progress Series*, 45, 137–144.

361 Lüning, K., Wagner, A., & Buchholz, C. (2000). Evidence for inhibitors of sporangium
362 formation in *Laminaria digitate* (Phaeophyceae) during the season of rapid growth. *Journal of*
363 *Phycology*, 36, 1 129-1 134. <https://doi.org/10.1046/j.1529-8817.2000.00017.x>.

364 Mizuta, H., Nimura, K., & Yamamoto, H. (1999). Sorus development on median and marginal
365 parts of the sporophyte of *Laminaria japonica* Areschoug (Phaeophyceae). *Journal of Applied*
366 *Phycology*, 11, 585-591.

367 Nimura K, Mizuta H, Yamamoto H (2002a) Critical contents of nitrogen and phosphorus for
368 sorus formation in four *Laminaria* species. *Botanica Marina*, 45, 184–188.

369 Nimura, K., & Mizuta, H. (2002b). Pattern of sori on the blade segments of *Laminaria japonica*
370 Areschoug (Laminariales, Phaeophyceae). *SUISANZOUSHOKU*, 50, 157-162. (in Japanese)

371 Nimura, K., & Mizuta, H. (2002c). Inducible effects of abscisic acid on sporophyte discs from
372 *Laminaria japonica* Areschoug (Laminariales, Phaeophyceae). *Journal of Applied Phycology*,
373 14, 159-163.

374 Oishi, K. (1987). *Kombu no michi*. Tokyo: Daiichi shobo. (in Japanese)

375 Pang, S. J., & Lüning, K. (2004). Breaking seasonal limitation: year-round sporogenesis in the
376 brown alga *Laminaria saccharina* by blocking the transport of putative sporulation inhibitors.
377 *Aquaculture*, 240, 531-541. <https://doi.org/10.1016/j.aquaculture.2004.06.034>.

378 Shan, T., Liu, F., Liu, Q., & Pang, S. (2011). Review and prospect of “summer sporeling”
379 technique of *Saccharina japonica* in China. *Journal of Agricultural Science and Technology*, 13:
380 129–134. (in Chinese)

381 Shan ,T, Yotsukura, N., & Pang, S. (2017). Novel implications on the genetic structure of
382 representative populations of *Saccharina japonica* (Phaeophyceae) in the Northwest Pacific as
383 revealed by highly polymorphic microsatellite markers. *Journal of Applied Phycology*, 29, 631-
384 638. DOI 10.1007/s10811-016-0888-2.

385 Skriptsova, A.V., and Titlyanov E.A. (2003). Effect of the meristem on sporification of
386 *Laminaria cichorioides*. Russian Journal of Marine Biology, 29, 372-377.

387 Su, L., Pang, S. J., Shan, T. F., & Li, X. (2017). Large-scale hatchery of the kelp *Saccharina*
388 *japonica*: a case study experience at Lvshun in northern China. Journal of Applied Phycology,
389 29, 3 003-3 013. DOI 10.1007/s10811-017-1154-y.

390 Su, L., Shan, T. F., Jing, L., Pang, S. J., Leng, X. F., Zhang, Y., & Gao, H. T. (2020).
391 Aquaculture of the hybrid cultivars of *Saccharina japonica*: Removing the obstacle of sori
392 production by photoperiodic control. Aquaculture, 519, 734 917.
393 <https://doi.org/10.1016/j.aquaculture.2019.734917>.

394 Sohn, C. H. (1998). The seaweed resources of Korea. In A.T. Critchley & M. Ohno (Eds.),
395 Seaweed resources of the world (pp. 15-33). Tokyo: JICA.

396 Tatewaki, M. (1966). Formation of a crustaceous sporophyte with unilocular sporangia in
397 *Scytosiphon lomentaria*. Phycologia, 6, 62-66.

398 Tseng, C. K., Sun, K. Y., & Wu, C. Y. (1955). On the cultivation of Haidai (*Laminaria japonica*
399 Aresch) by summering young sporophytes at low temperature. Acta Botanica Sinica, 4, 255–
400 264. (in Chinese)

401 Yotsukura, N., Kawashima, S., Kawai, T., Abe, T., & Druehl, L. D. (2008). A systematic re-
402 examination of four *Laminaria* species: *L. japonica*, *L. religiosa*, *L. ochotensis* and *L. diabolica*.
403 Journal of Japanese Botany, 83, 165-176.

404 Zhao, X. B., Pang, S. J., Liu, F., Shan, T. F., Li, J., Gao, S. Q., & Kim, H. G. (2016).
405 Intraspecific crossing of *Saccharina japonica* using distantly related unialgal gametophytes
406 benefits kelp farming by improving blade quality and productivity at Sanggou Bay, China.
407 Journal of Applied Phycology, 28, 449-455. DOI 10.1007/s10811-015-0597-2.

408

409

410

411

412

413 **Table**

Table 1 The sporophytes of *Saccharina japonica* that was conducted maturation induction in this study.

Sample No.	Collection date	Collection site	Age	Maturity at the time of collection	Artificial cutting of blade	note
No. 1	July 24. 2018	Oyasu, Hakodate City	1st year	*		originated from 2-year-old wild kelp collected the year before and cultured in the sea
No. 2	July 24. 2018	Oyasu, Hakodate City	1st year	*		
No. 3	July 24. 2018	Oyasu, Hakodate City	1st year			wild
No. 4	July 24. 2018	Oyasu, Hakodate City	1st year			
No. 5	June 24. 2019	Oyasu, Hakodate City	1st year		*	
No. 6	June 24. 2019	Oyasu, Hakodate City	1st year		*	
No. 7	June 24. 2019	Oniwaki, Rishirifuji Town	2nd year			
No. 8	July 6. 2019	Kuzurehama, Rausu Town	2nd year			

414

415

416

417

Table 2 The percentages of sori area to blade area on each part of blade of experimental thallus (No. 1-8) with maturation induction.

Side	Part of blade	Elapsed days	Sample No.				Side	Part of blade	Elapsed days	Sample No.			
			No. 1	No. 2	No. 3	No. 4				No. 5	No. 6	No. 7	No. 8
Upper side	Top part	13		1.4			Upper side	Top part	16	-	-	-	-
		19	6.3	10.1					21	2.0	2.0	-	-
		26	16.7	4.5	8.9	3.2			25	-	-	-	-
		33	21.0	4.5	9.3	3.5			28	3.2	2.9	0.8	-
	Middle part	13		0.1				37	0.1	3.1	5.0	-	
		19	5.3	0.1		0.1		16	-	-	-	0.6	
		26	7.4	4.5	4.0	1.4		21	-	-	-	-	
		33	8.2	5.4	2.8	1.2		25	-	-	-	1.2	
	Base part	13		0.7				28	1.4	3.9	0.2	-	
		19	16.2	1.2	0.3			37	2.0	2.9	0.4	-	
		26	19.9	1.7	7.4	6.6		16	-	-	-	17.6	
		33	21.3	1.3	9.9	4.9		21	-	-	-	-	
Underside	Top part	13		8.1			Underside	Base part	25	-	-	-	34.4
		19	36.0	12.1	0.4	0.5			28	0.5	0.8	-	-
		26	46.2	18.3	26.0	20.1			37	0.2	2.9	-	-
		33	53.8	19.1	25.4	18.6			16	-	-	-	-
	Middle part	13		4.7				21	11.8	5.4	-	-	
		19	16.4	10.2		0.8		25	-	-	-	1.6	
		26	24.4	23.8	26.4	12.0		28	14.4	8.7	-	-	
		33	23.3	23.2	27.2	14.0		37	14.2	7.7	11.5	-	
	Base part	13		18.1				16	-	-	-	11.9	
		19	51.6	44.3	14.5	24.3		21	-	8.2	-	-	
		26	55.8	52.6	59.5	47.1		25	-	-	-	19	
		33	54.3	49.6	59.5	48.2		28	17.4	19.0	13.2	-	
						37	25.2	24.1	18.1	-			
						Base part	16	-	-	-	37.7		
					21		-	8.7	-	-			
					25		-	-	-	67.7			
					28		31.1	23.1	37.8	-			
					37		50.7	44.4	46.5	-			

418

419

420

421

422

423

424 **Figure legends**

425 Figure 1 Sori formed on the underside of experimental thalli (A-E) and of a wild
426 thallus (F) of *Saccharina japonica*. The sori were processed green with
427 TouchDeMeasure. A: No. 1 after 19th day of the cultivation, B: No. 5 after 21th day of
428 the cultivation, C: No. 7 after 37th day of the cultivation, D: The underside of No. 8 after
429 16th day of the cultivation, E: No. 8 after 25th day of the cultivation, F: Wild thallus
430 collected on December 16, 2019, in the Oniwaki district of Rishirifuji Town.

431 Scale=10cm.

432 Figure 2 Averages of total length (A), blade length (B), blade width (C), and wet
433 weight of thalli of *Saccharina japonica* brought up in the sea. a: Experimental thalli
434 derived from seedlings obtained using maturation induction, b: Thalli derived from
435 seedlings obtained by the conventional method. **: $p < 0.01$.

Figure 1



