

HOKKAIDO UNIVERSITY

Title	Classification of Irreducible (g, k)-Modules Associated to the Ideals of Minimal Nilpotent Orbits for Simple Lie Groups of Type A
Author(s)	Tamori, Hiroyoshi
Citation	IMRN: International Mathematics Research Notices, 2023(5), 3814-3847 https://doi.org/10.1093/imrn/rnab356
Issue Date	2021-12-21
Doc URL	http://hdl.handle.net/2115/88245
Rights	This is a pre-copyedited, author-produced version of an article accepted for publication in IMRN: International Mathematics Research Notices following peer review. The version of record: Hiroyoshi Tamori, Classification of Irreducible (g, k)-Modules Associated to the Ideals of Minimal Nilpotent Orbits for Simple Lie Groups of Type A. IMRN, 2023, 5, 3814–3847 (March 2023) is available online at: https://doi.org/10.1093/imrn/rnab356
Туре	article (author version)
File Information	Int. Math. Res. Notices_rnab356.pdf

6

Instructions for use

CLASSIFICATION OF IRREDUCIBLE (g, \mathfrak{k}) -MODULES ASSOCIATED TO THE IDEALS OF MINIMAL NILPOTENT ORBITS FOR SIMPLE LIE GROUPS OF TYPE A

HIROYOSHI TAMORI

ABSTRACT. We classify completely prime primitive ideals whose associated varieties are the closure of the minimal nilpotent orbit of $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$, and classify irreducible $(\mathfrak{g}, \mathfrak{k})$ -modules which have those ideals as annihilators. Moreover, we irreducibly decompose them as \mathfrak{k} -modules.

Mathematics Subject Classification 2010: 22E46.

Key Words and Phrases: Minimal representation; Completely prime primitive ideal; Simple Lie group; Type A.

1. INTRODUCTION

Let G be a simple real Lie group and $\mathfrak{g}_0 := \operatorname{Lie}(G)$. Fix a maximally compact subalgebra \mathfrak{k}_0 of \mathfrak{g}_0 . Assume that the complexification \mathfrak{g} of \mathfrak{g}_0 is simple. Joseph proved that if \mathfrak{g} is not of type A, then the universal enveloping algebra $U(\mathfrak{g})$ has a unique completely prime ideal J_0 whose associated variety, which is a subset of $\mathfrak{g}^{\vee} := \operatorname{Hom}_{\mathbb{C}}(\mathfrak{g}, \mathbb{C})$, is the closure of the minimal nilpotent coadjoint orbit \mathcal{O}^{\min} [10], [4, Theorem 3.1]. Minimal representations of G are defined as admissible irreducible representations where the annihilators of the associated $(\mathfrak{g}, \mathfrak{k})$ -modules are equal to the Joseph ideal J_0 .

Minimal representations are classified and are known to be infinitesimally equivalent to unitary representations [20, Corollary 5.1]. In the Kirillov-Kostant orbit philosophy, they are considered to be attached to *G*-orbits in $\mathcal{O}^{\min} \cap \{X \in \mathfrak{g}^{\vee} \mid X(\mathfrak{g}_0) \subset \mathbb{R}\}$, and are considered to be a part of building blocks of the unitary dual of *G*.

Moreover, minimal representations are of interest in physics. For example, the oscillator representation of the metaplectic group (whose irreducible components are minimal) has a realization as the bound states of the quantum harmonic oscillator, and the minimal representation of the indefinite orthogonal group $O(p, 2)(p \ge 6)$ has a realization as some solution space of the wave equation in the Minkowski space (see [13, Theorem 1.4] for example).

On the other hand, there exist representations called "minimal" even when \mathfrak{g} is simple of type A, that is, when the definition of the Joseph ideal (hence the one of minimality) is not given. Here the vague term "minimal" means that there is an interest in physics as in the previous examples, the \mathfrak{k} -types are as simple as possible (called pencil, see Corollary 3.5 (2) for the definition), or there is a relation with the minimal nilpotent orbit. Such representations include the ladder representation of O(2, 4) (which is locally isomorphic to SU(2, 2)) expressing the bound states of the Hydrogen atom [12, Remark 3.6.2 (3)] and the irreducible unitary representations of the double cover $\widetilde{SL}(3,\mathbb{R})$ of SL(3, \mathbb{R}) given by Torasso (see [21, Theoreme VII.1], [19]).

The aim of this paper is to define minimality in terms of annihilators of $(\mathfrak{g}, \mathfrak{k})$ modules so that the above representations are minimal, and to classify minimal
representations for connected simple real Lie groups of type A_{n-1} $(n \geq 2)$. Hence

this study can be regarded as a small step of classifying irreducible $(\mathfrak{g}, \mathfrak{k})$ -modules associated with a fixed completely prime primitive ideal.

Our first claim is that the two-sided ideals in $U(\mathfrak{g})$ whose associated graded ideals coincide with the ideal defined by \mathcal{O}^{\min} are parametrized by the complex numbers \mathbb{C} (see Theorem 2.8). If we consider a simple Lie algebra not of type A, that condition characterizes the Joseph ideal [4, Theorem 3.1].

Let us write J_a for the ideal corresponding to $a \in \mathbb{C}$ (see Definition 2.7 for the precise definition). We say that an irreducible $(\mathfrak{g}, \mathfrak{k})$ -module is *a*-minimal if its annihilator equals J_a , and is minimal if it is *a*-minimal for some $a \in \mathbb{C}$ (see Definition 3.1).

The associated variety gives us a necessary condition for the existence of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules. If a minimal $(\mathfrak{g}, \mathfrak{k})$ -module exists, then $\mathcal{O}^{\min} \cap \{X \in \mathfrak{g}^{\vee} \mid X(\mathfrak{g}_0) \subset \mathbb{R}\}$ is not empty. Hence it suffices to consider the real forms $\mathfrak{su}(p,q)$ (p,q>0) and $\mathfrak{sl}(n,\mathbb{R})$ by [18, Proposition 4.1].

The main theorems in this paper are Theorems 5.1 and 5.3, which classify *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -modules for any $a \in \mathbb{C}$ and $\mathfrak{g}_0 = \mathfrak{su}(p,q)$ (p,q > 0), $\mathfrak{sl}(n,\mathbb{R})$. The number of the isomorphism classes of *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -modules is given in Table 1.1. We remark that many of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules do not admit nondegenerate in-

TABLE 1.1. The number of the isomorphism classes of *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -modules

\mathfrak{g}_0	a	number
$\mathfrak{su}(p,1)(p\geq 2)$	\mathbb{C}	2
$\boxed{ \mathfrak{su}(1,q)(q\geq 2)}$	\mathbb{C}	2
$\mathfrak{su}(p,q)(p,q\geq 2)$	$(p+q)/2 + \mathbb{Z}$	2
	$\mathbb{C} \setminus \left((p+q)/2 + \mathbb{Z} \right)$	0
$\mathfrak{sl}(n,\mathbb{R})(n\geq 4)$	\mathbb{C}	2
$\mathfrak{sl}(3,\mathbb{R})$	Z	3
	$\mathbb{C}\setminus\mathbb{Z}$	2

variant Hermitian forms and are not unitarizable, which does not occur when \mathfrak{g}_0 is not of type A. For the classification for unitarizable minimal $(\mathfrak{g}, \mathfrak{k})$ -modules, see Remarks 5.2 (3) and 5.4 (2).

For $\mathfrak{su}(1,q)$ (q > 0) and $\mathfrak{sl}(n,\mathbb{R})$, there exist *a*-minimal $(\mathfrak{g},\mathfrak{k})$ -modules for any $a \in \mathbb{C}$. They are related with the induction from maximal $(\theta$ -stable or real) parabolic subalgebras of \mathfrak{g} corresponding to the partition (1, n - 1). From the viewpoint of the orbit method, the situation matches to the fact that the nonzero coadjoint orbits of minimal dimension consist of \mathcal{O}^{\min} and semisimple orbits through nonzero characters of the parabolic subalgebras.

Our construction of a minimal $(\mathfrak{g}, \mathfrak{k})$ -module for $\mathfrak{g}_0 = \mathfrak{su}(p, q)$ (p, q > 0) and the one for $\mathfrak{sl}(n, \mathbb{R})$ are different. We realize it as the irreducible quotient of a Verma module for $\mathfrak{su}(p,q)$, while we construct it as the infinite-dimensional irreducible subquotient of the kernel of an intertwining differential operator between parabolically induced modules for $\mathfrak{g}_0 = \mathfrak{sl}(n, \mathbb{R})$. The latter construction for Torasso's representation (n = 3) was given in [15, Theorem 1.6], and the one for genuine *a*minimal representations of the universal cover of SL(3, \mathbb{R}) was recently given in [16, Theorem 1.7 (5)]. We give a general framework for the construction in Section 4.

The proof of exhaustion uses the facts that highest weights of \mathfrak{k} -types of a minimal $(\mathfrak{g}, \mathfrak{k})$ -module lie in some specified lattice, and that *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -modules are isomorphic to each other if they have a common \mathfrak{k} -type (see Proposition 3.3). The argument is the same as the classification when \mathfrak{g}_0 is not of type A [20]. We also need to show the injectivity of some intertwining differential operator for the nonexistence of minimal $(\mathfrak{sl}(3,\mathbb{C}),\mathfrak{so}(3,\mathbb{R}))$ -modules with some \mathfrak{k} -types (see Proposition 6.1).

Notation: We set $\mathbb{N} := \{0, 1, 2, \ldots\}$. Lie groups will be denoted by Latin capital letters, their Lie algebras by corresponding lower case German letters with subscript zero. We omit the subscript zero to denote their complexifications. Given a simple Lie algebra \mathfrak{g} , a Cartan subalgebra \mathfrak{h} , and a finite-dimensional \mathfrak{g} -module V, we write V_{μ} for the weight space of V of weight μ with respect to \mathfrak{h} .

2. Primitive ideals and irreducible highest weight modules

Let $n \geq 2$ and $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$. In this section, we classify two-sided ideals J in the enveloping algebra $U(\mathfrak{g})$ whose associated graded ideals gr J are the ideal $\mathcal{I}(\mathcal{O}^{\min})$ defined by the minimal nilpotent coadjoint orbit \mathcal{O}^{\min} of \mathfrak{g}^{\vee} . Moreover, we obtain the classification of irreducible highest weight modules whose annihilators are equal to one of these ideals.

For $A \in \mathfrak{g}$ and $1 \leq i, j \leq n$, we write $A_{i,j}$ for the (i, j)-th component of the matrix A. We write \mathfrak{h} for the Cartan subalgebra of \mathfrak{g} consisting of diagonal matrices. Let e_i $(1 \leq i \leq n)$ be an element of the dual \mathfrak{h}^{\vee} defined by $e_i(H) = H_{i,i}$ for $H \in \mathfrak{h}$. Then the set of roots for $(\mathfrak{g}, \mathfrak{h})$ is written as $\Sigma = \{ e_i - e_j \mid 1 \leq i, j \leq n \}$. We fix a set of positive roots $\Sigma^+ = \{ e_i - e_j \mid i < j \}$. For an element λ in \mathfrak{h}^{\vee} , we define $\lambda_i \in \mathbb{C}$ $(1 \leq i \leq n)$ by $\lambda = \sum_i \lambda_i e_i$ and $\sum_i \lambda_i = 0$. Let us write $(\lambda_1, \ldots, \lambda_n)$ for λ .

We first describe the irreducible decomposition of the second symmetric power $S^2(\mathfrak{g})$. Let I_n be the unit *n*-by-*n* matrix, $E_{i,j}$ the matrix whose (i, j)-th component is one and the others are zero. Set

$$T_{i,j} := E_{i,j} - \frac{\delta_{i,j}}{n} I_n$$

Here $\delta_{i,j}$ denotes the Kronecker delta. We remark that $A = \sum_{1 \leq i,j \leq n} A_{i,j} T_{i,j}$ holds for every $A \in \mathfrak{g}$. We define a nondegenerate invariant bilinear form B on \mathfrak{g} by

(2.1)
$$B(X,Y) := \operatorname{Tr}(XY) \quad (X,Y \in \mathfrak{g}).$$

Put

$$\Omega := \sum_{1 \le i,j \le n} T_{i,j} T_{j,i} \in S^2(\mathfrak{g}).$$

Proposition 2.1. (1) As a \mathfrak{g} -module, the second symmetric power $S^2(\mathfrak{g})$ irreducibly decomposes to

(2.2)
$$\begin{cases} F(2e_1 - 2e_n) \oplus F(e_1 + e_2 - e_{n-1} - e_n) \oplus F(e_1 - e_n) \oplus F(0) & \text{if } n \ge 4, \\ F(2e_1 - 2e_n) \oplus F(e_1 - e_n) \oplus F(0) & \text{if } n = 3, \\ F(2e_1 - 2e_n) \oplus F(0) & \text{if } n = 2. \end{cases}$$

Here $F(\lambda)$ denotes the irreducible g-submodule of $S^2(\mathfrak{g})$ with highest weight λ and we have $F(0) = \mathbb{C}\Omega$.

(2) For $n \ge 4$, the g-submodule $F(e_1 + e_2 - e_{n-1} - e_n) \oplus F(e_1 - e_n) \oplus F(0)$ is equal to

(2.3)
$$\operatorname{span} \{ T_{i,j} T_{k,l} - T_{i,l} T_{k,j} \mid 1 \le i, j, k, l \le n \}.$$

(3) For $n \geq 3$, the linear map

$$\mathfrak{g} \to F(e_1 - e_n); A \mapsto \sum_{i,j,k} A_{i,j} T_{i,k} T_{k,j}$$

is a g-isomorphism.

Proof. By $[T_{r,s}, T_{i,j}] = \delta_{i,s}T_{r,j} - \delta_{r,j}T_{i,s}$, we have

$$\begin{split} [T_{r,s}, T_{i,j}T_{k,l} - T_{i,l}T_{k,j}] &= (\delta_{i,s}T_{r,j} - \delta_{r,j}T_{i,s})T_{k,l} + T_{i,j}(\delta_{k,s}T_{r,l} - \delta_{r,l}T_{k,s}) \\ &- (\delta_{i,s}T_{r,l} - \delta_{r,l}T_{i,s})T_{k,j} - T_{i,l}(\delta_{k,s}T_{r,j} - \delta_{r,j}T_{k,s}) \\ &= \delta_{i,s}(T_{r,j}T_{k,l} - T_{r,l}T_{k,j}) - \delta_{r,j}(T_{i,s}T_{k,l} - T_{i,l}T_{k,s}) \\ &+ \delta_{k,s}(T_{i,j}T_{r,l} - T_{i,l}T_{r,j}) - \delta_{r,l}(T_{i,j}T_{k,s} - T_{i,s}T_{k,j}). \end{split}$$

Therefore the subspace (2.3) is invariant under the action of \mathfrak{g} and has a highest weight vector $T_{1,n-1}T_{2,n} - T_{1,n}T_{2,n-1}$ of weight $e_1 + e_2 - e_{n-1} - e_n$ for $n \ge 2$.

Let us consider a vector $\sum_{k} T_{i,k} T_{k,j} = \sum_{k} (T_{i,k} T_{k,j} - T_{i,j} T_{k,k})$, which belongs to (2.3). We see

$$[T_{r,s}, \sum_{k} T_{i,k} T_{k,j}] = \sum_{k} \{ (\delta_{i,s} T_{r,k} - \delta_{r,k} T_{i,s}) T_{k,j} + T_{i,k} (\delta_{k,s} T_{r,j} - \delta_{r,j} T_{k,s}) \}$$
$$= \delta_{i,s} \sum_{k} T_{r,k} T_{k,j} - \delta_{r,j} \sum_{k} T_{i,k} T_{k,s}.$$

Therefore $[\mathfrak{g}, \Omega] = 0$ and the map in (3) is a \mathfrak{g} -homomorphism into $S^2(\mathfrak{g})$. For $n \geq 3$, the submodule (2.3) has highest weight $e_1 - e_n$ since $\sum_k T_{1,k} T_{k,n}$ is nonzero.

Furthermore, (2.3) has highest weight 0 as it contains Ω , and it does not contain a nonzero multiple of $T_{1,n}T_{1,n}$, which is a highest weight vector of weight $2e_1 - 2e_n$ in $S^2(\mathfrak{g})$.

By the Weyl dimension formula, we see the dimension of (2.2) is equal to the square of n(n-1)/2, which is the dimension of $S^2(\mathfrak{g})$. Hence (2.2) equals the direct sum of $F(2e_1 - 2e_n)$ and (2.3), and our claim follows.

For
$$1 \le i < j < k < l \le n$$
, we put
$$\mathcal{M}_{i,j,k,l} := \left\{ a_1(T_{i,j} + T_{k,l}) + a_2(T_{i,k} + T_{j,l}) + a_3(T_{i,l} + T_{j,k}) \in \mathfrak{g} \mid \sum_{r=1}^3 a_r = 0 \right\}.$$

Define a injective linear map

$$T: \{ A \in \mathfrak{gl}(n,\mathbb{C}) \mid A_{i,j} = 0 \text{ for } i \ge j \} \to S^2(\mathfrak{g})_0; A \mapsto \sum_{i,j} A_{i,j}(T_{i,i}T_{j,j} - T_{i,j}T_{j,i}).$$

Proposition 2.2. Assume $n \ge 4$. Then the zero weight space $F(e_1+e_2-e_{n-1}-e_n)_0$ is equal to

span {
$$T(\mathcal{M}_{i,j,k,l}) \mid 1 \le i < j < k < l \le n$$
 }.

Proof. Define an $\mathfrak{su}(n)$ -invariant Hermitian inner product (\cdot, \cdot) on \mathfrak{g} by $(X, Y) := B(X, {}^t\overline{Y}) = \operatorname{Tr}(X^t\overline{Y}) \ (X, Y \in \mathfrak{g})$, where \overline{Y} denotes the complex conjugate of Y. It induces an $\mathfrak{su}(n)$ -invariant Hermitian inner product on $\mathfrak{g} \otimes \mathfrak{g}$ and one on $S^2(\mathfrak{g})$ via the symmetrization, which we write (\cdot, \cdot) by abuse of notation. By the assumption $n \geq 4$ and Proposition 2.1 and its proof, the zero weight space $F(e_1 + e_2 - e_{n-1} - e_n)_0$ is the orthogonal complement of $F(e_1 - e_n)_0 + F(0) = \operatorname{span} \{\sum_k T_{i,k} T_{k,i} \mid 1 \leq i \leq n\}$ in the zero weight space span $\{T_{i,i}T_{j,j} - T_{i,j}T_{j,i} \mid i < j\}$ of (2.3). In particular, dim $F(e_1 + e_2 - e_{n-1} - e_n)_0 = n(n-1)/2 - n = n(n-3)/2$.

Let us prove that $T(\mathcal{M}_{i,j,k,l})$ is contained in $F(e_1 + e_2 - e_{n-1} - e_n)_0$ for any i < j < k < l. By

$$(T_{i,j}, T_{r,s}) = \operatorname{Tr}\left(\left(E_{i,j} - \frac{\delta_{i,j}}{n}I_n\right)\left(E_{s,r} - \frac{\delta_{r,s}}{n}I_n\right)\right) = \delta_{i,r}\delta_{j,s} - \frac{\delta_{i,j}\delta_{r,s}}{n},$$

we have

$$2\Big(T_{i,i}T_{j,j} - T_{i,j}T_{j,i}, \sum_{s} T_{r,s}T_{s,r}\Big)$$

$$= \sum_{s} \{ (T_{i,i}, T_{r,s})(T_{j,j}, T_{s,r}) + (T_{j,j}, T_{r,s})(T_{i,i}, T_{s,r}) \\ - (T_{i,j}, T_{r,s})(T_{j,i}, T_{s,r}) - (T_{j,i}, T_{r,s})(T_{i,j}, T_{s,r}) \} \\ = \sum_{s} \left\{ 2 \left(\delta_{i,r} \delta_{i,s} - \frac{\delta_{r,s}}{n} \right) \left(\delta_{j,r} \delta_{j,s} - \frac{\delta_{r,s}}{n} \right) - \delta_{i,r} \delta_{j,s} - \delta_{j,r} \delta_{i,s} \right\} \\ = 2 \left(\delta_{i,r} - \frac{1}{n} \right) \left(\delta_{j,r} - \frac{1}{n} \right) - \delta_{i,r} - \delta_{j,r} \\ = \frac{2}{n^2} - \left(\frac{2}{n} + 1 \right) \left(\delta_{i,r} + \delta_{j,r} \right)$$

for any i < j and $1 \le r \le n$. Since $\sum_{r,s} A_{r,s} = 0$ and $\sum_r (A_{r,r_0} + A_{r_0,r}) = 0$ for any $A \in \mathcal{M}_{i,j,k,l}$ and $1 \le r_0 \le n$, it follows that $T(\mathcal{M}_{i,j,k,l})$ is orthogonal to $F(e_1 - e_n)_0 + F(0)$, and is included in $F(e_1 + e_2 - e_3 - e_4)_0$.

Let us prove that the linear map Π from $\sum_{i < j < k < l} \mathcal{M}_{i,j,k,l}$ to $\bigoplus_{1 \le i < j \le n-2} \mathbb{C} \oplus \bigoplus_{1 \le i \le n-3} \mathbb{C}$ defined by $\Pi(A) := ((A_{i,j})_{(i,j)}, (A_{i,n-1})_i)$ is bijective. If it holds, we obtain dim $\sum_{i < j < k < l} \mathcal{M}_{i,j,k,l} = (n-2)(n-3)/2 + n - 3 = n(n-3)/2 = \dim F(e_1 + e_2 - e_3 - e_4)_0$, and the proof is complete.

For any $(a_{i,j}) \in \bigoplus_{1 \le i \le j \le n-2} \mathbb{C}$ and $(a_{i,n-1}) \in \bigoplus_{1 \le i \le n-3} \mathbb{C}$, the image of

$$\sum_{1 \le i < j \le n-2} a_{i,j} (T_{i,j} + T_{n-1,n} - T_{i,n} - T_{j,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n-1} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n-1} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{1 \le i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n-1} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n-1} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{i \le n-3} a_{i,n-1} (T_{i,n-1} + T_{n-2,n-1} - T_{i,n-1} - T_{n-2,n-1}) + \sum_{i \le n-3} a_{i,n-1} (T_{i,n-1} - T_{i,n-1} -$$

is equal to $((a_{i,j}), (a_{i,n-1}))$. Hence Π is surjective.

What is left is to prove the injectivity of Π . Assume $A \in \sum_{i < j < k < l} \mathcal{M}_{i,j,k,l}$ and $\Pi(A) = 0$. We see $\sum_{j=1}^{i-1} A_{j,i} + \sum_{j=i+1}^{n} A_{i,j} = 0$ for any $1 \le i \le n$. Hence for any $1 \le i \le n-3$, we have $A_{i,n} = -\sum_{j=i+1}^{n-1} A_{i,j} - \sum_{j=1}^{i-1} A_{j,i} = 0$. Moreover, the sum of any two of $A_{n-2,n-1}, A_{n-2,n}, A_{n-1,n}$ is zero. Therefore A = 0, and Π is injective. \Box

We next classify irreducible highest weight modules annihilated by some \mathfrak{g} -submodules of $\bigoplus_{0 \leq i \leq 2} S^i(\mathfrak{g})$. For $\lambda \in \mathfrak{h}^{\vee}$, we write $L(\lambda)$ for the irreducible highest weight \mathfrak{g} -module with highest weight λ . Let l_{λ} be a nonzero highest weight vector in $L(\lambda)$. We remark that the infinitesimal character of $L(\lambda)$ is the orbit of the Weyl group through $\lambda + \rho$, where $\rho = \sum_{1 \leq i \leq n} ((n+1-2i)/2)e_i$ is half the sum of positive roots for Σ^+ .

For $a \in \mathbb{C}$, let F^a be the \mathfrak{g} -submodule of $\mathfrak{g} + S^2(\mathfrak{g})$ defined by

(2.4)
$$F^{a} := \begin{cases} F(e_{1} + e_{2} - e_{n-1} - e_{n}) + F(e_{1} - e_{n})^{a} & \text{if } n \ge 4, \\ F(e_{1} - e_{n})^{a} & \text{if } n = 2, 3, \end{cases}$$

where

$$F(e_1 - e_n)^a := \left\{ \left| \sum_{i,j,k} A_{i,j} T_{i,k} T_{k,j} - \frac{a(n-2)}{n} A \right| A \in \mathfrak{g} \right\} \subset \mathfrak{g} + S^2(\mathfrak{g}).$$

The normalizing factor (n-2)/n is put in order to simplify our assertions. We remark that $F^a = 0$ when n = 2.

For a positive integer m, put $\mathbf{1}_m$ to be the m-tuple

$$\mathbf{1}_m := (\underbrace{1,\ldots,1}_m).$$

We regard $\mathbf{1}_0$ as the empty tuple. For $1 \leq i \leq n$ and $a \in \mathbb{C}$, let us define an element of \mathfrak{h}^{\vee} by

$$(2.5)\lambda(i,a) := \frac{1}{n} \left(\left(-a - \frac{n}{2} \right) \mathbf{1}_{i-1}, (n-1)a - \frac{n(n+1-2i)}{2}, \left(-a + \frac{n}{2} \right) \mathbf{1}_{n-i} \right).$$

Let sym be the symmetrization map, that is, the \mathfrak{g} -isomorphism defined by

sym:
$$S(\mathfrak{g}) \to U(\mathfrak{g}); A_1 A_2 \cdots A_m \mapsto \frac{1}{m!} \sum_{\eta \in \mathfrak{S}_m} A_{\eta(1)} A_{\eta(2)} \cdots A_{\eta(m)},$$

where $m \in \mathbb{N}$ and \mathfrak{S}_m denotes the group of permutations of $\{1, 2, \ldots, m\}$.

We use the following lemma to compute annihilators of highest weight modules:

Lemma 2.3. Let $\lambda \in \mathfrak{h}^{\vee}$, and F be a \mathfrak{g} -submodule of $U(\mathfrak{g})$ with respect to the adjoint action. Then $\operatorname{sym}(F)$ is contained in the annihilator $\operatorname{Ann} L(\lambda)$ if and only if the zero weight space $\operatorname{sym}(F_0)$ annihilates l_{λ} .

Proof. It suffices to show the "if" part. Suppose $\operatorname{sym}(F_0)l_{\lambda} = 0$. We write \mathbb{C}_{λ} for the character of the Borel subalgebra \mathfrak{b} where \mathfrak{h} acts by λ and the nilradical acts trivially. Then $L(\lambda)$ is isomorphic to the quotient of the Verma module $M(\lambda) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} \mathbb{C}_{\lambda}$ by the maximal proper submodule. Let $m_{\lambda} \in M(\lambda)$ be a nonzero highest weight vector of weight λ . Let proj_2 be the projection to the second component of the decomposition $U(\mathfrak{g}) = U(\mathfrak{g})\mathfrak{n} \oplus U(\mathfrak{b}^-)$. Since F is a \mathfrak{g} -module, $\operatorname{sym}(F)M(\lambda)$ is a \mathfrak{g} -submodule of $M(\lambda)$. By the Poincaré-Birkoff-Witt theorem, we see $\operatorname{sym}(F)M(\lambda) = \operatorname{sym}(F)U(\mathfrak{n}^-)m_{\lambda} = U(\mathfrak{n}^-)\operatorname{sym}(F)m_{\lambda} = U(\mathfrak{n}^-)\operatorname{proj}_2(\operatorname{sym}(F))m_{\lambda}$. Hence the λ weight space of $\operatorname{sym}(F)M_{\lambda}$ is $\operatorname{sym}(F_0)m_{\lambda}$, which is zero by our assumption. Therefore the submodule $\operatorname{sym}(F)M(\lambda)$ is proper, and $\operatorname{sym}(F)L(\lambda)$ is zero.

Proposition 2.4. Let $a \in \mathbb{C}, \lambda \in \mathfrak{h}^{\vee}$. Then $\operatorname{sym}(F^a)$ annihilates the highest weight module $L(\lambda)$ if and only if $\lambda = \lambda(i, a)$ for some $1 \le i \le n$.

For the proof of Proposition 2.4, we use the following

Lemma 2.5. Assume $n \ge 4$. The symmetrization sym $(F(e_1 + e_2 - e_{n-1} - e_n))$ annihilates $L(\lambda)$ if and only if $\lambda = \lambda(i, a)$ for some $1 \le i \le n$ and $a \in \mathbb{C}$.

Proof. By Lemma 2.3, Proposition 2.2 and the definition of $\mathcal{M}_{i < j < k < l}$, we see that $\operatorname{sym}(F(e_1 + e_2 - e_{n-1} - e_n)) \subset \operatorname{Ann} L(\lambda)$ is equivalent to $\operatorname{sym}(T(\mathcal{M}_{i,j,k,l}))l_{\lambda} = 0$ for i < j < k < l, and to

(2.6)

$$sym(T_{i,i}T_{j,j} - T_{i,j}T_{j,i} + T_{k,k}T_{l,l} - T_{k,l}T_{l,k})l_{\lambda}$$

$$= sym(T_{i,i}T_{k,k} - T_{i,k}T_{k,i} + T_{j,j}T_{l,l} - T_{j,l}T_{l,j})l_{\lambda}$$

$$= sym(T_{i,i}T_{l,l} - T_{i,l}T_{l,i} + T_{j,j}T_{k,k} - T_{j,k}T_{k,j})l_{\lambda}$$

for i < j < k < l. Since we have

$$sym(T_{r,r}T_{s,s} - T_{r,s}T_{s,r})l_{\lambda} = \{\lambda_r\lambda_s - (\lambda_r - \lambda_s)/2\}l_{\lambda} = \{(\lambda_r + 1/2)(\lambda_s - 1/2) + 1/4\}l_{\lambda}$$
for $1 \le r < s \le n$, the equations (2.6) are equivalent to

(2.7)

$$(\lambda_i + 1/2)(\lambda_j - 1/2) + (\lambda_k + 1/2)(\lambda_l - 1/2)$$

$$= (\lambda_i + 1/2)(\lambda_k - 1/2) + (\lambda_j + 1/2)(\lambda_l - 1/2)$$

$$= (\lambda_i + 1/2)(\lambda_l - 1/2) + (\lambda_j + 1/2)(\lambda_k - 1/2)$$

for i < j < k < l. The equations (2.7) hold if and only if

$$(\lambda_i - \lambda_l + 1)(\lambda_j - \lambda_k) = (\lambda_i - \lambda_j)(\lambda_k - \lambda_l) = 0 \quad \text{for } i < j < k < l,$$

which implies the assertion by the condition $\sum_i \lambda_i = 0$.

Proof of Proposition 2.4. By Lemma 2.3, $\operatorname{sym}(F^a) \subset \operatorname{Ann} L(\lambda)$ is equivalent to $\operatorname{sym}(F_0^a)l_{\lambda} = 0$. We see $F(e_1 - e_1)^a = \{\sum_{i=1}^{n} A_i T_{i,i}, T_{i,i} - a(n-2)/n \sum_{i=1}^{n} A_i T_{i,i} \mid \sum_{i=1}^{n} A_i = 0\}$ and

We see
$$F(e_1 - e_n)_0^a = \{\sum_{i,k} A_i T_{i,k} T_{k,i} - a(n-2)/n \sum_i A_i T_{i,i} \mid \sum_i A_i = 0\}$$
 and

$$\sum_{i=1}^{n} \sum_{j=1}^{n} A_j T_{i,j} = 0$$

$$\operatorname{sym}\left(\sum_{i,k} A_i T_{i,k} T_{k,i} - \frac{\langle \cdot \cdot \rangle}{n} \sum_i A_i T_{i,i}\right) l_{\lambda}$$

$$= \left(\sum_i A_i \lambda_i^2 + \frac{1}{2} \sum_{i < k} A_i [T_{i,k}, T_{k,i}] + \frac{1}{2} \sum_{i > k} A_i [T_{k,i}, T_{i,k}] - \frac{a(n-2)}{n} \sum_i A_i \lambda_i\right) l_{\lambda}$$

$$= \sum_i A_i C_i l_{\lambda},$$

where we put

$$C_i := \lambda_i^2 + \left(\frac{n+1}{2} - i - \frac{a(n-2)}{n}\right)\lambda_i + \frac{1}{2}\sum_{k=1}^{i-1}\lambda_k - \frac{1}{2}\sum_{k=i+1}^n \lambda_k$$

Therefore sym $(F(e_1 - e_n)_0^a)l_{\lambda} = 0$ if and only if the scalar C_i is independent of $1 \le i \le n$, that is,

(2.8)
$$(\lambda_{i+1} - \lambda_i) \left(\lambda_{i+1} + \lambda_i + \frac{n}{2} - i - \frac{a(n-2)}{n} \right) = 0$$
 for $1 \le i \le n-1$.

When n = 2, (2.8) holds and any element in \mathfrak{h}^{\vee} can be written as $\lambda(1, a)$ for some $a \in \mathbb{C}$, and the assertion follows. When n = 3, the condition $\sum \lambda_i = 0$ implies that (2.8) is equivalent to $\lambda = (-1 + 2a/3, (1/2 - a/3)\mathbf{1}_2), (-1/2 - a/3, 2a/3, 1/2 - a/3), ((-1/2 - a/3)\mathbf{1}_2, 1 + 2a/3)$, which is our claim by $F_0^a = F(e_1 - e_3)_0^a$ and (2.5).

When $n \ge 4$, the definition of F^a , Lemma 2.5 and the above argument imply that $\operatorname{sym}(F_0^a)l_{\lambda} = 0$ if and only if λ is equal to $\lambda(r, c)$ satisfying (2.8) for some $1 \le r \le n$ and $c \in \mathbb{C}$. Since $\lambda(r, c)$ satisfies (2.8) exactly when c = a, we obtain the desired conclusion.

Remark 2.6. The infinitesimal character of $L(\lambda(i, a))$ is the orbit through $\lambda(i, a) + \rho = \sum_{k=1}^{i-1} (n/2 - k - a/n)e_k + a(n-1)/ne_i + \sum_{k=i+1}^{n} (n/2 - k + 1 - a/n)e_k$ and does not depend on $1 \le i \le n$. In particular, the Casimir element sym(Ω) acts by

$$\begin{split} \|\lambda(n,a) + \rho\|^2 &- \|\rho\|^2 \\ &= \sum_{i=1}^{n-1} \left\{ \left(\frac{n+1}{2} - i - \frac{2a+n}{2n}\right)^2 - \left(\frac{n+1}{2} - i\right)^2 \right\} + \frac{a^2(n-1)^2}{n^2} - \frac{(n-1)^2}{4} \\ &= \frac{(n-1)(2a+n)}{2n} \left(\frac{2a+n}{2n} - 2\left(\frac{n+1}{2} - \frac{n}{2}\right)\right) + \frac{(n-1)^2(2a+n)(2a-n)}{4n^2} \\ &= \frac{(n-1)(2a+n)(2a-n)}{4n}, \end{split}$$

where the norm $\|\cdot\|$ on \mathfrak{h}^{\vee} is induced from the invariant bilinear form *B* defined in (2.1).

Definition 2.7. For $a \in \mathbb{C}$, we define J_a to be the two-sided ideal in $U(\mathfrak{g})$ generated by the subspace sym (F^a) and the element sym $(\Omega) - (n-1)(a+n/2)(a-n/2)/n$:

$$J_a := \left\langle \operatorname{sym}(X), \operatorname{sym}(\Omega) - \frac{(n-1)(2a+n)(2a-n)}{4n} \mid X \in F^a \right\rangle$$

Theorem 2.8. (1) Let J be a two-sided ideal of $U(\mathfrak{g})$. The following are equivalent:

(i) $J = J_a$ for some $a \in \mathbb{C}$.

H. TAMORI

(ii) the associated graded ideal gr J is equal to the ideal $\mathcal{I}(\overline{\mathcal{O}^{\min}})$ defined by the closure $\overline{\mathcal{O}^{\min}}$ of the minimal nilpotent coadjoint orbit in \mathfrak{g}^{\vee} .

(iii) J is completely prime, primitive and has $\overline{\mathcal{O}^{\min}}$ as its associated variety. Moreover, we have $J_a = J_{a'}$ if and only if a = a' when $n \ge 3$, and $a = \pm a'$ when n = 2.

(2) Let $\lambda \in \mathfrak{h}^{\vee}$ and $a \in \mathbb{C}$. Then the annihilator of $L(\lambda)$ is equal to J_a if and only if $\lambda = \lambda(i, a)$ for some $1 \leq i \leq n$ satisfying

(2.9)
$$a \notin \frac{n}{2} + \mathbb{N} \text{ if } i = 1 \text{ and } a \notin -\frac{n}{2} - \mathbb{N} \text{ if } i = n.$$

- **Remark 2.9.** (1) When \mathfrak{g} is simple Lie algebra not of type A, there uniquely exists the two-sided ideal J satisfying (ii) (or equivalently, (iii)) [4, Theorem 3.1]. The ideal is called the Joseph ideal.
 - (2) The equivalence of (i) and (ii), and (2) for the case a = n/2 i, n/2 i + 1 or i = 1, n (that is, the case where λ extends to a character of a standard maximal parabolic subalgebra) can be deduced form the argument in [2, Section 7.8] (there is some typo: c' should be c'(n+2)/n except for " $\phi = \phi^a + c'\phi^{s}$ ").

Proof of Theorem 2.8. Let $a \in \mathbb{C}$. We first claim gr $J_a = \mathcal{I}(\overline{\mathcal{O}^{\min}})$ and (2). By Proposition 2.4 and Remark 2.6, we have $J_a \subset \operatorname{Ann} L(\lambda)$ if and only if $\lambda = \lambda(i, a)$ for $1 \leq i \leq n$. Hence the proof of (2) is reduced to show that $J_a = \operatorname{Ann} L(\lambda(i, a))$ exactly when the condition (2.9) holds.

By [6, Theorem III.2.1] and Proposition 2.1, the ideal $\mathcal{I}(\overline{\mathcal{O}^{\min}})$ is generated by $F(e_1 + e_2 - e_{n-1} - e_n)$ (if $n \ge 4$), $F(e_1 - e_n)$ (if $n \ge 3$) and F(0). Therefore the definition of J_a and $J_a \subset \operatorname{Ann} L(\lambda(i, a))$ shows $\mathcal{I}(\overline{\mathcal{O}^{\min}}) \subset \operatorname{gr} J_a \subset \operatorname{gr} \operatorname{Ann} L(\lambda(i, a))$.

Any ideal strictly containing $\mathcal{I}(\overline{\mathcal{O}^{\min}})$ has finite-codimension in $S(\mathfrak{g})$ since $S^k(\mathfrak{g})$ decomposes to the direct sum of the k-th Cartan power of \mathfrak{g} and $\mathcal{I}(\overline{\mathcal{O}^{\min}}) \cap S^k(\mathfrak{g})$ as a \mathfrak{g} -module for $k \in \mathbb{N}$ [6, Proposition III.1.1]. Moreover, we see that gr Ann $L(\lambda(i, a))$ has infinite-codimension in $S(\mathfrak{g})$ if and only if $L(\lambda(i, a))$ is infinite-dimensional, or the condition (2.9) holds. In particular, the ideal J_a has infinite-codimension by $J_a \subset \operatorname{Ann} L(1, a) \cap \operatorname{Ann} L(n, a)$ and $(n/2 + \mathbb{N}) \cap (-n/2 + \mathbb{N}) = \emptyset$. Therefore we obtain $\mathcal{I}(\overline{\mathcal{O}^{\min}}) = \operatorname{gr} J_a \subset \operatorname{gr} \operatorname{Ann} L(\lambda(i, a))$ and the last inclusion becomes an equality exactly in the case (2.9), which proves our claims.

We next prove (1). By $\mathcal{I}(\overline{\mathcal{O}^{\min}}) = \operatorname{gr} J_a$, the ideal J_a is completely prime since $\mathcal{I}(\overline{\mathcal{O}^{\min}})$ is prime in $S(\mathfrak{g})$, and the associated variety Ass J_a is $\overline{\mathcal{O}^{\min}}$. Hence (i) implies (ii) and (iii).

Let us prove (ii) \Rightarrow (i). Assume (ii). By the above result by Garfinkle, there exist scalars $a, c \in \mathbb{C}$ such that F^a and $\operatorname{sym}(\Omega) - c$ is included in J. Take a maximal (hence primitive) ideal J^{\max} containing J. By the work of Duflo [9, Satz 7.3], J^{\max} is the annihilator of some irreducible highest weight module. It follows from $F^a \subset J^{\max}$ and Proposition 2.4 that $J^{\max} = \operatorname{Ann} L(\lambda(i, a))$ for some $1 \leq i \leq n$. Since J^{\max} contains $\operatorname{sym}(\Omega) - (n-1)(a-n/2)(a+n/2)/n$ and does not contain scalars, we see c = (n-1)(a-n/2)(a+n/2)/n and $J_a \subset J$. Then $\operatorname{gr} J_a = \mathcal{I}(\overline{\mathcal{O}^{\min}}) = \operatorname{gr} J$ implies $J = J_a$, and (i) follows.

Let us prove (iii) \Rightarrow (i). Assume (iii). By [17, Théorème IV.1], there exist a standard parabolic subalgebra \mathfrak{q} and a character χ of \mathfrak{q} such that $J = \operatorname{Ann} U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \chi$. By [9, Satz 10.9], we have

 $\dim \mathfrak{g}/\mathfrak{q} = \operatorname{Dim}(U(\mathfrak{g}) \otimes_{U(\mathfrak{g})} \chi) = \operatorname{Dim}(U(\mathfrak{g})/J)/2 = \dim \overline{\mathcal{O}^{\min}}/2 = n-1,$

where Dim denotes the Gelfand-Kirillov dimension. Therefore \mathfrak{q} is the maximal parabolic subalgebra corresponding to the partition (1, n-1) or the one corresponding

to (n-1,1), and χ is equal to $\lambda(1,a)$ or $\lambda(n,a)$ on \mathfrak{h} for some $a \in \mathbb{C}$, respectively. By Lemma 2.10, we see $J_a \subset J$, and (i) follows from $\operatorname{gr} J_a = \mathcal{I}(\mathcal{O}^{\min}) \supset \operatorname{gr} J$.

Finally, we prove the last assertion of (1). Suppose $J_a = J_{a'}$. It suffices to show a = a' when $n \ge 3$, and $a = \pm a'$ when n = 2. By gr $J_a = \mathcal{I}(\mathcal{O}^{\min})$, we see that J_a does not contain a nonzero element in \mathfrak{g} nor a nonzero scalar. Therefore the nonzeroness of F^a and $F^{a'}$ shows a = a' when $n \ge 3$. When n = 2, the ideal J_a is generated by $\operatorname{sym}(\Omega) - (a^2 - 1)/2$ and we see $a = \pm a'$.

The next lemma is used not only in the proof of Theorem 2.8 but also in the construction of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules in Theorem 5.3.

Lemma 2.10. Let $a \in \mathbb{C}$ and \mathfrak{q} be the standard parabolic subalgebra $\mathfrak{q}_{(1,n-1)}$ (resp. $\mathfrak{q}_{(n-1,1)}$) corresponding to the partition (1, n-1) (resp. (n-1, 1)). Then the ideal J_a annihilates the generalized Verma module $U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \mathbb{C}_{\lambda(1,a)}$ (resp. $U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \mathbb{C}_{\lambda(n,a)}$).

Proof. Let J be the anti-diagonal n-by-n matrix whose anti-diagonal entries are one. The involution on \mathfrak{g} defined by $X \mapsto -J^t X J$ for $X \in \mathfrak{g}$ preserves \mathfrak{h} . It maps $\mathfrak{q}_{(1,n-1)}$ onto $\mathfrak{q}_{(n-1,1)}$, and induces an isomorphism on \mathfrak{h}^{\vee} sending $\lambda(1,a)$ to $\lambda(n,-a)$ for $a \in \mathbb{C}$. Hence it suffices to show our assertion for $\mathfrak{q} = \mathfrak{q}_{(1,n-1)}$.

Since J_a and $\operatorname{Ann}(U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \mathbb{C}_{\lambda(1,a)})$ have the same infinitesimal character, it suffices to prove $\operatorname{sym}(F^a) \subset \operatorname{Ann}(U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \mathbb{C}_{\lambda(1,a)})$. Let $m_{\lambda(1,a)}$ be a nonzero highest weight vector of weight $\lambda(1, a)$ in $U(\mathfrak{g}) \otimes_{U(\mathfrak{q})} \mathbb{C}_{\lambda(1,a)}$, and write $(F^a)^-$ for the space of lowest weight vectors in F^a . We have

 $\operatorname{sym}(F^a)U(\mathfrak{g})m_{\lambda(1,a)} = U(\mathfrak{g})\operatorname{sym}(F^a)m_{\lambda(1,a)} = U(\mathfrak{g})\operatorname{sym}((F^a)^-)m_{\lambda(1,a)}.$

In the last equality, we used $\operatorname{sym}([\mathfrak{n}, (F^a)^-])m_{\lambda(1,a)} = [\mathfrak{n}, \operatorname{sym}((F^a)^-)]m_{\lambda(1,a)} = \mathfrak{n}\operatorname{sym}((F^a)^-)m_{\lambda(1,a)}$, where \mathfrak{n} denotes the subalgebra of \mathfrak{g} consisting of all positive root vectors. Therefore we are reduced to see $\operatorname{sym}((F^a)^-)m_{\lambda(1,a)} = 0$. From Proposition 2.1, the subspace $(F^a)^-$ is spanned by $T_{n,1}T_{n-1,2} - T_{n,2}T_{n-1,1}$ (if $n \geq 4$) and $\sum_k T_{n,k}T_{k,1} - a(n-2)/nT_{n,1}$. By $T_{n-1,2}$ (if $n \geq 4$), $T_{n,k} \in [\mathfrak{q},\mathfrak{q}]$ for $2 \leq k \leq n-1$, we obtain

$$\begin{aligned} \operatorname{sym}(T_{n,1}T_{n-1,2} - T_{n,2}T_{n-1,1})m_{\lambda(1,a)} &= 0 \quad \text{if } n \ge 4, \\ \operatorname{sym}\left(\sum_{k} T_{n,k}T_{k,1} - \frac{a(n-2)}{n}T_{n,1}\right)m_{\lambda(1,a)} \\ &= \left(\lambda(1,a)_1 - \frac{1}{2} + \lambda(1,a)_n + \frac{n-1}{2} - \frac{a(n-2)}{n}\right)T_{n,1}m_{\lambda(1,a)} = 0, \end{aligned}$$

and the proof is complete.

3. MINIMAL REPRESENTATIONS

In this section, we extend the definition of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules to real simple Lie algebras including type A, and describe some properties on their \mathfrak{k} -types and a criterion for isomorphism. In this section, we do not restrict ourselves to simple Lie algebras of type A.

We will write \mathfrak{g}_0 for a real simple Lie algebra whose complexification \mathfrak{g} is simple. Fix a Cartan involution θ on \mathfrak{g}_0 , and write $\mathfrak{g}_0 = \mathfrak{k}_0 + \mathfrak{p}_0$ for the Cartan decomposition with respect to θ .

- **Definition 3.1.** (1) An irreducible $(\mathfrak{g}, \mathfrak{k})$ -module is called *minimal* if the associated graded ideal of the annihilator is the ideal defined by the closure of the minimal nilpotent coadjoint orbit in \mathfrak{g}^{\vee} .
 - (2) Let $a \in \mathbb{C}$ and assume $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$ $(n \geq 2)$. An irreducible $(\mathfrak{g}, \mathfrak{k})$ -module is called *a-minimal* if the annihilator equals the ideal J_a (see Definition 2.7).

H. TAMORI

(3) An irreducible admissible representation of a simple Lie group G is called minimal (resp. a-minimal) if the underlying (g, t)-module is minimal (resp. a-minimal).

Remark 3.2. Let V be an irreducible $(\mathfrak{g}, \mathfrak{k})$ -module.

- (1) When \mathfrak{g} is not of type A, the $(\mathfrak{g}, \mathfrak{k})$ -module V is minimal if and only if the annihilator of V is the Joseph ideal (see Remark 2.9 (1)). When $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$ $(n \geq 2), V$ is minimal if and only if V is *a*-minimal for some $a \in \mathbb{C}$, which holds if and only if Ann V is a completely prime primitive ideal whose associated variety is the closure of the minimal nilpotent orbit by Theorem 2.8 (1).
- (2) Assume V is minimal. Write $\mathfrak{k}^{\perp} := \{X \in \mathfrak{g}^{\vee} \mid X(\mathfrak{k}) = 0\}$. Since V is infinite-dimensional, we have $0 \subsetneq \operatorname{Ass} V \subset (\operatorname{Ass} \operatorname{Ann} V) \cap \mathfrak{k}^{\perp} = \overline{\mathcal{O}^{\min}} \cap \mathfrak{k}^{\perp} = (\mathcal{O}^{\min} \cap \mathfrak{k}^{\perp}) \cup \{0\}$. Hence, for the existence of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules, we have a necessary condition

$$(3.1) \qquad \qquad \mathcal{O}^{\min} \cap \mathfrak{k}^{\perp} \neq \emptyset.$$

In particular, there exist no minimal $(\mathfrak{g}, \mathfrak{k})$ -modules for $\mathfrak{g}_0 = \mathfrak{su}(n), \mathfrak{sl}(n, \mathbb{H})$ $(n \geq 2)$ by [18, Proposition 4.1].

(3) Assume g = sl(n, C) (n ≥ 2) and V is a-minimal. Then the contragredient (g, t)-module of V is (-a)-minimal. When g₀ is su(p,q) with p+q = n (resp. sl(n, R)), the complex conjugate of V is (-ā)-minimal (resp. ā-minimal). In particular, if n ≥ 3 and V admits a nondegenerate invariant Hermitian form, then a ∈ R (resp. a ∈ √-1R) by Theorem 2.8 (1).

For a reductive Lie algebra \mathfrak{l} , we write \mathfrak{l}_{ss} and $\mathfrak{z}(\mathfrak{l})$ for the derived subalgebra $[\mathfrak{l}, \mathfrak{l}]$ and the center, respectively.

From now, we also assume (3.1). Fix a Cartan subalgebra \mathfrak{t}_0 of \mathfrak{k}_0 . Define \mathfrak{h}_0^c to be the centralizer $\mathfrak{z}_{\mathfrak{g}_0}(\mathfrak{t}_0)$ of \mathfrak{t}_0 in \mathfrak{g}_0 . Then \mathfrak{h}_0^c is a maximally compact Cartan subalgebra of \mathfrak{g}_0 [11, Proposition 6.60]. By the assumption (3.1), we can take a positive system $\Sigma(\mathfrak{g},\mathfrak{h}^c)^+$ of the root system $\Sigma(\mathfrak{g},\mathfrak{h}^c)$ such that the highest root ψ is noncompact imaginary and $\theta\phi \in \Sigma(\mathfrak{g},\mathfrak{h}^c)^+$ for any $\phi \in \Sigma(\mathfrak{g},\mathfrak{h}^c)^+$. Then $\Sigma(\mathfrak{g},\mathfrak{h}^c)^+$ defines a positive system for $(\mathfrak{k}_{ss},\mathfrak{k}_{ss} \cap \mathfrak{t})$ by

$$\Sigma(\mathfrak{k}_{\mathrm{ss}},\mathfrak{k}_{\mathrm{ss}}\cap\mathfrak{t})^{+} := \left\{ \mu \mid \phi \in \Sigma(\mathfrak{g},\mathfrak{h}^{\mathrm{c}})^{+}, \phi = 0 \text{ on } \mathfrak{z}(\mathfrak{k}), \mu = \phi|_{\mathfrak{k}_{\mathrm{ss}}\cap\mathfrak{t}} \right\}.$$

Let (\cdot, \cdot) be the inner product on the real vector space spanned the set of roots $\Sigma(\mathfrak{g}, \mathfrak{h}^c)$ induced by a nondegenerate invariant bilinear form B on \mathfrak{g} . We normalize the invariant form so that $(\psi, \psi) = 2$. Let $\mathfrak{k}^1, \mathfrak{p}^1$ be the subspaces of $\mathfrak{k}, \mathfrak{p}$ spanned by weight spaces of \mathfrak{t} -weights whose inner product with ψ are $(\psi, \psi)/2 = 1$, respectively. We will write $\rho(\mathfrak{k}^1, \mathfrak{t})$ for half the sum of such roots in $\Sigma(\mathfrak{k}, \mathfrak{t})$. Set $(\mathbb{C}\psi)^{\perp} := \mathbb{C} \{ H_{\phi} \in \mathfrak{h}^c \mid \phi \in \Sigma(\mathfrak{g}, \mathfrak{h}^c), (\phi, \psi) = 0 \}$, where H_{ϕ} denotes the element in \mathfrak{h}^c corresponding to $\phi \in (\mathfrak{h}^c)^{\vee}$ under B. Let $\mathfrak{t}^{\perp}_{\text{Heis}}$ be the set of elements in \mathfrak{t}^{\vee} annihilating $(\mathbb{C}\psi)^{\perp} \cap \mathfrak{t}$. Here we regard \mathfrak{t}^{\vee} as a subspace of \mathfrak{h}^c via the invariant form.

Let us check

(3.2)
$$\mathfrak{t}_{\mathrm{Heis}}^{\perp} = \begin{cases} \mathbb{C}\psi & \text{if } \mathfrak{g}_0 \not\cong \mathfrak{su}(p,q), \\ \mathbb{C}(e_1 - e_n) + \mathbb{C}(e_1 + e_n - 2/n\sum_i e_i) & \text{if } \mathfrak{g}_0 = \mathfrak{su}(p,q). \end{cases}$$

Here we put n = p + q and $\{e_i - e_{i+1}\}_{1 \le i < n}$ denotes the simple roots of $\Sigma(\mathfrak{g}, \mathfrak{h}^c)$ in the latter case. If \mathfrak{g} is not of type A, the extended Dynkin diagram of \mathfrak{g} shows that the θ -stable subspace $(\mathbb{C}\psi)^{\perp}$ contains vectors corresponding to $(\operatorname{rank} \mathfrak{g} - 1)$ simple roots. Hence $\mathfrak{t}_{\operatorname{Heis}}^{\perp} = \mathbb{C}\psi$. By the assumption (3.1), the remaining cases are $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{R}), \mathfrak{su}(p, q)$. For these cases, the extended Dynkin diagram of \mathfrak{g} shows

 $\mathbb{C}(e_1 - e_n) \subset \mathfrak{t}_{\mathrm{Heis}}^{\perp} \subset \mathbb{C}(e_1 - e_n) + \mathbb{C}(e_1 + e_n - 2/n\sum_i e_i). \text{ Calculating dim}((\mathbb{C}\psi)^{\perp} \cap \mathfrak{t})$ for these cases, we obtain (3.2).

Let us see some basic properties of minimal $(\mathfrak{g}, \mathfrak{k})$ -modules:

Proposition 3.3. Let V be a minimal $(\mathfrak{g}, \mathfrak{k})$ -module. Then the following hold.

- (1) V is \mathfrak{k} -multiplicity free.
- (2) Assume that a nonzero root vector of root ψ annihilates no nonzero element in V. Then the highest weights of \mathfrak{k} -types in V belong to the subspace

$$-
ho(\mathfrak{k}^{\perp},\mathfrak{t}) + \mathfrak{t}_{\mathrm{Heis}}^{\perp}.$$

(3) Let V' be a $(\mathfrak{g}, \mathfrak{k})$ -module. Assume V' has a common \mathfrak{k} -type as V and Ann $V \subset$ Ann V'. Then V' is isomorphic to V.

Proof. (1) and (3) follows from the same argument as [20, Proposition 3.1].

We next prove (2). Let v be a highest weight vector in the \mathfrak{k} -module V of weight $\mu \in \mathfrak{t}^{\vee}$. Put $d := \dim \mathfrak{k}^1$ and write $\{\phi \in \Sigma(\mathfrak{k}, \mathfrak{t}) \mid (\phi, \psi) = 1\} = \{\phi_i\}_{i=1}^d$. Since the Lie subalgebra $\mathfrak{k}^1 + \mathfrak{p}^1 + \mathfrak{g}_{\psi}$ is a Heisenberg subalgebra, we can take \mathfrak{t} -weight vectors $\begin{aligned} z \in \mathfrak{p}_{\psi}, \, x_i \in \mathfrak{k}^1_{\phi_i} \text{ and } y_i \in \mathfrak{p}^1_{-\psi-\phi_i} \text{ satisfying } [x_i, y_j] = \delta_{i,j} z \text{ for } 1 \leq i, j \leq d. \\ \text{Set } \mathfrak{r} := \mathfrak{z}_{\mathfrak{g}}(\mathfrak{g}_{\psi} + \mathfrak{g}_{-\psi} + \mathbb{C}H_{\psi}) \text{ and define } \Xi \colon \mathfrak{r} \to S^2(\mathfrak{g}) \text{ by} \end{aligned}$

$$\Xi(X) := Xz + \frac{1}{2} \sum_{1 \le i \le d} ([X, x_i]y_i - [X, y_i]x_i).$$

By [5, Proposition 4.3], the linear map Ξ is an injective r-homomorphism and the image of Ξ is contained in the sum E of irreducible g-submodules of $S^2(g)$ which are not isomorphic to the trivial \mathfrak{g} -module and \mathfrak{g} . Since the Joseph ideal and the ideals J_a ($a \in \mathbb{C}$) contain the symmetrization of E by [6] and Definition 2.7, any element in the image of sym $\circ \Xi$ annihilates the minimal (\mathfrak{g}, K) -module V. Hence, for $H \in (\mathbb{C}\psi)^{\perp} \cap \mathfrak{t} \subset \mathfrak{r}$, we have

$$0 = \operatorname{sym} \circ \Xi(H)v = \operatorname{sym} \left(zH + \frac{1}{2} \sum_{i=1}^{d} ([H, x_i]y_i - [H, y_i]x_i) \right) v$$
$$= zHv + \frac{1}{4} \sum_{i=1}^{d} (2\phi_i - \psi)(H)(x_iy_i + y_ix_i)v = (\mu + \rho(\mathfrak{k}^1))(H)zv$$

Here we used $\psi(H) = 0$, $[x_i, y_i] = z$ and $x_i v = 0$ at the last equality. Therefore $\mu + \rho(\mathfrak{k}^1) \in \mathfrak{t}_{\text{Heis}}^{\perp}$ by the assumption.

Remark 3.4. We write $V^{\mathfrak{g}_{\psi}}$ for the set of elements in V annihilated by \mathfrak{g}_{ψ} . Since a minimal $(\mathfrak{g}, \mathfrak{k})$ -module V is irreducible and infinite-dimensional, the subspace $V^{\mathfrak{g}_{\psi}}$ or $V^{\mathfrak{g}_{-\psi}}$ is zero from [22, Lemma 3.2]. Hence by reversing the positivity and making $-\psi$ the highest weight of g if necessary, we obtain the assumption $V^{\mathfrak{g}_{\psi}}=0$ of Proposition 3.3 (2). We remark that if the center of \mathfrak{k} is trivial, then $V^{\mathfrak{g}_{\psi}} = V^{\mathfrak{g}_{-\psi}} = 0.$

The following properties are used to classify minimal $(\mathfrak{g}, \mathfrak{k})$ -modules for $\mathfrak{g} \cong$ $\mathfrak{sl}(n,\mathbb{C})$ in Section 5:

Corollary 3.5. Let V be a minimal $(\mathfrak{g}, \mathfrak{k})$ -module. Then the following hold.

- (1) Assume \mathfrak{g}_0 is Hermitian and $\mathfrak{g}_0 \not\cong \mathfrak{sl}(2,\mathbb{R})$. Then V is an irreducible highest or lowest weight module.
- Assume $V^{\mathfrak{g}_{\psi}} = 0$. If \mathfrak{g}_0 is not isomorphic to $\mathfrak{sl}(n,\mathbb{R})(n \geq 3 \text{ odd or } n = 2)$, (2)then V has pencil \mathfrak{k} -types: there exist a dominant integral weight $\mu_0 \in \mathfrak{t}^{\vee}$ with respect to $\Sigma(\mathfrak{k}_{ss},\mathfrak{k}_{ss}\cap\mathfrak{t})^+$ such that the \mathfrak{k} -type decomposition of V is multiplicity free and is given by the direct sum of the irreducible \mathfrak{k} -modules with highest weight $\mu_0 + k\psi$ for $k \in \mathbb{N}$.

Proof. We may assume (3.1). Moreover, let us first assume $\mathfrak{g} \not\cong \mathfrak{su}(1,2)$. For (1), it suffices to show that $V^{\mathfrak{g}_{\psi}}$ or $V^{\mathfrak{g}_{-\psi}}$ is nonzero. Suppose, contrary to our claim, $V^{\mathfrak{g}_{\psi}} = V^{\mathfrak{g}_{-\psi}} = 0$. By applying Proposition 3.3 (2) to the fixed positive system and the opposite positive system where $-\psi$ is the highest noncompact imaginary root, highest weights of \mathfrak{k} -types in V belong to

$$\{-
ho(\mathfrak{k}^1,\mathfrak{t})+\mathfrak{t}_{\mathrm{Heis}}^{\perp}\}\cap w_l\{
ho(\mathfrak{k}^1,\mathfrak{t})+\mathfrak{t}_{\mathrm{Heis}}^{\perp}\},$$

where w_l denotes the longest element in the Weyl group of $(\mathfrak{k}, \mathfrak{t})$. Therefore $w_l \psi$ belongs to $\mathfrak{t}_{\text{Heis}}^{\perp}$.

When $\mathfrak{g}_0 \not\cong \mathfrak{su}(p,q)$, the equality (3.2) shows $w_l \psi \in \mathbb{C}\psi$ and irreducible components of \mathfrak{k} -module \mathfrak{p} must be one-dimensional. Then \mathfrak{g} is three-dimensional, and is isomorphic to $\mathfrak{sl}(2,\mathbb{C})$, a contradiction.

When $\mathfrak{g}_0 = \mathfrak{su}(p,q)$, the equality (3.2) shows that $e_p - e_{p+1}$ belongs to $\mathbb{C}(e_1 - e_n) + \mathbb{C}(e_1 + e_n - 2/n \sum_i e_i)$, which shows p + q = 2, 3. It contradicts to our assumptions $\mathfrak{g}_0 \not\cong \mathfrak{su}(1,2)$ and (3.1).

We next prove (2) for $\mathfrak{g}_0 \ncong \mathfrak{su}(1,2)$. By (3.2), the set of weights of \mathfrak{p} in $\mathfrak{t}_{\text{Heis}}^{\perp}$ are

(3.3)
$$\begin{cases} \{\pm\psi\} & \text{if } \mathfrak{g} \not\cong \mathfrak{sl}(n,\mathbb{R}) (n \ge 3, \text{ odd}), \mathfrak{su}(1,2), \\ \{\pm\psi/2, \pm\psi\} & \text{if } \mathfrak{g} \cong \mathfrak{sl}(n,\mathbb{R}) \ (n \ge 3, \text{ odd}). \end{cases}$$

Let $V(\mu)$ be the \mathfrak{k} -isotypic component in V of the irreducible \mathfrak{k} -module with highest weight μ . By the assumption and Proposition 3.3 (2) and (3.3), the subspace $\mathfrak{g}V(\mu)$ is included in $V(\mu - \psi) + V(\mu) + V(\mu + \psi)$. Then it follows from the infinitedimensionality of V and Proposition 3.3 (1) that the minimal $(\mathfrak{g}, \mathfrak{k})$ -module V has pencil \mathfrak{k} -types.

It remains to prove assertions for $\mathfrak{g} = \mathfrak{su}(1,2)$. By taking the contragredient if necessary, we may assume that the action of $\mathfrak{g}_{e_1-e_3}$ on V is faithful (see Remark 3.4). From Remark 3.2 (1), there exists some $a \in \mathbb{C}$ such that V is *a*-minimal. Let vbe a nonzero highest weight vector of a \mathfrak{k} -type in V with highest weight μ . By $0 = \operatorname{sym}(\sum_i T_{1,i}T_{i,3} - a/2T_{1,3})v = (\mu_1 + \mu_3 - 1/2 - a/2)T_{1,3}v$ and $V^{\mathfrak{g}_{e_1-e_3}} = 0$, we obtain $\mu_2 = -(a+1)/2$. Hence the difference of two highest weights of \mathfrak{k} -types in V belongs to the line $\mathbb{R}(e_1 - e_3)$. Since the set of dominant integral weights in $\mathbb{R}(e_1 - e_3)$ with respect to $\Sigma(\mathfrak{k}_{ss}, \mathfrak{k}_{ss} \cap \mathfrak{t})^+$ is $\mathbb{N}(e_1 - e_3)$, the minimal $(\mathfrak{g}, \mathfrak{k})$ -module V has pencil \mathfrak{k} -types and V is a lowest weight module. \Box

- **Remark 3.6.** (1) There is another proof of Corollary 3.5 (1) given by Vogan. Since \mathfrak{g} is not of type A_1 , the dimension of every nonzero $\operatorname{Int}(\mathfrak{k})$ -orbit in \mathfrak{p} is greater than one. Then by [23, Theorem 4.6], the associated variety of a minimal $(\mathfrak{g}, \mathfrak{k})$ -module V is irreducible, and included in an irreducible component of \mathfrak{p} . Therefore if \mathfrak{g}_0 is Hermitian, then V is a highest (or lowest) weight module.
 - (2) Even when g₀ ≈ sl(n, R)(n ≥ 3 odd), minimal (g, t)-modules has pencil ttypes, as we can see from the classification of minimal (g, t)-modules given in Theorem 5.3.

4. Covariant differentials

In this section, we construct an intertwining differential operator between parabolically induced representations where the infinite-dimensional composition factors of the kernel are all minimal. When $\mathfrak{g}_0 = \mathfrak{sl}(n,\mathbb{R})$, the \mathfrak{k} -finite vectors in the kernel will be determined in Section 5 for some specific cases. Since the method will be applied to a new construction of many minimal representations in a subsequent paper, we do not restrict ourselves to simple Lie algebras of type A.

Let G be a connected simple Lie group with finite center, and fix an Iwasawa decomposition $G = KA_{\min}N_{\min}$. Assume that the complexification \mathfrak{g} is simple and

 $\mathfrak{g} \cong \mathfrak{sl}(2, \mathbb{C})$. Write $\Sigma(\mathfrak{g}, \mathfrak{a}_{\min})$ for the restricted root system for $(\mathfrak{g}_0, \mathfrak{a}_{\min,0})$, and $\Sigma(\mathfrak{g}, \mathfrak{a}_{\min})^+$ for the positive system defined as the set of \mathfrak{a}_{\min} -weights of \mathfrak{n}_{\min} . Let Q = MAN be a standard parabolic subgroup of G and its Langlands decomposition. Take a maximally split Cartan subalgebra \mathfrak{h}_0^s of \mathfrak{g}_0 containing $\mathfrak{a}_{\min,0}$. Then the center $\mathfrak{z}(\mathfrak{m})$ of \mathfrak{m} is contained in \mathfrak{k} , and we have $\mathfrak{h}^s = (\mathfrak{m}_{ss} \cap \mathfrak{h}^s) \oplus \mathfrak{z}(\mathfrak{m}) \oplus \mathfrak{a}$. Via this direct sum decomposition, we will regard $\mathfrak{z}(\mathfrak{m})^{\vee}$ as a subspace of $(\mathfrak{h}^s)^{\vee}$. Moreover, any element $\mu \in \mathfrak{z}(\mathfrak{m})^{\vee}$ extends to a character of \mathfrak{m} via $\mathfrak{m} = \mathfrak{m}_{ss} \oplus \mathfrak{z}(\mathfrak{m})$. We use the same symbol μ for the character.

Let (σ, V) be a finite-dimensional irreducible representation of M and ν an element in the dual \mathfrak{a}^{\vee} . We further assume that the \mathfrak{m} -module V irreducibly decomposes to a multiple of a character. By abuse of notation, we write $d\sigma \in \mathfrak{z}(\mathfrak{m})^{\vee}$ for the character. Set

$$\lambda := -d\sigma - \nu \in \mathfrak{z}(\mathfrak{m})^{\vee} + \mathfrak{a}^{\vee} \subset (\mathfrak{h}^{\mathrm{s}})^{\vee}.$$

We use the same symbol for the character of \mathfrak{q} via $\mathfrak{q}/(\mathfrak{m}_{ss} + \mathfrak{n}) \cong \mathfrak{z}(\mathfrak{m}) \oplus \mathfrak{a}$. By letting M act by σ , A act by $\exp(\nu)$ and N act trivially, we obtain an irreducible representation (σ_{ν}, V) of Q. We write $C^{\infty}(G, \sigma_{\nu})$ to be the space consisting of all smooth functions from G to V. The representation of G induced by σ_{ν} is defined as

$$C^{\infty}(G, \sigma_{\nu})^{Q} := \{ f \in C^{\infty}(G, \sigma_{\nu}) \mid f(gq) = \sigma_{\nu}(q^{-1})f(g) \text{ for } g \in G, q \in Q \},\$$

where the action of G is induced by the left transition L on $C^{\infty}(G, \sigma_{\nu})$:

$$(L(h)f)(g) := f(h^{-1}g) \text{ for } h, g \in G, f \in C^{\infty}(G, \sigma_{\nu}).$$

We define a bilinear map $\Psi: U(\mathfrak{g}) \times C^{\infty}(G, \sigma_{\nu}) \to C^{\infty}(G, \sigma_{\nu})$ by the right differentiation:

$$\Psi(X_1 X_2 \cdots X_n, f)(g) := \left. \frac{d^n}{dt_1 \cdots dt_n} \right|_{t_1 = \cdots = t_n = 0} f(g e^{t_1 X_1} \cdots e^{t_n X_n})$$

for $n \in \mathbb{N}, X_1, \dots, X_n \in \mathfrak{g}_0, f \in C^{\infty}(G, \sigma_{\nu}), g \in G.$

We write $(\sigma_{\nu})^{\vee}$ for the contragredient representation of Q. Then the left ideal generated by $\operatorname{Ann}_{U(\mathfrak{q})}(\sigma_{\nu})^{\vee}$ is given by

(4.1)
$$I(\mathfrak{q},\lambda) := U(\mathfrak{g})\operatorname{Ann}_{U(\mathfrak{q})}(\lambda).$$

We remark that $I(\mathfrak{q}, \lambda)$ is stable under the adjoint action of Q. Let proj be the canonical projection

(4.2)
$$\operatorname{proj}: U(\mathfrak{g}) \to U(\mathfrak{g})/I(\mathfrak{q},\lambda)$$

Then Ψ factors through proj to induce a $(G \times Q)$ -intertwining operator

(4.3)
$$U(\mathfrak{g})/I(\mathfrak{g},\lambda) \otimes C^{\infty}(G,\sigma_{\nu})^{Q} \to C^{\infty}(G,\sigma_{\nu})$$

by [14, Lemma 2.14]. Here the $(G \times Q)$ -actions on $U(\mathfrak{g})/I(\mathfrak{q}, \lambda)$ and $C^{\infty}(G, \sigma_{\nu})$ are given by

$$(h,q)\operatorname{proj}(X) = \operatorname{proj}(\operatorname{Ad}(q)(X)), \quad (h,q)f(g) = \sigma_{\nu}(q)f(h^{-1}gq)$$

for $g, h \in G, q \in Q, X \in U(\mathfrak{g})$ and $f \in C^{\infty}(G, \sigma_{\nu})$.

Define ι to be the anti-involution of $U(\mathfrak{g})$ by

$$\iota(X) := -X \text{ for } X \in \mathfrak{g}$$

Definition 4.1. We define a \mathfrak{g} -submodule F of $\bigoplus_{0 \le i \le 2} S^i(\mathfrak{g})$ as follows.

- When \mathfrak{g} is not of type A, put F to be the \mathfrak{g} -module complement of the trivial \mathfrak{g} -submodule in $S^2(\mathfrak{g})$.
- When $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$, fix $a \in \mathbb{C}$ and put F to be F^a (see (2.4)).

H. TAMORI

Define W to be the Q-submodule of $U(\mathfrak{g})/I(\mathfrak{q},\lambda)$ defined by

$$W := \operatorname{proj} \circ \iota \circ \operatorname{sym}(F) \subset U(\mathfrak{g})/I(\mathfrak{q},\lambda).$$

From (4.3), the tensor-hom adjunction and the triviality of the action of Q on $C^{\infty}(G, \sigma_{\nu})^{Q}$, we obtain a G-intertwining second-order differential operator

$$D = D(Q, \sigma_{\nu}) \colon C^{\infty}(G, \sigma_{\nu})^Q \to C^{\infty}(G, W^{\vee} \otimes \sigma_{\nu})^Q$$

defined by $\langle w, Df(g) \rangle := \Psi(w, f)(g)$ for $w \in W, f \in C^{\infty}(G, \sigma_{\nu})^Q, g \in G$. Here $\langle \cdot, \cdot \rangle$ denotes the pairing of W and its dual W^{\vee} .

In the rest of this section, we study properties of the kernel of D. The kernel of D picks out functions annihilated by sym(F):

Lemma 4.2. In the setting of Definition 4.1, we have $\text{Ker } D = \{f \in C^{\infty}(G, \sigma_{\nu})^Q \mid dL(\text{sym}(F))f = 0\}$. Here dL denotes the differential of the left transition L.

Proof. Let $f \in C^{\infty}(G, \sigma_{\nu})^Q$. Then Df = 0 is equivalent to $\Psi(\iota(X), f)(g) = 0$ for $X \in \text{sym}(F), g \in G$. For $n \in \mathbb{N}, X_1, \ldots, X_n \in \mathfrak{g}_0, f \in C^{\infty}(G, \sigma_{\nu})$ and $g \in G$, we have

$$\Psi(\iota(X_1X_2\cdots X_n), f)(g) = \frac{d^n}{dt_l\cdots dt_n}\Big|_{t_1=\cdots=t_n=0} f(ge^{-t_nX_n}\cdots e^{-t_1X_1})$$

= $\frac{d^n}{dt_1\cdots dt_n}\Big|_{t_1=\cdots=t_n=0} f(e^{-t_n\operatorname{Ad}(g)(X_n)}\cdots e^{-t_1\operatorname{Ad}(g)(X_1)}g)$
= $dL(\operatorname{Ad}(g)(X_1X_2\cdots X_n))f(g).$

Therefore $\Psi(\iota(X), f)(g) = dL(\operatorname{Ad}(g)(X))f(g)$ for $X \in U(\mathfrak{g}), f \in C^{\infty}(G, \sigma_{\nu}), g \in G$. Hence Df = 0 is equivalent to dL(Y)f(g) = 0 for $Y \in \operatorname{sym}(F), g \in G$, and the proof is complete.

In the following proposition, the term "(*a*)-minimal" means "minimal" if \mathfrak{g} is not of type *A*, and "*a*-minimal" if $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$ $(n \geq 3)$.

Proposition 4.3. In the setting of Definition 4.1, the following hold.

- (1) Any (a)-minimal subrepresentation of $C^{\infty}(G, \sigma_{\nu})^Q$ is contained in Ker D.
- (2) Any infinite-dimensional irreducible subquotient of Ker D is an (a)-minimal representation.

Proof. Since sym(F) is contained in the Joseph ideal by [6] (when \mathfrak{g} is not of type A) and in J_a by Definition 2.7 (when $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$), the assertion (1) follows from Lemma 4.2.

Let us prove (2). Let τ be an infinite-dimensional irreducible subquotient of Ker *D*. Write *F'* for the image of *F* under the projection from $\bigoplus_{0 \le i \le 2} S^i(\mathfrak{g})$ to $S^2(\mathfrak{g})$. Then *F'* and the trivial \mathfrak{g} -submodule of $S^2(\mathfrak{g})$ generates $\mathcal{I}(\overline{\mathcal{O}^{\min}})$ by [6].

By Lemma 4.2, the annihilator $\operatorname{Ann} \tau$ contains $\operatorname{sym}(F)$. Hence $\operatorname{gr}(\operatorname{Ann} \tau)$ contains F'. Since $\operatorname{Ann} \tau$ has an infinitesimal character, the associated graded ideal $\operatorname{gr}(\operatorname{Ann} \tau)$ contains the trivial \mathfrak{g} -submodule of $S^2(\mathfrak{g})$. Therefore $\operatorname{gr}(\operatorname{Ann} \tau)$ contains $\mathcal{I}(\overline{\mathcal{O}^{\min}})$. Since the ideal $\operatorname{Ann} \tau$ has infinite codimension in $U(\mathfrak{g})$, we have $\operatorname{gr}(\operatorname{Ann} \tau) = \mathcal{I}(\overline{\mathcal{O}^{\min}})$. Therefore τ is (a)-minimal by Theorem 2.8 (1) and Remark 2.9 (1). Here we used the fact that $\operatorname{sym}(F^a) \subset J_b$ ($b \in \mathbb{C}$) implies b = a for $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$ ($n \geq 3$).

The following two lemmas give necessary conditions for Ker $D \neq 0$.

Lemma 4.4. When Ker $D \neq 0$, the annihilator of the irreducible highest weight module $L(\lambda)$ equals the Joseph ideal or the augmentation ideal $U(\mathfrak{g})\mathfrak{g}$ if \mathfrak{g} is not of type A, and contains J_{-a} if $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$.

Proof. Assume Ker $D \neq 0$. Then there exist $f \in \text{Ker } D$ and $g_0 \in G$ with $f(g_0) \neq 0$. Let us show that $\iota \circ \text{sym}(F)$ annihilates $L(\lambda)$. As Lemma 2.3, it suffices to show $X \in I(\mathfrak{q}, \lambda)$ for any weight vector $X \in \iota \circ \text{sym}(F)$ of weight zero. We write $\overline{\mathfrak{n}}$ for the nilradical of the opposite parabolic of \mathfrak{q} . By the Poincaré-Birkhoff-Witt theorem and the decomposition $\mathfrak{g} = \overline{\mathfrak{n}} + \mathfrak{q}$, we see $X \in c + I(\mathfrak{q}, \lambda)$ for some $c \in \mathbb{C}$. By $f \in \text{Ker } D$ and the proof of Lemma 4.2, we have $0 = dL(\text{Ad}(g_0) \circ \iota(X))f(g_0) = \Psi(X, f)(g_0) = cf(g_0)$. Therefore $X \in I(\mathfrak{q}, \lambda)$, and $\iota \circ \text{sym}(F)$ annihilates $L(\lambda)$.

When \mathfrak{g} is not of type A, it follows from [2, Proposition 5.3] that a proper primitive ideal containing $\iota \circ \operatorname{sym}(F) = \operatorname{sym}(F)$ is the Joseph ideal or $U(\mathfrak{g})\mathfrak{g}$, and the assertion follows. When $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$, it follows from the above argument, $\iota \circ \operatorname{sym}(F) = \operatorname{sym}(F^{-a})$ and Proposition 2.4 that λ equals $\lambda(i, -a)$ for some *i*. By Remark 2.6, the ideal J_{-a} annihilates $L(\lambda)$.

Let $\Delta(\mathfrak{g}, \mathfrak{a}_{\min})$ be the set of simple restricted roots, Δ_Q the subset of $\Delta(\mathfrak{g}, \mathfrak{a}_{\min})$ defining the standard parabolic subgroup Q. When Q is minimal, we have $\Delta_Q = \emptyset$. Define

$$\Delta_Q^{\nu} := \{ \alpha \in \Delta(\mathfrak{g}, \mathfrak{a}^{\mathrm{s}}) \mid \operatorname{proj}(\mathfrak{g}_{-\alpha}) = \operatorname{proj}(F_{-\alpha}) \},\$$

where proj is defined in (4.2) and the subscript $-\alpha$ means the \mathfrak{a}_{\min} -weight space of weight $-\alpha$. We write $Q^{\nu} = M^{\nu}A^{\nu}N^{\nu}$ for the standard parabolic subgroup corresponding to Δ_Q^{ν} and its Langlands decomposition. By $\Delta_Q \subset \Delta_Q^{\nu}$, we see $Q \subset Q^{\nu}$. Therefore we have the decomposition $\mathfrak{a} = \mathfrak{a}^{\nu} \oplus (\mathfrak{a} \cap \mathfrak{m}^{\nu})$ and $\mathfrak{z}(\mathfrak{m}) =$ $(\mathfrak{z}(\mathfrak{m}) \cap \mathfrak{m}_{ss}^{\nu}) \oplus \mathfrak{z}(\mathfrak{m}^{\nu})$.

Lemma 4.5. Assume that Ker D is nonzero. Then the character ν (resp. $d\sigma$) is zero on $\mathfrak{a} \cap \mathfrak{m}^{\nu}$ (resp. $\mathfrak{z}(\mathfrak{m}) \cap \mathfrak{m}_{ss}^{\nu}$) to define a character of \mathfrak{a}^{ν} (resp. $\mathfrak{z}(\mathfrak{m}^{\nu})$), and there exists an irreducible representation σ' of M^{ν} satisfying the following conditions.

- (1) The representation space of σ' agrees with that of σ .
- (2) The differentiated action of $(\mathfrak{m}_0^{\nu})_{ss}$ is trivial.
- (3) $\sigma'(m) = \sigma(m)$ for $m \in M$.

Moreover, we have $\operatorname{Ker} D(Q, \sigma_{\nu}) \subset C^{\infty}(G, \sigma'_{\nu})^{Q^{\nu}} \subset C^{\infty}(G, \sigma_{\nu})^{Q}$.

Proof. By the assumption, we can take $f \in \text{Ker } D$, $g_0 \in G$ with $f(g_0) \neq 0$. Then

(4.4)
$$f(gm) = \sigma(m^{-1})f(g), \quad \frac{d}{dt}\Big|_{t=0} f(ge^{tX}) = 0$$

for $g \in G, m \in M, X \in (\mathfrak{m}_0)_{\mathrm{ss}} \cup \mathfrak{n}_0 \cup (\mathfrak{g}_0)_{-\alpha} \ (\alpha \in \Delta_Q^{\nu})$. Moreover, the Lie subalgebra of \mathfrak{g}_0 generated by $(\mathfrak{m}_0)_{\mathrm{ss}}, \mathfrak{n}_0$ and $(\mathfrak{g}_0)_{-\alpha} \ (\alpha \in \Delta_Q^{\nu})$ contains $(\mathfrak{m}_0^{\nu})_{\mathrm{ss}}$:

(4.5)
$$(\mathfrak{m}_0^{\nu})_{\rm ss} \subset \langle (\mathfrak{m}_0)_{\rm ss}, \mathfrak{n}_0, (\mathfrak{g}_0)_{-\alpha} \mid \alpha \in \Delta_Q^{\nu} \rangle.$$

By (4.4) and (4.5), we obtain $(d\sigma + \nu)(H)f(g_0) = \frac{d}{dt}\Big|_{t=0} f(g_0 \exp(-tH)) = 0$ for $H \in (\mathfrak{a}_0 \cap \mathfrak{m}_0^{\nu}) + (\mathfrak{z}(\mathfrak{m}_0) \cap (\mathfrak{m}_0^{\nu})_{ss}) \subset (\mathfrak{m}_0^{\nu})_{ss}$. Therefore ν (resp. $d\sigma$) can be seen as an element in $(\mathfrak{a}^{\nu})^{\vee}$ (resp. $\mathfrak{z}(\mathfrak{m}^{\nu})^{\vee}$).

We next define a representation σ' of M^{ν} . Let $g \in M^{\nu}, v \in V$. Since $f \neq 0$ and σ is finite-dimensional and irreducible, the representation space V is spanned by f(G). Hence we can write $v = \sum_{j=1}^{m} a_j f_j(h_j)$ for some $m \in \mathbb{N}, a_j \in \mathbb{C}, f_j \in \text{Ker } D$ and $h_j \in G$ $(1 \leq j \leq m)$. Then we define $\sigma'(g)v$ to be $\sum_{j=1}^{m} a_j f_j(h_j g^{-1})$. Let us check that $\sigma'(g)v$ is well-defined. It suffices to show $\sum_{j=1}^{m} a_j f_j(h_j g^{-1}) = 0$

Let us check that $\sigma'(g)v$ is well-defined. It suffices to show $\sum_{j=1}^{m} a_j f_j(h_j g^{-1}) = 0$ assuming $\sum_{j=1}^{m} a_j f_j(h_j) = 0$. Since \mathfrak{m}_0^{ν} is generated by \mathfrak{m}_0 and $(\mathfrak{g}_0)_{\pm \alpha}$ ($\alpha \in \Delta_Q^{\nu}$), we can write $g = g_1 \cdots g_n$ for some $n \in \mathbb{N}$ and

$$g_i \in M \cup \exp(\mathfrak{n}_0) \cup \bigcup_{\alpha \in \Delta_Q^{\nu}} \exp((\mathfrak{g}_0)_{-\alpha}) \quad (1 \le i \le n)$$

By (4.4), we see $\sum_{j=1}^{m} a_j f_j(h_j g^{-1}) = \sigma(\prod_{1 \le i \le n, g_i \in M} g_i) \sum_{j=1}^{m} a_j f_j(h_j) = 0$. Hence $\sigma'(g)v$ does not depend on f and the expression of v.

We easily see that $\sigma'(g) \in \operatorname{GL}(V)$ and σ' is a group homomorphism from M^{ν} to $\operatorname{GL}(V)$. In a small neighborhood of the neutral element in M^{ν} , any element g can be smoothly decomposed into g_1g_2 for some $g_1 \in \exp(\mathfrak{g}(\mathfrak{m}_0^{\nu})), g_2 \in \exp((\mathfrak{m}_0^{\nu})_{ss}),$ and $\sigma'(g) = \sigma(g_1)$ by (4.4) and (4.5). Therefore σ' is a smooth representation of M^{ν} . The conditions (1), (2) and (3) and $f \in C^{\infty}(G, \sigma'_{\nu})^{Q^{\nu}}$ follow directly from the definition of σ' .

5. Classification of minimal representations

Let \mathfrak{g}_0 be a real form of $\mathfrak{sl}(n,\mathbb{C})$ $(n \geq 2)$. By Remark 3.2 (2), if there is a minimal $(\mathfrak{g},\mathfrak{k})$ -module, then \mathfrak{g}_0 is isomorphic to $\mathfrak{su}(p,q)$ (p,q > 0, p + q = n) or $\mathfrak{sl}(n,\mathbb{R})$. In this section, we classify *a*-minimal $(\mathfrak{g},\mathfrak{k})$ -modules for $\mathfrak{g}_0 = \mathfrak{su}(p,q)$ $(p,q > 0, p + q = n), \mathfrak{sl}(n,\mathbb{R})$ $(n \geq 3)$ for any $a \in \mathbb{C}$.

We remark that when $\mathfrak{g}_0 = \mathfrak{sl}(2, \mathbb{R}) \cong \mathfrak{su}(1, 1)$, any primitive ideal with infinitecodimension can be written as J_a for some $a \in \mathbb{C}$. Therefore an irreducible $(\mathfrak{g}, \mathfrak{k})$ module is minimal if and only if it is infinite-dimensional, and the classification of minimal $(\mathfrak{g}, \mathfrak{k})$ -module is well-known (see [7, Sections II.1.2–3], for example). As for the classification of unitarizable ones, see [7, Theorems II.1.1.3 and 1.1.5].

Let us first consider the case $\mathfrak{g}_0 = \mathfrak{su}(p,q)$ $(p,q > 0, n = p + q \ge 3)$ and $\mathfrak{k}_0 = \mathfrak{su}(p) \oplus \mathfrak{su}(q)$. We take a compact Cartan subalgebra \mathfrak{t}_0 of \mathfrak{g}_0 as diagonal matrices. We use the notation in Section 2.

Theorem 5.1. Let $\mathfrak{g}_0 = \mathfrak{su}(p,q)$ $(p,q > 0, n = p + q \ge 3)$, $a \in \mathbb{C}$. The following $(\mathfrak{g},\mathfrak{k})$ -modules are a-minimal and any a-minimal $(\mathfrak{g},\mathfrak{k})$ -module is isomorphic to some of them.

(1) In the case
$$p = 1$$
,
 $L(\lambda(1, a)) \quad a \notin n/2 + \mathbb{N}, \qquad L(\lambda(1, -a))^{\vee} \quad a \notin -n/2 - \mathbb{N},$
 $L(\lambda(2, a)) \quad a \in (n - 2)/2 + \mathbb{N}, \qquad L(\lambda(2, -a))^{\vee} \quad a \in -(n - 2)/2 - \mathbb{N}.$
(2) In the case $q = 1$,
 $L(\lambda(n, a)) \quad a \notin -n/2 - \mathbb{N}, \qquad L(\lambda(2, -a))^{\vee} \quad a \notin n/2 + \mathbb{N},$
 $L(\lambda(n - 1, a)) \quad a \in -(n - 2)/2 - \mathbb{N}, \qquad L(\lambda(n - 1, -a))^{\vee} \quad a \in (n - 2)/2 + \mathbb{N}.$
(3) In the case $p, q > 1$,
 $L(\lambda(p, a)) \quad a \in -(p - q)/2 - \mathbb{N}, \qquad L(\lambda(p, -a))^{\vee} \quad a \in (p - q)/2 + \mathbb{N},$
 $L(\lambda(p + 1, a)) \quad a \in -(p - q)/2 + \mathbb{N}, \qquad L(\lambda(p + 1, -a))^{\vee} \quad a \in (p - q)/2 - \mathbb{N}.$

Furthermore, these minimal a-modules are not isomorphic to each others except for the ones caused by

$$\lambda(p, -(p-q)/2) = \lambda(p+1, -(p-q)/2).$$

The \mathfrak{t} -types of an a-minimal $(\mathfrak{g}, \mathfrak{k})$ -module $L(\lambda(i, a))$ (resp. $L(\lambda(i, -a))^{\vee}$) are multiplicity free and the set of highest weights is given by $\lambda(i, a) + \mathbb{N}(-e_p + e_{p+1})$ (resp. $-w_l\lambda(i, -a) + \mathbb{N}(e_1 - e_n)$). Here w_l denotes the longest element of the Weyl group for $(\mathfrak{t}, \mathfrak{t})$.

Remark 5.2. (1) Since $-\lambda(i,a)_j = -\overline{\lambda(i,\overline{a})_j}$ for any $1 \leq j \leq n$, we have $L(\lambda(i,a))^{\vee} \cong \overline{L(\lambda(i,\overline{a}))}$.

(2) As in the proof of Lemma 2.10, there exists an isomorphism $\mathfrak{su}(p,q) \cong \mathfrak{su}(q,p)$ where the $\mathfrak{su}(p,q)$ -module $L(\lambda(i,a))$ maps to the $\mathfrak{su}(q,p)$ -module $L(\lambda(n+1-i,-a))$.

- (3) By [3, Theorem 7.4] or [8], when $L(\lambda(i, a))$ is a-minimal $(\mathfrak{g}, \mathfrak{k})$ -module, it is unitarizable if and only if

 - (i) p = 1, i = 1 and $a \in (-\infty, 2 n/2] \cup \{ n/2 i \mid 1 \le i \le n 3 \}$, (ii) q = 1, i = n and $a \in \{ -n/2 + i \mid 1 \le i \le n 3 \} \cup (-2 + n/2, \infty]$ or (iii) $i \neq 1, n$.
- (4) Recall that O(2,4) is locally isomorphic to SU(2,2). The $(\mathfrak{g},\mathfrak{k})$ -module of the irreducible unitary representation of O(2,4) constructed in [1, 12] is isomorphic to $L(\lambda(2,0)) \oplus L(\lambda(2,0))^{\vee}$, and the two irreducible components are 0-minimal.

Proof of Theorem 5.1. By Corollary 3.5 (1), any *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module is a highest or lowest weight module. From Remark 3.2 (3), it suffices to classify *a*-minimal highest weight $(\mathfrak{g}, \mathfrak{k})$ -modules. Theorem 2.8 (2) shows that the irreducible highest weight \mathfrak{g} -module $L(\lambda)$ is an *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module if and only if the highest weight λ is written as $\lambda(i,a)$ for some $1 \leq i \leq n$, satisfies (2.9), and is dominant integral with respect to $\Sigma(\mathfrak{k}_{ss},\mathfrak{k}_{ss}\cap\mathfrak{t})^+$. From the expression (2.5) of $\lambda(i,a)$ and the fact that $\lambda(i,a) = \lambda(i',a')$ if and only if a = a' and $(i',a') = \lambda(i',a')$ (i,a), (i-1, n/2 - i + 1), (i+1, n/2 - i) hold, we obtain the classification of aminimal $(\mathfrak{g}, \mathfrak{k})$ -modules.

The assertion on \mathfrak{k} -types follows from Corollary 3.5 (2).

The rest of this section is devoted to the case $\mathfrak{g}_0 = \mathfrak{sl}(n, \mathbb{R}) (n \geq 3)$. We will construct an *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module as the \mathfrak{k} -finite vectors of the kernel of an intertwining differential operator between parabolically induced representations. Furthermore, the construction will be applied to show the nonexistence of some *a*-minimal $(\mathfrak{sl}(3,\mathbb{C}),\mathfrak{so}(3,\mathbb{C}))$ -modules (see Proposition 6.1).

Let \mathfrak{h}_0 be the split Cartan subalgebra consisting of diagonal matrices in \mathfrak{g}_0 . Take a positive system $\Sigma(\mathfrak{g},\mathfrak{h})^+$ of restricted roots as in Section 2. Fix a standard parabolic subalgebra \mathfrak{q}_0 of \mathfrak{g}_0 . Let G be the universal cover $SL(n,\mathbb{R})$ of $SL(n,\mathbb{R})$, K the inverse image of SO(n), Q = MAN the standard parabolic subgroup with Lie algebra \mathfrak{q}_0 and its Langlands decomposition, σ an irreducible representation of the component group of M, and ν a character of the Lie algebra \mathfrak{a} . In Section 4, we defined an intertwining differential operator

$$D = D(Q, \sigma_{\nu}) \colon C^{\infty}(G, \sigma_{\nu})^Q \to C^{\infty}(G, (\operatorname{proj} \circ \iota \circ \operatorname{sym}(F^a))^{\vee} \otimes \sigma_{\nu})^Q.$$

We write $(\text{Ker } D)_K, C^{\infty}(G, \sigma_{\nu})^Q_K$ for the space of K-finite vectors of $\text{Ker } D, C^{\infty}(G, \sigma_{\nu})^Q$, respectively.

Let us state the classification result for $\mathfrak{g}_0 = \mathfrak{sl}(n,\mathbb{R})$. Put a subset Z of N to be

$$Z := \begin{cases} \mathbb{N} \cap (|a| - n/2 - 2\mathbb{N}) & \text{if } a \in \mathbb{R}, \\ \emptyset & \text{otherwise} \end{cases}$$

Theorem 5.3. (1) Let $Q_1 = M_1 A_1 N_1$ be the standard parabolic subgroup of G whose Lie algebra is a real form of $q_{(1,n-1)}$. Write triv for the trivial character of M_1 , and sgn for the nontrivial character of M_1 . Then the infinite-dimensional subquotients of $C^{\infty}(G, \operatorname{triv}_{-\lambda(1,-a)})_K^{Q_1}, C^{\infty}(G, \operatorname{sgn}_{-\lambda(1,-a)})_K^{Q_1}$, whose \mathfrak{k} -type decompositions are

(5.1)
$$\bigoplus_{k \in 2\mathbb{N} \setminus Z} \mathcal{H}^k(\mathbb{R}^n), \bigoplus_{k \in (2\mathbb{N}+1) \setminus Z} \mathcal{H}^k(\mathbb{R}^n),$$

are a-minimal, respectively. Here $\mathcal{H}^k(\mathbb{R}^n)$ denotes the space of harmonic polynomials on \mathbb{R}^n of homogeneous degree k, which is irreducible as a \mathfrak{k} module.

H. TAMORI

(2) Assume n = 3 and $a \in \mathbb{Z}$. Let B = MAN be the standard Borel subgroup of G. Write σ^0 for the irreducible two-dimensional representation of $M \cong$ $\{\pm 1, \pm \mathbf{i}, \pm \mathbf{j}, \pm \mathbf{k}\}$. Then Ker $D(B, \sigma^0_{-\lambda(2,-a)})_K$ is a-minimal and isomorphic to

$$\bigoplus_{k\geq 0} S^{2|a|+1+4k} \mathbb{C}$$

as \mathfrak{k} -modules, where $S^k \mathbb{C}^2$ denotes the k-th symmetric power of the twodimensional irreducible \mathfrak{k} -module \mathbb{C}^2 .

Moreover, any a-minimal (g, \mathfrak{k}) -module is isomorphic to one of the above.

Remark 5.4. (1) When n = 3 and a = 0, the representation Ker $D(B, \sigma^0_{-\lambda(2,0)})$ is infinitesimally equivalent to the one given by Torasso [21], and the realization as the kernel of an intertwining differential operator was given by Kubo and Ørsted [15]. The *K*-type formula in Theorem 5.3 (2) (more precisely, Proposition 6.1 in a different parametrization) was obtained by their recent work [16, Theorem 1.7 (5)].

For $a = \pm 1$, the *a*-minimal representation of $SL(3, \mathbb{R})$ which does not descend to the representation of $SL(3, \mathbb{R})$ appeared in [23, Example 12.4].

(2) If an a-minimal (g, ℓ)-module is unitarizable, then a ∈ √-1R by Remark 3.2
(3). Conversely, assume a ∈ √-1R. Theorem 5.3 says that an a-minimal (g, ℓ)-module is associated to a composition factor of unitary principal series or Torasso's representation, which is unitary. Therefore it is unitarizable.

Proof of Theorem 5.3 (1). By Theorem 2.8 (2), the finite-dimensional \mathfrak{g} -modules whose annihilators include J_a are the space of homogeneous polynomials $\mathcal{P}^{-a-n/2}(\mathbb{R}^n)$ of degree -a - n/2 if $-a - n/2 \in \mathbb{N}$ and the contragredient of $\mathcal{P}^{a-n/2}(\mathbb{R}^n)$ if $a - n/2 \in \mathbb{N}$, and do not exist otherwise. Here the \mathfrak{g} -action on $\mathcal{P}^{|a|-n/2}(\mathbb{R}^n)$ is induced by the symmetric power of the contragredient of the *n*-dimensional defining representation of \mathfrak{g} when $|a| - n/2 \in \mathbb{N}$. As \mathfrak{k} -modules, they are isomorphic to

(5.2)
$$\bigoplus_{k\in Z} \mathcal{H}^k(\mathbb{R}^n)$$

by the theory of harmonic polynomials.

By Lemma 2.10, $\iota \circ \operatorname{sym}(F^a) = \operatorname{sym}(F^{-a})$ annihilates the generalized Verma module $U(\mathfrak{g}) \otimes_{U(\mathfrak{q}_1)} \mathbb{C}_{\lambda(1,-a)}$. Hence we have Ker $D(Q_1, \sigma_{-\lambda(1,-a)}) = C^{\infty}(G, \sigma_{-\lambda(1,-a)})^{Q_1}$ for any $\sigma = \operatorname{triv}$, sgn and Proposition 4.3 (2) shows that any infinite-dimensional composition factor is *a*-minimal. By the Frobenius reciprocity and the branching law for the pair $(\mathfrak{so}(n), \mathfrak{so}(n-1))$ (see [11, Proposition 9.16] for example), the \mathfrak{k} -type decomposition of $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_K^{Q_1}$ is given by

(5.3)
$$\bigoplus_{k \in 2\mathbb{N} + \epsilon} \mathcal{H}^{k}(\mathbb{R}^{n}), \text{ where } \epsilon = \begin{cases} 0 & \text{if } \sigma = \text{triv}, \\ 1 & \text{if } \sigma = \text{sgn.} \end{cases}$$

There exists an irreducible infinite-dimensional subquotient V of $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_{K}^{Q_{1}}$ as Z does not contain $2\mathbb{N} + \epsilon$. Let ψ be the highest weight of \mathfrak{k} -module \mathfrak{p} as in Section 3. Since the highest weight $\mathcal{H}^{k}(\mathbb{R}^{n})$ is $k\psi/2$, we see $\mathfrak{g} \cdot \mathcal{H}^{k}(\mathbb{R}^{n}) \subset$ $\mathcal{H}^{k-2}(\mathbb{R}^{n}) + \mathcal{H}^{k}(\mathbb{R}^{n}) + \mathcal{H}^{k+2}(\mathbb{R}^{n})$ as subspaces of $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_{K}^{Q_{1}}$. Therefore the \mathfrak{k} -type decomposition of V is written as $\bigoplus_{k \in 2\mathbb{N}+k_{0}} \mathcal{H}^{k}(\mathbb{R}^{n})$ for some $k_{0} \in \mathbb{N}$, and the multiplicity of V is one by (5.3).

Assume that $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_{K}^{Q_{1}}$ has a composition factor V' other than V. The \mathfrak{k} -type decomposition of V shows that V' is finite-dimensional. By the above argument, $||a|| - n/2 \in \mathbb{N}$ and V' is isomorphic to $\mathcal{P}^{-a-n/2}(\mathbb{R}^{n})$ if $-a - n/2 \in \mathbb{N}$

and the contragredient of $\mathcal{P}^{a-n/2}(\mathbb{R}^n)$ if $a - n/2 \in \mathbb{N}$. By (5.2) and (5.3), the multiplicity of V' is one and there are no composition factors other than V, V'.

Conversely, if $||a|| - n/2 \in \mathbb{N}$ and $C^{\infty}(G, \sigma_{-\lambda(1, -a)})_{K}^{Q_{1}}$ contains a \mathfrak{k} -type $\mathcal{H}^{k}(\mathbb{R}^{n})$ for some $k \in \mathbb{Z}$, then there exists a finite-dimensional composition factor by Proposition 3.3 (3) and (5.2).

By the above argument, if $(2\mathbb{N} + \epsilon) \cap Z = \emptyset$, $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_{K}^{Q_{1}}$ is irreducible (hence *a*-minimal). If $(2\mathbb{N}+\epsilon)\cap Z \neq \emptyset$, the composition factors of $C^{\infty}(G, \sigma_{-\lambda(1,-a)})_{K}^{Q_{1}}$ consist of an *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module whose \mathfrak{k} -type decomposition is as in (5.1) and one of the above finite-dimensional \mathfrak{g} -modules, and their multiplicities are one, which proves (1).

6. Genuine minimal representations of $\widetilde{SL}(3,\mathbb{R})$

Let $\mathfrak{g}_0 = \mathfrak{sl}(3,\mathbb{R})$. We use the notation in Sections 4 and 5. In this section, we explicitly describe the space of \mathfrak{k} -finite vectors of the kernel of $D(B, \sigma^0_{-\lambda(2,-a)})$ and prove the rest of Theorem 5.3.

We first fix notations for a covering map from SU(2) onto SO(3). Let $\mathbb{H} = \mathbb{R}1 + \mathbb{R}\mathbf{i} + \mathbb{R}\mathbf{j} + \mathbb{R}\mathbf{k}$ be the quaternion algebra. We regard \mathbb{H} as a vector space over $\mathbb{C} = \mathbb{R}1 + \mathbb{R}\mathbf{i}$ by the right multiplication, take $\{1, \mathbf{j}\}$ as a basis, and identify \mathbb{H} with \mathbb{C}^2 . The standard Hermitian inner product on \mathbb{C}^2 induces an inner product on \mathbb{H} given by $(\mathbf{x}, \mathbf{y}) = (\mathbf{x}\overline{\mathbf{y}} + \mathbf{y}\overline{\mathbf{x}})/2$ for $\mathbf{x}, \mathbf{y} \in \mathbb{H}$, where $\overline{\mathbf{y}}$ denotes the conjugate of \mathbf{y} . Since the left multiplication by a unit quaternion is a \mathbb{C} -vector space homomorphism and preserves the norm on \mathbb{H} , the group of unit quaternions are identified with SU(2). We also identify the imaginary part of \mathbb{H} with \mathbb{R}^3 by $b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \leftrightarrow {}^t(b, c, d)$ for $b, c, d \in \mathbb{R}$. Since the imaginary part of \mathbb{H} is stable under the conjugation by a unit quaternion, we obtain a two-to-one homomorphism

$$K = \mathrm{SU}(2) \to \mathrm{SO}(3); \ \mathbf{y} \mapsto (\mathbf{x} \mapsto \mathbf{y} \mathbf{x} \overline{\mathbf{y}}).$$

The differentiation and the complexification give the isomorphism of Lie algebras (6.1)

$$\mathfrak{sl}(2,\mathbb{C}) \cong \mathfrak{so}(3,\mathbb{C}); \frac{1}{2} \begin{pmatrix} a\sqrt{-1} & -b+c\sqrt{-1} \\ b+c\sqrt{-1} & -a\sqrt{-1} \end{pmatrix} \leftrightarrow \begin{pmatrix} 0 & c & b \\ -c & 0 & -a \\ -b & a & 0 \end{pmatrix} \quad (a,b,c\in\mathbb{C}).$$

We next fix notation for the irreducible representations of K. Let $m \in \mathbb{N}$ and $\mathcal{P}_m[t]$ the space of polynomials of degree up to m with variable t. The linear action of K on $\mathcal{P}_m[t]$ defined by

(6.2)
$$g^{-1}q(t) := (\beta t + \overline{\alpha})^m q((\alpha t - \overline{\beta})/(\beta t + \overline{\alpha}))$$

for $g = \begin{pmatrix} \alpha & -\overline{\beta} \\ \beta & \overline{\alpha} \end{pmatrix} \in K, q \in \mathcal{P}_m[t]$ gives us an irreducible (m + 1)-dimensional representation. Then the differentiated action π_m of $\mathfrak{sl}(2,\mathbb{C})$ on $\mathcal{P}_m[t]$ is given by (6.3)

$$\pi_m\left(\begin{pmatrix}1&0\\0&-1\end{pmatrix}\right) = m - 2t\frac{d}{dt}, \ \pi_m\left(\begin{pmatrix}0&1\\0&0\end{pmatrix}\right) = -\frac{d}{dt}, \ \pi_m\left(\begin{pmatrix}0&0\\1&0\end{pmatrix}\right) = -mt + t^2\frac{d}{dt}$$

For $m \in \mathbb{N}$, we define a nondegenerate K-invariant pairing \langle , \rangle_m of the m-th symmetric power $S^m \mathbb{C}^2$ and $\mathcal{P}_m[t]$ by the linear extension of

$$\left\langle \begin{pmatrix} 1\\0 \end{pmatrix}^{i} \begin{pmatrix} 0\\1 \end{pmatrix}^{m-i}, t^{j} \right\rangle_{m} \coloneqq \begin{cases} i!(m-i)!/m! & \text{if } i=j, \\ 0 & \text{otherwise} \end{cases}$$

for $0 \leq i, j \leq m$.

We regard σ^0 as the restriction of the defining representation \mathbb{C}^2 of K to M. By the Frobenius reciprocity, the restriction map induces an isomorphism from $C^{\infty}(G, \sigma^0_{-\lambda(2,-a)})^B$ to $C^{\infty}(K, \sigma^0)^M$ as K-modules. Since $\mathcal{P}_m[t]$ is isomorphic to the dual of $S^m \mathbb{C}^2$ as K-modules, we have a linear injection (6.4)

$$\Phi_m \colon S^m \mathbb{C}^2 \otimes (\mathcal{P}_m[t] \otimes \sigma^0)^M \to C^\infty(K, \sigma^0)^M; \ \Phi_m\left(v \otimes \begin{pmatrix} q_1(t) \\ q_2(t) \end{pmatrix}\right)(k) := \begin{pmatrix} \langle v, kq_1(t) \rangle_m \\ \langle v, kq_2(t) \rangle_m \end{pmatrix}$$

for ${}^{t}(q_{1}(t), q_{2}(t)) \in (\mathcal{P}_{m}[t] \otimes \sigma^{0})^{M}, v \in S^{m}\mathbb{C}^{2}, k \in K$. Then the direct sum of Φ_{m} for $m \geq 0$ gives an isomorphism

$$C^{\infty}(G, \sigma^0_{-\lambda(2, -a)})^Q_K \cong \bigoplus_{m \ge 0} S^m \mathbb{C}^2 \otimes (\mathcal{P}_m[t] \otimes \sigma^0)^M.$$

For $a \in \mathbb{Z}$ and $k \in \mathbb{N}$, we put

$$m(a,k) := 2|a| + 1 + 4k$$

and define a polynomial $q(a, k : t^2) \in \mathcal{P}_{m(a,k)}[t]$ by

$$\begin{split} q(a,k;t^2) &:= {}_2F_1(-|a|-1/2-2k,-a/2-|a|/2-k:a/2-|a|/2+1/2-k;t^2) \\ &= \sum_{l=0}^{a/2+|a|/2+k} \frac{(-|a|-1/2-2k)_l(-a/2-|a|/2-k)_l}{l!(a/2-|a|/2+1/2-k)_l} t^{2l}. \end{split}$$

Here $_2F_1$ denotes the Gauss hypergeometric function and we use the Pochhammer symbol $(x)_l = x(x+1)\cdots(x+l-1)$.

The subspace $\operatorname{Ker} D(B, \sigma^0_{-\lambda(2,-a)})_K$ of $C^{\infty}(G, \sigma^0_{-\lambda(2,-a)})^B$ is described explicitly in the following

Proposition 6.1. When $a \notin \mathbb{Z}$, Ker $D(B, \sigma^0_{-\lambda(2,-a)})_K = 0$. When $a \in \mathbb{Z}$, Ker $D(B, \sigma^0_{-\lambda(2,-a)})_K$ equals

$$\begin{cases} \bigoplus_{k\geq 0} \Phi_{m(a,k)} \left(S^{m(a,k)} \mathbb{C}^2 \otimes \begin{pmatrix} -t^{m(a,k)} q(a,k;t^{-2}) \\ q(a,k;t^2) \end{pmatrix} \right) & \text{if a is even,} \\ \bigoplus_{k\geq 0} \Phi_{m(a,k)} \left(S^{m(a,k)} \mathbb{C}^2 \otimes \begin{pmatrix} q(a,k;t^2) \\ -t^{m(a,k)} q(a,k;t^{-2}) \end{pmatrix} \right) & \text{if a is odd,} \end{cases}$$

and is isomorphic to $\bigoplus_{k\geq 0}S^{m(a,k)}\mathbb{C}^2$ as $\mathfrak{k}\text{-modules}.$

We use the following lemma in the proof of Proposition 6.1. Let $\mathfrak n$ be the nilradical of $\mathfrak b.$

Lemma 6.2. Let $a \in \mathbb{C}, \lambda \in \mathfrak{h}^{\vee}$. Then $[\mathfrak{n}, \iota \circ \operatorname{sym}(F^a)]$ is included in $I(\mathfrak{b}, \lambda)$ (see (4.1) for the definition) if and only if $\lambda = \lambda(2, -a)$.

Proof. The lowest weight vector $\operatorname{sym}(\sum_{i=1}^{3} T_{3,i}T_{i,1} + a/3T_{3,1})$ in $\iota \circ \operatorname{sym}(F^{a}) = \operatorname{sym}(F^{-a})$ is $T_{3,2}T_{2,1} + (-\lambda_{2} + a/3 - 1/2)T_{3,1}$ modulo $I(\mathfrak{b}, \lambda)$. Since we have

$$\begin{split} [T_{1,2}, T_{3,2}T_{2,1} + (-\lambda_2 + a/3 - 1/2)T_{3,1}] &\equiv T_{3,2}(T_{1,1} - T_{2,2}) - (-\lambda_2 + a/3 - 1/2)T_{3,2} \\ &\equiv (\lambda_1 - a/3 + 1/2)T_{3,2} \bmod I(\mathfrak{b}, \lambda), \end{split}$$

$$\begin{split} [T_{2,3}, T_{3,2}T_{2,1} + (-\lambda_2 + a/3 - 1/2)T_{3,1}] &\equiv (T_{2,2} - T_{3,3})T_{2,1} + (-\lambda_2 + a/3 - 1/2)T_{2,1} \\ &\equiv (-\lambda_3 + a/3 + 1/2)T_{2,1} \bmod I(\mathfrak{b}, \lambda), \end{split}$$

the condition $[\mathfrak{n}, \iota \circ \operatorname{sym}(F^a)] \subset I(\mathfrak{b}, \lambda)$ is equivalent to $\lambda_1 = a/3 - 1/2$ and $\lambda_3 = a/3 + 1/2$, which says $\lambda = \lambda(2, -a)$.

Proof of Proposition 6.1. Since $I(\mathfrak{b}, \lambda(2, a))$ is closed under the adjoint action of \mathfrak{n} , Lemma 6.2 and its proof show that $\iota \circ \operatorname{sym}(F^a) + I(\mathfrak{b}, \lambda(2, -a))$ is spanned by $T_{3,2}T_{2,1} + (-\lambda(2, -a)_2 + a/3 - 1/2)T_{3,1}$, which equals

$$X := (T_{2,1} - T_{1,2})(T_{3,2} - T_{2,3}) + (a + 1/2)(T_{3,1} - T_{1,3}) \in U(\mathfrak{k})$$

modulo $I(\mathfrak{b}, \lambda(2, a))$.

Let ${}^{t}(q_{1}(t), q_{2}(t))$ be a nonzero element in $(\mathcal{P}_{m}[t] \otimes \sigma^{0})^{M}$. By (6.2), the condition ${}^{t}(q_{1}(t), q_{2}(t)) \in (\mathcal{P}_{m}[t] \otimes \sigma^{0})^{M}$ is equivalent to

$$\begin{pmatrix} i \\ & -i \end{pmatrix} \begin{pmatrix} q_1(t) \\ q_2(t) \end{pmatrix} = \begin{pmatrix} \mathbf{i}^{-1}q_1(t) \\ \mathbf{i}^{-1}q_2(t) \end{pmatrix} = \begin{pmatrix} (-i)^m q_1(-t) \\ (-i)^m q_2(-t) \end{pmatrix} + \\ \begin{pmatrix} & -1 \\ 1 \end{pmatrix} \begin{pmatrix} q_1(t) \\ q_2(t) \end{pmatrix} = \begin{pmatrix} \mathbf{j}^{-1}q_1(t) \\ \mathbf{j}^{-1}q_2(t) \end{pmatrix} = \begin{pmatrix} t^m q_1(-t^{-1}) \\ t^m q_2(-t^{-1}) \end{pmatrix}.$$

Hence our assumption on $q_1(t), q_2(t)$ is written as

- (1) m is odd,
- (2) $q_1(t)$ is odd when $m \equiv 1 \mod 4$, $q_1(t)$ is even when $m \equiv 3 \mod 4$, and (3) $q_2(t) = -t^m q_1(-t^{-1}) \neq 0$.
- Let $F \in C^{\infty}(G, \sigma_{-\lambda(2,-a)}^{0})^{B}$. By the decomposition G = KB, we see that $F \in \text{Ker } D$ if and only if DF = 0 on K. By the definitions of D (see Definition 4.1) and Φ_m (6.4), the irreducible K-submodule $\Phi_m(S^k(\mathbb{C}^2) \otimes {}^t(q_1(t), q_2(t)))$ is included in $\text{Ker } D(B, \sigma_{-\lambda(2,-a)}^0)$ if and only if $\pi_m(X)q_1(t) = 0$ and $\pi_m(X)q_2(t) = 0$.

Under the isomorphism (6.1), the vector 4X corresponds to

$$\begin{pmatrix} 0 & -\sqrt{-1} \\ -\sqrt{-1} & 0 \end{pmatrix} \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix} + (1+2a) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

By (6.3), the action $\pi_m(4X)$ on $\mathcal{P}_m[t]$ is given by the differential operator

$$(-mt+t^2\frac{d}{dt}-\frac{d}{dt})(m-2t\frac{d}{dt}) + (1+2a)(mt-t^2\frac{d}{dt}-\frac{d}{dt}).$$

Let $a(t) = \sum^k a_i t^l \in \mathcal{P}_{-}[t]$. Since

Let
$$q(l) = \sum_{l=0} a_l l^{l} \in \mathcal{P}_m[l]$$
. Since
 $\pi_m(4X)q(l) = \sum_l a_l(\{(m-2l)(-m+l) + (1+2a)(m-l)\}t^{l+1} + \{(m-2l)(-l) + (1+2a)(-l)\}t^{l-1})$
 $= \sum_l a_l\{(m-l)(-m+2l+1+2a)t^{l+1} - l(m-2l+1+2a)t^{l-1}\},$

the condition $\pi_m(X)q(t) = 0$ is equivalent to

$$a_{l}\frac{l}{2}\left(\frac{l}{2}+k_{0}-\frac{m}{2}\right) = a_{l-2}\left(\frac{l}{2}-\frac{m+2}{2}\right)\left(\frac{l}{2}-k_{0}-1\right)$$

for any l. Here we put $a_l = 0$ for any $l \neq 0, \ldots, k$ and

$$k_0 := (m - 1 - 2a)/4.$$

By the assumption that m is odd, there is a nonzero polynomial solution to $\pi_m(X)q(t) = 0$ if and only if $2k_0 \in \{0, 2, \ldots, m-1\}$ if and only if $m \in 2|a|+1+4\mathbb{N}$. In this case, the solution space is spanned by

$${}_{2}F_{1}(-m/2,-k_{0};k_{0}+1-m/2;t^{2}) = \sum_{k=0}^{k_{0}} \frac{(-m/2)_{k}(-k_{0})_{k}}{(1)_{k}(k_{0}+1-m/2)_{k}} t^{2k},$$

$$t^{m}{}_{2}F_{1}(-m/2,-k_{0};k_{0}+1-m/2;t^{-2}) = \sum_{k=0}^{k_{0}} \frac{(-m/2)_{k}(-k_{0})_{k}}{(1)_{k}(k_{0}+1-m/2)_{k}} t^{m-2k}.$$

By the above arguments, $q_1, q_2 \in \mathcal{P}_m[t]$ satisfies the assumptions (1), (2), (3) and $\pi_m(X)q_1(t) = \pi_m(X)q_2(X) = 0$ if and only if $a \in \mathbb{Z}$, m = m(a,k) for some $k \in \mathbb{N}$ and

$$(q_1(t), q_2(t)) \in \begin{cases} \mathbb{C}(-t^{m(a,k)}q(a,k:t^{-2}), q(a,k:t^2)) & \text{when } a \text{ is even,} \\ \mathbb{C}(q(a,k:t^2), -t^{m(a,k)}q(a,k:t^{-2})) & \text{when } a \text{ is odd,} \end{cases}$$

which is the desired conclusion.

Let us finish the proof of Theorem 5.3. Let $\mathfrak{g}_0 = \mathfrak{sl}(n, \mathbb{R})$.

Proof of the rest of Theorem 5.3. The assertion (2) follows from Proposition 6.1.

Let us prove the exhaustion of *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -modules. Let us take a positive system on $\Phi(\mathfrak{g}, \mathfrak{h}^c)$ such that the highest root ψ is noncompact imaginary as in Section 3. By $\mathfrak{p} \cong S^2(\mathbb{C}^n)$ as \mathfrak{k} -modules, we see $\rho(\mathfrak{k}^1, \mathfrak{t}) = n\psi/2$. Therefore the description (3.2) and Proposition 3.3 (2) show that the highest weight λ of a \mathfrak{k} -type of an *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module belongs to $\mathbb{N}\psi/2$ if $n \geq 4$ and to $\mathbb{N}\psi/4$ if n = 3. When λ belongs to $\mathbb{N}\psi/2$, the *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module has a common \mathfrak{k} -type with the one appearing in Theorem 5.3 (1) by (5.3). Therefore they are isomorphic by Proposition 3.3 (3).

Hence what is left is to prove that when $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{C})$, any *a*-minimal $(\mathfrak{g}, \mathfrak{k})$ -module V which does not exponentiate to a representation of SO(3) (called genuine) is isomorphic to the one in (2). By Casselman's subrepresentation theorem, we can embed V into $C^{\infty}(G, \sigma_{-\lambda})_K^B$ for some $\sigma \in \widehat{M}, \lambda \in \mathfrak{a}^{\vee}$. Proposition 4.3 (1) shows that V is contained in Ker $D(B, \sigma_{-\lambda})$. Since V is genuine, σ is the two-dimensional irreducible representation σ^0 of M. Moreover, since the identity component of any standard parabolic subgroup strictly larger than B contains the center of G, Lemma 4.5 shows that $[\mathfrak{n}, \iota \circ \operatorname{sym}(F^a)]$ is included in $I(\mathfrak{b}, \lambda)$ so that Ker $D(B, \sigma_{-\lambda})$ does not descend to the representation of SL(3, \mathbb{R}). By Lemma 6.2, we must have $\lambda = \lambda(2, -a)$. Therefore V is isomorphic to the minimal *a*-module in Proposition 6.1, and the proof of Theorem 5.3 is complete.

Acknowledgements

The author is deeply grateful to Prof. Toshiyuki Kobayashi for helpful comments and warm encouragement. The author expresses his sincere thanks to Prof. David Vogan for insightful comments and letting him know the work by Mœglin [17] and the alternative proof of Corollary 3.5. This work was supported by JSPS KAKENHI Grant Number 17J01075, 20J00024 and the Program for Leading Graduate Schools, MEXT, Japan.

FACULTY OF SCIENCE, HOKKAIDO UNIVERSITY, KITA 10, NISHI 8, KITA-KU, SAPPORO, HOKKAIDO, 060-0810, JAPAN

 $E\text{-}mail: \verb+tamori@math.sci.hokudai.ac.jp+$

References

- B. Binegar and R. Zierau. Unitarization of a singular representation of SO(p,q). Comm. Math. Phys., 138:245–258, 1991.
- [2] A. Braverman and A. Joseph. The minimal realization from deformation theory. J. Algebra, 205:13–36, 1998.
- [3] T. Enright, R. Howe, and N. Wallach. A classification of unitary highest weight modules. In *Representation theory of reductive groups*, volume 40 of *Progr. Math.*, pages 97–143. Birkhäuser Boston, 1983.
- [4] W. T. Gan and G. Savin. Uniqueness of Joseph ideal. Math. Res. Lett., 11: 589–597, 2004.

- [5] W. T. Gan and G. Savin. On minimal representations definitions and properties. *Represent. Theory*, 9:46–93, 2005.
- [6] D. Garfinkle. A new construction of the Joseph ideal. Phd. thesis, M.I.T., 1982.
- [7] R. Howe and E. C. Tan. Nonabelian harmonic analysis. Universitext. Springer-Verlag, New York, 1992. ISBN 0-387-97768-6.
- [8] H. P. Jakobsen. Hermitian symmetric spaces and their unitary highest weight modules. J. Funct. Anal., 52(3):385–412, 1983.
- [9] J. C. Jantzen. Einhüllende Algebren halbeinfacher Lie-Algebren, volume 3 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1983. ISBN 3-540-12178-1.
- [10] A. Joseph. The minimal orbit in a simple Lie algebra and its associated maximal ideal. Ann. Sci. École. Norm. Sup. (4), 9(1):1–29, 1976.
- [11] A. Knapp. *Lie Groups Beyond an Introduction*, volume 140 of *Progr. Math.* Birkhäuser, second edition, 2002.
- [12] T. Kobayashi and B. Ørsted. Analysis on the minimal representation of O(p,q), I. Realization via conformal geometry. *Adv. Math.*, 180:486–512, 2003.
- [13] T. Kobayashi and B. Ørsted. Analysis on the minimal representation of O(p, q), III. Ultrahyperbolic equations on $\mathbb{R}^{p-1,q-1}$. Adv. Math., 180:551–595, 2003.
- [14] T. Kobayashi and M. Pevzner. Differential symmetry breaking operators: I. General theory and F-method. Selecta Math. (N. S.), 22(2):801–845, 2016.
- [15] T. Kubo and B. Ørsted. On the space of K-finite solutions to intertwining differential operators. *Represent. Theory*, 23:213–248, 2019.
- [16] T. Kubo and B. Ørsted. Classification of k-type formulas for the Heisenberg ultrahyperbolic operator \Box_s for $\widetilde{SL}(3,\mathbb{R})$ and tridiagonal determinants for local Heun functions. arXiv:2101.06810, 2021.
- [17] C. Mœglin. Idéaux complètement premiers de l'algèbre enveloppante de $gl_n(\mathbb{C})$. J. Algebra, 106(2):287–366, 1987.
- [18] T. Okuda. Smallest complex nilpotent orbits with real points. J. Lie Theory, 25(2):507–533, 2015.
- [19] J. Rawnsley and S. Sternberg. On representations associated to the minimal nilpotent coadjoint orbit of SL(3, ℝ). Amer. J. Math., 104(6):1153–1180, 1982.
- [20] H. Tamori. Classification of minimal representations of real simple Lie groups. Math. Z., 292(1-2):387–402, 2019.
- [21] P. Torasso. Quantification géométrique, opérateurs d'entrelacement et représentations unitaires de $\widetilde{SL}_3(\mathbb{R})$. Acta Math., 150(3-4):153-242, 1983.
- [22] D. Vogan. Singular unitary representations. In Noncommutative harmonic analysis and Lie groups, volume 880 of Lecture Notes in Math., pages 506–535. Springer, 1981.
- [23] D. Vogan. Associated varieties and unipotent representations. In Harmonic analysis on reductive groups (Brunswick, ME, 1989), volume 101 of Progr. Math., pages 315–388. Birkhäuser Boston, Boston, MA, 1991.