

| Title | Assessing insect herbivory on broadleaf canopy trees at 19 natural forest sites across Japan |
|------------------|---|
| Author(s) | Takafumi, Hino; Kanno, Yuri; Abe, Shin; Abe, Tetsuto; Enoki, Tsutomu; Hirao, Toshihide; Hiura, Tsutom; Hoshizaki, Kazuhiko; Ida, Hideyuki; Ishida, Ken; Maki, Masayuki; Masaki, Takashi; Naoe, Shoji; Noguchi, Mahoko; Otani, Tatsuya; Sato, Takanori; Sakimoto, Michinori; Sakio, Hitoshi; Takagi, Masahiro; Takashima, Atsushi; Tokuchi, Naoko; Utsumi, Shunsuke; Hidaka, Amane; Nakamura, Masahiro |
| Citation | Ecological Research, 36(3), 562-572 https://doi.org/10.1111/1440-1703.12215 |
| Issue Date | 2021-03-25 |
| Doc URL | http://hdl.handle.net/2115/88734 |
| Rights | This is the peer reviewed version of the following article: Takafumi, H, Kanno, Y, Abe, S, et al. Assessing insect herbivory on broadleaf canopy trees at 19 natural forest sites across Japan. Ecological Research. 2021; 36: 562–572., which has been published in final form at https://doi.org/10.1111/1440-1703.12215. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley 's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited. |
| Туре | article (author version) |
| File Information | Takafumi et al. 2021 EcoRes.pdf |



1 **Title**

2 Assessing insect herbivory on broadleaf canopy trees at 19 natural forest sites across

- 3 Japan
- 4

5 Authors

- 6 Hino Takafumi^{1,a}, Yuri Kanno², Shin Abe³, Tetsuto Abe⁴, Tsutomu Enoki⁵, Toshihide
- 7 Hirao⁶, Tsutom Hiura^{7,b}, Kazuhiko Hoshizaki⁸, Hideyuki Ida^{9,c}, Ken Ishida^{10,d},
- 8 Masayuki Maki¹¹, Takashi Masaki³, Shoji Naoe^{3,e}, Mahoko Noguchi¹², Tatsuya Otani¹³,
- 9 Takanori Sato¹⁴, Michinori Sakimoto¹⁵, Hitoshi Sakio^{16,f}, Masahiro Takagi¹⁷, Atsushi
- 10 Takashima¹⁸, Naoko Tokuchi¹⁵, Shunsuke Utsumi^{19,g}, Amane Hidaka^{1,h}, Masahiro
- 11 Nakamura^{2*}
- 12

13 Affiliations

- ¹⁴ ¹ Network Center of the Forest and Grassland Survey of the Monitoring Sites 1000
- 15 Project, Japan Wildlife Research Center, c/o Tomakomai Experimental Forest, Field
- 16 Science Center for Northern Biosphere, Hokkaido University, Tomakomai, Japan
- ¹⁷ ² Wakayama Experimental Forest, Field Science Center for Northern Biosphere,
- 18 Hokkaido University, Wakayama, Japan
- 19³ Forestry and Forest Products Research Institute, Tsukuba, Japan
- ⁴ Kyushu Research Center, Forestry and Forest Products Research Institute, Kumamoto,
- 21 Japan
- ⁵ Faculty of Agriculture, Kyushu University, Fukuoka, Japan
- ⁶ The University of Tokyo Chichibu Forest, Graduate School of Agricultural and Life
- 24 Sciences, The University of Tokyo, Chichibu, Japan
- ⁷ Tomakomai Experimental Forest, Field Science Center for Northern Biosphere,
- 26 Hokkaido University, Tomakomai, Japan
- ⁸ Faculty of Bioresource Sciences, Akita Prefectural University, Akita, Japan
- ⁹ Institute of Nature Education in Shiga Heights, Faculty of Education, Shinshu
- 29 University, Yamanouchi, Japan

| 30 | ¹⁰ Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, |
|----|--|
| 31 | Japan |
| 32 | ¹¹ Botanical Gardens, Tohoku University, Sendai, Japan |
| 33 | ¹² Tohoku Research Center, Forestry and Forest Products Research Institute, Morioka, |
| 34 | Japan |
| 35 | ¹³ Shikoku Research Center, Forestry and Forest Products Research Institute, Kochi, |
| 36 | Japan |
| 37 | ¹⁴ Ecohydrology Research Institute, The University of Tokyo Forests, Graduate School |
| 38 | of Agricultural and Life Sciences, The University of Tokyo, Seto, Japan |
| 39 | ¹⁵ Field Science Education and Research Center, Kyoto University, Kyoto, Japan |
| 40 | ¹⁶ Field Center for Sustainable Agriculture and Forestry, Faculty of Agriculture, Niigata |
| 41 | University, Sado, Japan |
| 42 | ¹⁷ Faculty of Agriculture, University of Miyazaki, Miyazaki, Japan |
| 43 | ¹⁸ Yona Field, Subtropical Field Science Center, Faculty of Agriculture, University of |
| 44 | the Ryukyus, Kunigami, Japan |
| 45 | ¹⁹ Uryu Experimental Forest, Field Science Center for Northern Biosphere, Hokkaido |
| 46 | University, Horokanai, Japan |
| 47 | |
| 48 | * corresponding author |
| 49 | |
| 50 | Present address: |
| 51 | ^a The National Ainu Museum, Shiraoi, Japan |
| 52 | ^b Department of Ecosystem Studies, Graduate School of Agricultural and Life Sciences, |
| 53 | The University of Tokyo, Tokyo, Japan |
| 54 | ^c Faculty of Education, Shinshu University, Nagano, Japan |
| 55 | ^d Amami Ecosystem Study Group |
| 56 | ^e Tohoku Research Center, Forestry and Forest Products Research Institute, Morioka, |
| 57 | Japan |
| 58 | ^f Sado Island Center for Ecological Sustainability, Niigata University, Sado, Japan |
| 59 | ^g Field Science Center for Northern Biosphere, Hokkaido University, Sapporo, Japan |

60 ^h Graduate School of Agriculture, Kyoto University, Kyoto, Japan

61

62 Abstract

63 We present the largest freely available herbivory dataset for Japan representing data 64 collected from a network of 19 natural forest sites across the country. Sampled network 65 sites were part of the Monitoring Sites 1000 Project organized by the Ministry of the Environment. Sites were located across a range of climate zones, from subarctic to 66 subtropical, and broadleaf trees (both evergreen and deciduous) were targeted at each 67 68 site. Litterfall traps were used to assess leaf damage caused by leaf-chewing insects in 69 2014 and 2015. Using a standardized protocol, we assessed herbivory on 117,918 leaves 70 of 39 dominant tree species. Preliminary analyses suggest that insect herbivory 71 increases with increasing latitude for deciduous broadleaf species. In particular, oak 72 (Quercus crispula) and beech (Fagus crenata) were subject to increased insect 73 herbivory with increasing latitude. In contrast, insect herbivory decreased with 74 increasing latitude in evergreen broadleaf species. The latitudinal gradient of herbivory 75 differed according to leaf type (i.e., evergreen or deciduous). This dataset offers 76 excellent opportunities for meta-analysis and comparative studies of herbivory among 77 various forest types.

78

79 **Keywords**

80 evergreen species, deciduous species, insect-plant interactions, the Monitoring sites

- 81 1000 Project, latitudinal gradient
- 82

83 Metadata

84 **1. Introduction**

85 Herbivory, the act of eating plants, plays a fundamental role in carbon and nutrient 86 cycling (Metcalfe et al., 2013) and may affect diversity and structure in forest 87 communities by altering biomass (Cyr & Pace, 1993) and tree fitness (Coley & Barone, 88 1996). Exploring changes in herbivory that may reflect global climate change has 89 become increasingly popular in recent decades (Kozlov & Zvereva, 2015). However, 90 debate remains as to the mechanisms by which biotic and abiotic factors regulate 91 herbivory (e.g., Coley, Bryant, & Chapin, 1985; Endara & Coley, 2011; Moles, Bonser, 92 Poore, Wallis, & Foley, 2011). 93 It is believed that biotic interactions become stronger as latitude decreases 94 (Dobzhansky, 1950; Janzen 1970; Schemske et al. 2009). For insect-plant interactions, 95 ecologists have long examined how insect herbivory intensity varies with latitude (e.g., MacArthur, 1972; Pennings & Siliman, 2005; Schemske, Mittelbach, Cornell, Sobel, & 96 97 Roy, 2009; Hiura & Nakamura, 2013). In two seminal review papers published in the 98 1990s, Coley and colleagues found that tropical forests experienced significantly higher 99 herbivory than did temperate forests (Coley & Aide, 1991; Coley & Barone, 1996). These 100 studies served as the foundation for the latitudinal herbivory-defense hypothesis 101 (LHDH), which predicts that herbivory will be higher in warmer and more stable 102 climates (i.e., at low latitude or low elevation), which then leads to selective pressure 103 for the evolution of more effective plant defenses (Dobzhansky, 1950; MacArthur, 104 1972; Coley & Aide, 1991). Latitudinal gradients in insect herbivore abundance and 105 species richness are likely responsible for stronger herbivore pressure on plants at low 106 latitudes (Salazar & Marquis, 2012). It is widely accepted that forests in warmer 107 climates suffer greater herbivory than those in cooler climates (Kozlov, 2008; Kozlov, 108 van Nieukerken, Zverev, & Zvereva, 2013; Lim, Fine, & Mittelbach, 2015; Wang et al., 109 2016; Zhang, Zhang, & Ma, 2016; Moreira et al., 2018). Additionally, variation in the 110 production cost of defense compounds as predicted by optimal defense theory is an 111 important determinant of geographic patterns in plant defense (Kooyers et al. 2017). 112 However, some studies have shown that no latitudinal gradient in herbivory exists (e.g.,

Andrew & Hughes, 2005), that the trend is counter to what would generally be expected by the LHDH (e.g., Moles et al., 2011), or that the trend is nonlinear (e.g., Kozlov, Lanta, Zverev, & Zvereva, 2015). Therefore, despite widespread interest in this pattern, the empirical evidence comprises controversial results regarding latitudinal gradients in herbivory. The wide range of latitudes (20–46 degrees) and mountainous terrain in Japan, which include a wide range of subarctic and subtropical forest types themselves, gives rise to suitable conditions for addressing this question.

- 120 This study reports insect herbivory data collected across a network of 19 121 natural forest sites; this is the largest published herbivory dataset available for Japan 122 with sites spanning a wide latitudinal gradient (26.74–44.37°N). This study focuses on 123 the latitudinal gradient in insect herbivory in temperate regions of East Asia. The forest 124 site network is part of the Forest and Grassland Survey of the Monitoring Sites 1000 125 Project established by the Ministry of the Environment (Ishihara, Toyota, & Nakamura, 126 2007; Ishihara et al., 2010). Some Monitoring Sites 1000 locations are also part of the 127 Japan Long-Term Ecological Research Network (JaLTER), which further contributes to 128 the International Long-Term Ecological Research Network (ILTER). These herbivory 129 data, which were collected in 2014 and 2015, offer an opportunity for meta-analysis and 130 comparative studies among forest types and for testing community ecology and 131 ecosystem function hypotheses at broad spatial scales (e.g., testing for a latitudinal 132 gradient in insect-plant interactions and elucidating the relative importance of insect 133 herbivory in forest ecosystem function). 134
- 135 2. IDENTIFER
- 136 xxxx(Ecologial Research Data Paper Archives ID, describe after accepted)
- 137
- 138 3. CONTRIBUTER
- 139 A. Dataset Owner

Masahiro Nakamura, Wakayama Experimental Forest, Hokkaido University, Hokkaido,
Japan

142

- 143 B. Dataset Creator
- 144 Hino Takafumi, Network Center of the Forest and Grassland Survey of the Monitoring
- 145 Sites 1000 Project, Japan Wildlife Research Center
- 146 Masahiro Nakamura, Wakayama Experimental Forest, Field Science Center for
- 147 Northern Biosphere, Hokkaido University, Wakayama, Japan
- 148
- 149 4. GEOGRAPHIC COVERAGE
- 150 A. Geographic Description
- 151 Japan
- 152 B. Geographical Position
- 153 Longitude: 128° 23´ E 142° 28´ E
- 154 latitude: 26° 74′ N 44° 37′ N
- 155
- 156 5. TEMPORAL COVERAGE
- 157 A. Begin:
- 158 2014
- 159 B. End:
- 160 2015
- 161

162 6. TAXONOMIC COVERAGE

163 The data include 20 families, 27 genus and 39 species of broadleaf trees (see SpList.csv

164 for details). Castanopsis spp. in Tano site includes Castanopsis sieboldii and

165 *Castanopsis cuspidate*.

166

167 7. METHODS

168 A. Study sites and data acquisition

169 Herbivory data were obtained at 19 natural forest sites as part of the Monitoring Sites

- 170 1000 Project, with one to five permanent plots located within each site. These sites
- 171 covered the major climate zones and biogeographic regions in Japan (Ministry of the
- 172 Environment 2001, http://www.env.go.jp/press/press.php?serial=2908, last accessed on

173 October 6, 2020) (Fig. 1, Table 1; see also Ishihara et al., 2011) as well as the four 174 major forest types. Plots were classified to four forest types based on the dominant tree 175 species; evergreen conifer forest (EC), broadleaf and conifer mixed forest (BC), 176 deciduous broadleaf forest (DB), and evergreen broadleaf forest (EB) (Ishihara et al., 177 2011). The majority of the surveyed forest were old-growth or older secondary forests, 178 but some were secondary forests aged <100 years. We classified forests stands to three 179 age categories: old growth (OG), old secondary (OS), and secondary (S) (Ishihara et al., 180 2011; see Appendix 1 for details).

181 The mean annual temperature and precipitation and mean maximum snow 182 depth during 1981–2010 were extracted from the Mesh Climate Data 2010 database 183 distributed by the Japan Meteorological Agency (2012). The database provides climate 184 variables estimated at 1-km spatial resolution. Mean annual temperature was corrected 185 to account for altitudinal difference between a given plot and the 1-km cell mean using 186 a lapse rate of 0.55°C per 100 m. Database estimates for snow depth may contain 187 inaccuracies (i.e., over- or underestimation) given that snow depth is highly spatially 188 heterogeneous. Therefore, we provide snow depths reported in other publications and 189 personal observations in Appendix 1 and Table S1.

Additional ecological data collected at the 19 study sites were available from publications, including seasonal patterns and inter-annual dynamics of litterfall (Suzuki et al., 2012), forest stand structure, composition, and dynamics (Ishihara et al., 2010), as well as ground-dwelling beetle community and understory variables (Niwa et al., 2016). At the Shiiba site (SI), only tree census data (Ishihara et al., 2010) were available.

Additional ecological information including understory vegetation, disturbance history,
and soil parameters collected from study sites are provided in Appendix 1 and partially
summarized in Table S1.

198

199 B. Sampling design, field methodology, and preliminary analyses

200 B-1. Sampling design

201 One permanent plot at each of the 19 sites was selected for litter collection. Plots ranged

in size from 0.64 to 1.2 ha and were usually approximately 1 ha in size (see Appendix 1,

SiteList.csv). Plots were placed or their shapes were adjusted to avoid forest roads and
urban edges. Plots were typically divided into 10 x 10-m grid cells. Typically, 25
conical litter traps, each with a circular collection area of 0.5 m², were installed within
each plot (see Suzuki et al., 2012 for details of the litter traps). Traps were placed
approximately 1 m above the ground at grid cell corners and were spaced 20 m apart.
Litterfall that collected within the traps was collected monthly. Traps were either
removed or placed on the ground during winter at sites that received snow.

211 B-2. Field methodology

212 Litterfall from these traps was then used for a visual assessment of leaf damage by 213 leaf-chewing insects. Herbivory has been assessed visually in other publications (e.g., 214 review by Kozlov et al., 2015). Although few studies have used litter traps for the visual 215 assessment of herbivory (but see Hiura & Nakamura, 2013), litter traps are useful for 216 collecting litterfall, especially for the assessment of multiple tree species over long 217 periods. One technician visually assessed leaf damage blindly with respect to site. The 218 protocol was standardized across all sites, enabling comparisons among forest types. At 219 broadleaf forest sites, the 5 traps with the largest amount of litter were selected from all 220 25 traps. At BC sites, 5 traps located near broadleaf trees were selected. During the 221 4-month period when litterfall was highest (Table 2), we visually categorized the extent 222 of herbivory on fallen leaves to six classes: no damage; 1-10% of leaf area lost; 11-223 25% loss; 26–50% loss; 51–75% loss; and >76% loss (Nakamura, Asanuma, & Hiura, 224 2010; Nakamura, Nakaji, Muller, & Hiura, 2014). This herbivory assessment was 225 applied to all broadleaf tree species that were considered dominant at each site (up to 226 50% dominance, Table 3). In addition, oak (Quercus crispula, 8 sites in 2014, 8 sites in 227 2015) and beech (Fagus crenata, 8 sites in 2014, 9 sites in 2015) leaves were also 228 assessed regardless of their dominance at sites.

229

230 B-3. Preliminary analyses

231 We preliminarily analyzed the latitudinal gradient in insect herbivory as observed in

deciduous species, evergreen species, *Q. crispula*, and *F. crenata* in 2014 and 2015

233 using linear mixed models (LMM) in the R library lme4 package. In the models, the 234 median herbivory rate of each leaf (i.e., 0%, 5.5%, 18%, 38%, 63%, and 88%, 235 respectively) was treated as a response variable; latitude, year sampled, and their 236 interaction were treated as fixed effect variables. The year sampled was treated as a 237 factor. Plot ID was treated as a random effect variable. The likelihood-ratio chi-square 238 test was used to determine whether the herbivory rate was significantly related to 239 latitude. The preliminary analyses indicated that insect herbivory on deciduous species 240 significantly increased toward higher latitudes ($\chi^2 = 35.03$, P < 0.001; Fig. 2a, Table 4), whereas that on evergreen species significantly decreased ($\gamma^2 = 4.82$, P < 0.05; Fig. 2b, 241 242 Table 4). These results were congruent with the latitudinal pattern in defense strategy in 243 deciduous and evergreen species reported by Saihanna et al. (2018), who showed that 244 deciduous species used physical defenses at lower latitudes, whereas evergreen species 245 exhibited the opposite latitudinal defense patterns. In particular, insect herbivory on Q. *crispula* significantly increased with increasing latitude ($\chi^2 = 8.99$, P < 0.05; Fig. 3a, 246 247 Table 4), and herbivory on F. crenata tended to increase with latitude, although the 248 trend was not significant (Fig. 3b, Table 4). These results were congruent with the 249 latitudinal pattern in beech herbivory reported by Hiura & Nakamura (2013). 250 Our findings suggested that the latitudinal gradient of insect herbivory 251 differed depending on leaf type (i.e., evergreen or deciduous). Leaf type is an important 252 predictor of resource use; evergreen species typically show a more conservative

resource-use strategy and slower growth rates (Givinish, 2002), whereas deciduous

254 species show an exploitative resource-use strategy and higher growth rates (Reich,

Ellsworth, & Walters, 1998). Our preliminary results suggest that the resource-use
strategy may be an important determinant of the directionality of latitudinal gradients in
insect herbivory.

258

259 C. Data verification procedures

260 We note that our data collection at two sites, Kamigamo (2014) and Wakayama (2015),

261 was limited to a single year. The selected litter traps at some sites did not capture fallen

262 leaves of some target tree species. Specifically, we did not capture leaves from Acer

- 263 ukurunduense at Otanomosutaira in 2015, Castanea crenata at Ogawa in 2015, Illicium
- 264 anisatum at Shiiba in 2014, Quercus acuta at Aya in 2014, and Ilex macropoda at
- 265 Kamigamo in 2015. Thus, herbivory was not assessed for those species in those years.
- 266
- 267 8. DATA STATUS
- A. Latest update
- 269 January 15 2020
- 270
- B. Metadata status
- 272 Metadata are complete for this period and stored with the data.
- 273
- 274 9. ACCESSIBILITY
- A. License and usage right
- 276 This dataset is provided under a Creative Commons Attribution 4.0 International license
- 277 (CC-BY 4.0) (https://creativecommons.org/licenses/by/4.0/).
- 278
- B. Contact
- 280 Data Set Contact
- 281 Masahiro Nakamura
- 282 Wakayama Experimental Forest, Field Science Center for Northern Biosphere,
- 283 Hokkaido University
- 284 Hirai, Kozagawa, Wakayama 649-4563, Japan
- 285
- 286 10. DATA STRUCTURE
- A. Data tables

| Data file name | Description | | | |
|----------------|---|--|--|--|
| SiteList.csv | Site-wise information table. Sites are listed by latitude | | | |
| | from north to south. | | | |
| SpList.csv | Species-wise information table. | | | |

| Herbivory.csv | Herbivory data of each species in each site in 2014 and |
|---------------|--|
| | 2015. The herbivory of each leaf was classified into the |
| | six herbivory class, and a total number of the leaves of |
| | each the class is calculated (see method 7. B). |

288

- B. Format type
- 290 The data files are comma delimited (csv) in UTF-8 encoding.
- 291
- 292 C. Header information

Headers corresponding to variables names (see 10. D.) are included as the first row inthe data files.

- 295
- 296 D. Variable definitions

The variables are listed in the order they appear in each data file. Variable names are
headers included as the first row in the data files. All variables of SiteList.csv are
compliant with the data papers (Ishihara et al., 2010; Suzuki et al., 2012; Niwa et al.,
2016), although variables of Temp30yr, Rain30yr, and Snow30yr (1971-2000) are
updated to 1981-2010 due to updating the database Mesh Climate Data by the Japan
Meteorological Agency.

303

| Data file name | Variable name | Variable definition | | | |
|----------------|---------------|--|--|--|--|
| SiteList.csv | Code | Plot code. The third and fourth digits are | | | |
| | | unique to each site, and the fifth and sixth | | | |
| | | digits are plot numbers. | | | |
| | PlotID | Alphanumeric plot code. | | | |
| | | An alphanumeric code as [abbreviated site | | | |
| | | name]-[forest type code][plot number]. For | | | |
| | | example, TM-DB1 represents the first plot | | | |
| | | of deciduous broadleaf forest (DB) in the | | | |

| | site, Tomakomai (TM). |
|-----------|---|
| SiteName | Site name. |
| Туре | Alphabetical code for forest type: |
| | BC = Broadleaf conifer mixed forest. |
| | DB = Deciduous broadleaf forest. |
| | EB = Evergreen broadleaf forest. |
| | EC = Evergreen coniferous forest. |
| Status | Forest age classifications (See 7. A and |
| | Appendix 1.pdf). |
| | S: Secondary. |
| | OS: Old secondary. |
| | OG: Old growth. |
| Latitude | WGS84 latitudinal coordinates in decimal |
| | degrees (°) to 2 decimal places. |
| Longitude | WGS84 longitudinal coordinates in |
| | decimal degrees (°) to 2 decimal places. |
| Altitude | Elevation above mean sea level (m). |
| | The precision is 5 m. |
| Area | Plot area (ha). |
| | The precision is 0.01 ha. |
| Shape | Shape of plot. |
| Temp30yr | Mean of mean annual temperature during |
| | the period from 1981 to 2010 ($^{\circ}$ C). |
| | The precision is 0.1 $^{\circ}$ C. |
| Rain30yr | Mean of annual precipitation during the |
| | period from 1981 to 2010 (mm). |
| | The precision is 0.1 mm. |
| Snow30yr | Mean of annual maximum snow depth |
| | during the period from 1981 to 2010 (cm). |

| | | The precision is 1 cm. | | | | |
|---------------|---------------------|--|--|--|--|--|
| SpList.csv | SpJapan | Japanese common name of species, which | | | | |
| | | was romanized following the Kunrei-siki | | | | |
| | | romanization system (ISO3602). | | | | |
| | Family | Botanical family to which the species. | | | | |
| | Genus | Genus name. | | | | |
| | SpecificName | Specific name. | | | | |
| | Subspecies | "subsp." represents subspecies. | | | | |
| | SubspeciesName | Subspecies name. | | | | |
| | NameAuth | The scientific name of species with | | | | |
| | | authority name. Under-bars are used in | | | | |
| | | place of spaces. | | | | |
| | Synonym | Numeric code for a synonym in Japanese | | | | |
| | | common name. | | | | |
| | | 1 = accepted name. | | | | |
| | | 0 = synonym. | | | | |
| | Deciduous_Evergreen | Functional type of the species. | | | | |
| | | Deciduous_broadleaf or | | | | |
| | | Evergreen_broadleaf. | | | | |
| Herbivory.csv | SiteName | Site name. | | | | |
| | Year | Leaves sampled year. The leaf collection | | | | |
| | | period at Aichi-akazu, Kamigamo, and | | | | |
| | | Shiiba were two calendar years because | | | | |
| | | the peak month of litter fall occurred | | | | |
| | | during these calendar years (See Table 2). | | | | |
| | Genus | Genus name. | | | | |
| | SpecificName | Specific name. | | | | |
| | Subspecies | "subsp." represents subspecies. | | | | |
| | SubspeciesName | Subspecies name. | | | | |

| herbivory_0 | A total number of leaves as classified |
|-------------|---|
| | herbivory class 0 (no damage). |
| herbivory_1 | A total number of leaves as classified |
| | herbivory class 1 (1-10% loss). |
| herbivory_2 | A total number of leaves as classified |
| | herbivory class 2 (11-25% loss). |
| herbivory_3 | A total number of leaves as classified |
| | herbivory class 3 (26-50% loss). |
| herbivory_4 | A total number of leaves as classified |
| | herbivory class 4 (51-75% loss). |
| herbivory_5 | A total number of leaves as classified |
| | herbivory class 5 (> 76% loss). |
| Latitude | WGS84 latitudinal and longitudinal |
| | coordinates in decimal degrees (°) to 2 |
| | decimal places. |
| Longitude | Elevation above mean sea level (m). |

304

305 11. ACKNOWLEDGEMENT

We are grateful to Shigeru Niwa, Satoshi N. Suzuki, Hiroko Kurokawa, and YusukeOnoda

308 for valuable advice on the study, and Kazuo Doi, Kazuhiro Onishi, Fumiyo Konishi,

309 Hiroko Ikuma, Keiko Ono, and Hitomi Asano for the measuring the herbivory rate. We

thank also all individuals who have contributed to obtain the data reported here and

311 facilitate the Forest and Grassland Survey of Monitoring Sites 1000 Project of Ministry

312 of the Environment (see also Appendix.pdf). The study was partly supported by the

313 Environmental Research and Technology Development Fund (S-9-3) of the Ministry of

the Environment, Japan.

315

316 12. LITERATURE CITED

- 317 Andrew, N., & Hughes, L. (2005) Herbivore damage along a latitudinal gradient:
- 318 relative impacts of different feeding guilds. *Oikos* 108: 176-182. doi:
- 319 10.1111/j.0030-1299.2005.13457.x
- 320 Coley, P. D., & Aide, T. M. (1991) Comparison of herbivory and plant defenses in
- 321 temperate and tropical broad-leaved forests. In: Price P. W., Lewinsohn T. M.,
- 322 Fernandes, G. W. & Benson, W. W. (ed), *Plant-Animal Interactions: Evolutionary*
- 323 Ecology in Tropical and Temperate Regions. Wiley & Sons, New York, pp 25-49
- 324 Coley, P. D., & Barone, J. A. (1996) Herbivory and plant defences in tropical forests.

325 Annual Review of Ecology and Systematics 27: 305-335. doi:

- 326 10.1146/annurev.ecolsys.27.1.305
- 327 Coley, P. D., Bryant, J. P., & Chapin F. S. (1985) Resource availability and plant
- 328 antiherbivore defense. *Science* 230: 895-899. DOI 10.1126/science.230.4728.895
- 329 Cyr, H., & Pace, M. L. (1993) Magnitude and patterns of herbivory in aquatic and
- 330 terrestrial ecosystems. *Nature* 361: 148-150. doi: 10.1038/361148a0
- 331 Dobzhansky, T. (1950) Evolution in the tropics. *American Scientist* 38: 209-221. doi:
 332 10.2307/2989767
- 333 Endara, M., & Coley, P. D. (2011) The resource availability hypothesis revisited: a
- 334 meta-analysis. *Functional Ecology* 25: 389-398. doi:
- 335 10.1111/j.1365-2435.2010.01803.x
- Givinish, T. J. (2002) Adaptive significance of evergreen vs deciduous leaves: Solving
 the triple paradox. *Silva Fennica* 36: 703-734. doi: 10.14214/sf.535
- 338 Hiura, T. & Nakamura, M. (2013) Different mechanisms explain feeding type-specific
- 339 patterns of latitudinal variation in herbivore damage among diverse feeding types of
- 340 herbivorous insects. *Basic and Applied Ecology* 14: 480-488. doi:
- 341 10.1016/j.baae.2013.06.004
- 342 Ishihara, M., Toyota, A., & Nakamura, M. (2007) Monitoring Sites 1000 (Forest Survey)
- 343 (in Japanese). *Japanese Journal of Ecology* 57: 438-442. doi:
- 344 10.18960/seitai.57.3_438

- 345 Ishihara, M., Ishida, K., Ida, H., Itoh, A., Enoki, T., Ohkubo, T., Kaneko, T., Kaneko, N.,
- 346 Kuramoto, S., Sakai, T., Saito, S., Sakio, H., Sakimoto, M., Shibano, H., Sugita, H.,
- 347 Suzuki, M., Takagi, M., Takashima, A., Takyu, M., Tashiro, N., Tanaka, N., Tokuchi,
- 348 N., Namikawa, K., Niiyama, K., Nishimura, N., Noguchi, M., Nomiya, H., Hiura, T.,
- 349 Fujiwara, A., Hoshino, D., Homma, K., Makita, A., Masaki, T., Yoshioka, T., &
- 350 Yoshida, T. (2010) An introduction to forest permanent plot data at Core and Subcore
- 351 sites of the Forest and Grassland Survey of the Monitoring Sites 1000 Project (in
- 352 Japanese). Japanese Journal of Ecology 60: 111-123. doi: 10.18960/seitai.60.1_111
- 353 Ishihara, M. I., Suzuki, S. N., Nakamura, M., Enoki, T., Fujiwara, A., Hiura, T.,
- Homma, K., Hoshino, D., Hoshizaki, K., Ida, H., Ishida, K., Itoh, A., Kaneko, T.,
- 355 Kubota, K., Kuraji, K., Kuramoto, S., Makita, A., Masaki, T., Namikawa, K.,
- 356 Niiyama, K., Noguchi, M., Nomiya, H., Ohkubo, T., Saito, S., Sakai, T., Sakimoto,
- 357 M., Sakio H., Shibano, H., Sugita, H., Suzuki, M., Takashima, A., Tanaka, N.,
- 358 Tashiro, N., Tokuchi, N., Yoshida, T., & Yoshida, Y. (2011) Forest stand structure
- 359 composition and dynamics in 34 sites over Japan. *Ecological Research* 26:
- 360 1007-1008. doi: 10.1007/s11284-011-0847-y
- 361 Janzen, D. H. (1970) Herbivores and the number of tree species in tropical forests.
- 362 *American Naturalist* 104: 501-528.
- 363 Japan Meteorological Agency (2012) Mesh Climatic Data of Japan. Available at
- 364 <u>http://nlftp.mlit.go.jp/ksj/gml/datalist/KsjTmplt-G02.html</u>. (last accessed on March
 365 20, 2020)
- 366 Kooyers, N. J., Blackman, B. K., & Holeski, L. M. (2017) Optimal defense theory
- 367 explains deviation from latitudinal herbivory defense hypothesis. *Ecology* 98:
- 368 1036-1048. dio: 10.1002/ecy.1731
- 369 Kozlov, M. V. (2008) Losses of birch foliage due to insect herbivory along
- 370 geographical gradients in Europe: a climate-driven pattern? *Climatic Change* 87:371 107-117.
- 372 Kozlov, M. V., van Nieukerken, E. J., Zverev, V., & Zvereva, E. L. (2013) Abundance
- and diversity of birch-feeding leafminers along latitudinal gradients in northern
- 374 Europe. *Ecography* 16: 1138-1149. doi: 10.1111/j.1600-0587.2013.00272.x

- 375 Kozlov, M. V., & Zvereva, E. L. (2015) Changes in the background losses of woody
- 376 plant foliage to insects during the past 60 years: are the predictions fulfilled? *Biology*
- 377 Letters 11: 20150480. doi: 10.1098/rsbl.2015.0480
- 378 Kozlov, M. V., Lanta, V., Zverev, V., & Zvereva E. L. (2015) Global patterns in
- 379 background losses of woody plant foliage to insects. *Global Ecology and*
- 380 *Biogeography* 24: 1126-1135. doi: 10.1111/geb.12347
- 381 Lim, J. Y., Fine, P. V. A., & Mittelbach, G. G. (2015) Assessing the latitudinal gradient
- in herbivory. *Global Ecology and Biogeography* 24: 1106-1112. doi:
- 383 10.1111/geb.12336
- 384 MacArthur, R. H. (1972) Geographical ecology: patterns in the distribution of species.
- Harper and Row. New York, New York, USA.
- 386 Metcalfe, D. B., Gregory, P. A., Martin, R. E., Espejo, E. S., Huasco, W. H., Amezquita,
- 387 F. F., Carranza-Jimenez, L., Galiano-Cabrera, D. F., Baca, L. D., Sinca, F., Quispe, L.
- 388 P. H., Taype, I. A., Mora, L. E., Davila, A. R., Solorzano, M. M., Vilca, B. P., Roman,
- J. M. L., Bustios, P. C. G., Revilla, N. S., Tupayachi, R., Girardin, C. A. J., Doughty,
- 390 C. E., & Malhi, Y. (2013) Herbivory makes major contributions to ecosystem carbon
- and nutrient cycling in tropical forests. *Ecology Letters* 17: 324-332. doi:
- 392 10.1111/ele.12233
- 393 Moles, A., Bonser, S., Poore, A., Wallis, I., & Foley, W. J. (2011) Assessing the evidence
- for latitudinal gradients in plant defence and herbivory. *Functional Ecology* 25:
- 395 380-388. doi: 10.1111/j.1365-2435.2010.01814.x
- 396 Moreira, X., Castagneyrol, B., Abdala-Roberts, L., Teran, J. C. B., Timmermans, B. G.
- 397 H., Bruun, H. H., Covelo, F., Glauser, G., Rasmann, S., & Tack, A. J. M. (2018)
- 398 Latitudinal variation in plant chemical defences drives latitudinal patterns of leaf
- 399 herbivory. *Ecography* 41: 1124-1134. doi: 10.1111/ecog.03326
- 400 Nakamura, M., Nakaji, T., Muller, O., & Hiura, T. (2014) Different initial responses of
- 401 the canopy herbivory rate in mature oak trees to experimental soil and branch warming
- 402 in a soil-freezing area. *Oikos* 124:1071-1077. doi: 10.1111/oik.01940

- 403 Nakamura, M., Asanuma, M., & Hiura, T. (2010) Differential effects of host plant
- 404 hybridization on herbivore community structure and grazing pressure on forest
- 405 canopies. *Oikos* 119: 1445-1452. doi: 10.1111/j.1600-0706.2010.18255.x
- 406 Niwa, S., Toyota, A., Kishimoto, T., Sasakawa, K., Abe, S., Chishima, T., Higa, M.,
- 407 Hiura, T., Homma, K., Hoshino, D., Ida, H., Kamata, N., Kaneko, Y., Kawanishi, M.,
- 408 Kobayashi, K., Kubota, K., Kuraji, K., Masaki, T., Niiyama, K., Noguchi, M.,
- 409 Nomiya, H., Saito, S., Sakimoto, M., Sakio, H., Sato, S., Shibata, M., Takashima, A.,
- 410 Tanaka, H., Tashiro, N., Tokuchi, N., Torikai, H. & Yoshida, T. (2016) Monitoring
- 411 of the ground-dwelling beetle community and forest floor environment in 22
- 412 temperate forests across Japan. *Ecological Research*: 31. 607-608. doi:
- 413 10.1007/s11284-016-1379-2
- 414 Pennings, S. C, & Silliman, B. R. (2005) Linking biogeography and community
- 415 ecology: Latitudinal variation in plant-herbivore interaction strength. *Ecology* 86:
- 416 2310-2319. doi: 10.1890/04-1022
- 417 Reich, P. B., Ellsworth, D. S., & Walters, M. B. (1998) Leaf structure (specific leaf
- 418 area) modulates photosynthesis-nitrogen relations: Evidence from within and across
- 419 species and functional groups. *Functional Ecology* 12: 948-958. doi:
- 420 10.1046/j.1365-2435.1998.00274.x
- 421 Saihanna, S., Tanaka, T., Okamura, Y., Kusumoto, B., Shiono, T., Hirao, T., Kubota, Y.
- 422 & Murakami M. (2018) A paradox of latitudinal leaf defense strategies in deciduous
- 423 and evergreen broadleaved trees. *Ecological Research*: 33. 1011-1017. doi:
- 424 10.1007/s11284-018-1641-x
- 425 Schemske, D. W., Mittelbach, G. G., Cornell, HV., Sobel, J. M., & Roy, K. (2009) Is
- 426 there a latitudinal gradient in the importance of biotic interactions? Annual Review of
- 427 *Ecology, Evolution, and Systematics* 40: 245–269. doi:
- 428 10.1146/annurev.ecolsys.39.110707.173430
- 429 Salazar, D. & Marquis, R.J. (2012) Herbivore pressure increases toward the equator.
- 430 *Proceedings of the National Academy of Sciences* 109: 12616-12620. Doi:
- 431 https://doi.org/10.1073/pnas.1202907109

- 432 Suzuki, S. N., Ishihara, M. I., Nakamura, M., Abe, S., Hiura, T., Homma, K., Higa, M.,
- 433 Hoshino, D., Hoshizaki, K., Ida, H., Ishida, K., Kawanishi, M., Kobayashi, K., Kuraji,
- 434 K., Kuramoto, S., Masaki, T., Niiyama, K., Noguchi, M., Nomiya, H., Saito, S., Sakai,
- 435 T., Sakimoto, M., Sakio, H., Sato, T., Shibano, H., Shibata, M., Suzuki, M., Takashima,
- 436 A., Tanaka, H., Takagi, M., Tashiro, N., Tokuchi, N., Yoshida, T., & Yoshida, Y.
- 437 (2012) Nation-wide litter fall data from 21 forests of the Monitoring Sites 1000 Project
- 438 in Japan. *Ecological Research* 27: 989-990. doi: 10.1007/s11284-012-0980-2
- 439 Wang, X. F., Liu, J. F., Gao, W. Q., Deng, Y. P., Ni, Y. Y., Xiao, Y. H., Kang, F. F.,
- 440 Wang, Q., Lei, J. P., & Jiang, Z. P. (2016) Defense pattern of Chinese cork oak
- 441 across latitudinal gradients: influences of ontogeny, herbivory, climate and soil
- 442 nutrients. Scientific Reports 6: 27269. doi: 10.1038/srep27269
- 443 Zhang, S., Zhang, Y., & Ma, K. (2016) latitudinal variation in herbivory: hemispheric
- 444 asymmetries and the role of climatic drivers. *Journal of Ecology* 104: 1089-1095.
- 445 doi: 10.1111/1365-2745.12588
- 446

447 13. FIGURE LEGEND

- 448 Figure 1 Location of the 19 natural forest sites across Japan.
- 449 Figure 2 Latitudinal gradient of insect herbivory in (a) deciduous and (b) evergreen
- 450 broadleaf tree species in 2014 and 2015. Symbols represent the mean herbivory rate
- 451 of each site. Lines indicate fitted lines from the prediction from the linear mixed452 models.
- 453 Figure 3 Latitudinal gradient of insect herbivory in (a) *Quercus crispula* and (b) *Fagus*
- 454 *crenata* in 2014 and 2015. Symbols represent the mean herbivory rate of each site.
- 455 Lines indicate fitted lines from the prediction from linear mixed models.
- 456
- 457 Storage location URL
- 458 JaLTER Database URL...(describe after accepted)
- 459 The corresponding author stores the original data.
- 460

- 461 The English in this document has been checked by at least two professional editors,
- 462 both native speakers of English. For a certificate, please see:
- 463 <u>http://www.textcheck.com/certificate/pyW86q</u>
- 464







Table 1. Major climatic zones and biogeographic regions of the 19 sites. The blank cells incidate our study sites is not located on these climatic zones and biogeographic regions, although these distribute in Japan. "Rare" and "Not distributed" in the cell means that the combination of climatic zones and biogeographic regions is rarely or not distributed in Japan.

| Dia ang ang hia ang ian | Climatic zone | | | | | | | | |
|--|---|---|-----------------------------|-----------------|--|--|--|--|--|
| Biogeographic region | Subalpine or subarctic | Cool temperate | Warm temperate | Subtropical | | | | | |
| Eastern Hokkaido Island | | Uryu (UR) | Not distributed | Not distributed | | | | | |
| Western Hokkaido Island | Vestern Hokkaido land Tomakomai (TM) <i>Not distribute</i> | | Not distributed | Not distributed | | | | | |
| The Pacific Ocean side of the Northern Honshu Island | Pacific Ocean of the Northern hu Island Aobayama (AO) ¹ Ogawa (OG) Chichibu (CC) Ooyamazawa (OY) | | Not distributed | | | | | | |
| The Sea of Japan side of the Northern Honshu Island | Otanomosutaira (OT) | Kanumazawa (KM) Kayanodaira (KY) | mazawa (KM) nodaira (KY) | | | | | | |
| Hokuriku and Sanin region | | Ashiu (AU) | Kamigamo (KG) | Not distributed | | | | | |
| The Pacific Ocean side of the Chubu region | | | Aichi-akazu (AI) | Not distributed | | | | | |
| The Inland Sea | Not distributed | Rare | | Not distributed | | | | | |
| The Kii Peninsula, Shikoku and Kyushu Islands | Rare | Wakayama (Ichinomata (Aya (AY) Tano (TN) | | Not distributed | | | | | |
| The Amami and Ryukyu Islands | Not distributed | Not distributed | Not distributed Rare | | | | | | |
| The Bonin Islands | Not distributed | Not distributed | Not distributed | | | | | | |

¹ Sites located between the cool and warm temperate zones.

Table 2. The four months when these traps caught the largest amount of fallen leaves in each site. The visual measurement of leaf damage by leaf-chewing insects was applied to these leaves.

| Site name | Abbreviated site name | Leaf collection period |
|-----------------|-----------------------|--|
| Uryu | UR | August–November |
| Tomakomai | ТМ | August–November |
| Kanumazawa | KM | August–November |
| Aobayama | AO | September–December |
| Ogawa | OG | September–December |
| Kayanodaira | KY | August–November |
| Otanomousutaira | OT | July-November |
| Ooyamazawa | OY | August–November |
| Chichibu | CC | September–December |
| Ashiu | AU | August–November |
| Aichi-akazu | AI | October–January |
| Kamigamo | KG | October–January |
| Wakayama | WK | September–December |
| Ichinomata | IC | April–July |
| Shiiba | SI | September–January in 2014 (sampled year), September–December in 2015 (sampled year) |
| Aya | AY | April–July |
| Tano | TN | April–July |
| Amami | AM | March–June |
| Yona | YN | February–May |

| SiteName | Broadleaf tree species | | | | | | | | | |
|----------------|---|---------------------|-------------------------------|----------------------|--------------------|------------------|-------------------|-------------------|------------------|----------------|
| Uryu | Betula ermanii | Quercus crispula | Acer mono | | | | | | | |
| Tomakomai | Cercidiphyllum japonicum | Ostrya japonica | Quercus crispula | Tilia japonica | Acer mono | | | | | |
| Kanumazawa | Cercidiphyllum japonicum | Fagus crenata | Quercus crispula | Pterocarya rhoifolia | Aesculus turbinata | | | | | |
| Aobayama | Fagus crenata | Quercus crispula | Quercus serrata | Idesia polycarpa | Prunus verecunda | | | | | |
| Ogawa | Castanea crenata | Fagus crenata | Fagus japonica | Quercus crispula | Quercus serrata | Acer mono | | | | |
| Kayanodaira | Fagus crenata | | | | | | | | | |
| Otanomosutaira | Betula ermanii | Acer ukurunduense | | | | | | | | |
| Ooyamazawa | Cercidiphyllum japonicum | Fagus crenata | Fraxinus platypoda | | | | | | | |
| Chichibu | Carpinus cordata | Fagus crenata | Fagus japonica | Quercus crispula | | | | | | |
| Ashiu | Magnolia obvata | Betula grossa | Fagus crenata | Quercus crispula | Acer sieboldianum | | | | | |
| Aichi-akazu | Quercus serrata | Acer sieboldianum | Cleyera japonica | | | | | | | |
| Kamigamo | Castanopsis cuspidata | Clethra barbinervis | Ilex macropoda | Ilex pedunculosa | | | | | | |
| Wakayama | Fagus crenata | Quercus acuta | Stewartia monadelpha | | | | | | | |
| Ichinomata | Quercus salicina | Cleyera japonica | | | | | | | | |
| Shiiba | Illicium anisatum | Betula grossa | Carpinus laxiflora | Carpinus tschonoskii | Fagus crenata | Quercus crispula | Acer sieboldianum | Swida controversa | Kalopanax pictus | Acer rufinerve |
| Aya | Machilus thunbergii | Distylium racemosum | Quercus acuta | Quercus salicina | | | | | | |
| Tano | Machilus thunbergii | Distylium racemosum | Castanopsis spp. ¹ | Quercus salicina | | | | | | |
| Amami | Castanopsis sieboldii subsp. lutchuensis | | | | | | | | | |
| Yona | Castanopsis sieboldii subsp. lutchuensis | Schima wallichii | | | | | | | | |

Table 3. The measured species in this study. Broadleaf tree species with a dominant ranking of five or species with a total of up to 50% of dominance and broad distributed species Quercus crispula and Fagus crenata were measured.

¹ Castanopsis sieboldii and Castanopsis cuspidata were included.

Table 4. Results of generalized linear mixed-effect models for estimating the latitudinal gradient of insect herbivory of deciduous broadleaf species, evergreen broadleaf species, *Q. crispula*, and *F. crenata* in 2014 and 2015. Estimates of parameters, Chi-square value, and statistical significance (*P*-value) are given. The likelihood ratio test was used for the statistical significance.

| Functional type/Species | Parameter | Estimate | Chi-square | P-value |
|-------------------------|---------------|----------|------------|-----------|
| Deciduous broadleaf | Latitude | 2.91 | 35.03 | P < 0.001 |
| | Year | -16.66 | 76.29 | P < 0.001 |
| | Latitude*Year | 0.49 | 69.22 | P < 0.001 |
| Evergreen broadleaf | Latitude | -2.15 | 4.82 | P < 0.05 |
| | Year | -4.84 | 25.98 | P < 0.001 |
| | Latitude*Year | 0.11 | 0.58 | 0.45 |
| Quercus crispula | Latitude | 3.42 | 8.99 | P < 0.01 |
| | Year | 12.69 | 5.92 | P < 0.05 |
| | Latitude*Year | -0.25 | 1.01 | 0.32 |
| Fagus crenata | Latitude | 1.75 | 2.91 | 0.09 |
| | Year | -16.01 | 641.48 | P < 0.001 |
| | Latitude*Year | 0.66 | 18.11 | P < 0.001 |

Table S1. Detailed ecological information of each plot collected from study sites. Soil parameters, snow depth, and maximum canopy height collected from study sites. This information was described in Appendix of Ishihara et al. (2011), Suzuki et al. (2012), and Niwa et al. (2011). Plot ID indicates an alphanumeric code of each plot (see SiteList.esv and Ishihara et al. (2011)). Soil types based on the soil classification survey conducted by the Ministry of Land, Infrastructure, Transport and Tourism, Japan (http://mb-www.mlit.go.jpkokjo/inspect/landclassification/download/index.html, last accessed on March 13, 2020). In addition, soil types based on the Classification of Forest Soil in Japan (Forest Soil Division 1976) were also shown, which are according to related literatures and personal observations of researchers. 'NA' means data not available. See appendix 1 for details and references, and other ecological information.

| Site name I | PlotID | Soil type | | | Soil pH | Bedrock | | Snow depth | | Maximum canopy height | |
|-------------------|--|---|------------------------------------|-----------|-----------------------------|--|---|------------------|---|-----------------------|--|
| Site name 1 | FAO | Forest Soil Divisior | Reference | Value | Reference | Туре | Reference | Value (m |) Reference | Value (m) | Reference |
| Uryu U | JR-BC1 Humic Cambisols | Brown forest soil | Shibata et al. 2002 | 3.9-4.5 | Shibata et al . 2002 | Andesite tuff-breccia | Shibata et al . 2002 | 2 | Shibata et al. 2002 | 28 | Yoshida T. unpublished data |
| Tomakomai TI | M-DB1 (Andic) Rhegosols | Shallow top soil | Hiura et al. 1998 | 5.3-6.2 | Shibata et al. 1998 | Volcanic ejecta of 1-2 m depth | Igarashi 1987 | 0.5 | Shibata et al. 1998 | 26.5 | Ishihara M. personal observation |
| Kanumazawa K | M-DB1 Residual Regosols | Gravel (large and sandy), brown forest soil | Masaki et al. 1999 | 5.2 | - | Igneous rock (green tuff and others) | - | 1.8 | Suzuki et al. 2002 | 30 | Suzuki et al. 2002 |
| Aobayama A | AO-BC1 Humic Cambisols | Brown forest soil ² | - | NA | | Aobayama formation on tuff | - | 0.1 | - | 20 | Kobayashi K. personal observation |
| Ogawa O | OG-DB1 Ochric Cambisols | Brown forest soil partly black or gley soi | Masaki et al. 1999 | 4.7-6.2 | Yoshinaga et al. 2002 | Metamorphic rock, volcanic ejacta | Yoshinaga et al. 2002 | 0.5 | Masaki et al. 1999 | About 35 | Nakashizuka 2002 |
| Kayanodaira K | Y-DB1 Humic Cambisols | Brown forest soil | Ida H. personal observatior | NA | | Plateau originated from lava flow | Ida 2013 | 3-4 | Ida 2013 | 25 | Watanabe 1994 |
| Otanomousutaira O | DT-EC1 Humo-Ferric (Gleyic) Podzols | Wet humus podzolic partly dry podzolic or moderately moist brown forest soil | Takai et al. 1976 | 3.8-4.5 | Takai <i>et al.</i> 1976 | Deposition of andesite and volcanic mudflow | Takai et al. 1976 | 2 | Ida H. personal observation | 22 | Kuroiwa and Watanabe 1997 |
| Ooyamazawa O | OY-DB1 Humo-Ferric Podzols | Sand, gravel, rock | Sakio 1997 | NA | | Greywacke, sandstone | Sakio 1997 | 0.3 | Sakio 1997 | 35 | Sakio H. unpublished data |
| Chichibu C | C-DB1 Humic Cambisols | Moderately moist brown forest soi | University Forest in Chichibu 2000 | NA | | Sedimentary rock | University Forest in Chichibu 2000 | 0.2-0.3 | Sawada et al. 2005 | 29 | University Forest in Chichibu unpublished dat |
| Ashiu A | AU-EC1 Humic Cambisols | Brown forest soil | Ueda et al. 1993 | 4.5 | Ueda et al. 1993 | Sandstone, slate, mudstone, shale, cher | Ueda et al. 1993; Yamanaka et al. 1993 | 2-3 | Yamanaka et al. 1993 | 25 | Kawanabe et al. 1994; Sakimoto M. personal observation |
| Aichi-akazu A | AI-BC1 Humic Cambisols | Moderately moist brown forest soil | Moroto et al. 1987 | 4.5-5.1 | Moroto et al. 1987 | Deeply weathered granite | Moroto et al. 1987 | 0.1014 | University Forest in Aichi, the University of Tokyo unpublished data | 20 | Ariyakanonet al. 2000 |
| Kamigamo K | G-EC1 Gleysols | Dry brown forest soil | Tokuchi et al. 2002 | NA | | Bedded chert with siliceous shale | Kimura et al. 1998 | Few ⁵ | - | 20 | Sakimoto M. personal observation |
| Wakayama W | WK-EC1 Humic Cambisols | Moderately moist brown forest soil | Ueda et al. 1994 | 4.8-4.9 | Ueda et al. 1994 | Sandstone, shale | Toda et al. 2000 | 0.36 | - | 25-30 | Sakimoto M. personal observation |
| Ichinomata I | IC-BC1 Humic Cambisols | Moderately moist to weakly dried brown forest soil, dry podzolic soil | Hirai et al. 2007 | 3.6-5.1 | Hirai et al. 2007 | Sandstone, mudstone | Sakai et al. 2006 | 0.15 | Sakai T. personal observation | 41 | Sakai et al. 2006 |
| Shiiba S | SI-DB1 NA | Brown forest soil | Enoki T. personal observation | 4.97-5.29 | Shibata H. unpublished data | Granite | Hatsushima 1970 | 0.2 | Enoki T. unpublished data | 27 | Enoki T. unpublished data |
| Aya A | AY-EB1 Humic Cambisols | Dry, moderately moist, or moderately moist drier subtype brown forest soil | Sato et al. 1999 | NA | | Shale, sandstone, partly covered by pumice stone from volcanic eruption | ^e Ohnuki et al. 1998; Sato et al. 1999 | 0 | Masaki et al. 1999 | 30 | Saito and Sato 2007 |
| Tano T | FN-EB1 Andosols | Moderately moist brown soi | Takagi M. unpublished data | 5.7 | Takagi M. unpublished data | Shale | Endo 1958 | 0 | Takagi M. personal observatior | 25 | Takagi M. unpublished data |
| Amami A | AM-EB1 Humic Cambisols | Weakly dried to moderately moist yellow soil at the valley | Ishida K. personal observation | NA | | Shale partly sandstone | Ishida K. personal observation | 0 | Ishida K. personal observation | 25 | Kumamoto Forest Office and Japan Forest Technology Association 1997 |
| Yona Y | N-EB1 Helvic Acrisols | Weakly dried to moderately moist yellow soi | Yamamori et al. 1986 | 4.1-4.3 | Yamamori et al. 1986 | Sandstone and slate | Enoki 2003 | 0 | - | 20 | Shinzato et al. 1986 |

¹ Cambie Red-Yellow soils (based on Japan Soil Inventory https://soil-inventory.dc.affrc.go.jp), with boulders and gravels.

² Scale 1:50,000 Fundamental Land Classification Survey in Miyagi, Sendai, 1976. 3 http://www.biology.tohoku.ac.jp/garden/geology.htm

⁴ Average between 1966 and 1999.

⁵ Few cm (Kamigamo Experimental Station, Kyoto University http://fserc.kyoto-u.ac.jp/kami/). 6 Wakayama Forest Research Station, Kyoto University (http://fserc.kyoto-u.ac.jp/waka/).

7 27 m for Abies firma and 24 m for Fagus crenata.

Appendix 1 Detailed information of each plot.

This material gives detailed information of each plot: forest age, disturbance history, soil type, soil pH, bedrock, snow depth, dwarf bamboo as understory vegetation, maximum canopy height, layout of the plot and subplots, remarks (optional), and acknowledgements (optional). This material includes information which was described in Appendix of Ishihara et al., (2011), Suzuki et al., (2012), and Niwa et al., (2016). Plot ID indicates an alphanumeric code of each plot (see SiteList.csv and Ishihara et al., (2011)). For definition of forest age classifications, see 7. A and Ishihara et al., (2011). Forest age or maximum tree age is the age in 2010 unless specified. Soil types based on the soil classification system of the Food and Agriculture Organization of the United Nations (FAO) (Dudal, 1968), were extracted from the 1:200,000 scale soil map of the Land Classification Survey conducted by the Ministry of Land, Infrastructure, Transport and Tourism, Japan (http://nrb-www.mlit.go.jp/kokjo/inspect/landclassification/download/index.html, last accessed on March 13, 2020). In addition, soil types based on the Classification of Forest Soil in Japan (Forest Soil Division, 1976) were also shown, which are according to related literatures and personal observations of researchers. 'NA' means data not available. References with * are those conducted in the plot.

Site name: Uryu

Plot ID: UR-BC1

Forest age: OG.

Disturbance: No record of human disturbance (Yoshida T. personal communication).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Brown forest soil (Shibata et al., 2002).

Soil pH: 3.9-4.5 (Ozawa, Shibata, Satoh, & Sasa 2001).

Bedrock: Andesite tuff-breccia (Shibata et al., 2002).

Snow depth: 2 m (Shibata et al., 2002).

Dwarf bamboo as understory vegetation: Understory is covered by dwarf bamboo (Yoshida T.

personal observation).

Maximum canopy height: 28 m (Yoshida T. unpublished data).

Plot & Subplots: The shape of plot is 70×150 m. The direction of Y-axis is 71° west from true north. *Remarks*: Croplands are 600 m northwest and 300 m southwest of the plot.

Acknowledgements: We thank the staff of Uryu Experimental Forests of Hokkaido University for the field work.

Site name: Tomakomai

Plot ID: TM-DB1

Forest age: OG. About 270-340 years old (Igarashi, 1987).

Disturbance: The forest regenerated after the volcanic eruption of Mt. Tarumae in 1669 and 1739 (Igarashi, 1987). The forest was disturbed by strong typhoons in 1954 (Mishima, Taniguchi, Taniguchi, & Hishinuma, 1958) and 2004.

Soil type FAO: (Andic) Rhegosols.

Soil type Forest Soil Division: Shallow top soil (Hiura et al., 1998*).

Soil pH: 5.3-6.2 (Shibata, Kirikae, Tanaka, Sakuma, & Hatano, 1998).

Bedrock: Volcanic ejecta of 1-2 m depth (Igarashi, 1987).

Snow depth: 0.5 m (Hiura et al., 1998*).

Dwarf bamboo as understory vegetation: Understory vegetation is partly dominated by *Sasamorpha borealis* (Hiura et al., 1998*).

Maximum canopy height: 26.5 m (Ishihara M. personal observation).

- *Plot & Subplots*: The 1-ha plot is a part of a 9-ha permanent plot. The direction of Y-axis is 31° west from true north.
- Acknowledgements: We thank the staff of Tomakomai Experimental Forests of Hokkaido University for the field work.

Site name: Kanumazawa

Plot ID: KM-DB1

- *Forest age*: OG. Maximum tree age is about 1000 years old according to Suzuki, Osumi, Masaki, Takahashi, Daimaru, & Hoshizaki (2002*).
- *Disturbance*: Canopy gaps and more infrequent, debris flows. No sign of human disturbance although selective cuttings were conducted at surrounding forests until 20–30 years ago (Masaki, Tanaka, Tanouchi, Sakai, & Nakashizuka, 1999*; Suzuki et al., 2002*).

Soil type FAO: Residual Regosols.

Soil type Forest Soil Division: Cambic Red-Yellow soils (based on Japan Soil Inventory https://soil-inventory.dc.affrc.go.jp), with boulders and gravels; brown forest soil (Masaki et al., 1999*).

Soil pH: 5.2.

Bedrock: Igneous rock (green tuff and others).

Snow depth: 1.8 m (Suzuki et al., 2002*).

Dwarf bamboo as understory vegetation: S. kurilensis and S. palmata are distributed widely but dominant only patchily. Instead, evergreen shrub (*Camellia japonica var. decumbens*), tall herbs (e.g. genera Laportea, Elatostema and Petasites) and ferns (genera Polystichum, Dyropteris and Arachniodes) dominate the understory layer (Hoshizaki, Suzuki, & Sasaki, 1997*). Maximum canopy height: 30 m (Suzuki et al., 2002*).

Plot & Subplots: The 1-ha plot is a part of a 4.71-ha permanent plot. The direction of Y-axis is 11° east from true north.

- *Remarks*: A stream flows in the plot. A road is 50 m east and 100 m northwest of the plot. Rivers are 250 m southeast and 400 m southwest of the plot.
- Acknowledgements: We thank Wajirou Suzuki, Katsuhiro Osumi and Kazunori Takahashi for early setup of the plot; Ohta Kazuhide for soil descriptions.

Site name: Aobayama

Plot ID: AO-BC1

Forest age: OG.

- *Disturbance*: Human usage of the forest has been restricted for the past 400 years (Suzuki Mitsuo personal communication).
- Soil type FAO: Humic Cambisols.
- Soil type Forest Soil Division: Brown forest soil (Scale 1:50,000 Fundamental Land Classification Survey in Miyagi, Sendai, 1976).
- Soil pH: NA.

Bedrock: Aobayama formation on tuff (http://www.biology.tohoku.ac.jp/garden/geology.htm). *Snow depth*: 0.1 m.

- *Dwarf bamboo as understory vegetation*: Patchy distribution of *Sasa borealis* (Kobayashi K. personal observation).
- Maximum canopy height: 20 m (Kobayashi K. personal observation).

Plot & Subplots: The direction of Y-axis is 18° west from true north.

Remarks: Grasslands and urban areas are 150 m north and southeast of the plot.

Acknowledgements: We thank Kadzutaka Kobayashi, Motonari Oyama for the field work.

Site name: Ogawa

Plot ID: OG-DB1

Forest age: OG.

Disturbance: Although the forest is an old-growth forest, human disturbances such as fire, grazing, and selective cutting took place until 1930s at surrounding forests. Remains of charcoal making were found around the plot (Masaki et al., 1999*; Suzuki, 2002*).

Soil type FAO: Ochric Cambisols.

Soil type Forest Soil Division: Brown forest soil partly black or gley soil (Masaki et al., 1999*).

Soil pH: 4.7-6.2 (Yoshinaga, Takahashi, & Aizawa, 2002*).

Bedrock: Metamorphic rock, volcanic ejacta (Yoshinaga et al., 2002*).

Snow depth: 0.5 m (Masaki et al., 1999*).

Dwarf bamboo as understory vegetation: Patchy distribution of Sasamorpha borealis and Sasa nipponica (Suzuki, 2002*).

Maximum canopy height: About 35 m (Nakashizuka, 2002*).

Plot & Subplots: The 1.2-ha (100×120 m) plot is a part of a 6-ha permanent plot (see Nakashizuka & Matsumoto, 2002). The direction of Y-axis is 100° west from true north.

Remarks: Grasslands and croplands are 400 m north and 500 m southwest of the plot.

Acknowledgements: Grants in support came from the Ministry of Agriculture, Forestry and Fishery, and the Ministry of Education, Science, Sports and Culture.

Site name: Kayanodaira

Plot ID: KY-DB1

Forest age: OG.

Disturbance: A light selective cutting probably occurred because remains of charcoal making were found around the plot (Watanabe, 1993; Ida, 2013).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Brown forest soil (Ida H. personal observation).

Soil pH: NA.

Bedrock: Plateau originated from lava flow (Ida, 2013).

Snow depth: 3-4 m (Ida, 2013).

Dwarf bamboo as understory vegetation: Understory is dominated by 1.5 m high Sasa kurilensis and Sasa senanensis (Peters, Nakashizuka, & Ohkubo, 1992; Ida, 2013).

Maximum canopy height: 25 m (Watanabe, 1994).

Plot & Subplots: The direction of Y-axis is 97° west from true north. Subplots were slightly moved after the census in 2006.

Remarks: Grasslands and pastures are 100 m west and 350 m south of the plot.

Site name: Otanomosutaira

Plot ID: OT-EC1

Forest age: OG.

Disturbance: No record of human disturbance (Ida H. personal observation).

Soil type FAO: Humo-Ferric (Gleyic) Podzols.

Soil type Forest Soil Division: Wet humus podzolic partly dry podzolic or moderately moist brown forest soil (Takai, Kanazawa, Asami, Takeshima, & Kawashima, 1976).

Soil pH: 3.8–4.5 (Takai et al., 1976).

Bedrock: Deposition of andesite and volcanic mudflow (Takai et al., 1976).

Snow depth: 2 m (Ida H. personal observation).

Dwarf bamboo as understory vegetation: Understory is dominated by 1 m high *Sasa kurilensis* (Kuroiwa & Watanabe, 1997*).

Maximum canopy height: 22 m (Kuroiwa & Watanabe, 1997*).

Plot & Subplots: The direction of Y-axis is 3° west from true north.

Remarks: Grasslands, wetlands, ski slopes and a road are 600 m west to southwest of the plot.

Site name: Ooyamazawa

Plot ID: OY-DB1

Forest age: OG. 254-year-old tree was recorded in 1988 (Sakio, 1997*).

Disturbance: Fraxinus platypoda established after a land slide caused by an earthquake in 1770 to

1790 (Sakio, 1997*). No record of logging (Kubo, Sakio, Shimano, & Ohno, 2005*).

Soil type FAO: Humo-Ferric Podzols.

Soil type Forest Soil Division: Sand, gravel, rock (Sakio, 1997*).

Soil pH: NA.

Bedrock: Greywacke, sandstone (Sakio, 1997*).

Snow depth: 0.3 m (Sakio, 1997*).

Dwarf bamboo as understory vegetation: 2 m height *Sasamorpha borealis* dominates at slope (Sakio H. personal observation).

Maximum canopy height: 35 m (Sakio H. unpublished data).

Plot & Subplots: The direction of Y-axis is 116° west from true north.

Remarks: The plot includes a stream and was established in a riparian forest dominated by *Fraxinus platypoda*, *Pterocarya rhoifolia*, and *Cercidiphyllum japonicum* (Sakio, Kubo, Shimano, & Ohno 2002*).

Acknowledgements: We thank Drs. Motohiro Kawanishi, Motoki Higa, and Masako Kubo for the field works of the research site. Thanks are also due to the members of Mori to Mizu no Genryu Bunkajuku for their various assistances.

Site name: Chichibu

Plot ID: CC-DB1

Forest age: OG.

Disturbance: No record of logging since the University forest was established in 1916.

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Moderately moist brown forest soil (University Forest in Chichibu, 2000).

Soil pH: NA.

Bedrock: Sedimentary rock (University Forest in Chichibu, 2000).

Snow depth: 0.2–0.3 m (Sawada, Ohkubo, Kaji, & Oomura, 2005*).

Dwarf bamboo as understory vegetation: None.

Maximum canopy height: 29 m (University Forest in Chichibu unpublished data).

Plot & Subplots: The direction of Y-axis is true north.

Remarks: A river flows 350 m southwest of the plot.

Acknowledgements: We thank the staff of the University of Tokyo Chichibu Forest for the field work.

Site name: Ashiu

Plot ID: AU-EC1

Forest age: OG. 230-year-old tree was recorded in 1980 (Tamai & Tempo, 1990).

Disturbance: Since the establishment of Ashiu Experimental Forest in 1924, no human disturbance occurred (Yamanaka, Matumoto, Oshima, & Kawanabe, 1993). Mass mortality of Fagaceae trees by Japanese oak wilt has occurred since 2002.

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Brown forest soil (Ueda, Ando, & Kanzaki, 1993).

Soil pH: 4.5 (Ueda et al., 1993).

Bedrock: Sandstone, slate, mudstone, shale, chert (Ueda et al., 1993; Yamanaka, Matumoto, Oshima, & Kawanabe, 1993).

Snow depth: 2–3 m (Yamanaka et al., 1993).

- *Dwarf bamboo as understory vegetation*: None since before sever herbivory by Sika deer occurred (Sakimoto M. personal observation).
- *Maximum canopy height*: 25 m (Kawanabe, Ando, Sakai, & Wada, 1994; Sakimoto M. personal observation).

Plot & Subplots: The direction of Y-axis is 69° west from true north.

Acknowledgements: We thank the technical staff of Aisu Forest Station, Kyoto University for their work.

Site name: Aichi-akazu

Plot ID: AI-BC1

Forest age: S. Less than 100 years old (Shibano, 2000*).

Disturbance: The forest established on the previously bare land due to fuel wood consumption (Shibano, 2000*). *Chamaecyparis obtusa* trees were planted in 1917–1918 to prevent soil erosion. At present, the forest is composed of pine tree and broadleaf tree species that have naturally established. Mass mortality of pine trees by Pine wilt disease occurred in 1980s and late 2000s. In

2010 and 2011, many oak trees were attacked by ambrosia beetle *Platypus quercivorus*, which transport the pathogenic fungi *Raffaelea quercivora* causing Japanese oak wilt.

Soil type FAO: Humic Cambisols.

- *Soil type Forest Soil Division*: Moderately moist brown forest soil (Moroto, Mashimo, & Haruta, 1987).
- Soil pH: 4.5–5.1 (Moroto et al., 1987).

Bedrock: Deeply weathered granite (Moroto et al., 1987).

Snow depth: 0.101 m on average between 1966 and 1999 (University Forest in Aichi, the University of Tokyo unpublished data).

Dwarf bamboo as understory vegetation: None.

Maximum canopy height: 20 m (Ariyakanon, Numamoto, & Suzuki, 2000).

- *Plot & Subplots*: The direction of Y-axis is true north. All subplots were relocated from outside to inside the plot after the census in 2004. Subplot 1, 3, 4 and 5 were moved several meters after the census in 2009.
- *Remarks*: Roads are 100 m south and 200 m north of the plot. Open lands and some buildings are 200 m west of the plot.

Acknowledgements: We thank Haruo Sawada, Yukiko Kamata, Masaki Matsui, Takehiko Tsukamoto, Yoshihide Hara, Kenji Yanase, Nobuyo Mizuno, Dr. Hirofumi Shibano, Dr. Noriko Iwai, and Dr. Koichiro Kuraji for the field work.

Site name: Kamigamo

Plot ID: KG-EC1

Forest age: S. About 90 years old (Sakimoto, Morishita, & Sakanoue, 2009b*).

Disturbance: After mass mortality of dominant pine trees by Pine wilt disease in 1970s,

Chamaecyparis obtusa that formed the middle and lower layers have become dominant (Sakimoto M. unpublished data).

Soil type FAO: Gleysols.

Soil type Forest Soil Division: Dry brown forest soil (Tokuchi, Fujimaki, & Terai, 2002*).

Soil pH: NA.

Bedrock: Bedded chert with siliceous shale (Kimura et al., 1998).

Snow depth: Few cm (Kamigamo Experimental Station, Kyoto University

http://fserc.kyoto-u.ac.jp/kami/).

Dwarf bamboo as understory vegetation: None (Sakimoto M. personal observation).

Maximum canopy height: 20 m (Sakimoto M. personal observation).

Plot & Subplots: The shape of plot is 80×80 m. The direction of Y-axis is 180° east from true north.

Remarks: A grassland and a golf course are 50 m south and 500 m southwest of the plot, respectively.

Urban areas are 150 m north, east and south of the plot.

Site name: Wakayama

Plot ID: WK-EC1

Forest age: OS. About 100 years old (Sakimoto et al., 2009a*).

Disturbance: Cut stumps created in 1920-1922 were found and the forest was used until the

establishment of the University Forest in 1926 (Furuno, Uenishi, & Uenishi, 1986).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Moderately moist brown forest soil (Ueda., Ando, & Takeuchi, 1994).

Soil pH: 4.8–4.9 (Ueda et al., 1994).

Bedrock: Sandstone, shale (Toda et al., 2000).

Snow depth: 0.3 m (Wakayama Forest Research Station, Kyoto University

http://fserc.kyoto-u.ac.jp/waka/).

Dwarf bamboo as understory vegetation: None (Sakimoto M. personal observation).

Maximum canopy height: 25-30 m (Sakimoto M. personal observation).

Plot & Subplots: The direction of Y-axis is 30° east from true north.

Site name: Ichinomata

Plot ID: IC-BC1

Forest age: OG. Maximum tree age is about 300 years old (Sakai T. unpublished data).

Disturbance: Chamaecyparis obtusa trees were cut selectively in 1985-1986 at the ridge (Sakai,

Sakai, Kuramoto, & Sato, 2006*; Sakai T. personal communication).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Moderately moist to weakly dried brown forest soil, dry podzolic soil (Hirai , Kaneko, & Takahashi, 2007*).

Soil pH: 3.6–5.1 (Hirai et al., 2007*).

Bedrock: Sandstone, mudstone (Sakai et al., 2006*).

Snow depth: 0.15 m (Sakai T. personal observation).

Dwarf bamboo as understory vegetation: None (Sakai T. personal observation).

Maximum canopy height: 41 m (Sakai et al., 2006*).

Plot & Subplots: The direction of Y-axis is 142° west from true north.

Remarks: A road is 450 m east of the plot.

Acknowledgements: We thank Tatsuro Kawasaki, Ryuichi Tabuchi and Atsushi Sakai for setup and maintenance of the research site. Thanks are also due to the members of Shikoku Research Center, Forestry and Forest Products Research Institute for their various assistances.

Site name: Shiiba

Plot ID: SI-DB1

Forest age: OS.

Disturbance: No record of logging since the establishment of University Forest at about 70 years ago.

No cut stump can be found although large trees are scarce (Enoki T. personal observation).

Soil type: Brown forest soil (Enoki T. personal observation).

Soil pH: 4.97–5.29 (Shibata H. unpublished data).

Bedrock: Granite (Hatsushima, 1970).

Snow depth: 0.2 m (Enoki T. unpublished data).

Dwarf bamboo as understory vegetation: Sasamorpha borealis dominates (Murata et al., 2009; Saruki, Inoue, Shiba, Nagasawa, Osaki, & Kubota, 2004).

Maximum canopy height: 27 m for *Abies firma* and 24 m for *Fagus crenata* (Enoki T. unpublished data).

Plot & Subplots: The shape of plot is 100×100 m.

Acknowledgements: We thank technical staff of the Shiiba Research Forest for the field work.

Site name: Aya

Plot ID: AY-EB1

Forest age: OG.

Disturbance: No record of human disturbance (Tanouchi and Yamamoto, 1995*). The forest experienced typhoon disturbance in 1993, 2004 and 2005 (Saito and Sato, 2007*).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Dry, moderately moist, or moderately moist drier subtype brown forest soil (Sato et al., 1999*).

Soil pH: NA.

Bedrock: Shale, sandstone, partly covered by pumice stone from volcanic eruption (Ohnuki, Sato, Fujimoto, & Inagaki 1998*; Sato et al., 1999*).

Snow depth: 0 m (Masaki et al., 1999*).

Dwarf bamboo as understory vegetation: None (Saito S. personal observation).

Maximum canopy height: 30 m (Saito and Sato, 2007*).

Plot & Subplots: The 1-ha plot is a part of a 4-ha permanent plot. The direction of Y-axis is 166° west from true north.

Remarks: A road and a river are 450 m northeast of the plot.

Acknowledgements: We thank Masahiro Kiyofuji, Dai Sugimoto, Mikiyo Hatomura, Yoshimi Narimatsu and the member of Kyushu Research Center, Forestry and Forest Products Research Institute for their various assistances. Site name: Tano

Plot ID: TN-EB1

Forest age: S. 87 years old (Kubota and Takagi, 2007*).

Disturbance: The forest regenerated in 1924 (Kubota and Takagi, 2007*).

Soil type FAO: Andosols.

Soil type Forest Soil Division: Moderately moist brown soil (Takagi M. unpublished data).

Soil pH: 5.7 (Takagi M. unpublished data).

Bedrock: Shale (Endo, 1958).

Snow depth: 0 m (Takagi M. personal observation).

Dwarf bamboo as understory vegetation: None (Takagi M. personal observation).

Maximum canopy height: 25 m (Takagi M. unpublished data).

Plot & Subplots: The direction of Y-axis is 6° east from true north.

Acknowledgements: We thank the staff of University of Miyazaki Tano Forest Science Station for the field work. We also thank Machiko Kuroki for processing litter trap samples.

Site name: Amami

Plot ID: AM-EB1

Forest age: OS. About 140 years old.

Disturbance: Remains of charcoal making were found in the plot. Protected from human disturbance for 100 years as a reserve (Ishida, Kawaguchi, Torikai, Takashi, & Kawaguchi, 2008).

Soil type FAO: Humic Cambisols.

Soil type Forest Soil Division: Weakly dried to moderately moist yellow soil at the valley (Ishida K. personal observation).

Soil pH: NA.

Bedrock: Shale partly sandstone (Ishida K. personal observation).

Snow depth: None (Ishida K. personal observation).

Dwarf bamboo as understory vegetation: None (Ishida K. personal observation).

Maximum canopy height: 20 m (Kumamoto Forest Office and Japan Forest Technology Association, 1997).

Plot & Subplots: The direction of Y-axis is 175° east from true north. Subplots were slightly moved after the first pitfall trapping in 2005.

Acknowledgements: We thank Hidemi Kawaguchi and the member of Amami Ecosystem Study Group.

Site name: Yona

Plot ID: YN-EB1

Forest age: OS.

Disturbance: Human disturbance such as selective cutting occurred until 1950s (Enoki, 2003*; Saito, 2011).

Soil type FAO: Helvic Acrisols.

Soil type Forest Soil Division: Weakly dried to moderately moist yellow soil (Yamamori, Hirata, Aramoto, Sunakawa, & Asato 1986).

Soil pH: 4.1–4.3 (Yamamori et al., 1986).

Bedrock: Sandstone and slate (Enoki, 2003*).

Snow depth: 0 m.

Dwarf bamboo as understory vegetation: Pleioblastus linearis distributed at ridges (Takashima A. personal observation).

Maximum canopy height: 20 m (Shinzato, Taba, Hirata, & Yamamori, 1986).

Plot & Subplots: The direction of Y-axis is 40° west from true north.

Acknowledgements: We thank Takanori Kinjo and the staff of Yona field Subtropical Field Science Center, Faculty of Agriculture, University of the Ryukyus.

Literature Cited

- Ariyakanon, N., Numamoto, S., & Suzuki, M. (2000) Sixty-year decreasing trend of bare land in Shirasaka watershed, University Forest in Aichi, revealed by aerial photography. *Bulletin of the* University of Tokyo Forests 103: 339-348
- Dudal, R. (1968) Definitions of soil units for the soil map of the world. *World Soil Resources Reports* No. 33, FAO, Rome
- Endo, T. (1958) Report of the geological survey of the Tano Instructional Forest of Miyazaki University (in Japanese). *Bulletin of the Miyazaki University Forests* 2: 1-25
- Enoki, T. (2003) Microtopography and distribution of canopy trees in a subtropical evergreen broad-leaved forest in the northern part of Okinawa Island, Japan. *Ecological Research* 18: 103-113
- Forest Soil Division. (1976) Classification of forest soil in Japan (1975) (in Japanese with English summary). *Bulletin of the Government Forest Experimental Station* 280: 1-28

- Furuno, T., Uenishi, Y., & Uenishi, K. (1986) Investigations on the productivity of Japanese fir (*Abies firma* Sieb. et Zucc.) and hemlock (*Tsuga sieboldii* Carr.) stands in Kyoto University
 Forest in Wakayama (VII) Profile of the natural reservation forest in compartment 9 (in Japanese). *Bulletin of the Kyoto University Forests* 57: 60-75
- Hatsushima, S. (1970) Preliminary reports on the forest flora of the Miyazaki Experimental Forest of Kyushu University (in Japanese with English summary). *Bulletin of the Kyushu University Forests* 44: 1-80
- Hirai, K., Kaneko, S., & Takahashi, M. (2007) Nitrogen mineralization of forest soil along the climate in Japan: estimation of rate of nitrogen mineralization in the field by soil properties, temperature and soil type (in Japanese with English summary). *Japanese Journal of Forest Environment* 49: 123-131
- Hiura, T., Fujito, E., Ishii, T., Naniwa, A., Sugata, S., Ishida, K., Murakami, M., Kato, E., Maeno, H.,
 Fukushima, Y., & Sakai, T. (1998) Stand structure of a deciduous broad-leaved forest in
 Tomakomai Experimental Forest, based on a large-plot data (in Japanese with English abstract). *Research Bulletin of Hokkaido University Forests* 55: 1-10
- Hoshizaki, K., Suzuki, W., & Sasaki, S. (1997) Impacts of secondary seed dispersal and herbivory on seedling survival in *Aesculus turbinata*. *Journal of Vegetation Science* 8: 735-742
- Ida, H. (2013) Forest structure in a beech (*Fagus crenata* Blume) stand on a 1-ha permanent plot for the Monitoring Sites 1000 Project in Kayanodaira, central Japanese snowbelt. *Bulletin of the Institute of Nature Education in Shiga Heights, Shinshu University* 50: 33-40
- Igarashi, Y. (1987) Vegetational succession in the Tomakomai Experiment Forest area (in Japanese with English summary). *Research Bulletins of the College Experiment Forests Hokkaido University* 44: 405-427
- Ishida, K., Kawaguchi, H., Torikai, H., Takashi, M., & Kawaguchi, K. (2008) Result of forest survey at Kinsakubaru national forest in Amami Island and ecosystem managements from the view point of mast seeding in *Castanopsis sieboldii* (in Japanese). *Japanese Forest Society Annual Meeting Database* 119: 4
- Ishihara, M. I., Suzuki, S. N., Nakamura, M., Enoki, T., Fujiwara, A., Hiura, T., Homma, K., Hoshino, D., Hoshizaki, K., Ida, H., Ishida, K., Itoh, A., Kaneko, T., Kubota, K., Kuraji, K., Kuramoto, S., Makita, A., Masaki, T., Namikawa, K., Niiyama, K., Noguchi, M., Nomiya, H.,

Ohkubo, T., Saito, S., Sakai, T., Sakimoto, M., Sakio, H., Shibano, H., Sugita, H., Suzuki, M., Takashima, A., Tanaka, N., Tashiro, N., Tokuchi, N., Yoshida, T., & Yoshida, Y. (2011) Forest stand structure composition and dynamics in 34 sites over Japan. *Ecological Research* 26: 1007-1008

- Kawanabe, S., Ando, M., Sakai, T., & Wada, S. (1994) The dynamics of natural forest on cool temperate deciduous forest zone mixing Sugi (*Cryptomeria japonica*) trees (II) - Studies on stand structure of Masukami A and B plots at Ashiu- (in Japanese). *Reports of the Kyoto University Forests* 26: 66-75
- Kimura, K., Yoshioka, T., Imoto, N., Tanaka, S., Musashino, M., & Takahashi, Y. (1998) Geology of the Kyoto-Tohokubu district (in Japanese with English summary). With geological Sheet Map at 1: 50,000. *Geological Survey of Japan*, Tsukuba, pp 89
- Kubo, M., Sakio, H., Shimano, K., & Ohno, K. (2005) Age structure and dynamics of *Cercidiphyllum japonicum* sprouts based on growth ring analysis. *Forest Ecology and Management* 213: 253-260
- Kubota, K., & Takagi, M. (2007) Long-term forest dynamic research site at an evergreen broadleaf secondary forest in Miyazaki University Forests (in Japanese). Annual Report of Field Science Center, Faculty of Agriculture, Miyazaki University 6: 57-61
- Kumamoto Forest Office, Japan Forest Technology Association. (1997) *Research report on the preservation of Scolopax mira and rare wildlife species* (in Japanese, title translated by the authors). Kumamoto Forest Office, Kumamoto
- Kuroiwa, Y., & Watanabe, R. (1997) Forest structure of a subalpine coniferous forest at Otanomousutaira, Shigakogen. *Institute of Nature Education in Shiga Heights, Shinshu University* 34: 11-22
- Masaki, T., Tanaka, H., Tanouchi, H., Sakai, T., & Nakashizuka, T. (1999) Structure, dynamics and disturbance regime of temperate broad-leaved forests in Japan. *Journal of Vegetation Science* 10: 805-814
- Mishima, T., Taniguchi, S., Taniguchi, M., & Hishinuma, Y. (1958) The actual states of wind damage in the Tomakomai Experimental Forest of Hokkaido University (II) (on the natural forest) (in Japanese with English summary). *Research Bulletins of the College Experiment Forests Hokkaido University* 19: 1-39

- Moroto, K., Mashimo, Y., & Haruta, Y. (1987) Soil properties and growth of Japanese red pine (*Pinus densiflora*) in the hilly and low-mountainous region of Central Japan (in Japanese with English summary). *Journal of the Japanese Forest Society* 69: 371-378
- Murata, I., Saruki, S., Kubota, K., Inoue, S., Tashiro, N., Enoki, T., Utsumi, Y., & Inoue, S. (2009) Effects of sika deer (*Cervus nippon*) and dwarf bamboo (*Sasamorpha borealis*) on seedling emergence and survival in cool-temperate mixed forests in the Kyushu Mountains. *Journal of Forest Research* 14: 296-301
- Nakashizuka, T. (2002) Disturbance regimes. In: Nakashizuka, T., & Matsumoto, Y. (eds) *Diversity* and interaction in a temperate forest community –Ogawa Forest Reserve of Japan. Springer, Tokyo, pp 67-80
- Nakashizuka, T., & Matsumoto, Y. (2002) *Diversity and interaction in a temperate forest community* -Ogawa Forest Reserve of Japan. Springer, Tokyo.
- Niwa, S., Toyota, A., Kishimoto, T., Sasakawa, K., Abe, S., Chishima, T., Higa, M., Hiura, T.,
 Homma, K., Hoshino, D., Ida, H., Kamata, N., Kaneko, Y., Kawanishi, M., Kobayashi, K.,
 Kubota, K., Kuraji, K., Masaki, T., Niiyama, K., Noguchi, M., Nomiya, H., Saito, S., Sakimoto,
 M., Sakio, H., Sato, S., Shibata, M., Takashima, A., Tanaka, H., Tashiro, N., Tokuchi, N., Torikai,
 H. & Yoshida, T. (2016) Monitoring of the ground-dwelling beetle community and forest floor
 environment in 22 temperate forests across Japan. *Ecological Research*, 31, 607-608.
- Ohnuki, Y., Sato T., Fujimoto, K., & Inagaki, M. (1998) Dynamics and physical properties of surficial soil and microtopography at Aya evergreen broad-leaved forest, southwestern Japan (in Japanese with English summary). *Japanese Journal of Forest Environment* 40: 67-74
- Ozawa, M., Shibata, H., Sato, F., & Sasa, K. (2001) Annual element budget of soil in snow-dominated forested ecosystem. *Water, Air, & Soil Pollution* 130: 703-708
- Peters, R., Nakashizuka, T., & Ohkubo, T. (1992) Regeneration and development in beech-dwarf bamboo forest in Japan. *Forest Ecology and Management* 55: 35-50
- Saito, K. (2011) Forest age distribution in Kunigami-village, Okinawa, based on forest register data (in Japanese). *Papers on environmental information science* 25: 245-250
- Saito, S., & Sato, T. (2007) Characteristics of typhoon damage to major tree species in a Lucidophyllous forest: a comparison of tree species responses to several typhoons at the Aya

long-term ecological research site (in Japanese with English summary). *Journal of the Japanese Forest Society* 89: 321-328

- Sakai, A., Sakai, T., Kuramoto, S., & Sato, S. (2006) Soil seed banks of an old-growth forest and an adjacent conifer plantation in a medium altitude region in Shikoku, Japan (in Japanese with English summary). *Japanese Journal of Forest Environment* 48: 85-90
- Sakimoto, M., Angeles-Perez, G., & Hirayama, K. (2009a) Spatial pattern, response to topology and coexistence of *Abies firma* and *Tsuga sieboldii* (in Japanese). *Abstract Annual Meeting of The Society of Vegetation Science* 14: 42
- Sakimoto, M., Morishita, K., & Sakanoue, N. (2009b) Spatial pattern and regeneration of *Chamaecyparis obtusa* in a natural forest of hilly area, Kyoto city (in Japanese). *Abstract Annual Meeting of Kansai Branch, Japan Forest Society* 60: 50
- Sakio, H. (1997) Effects of natural disturbance on the regeneration of riparian forests in a Chichibu Mountains, central Japan. *Plant Ecology* 132: 181-195
- Sakio, H., Kubo, M., Shimano, K., & Ohno, K. (2002) Coexistence of three canopy tree species in a riparian forest in the Chichibu Mountains, central Japan. *Folia Geobotanica* 37: 45-61
- Saruki, S., Inoue, S., Shiba, Y., Nagasawa, H., Osaki, S., & Kubota, K. (2004) Distribution and growth situation of Suzutake (*Sasamorpha borealis*) damaged by grazing of Shika deer (*Cervus nippon nippon*) in Miyazaki Forest of Kyushu University: Case study in 2003 (in Japanese with English summary). *Bulletin of the Kyushu University Forests* 85: 47-54
- Sato, T., Kominami, Y., Saito, S., Niiyama, K., Manabe, T., Tanouchi, H., Noma, N., & Yamamoto, S. (1999) An introduction to the Aya Research Site, a Long-Term Ecological Research site, in a warm temperate evergreen broad-leaved forest ecosystem in southwestern Japan: Research topics and design. *Bulletin of the Kitakyushu Museum of Natural History* 18: 157-180
- Sawada, H., Ohkubo, T., Kaji, M., & Oomura, K. (2005) Spatial distribution and topographic dependence of vegetation types and tree populations of natural forests in the Chichibu Mountains, Central Japan (in Japanese with English summary). *Journal of the Japanese Forest Society* 87: 293-303

- Shibano, H. (2000) Stand structure at Shirasaka watershed long-term research plot in University
 Forest in Aichi, the University of Tokyo (in Japanese). In: Kaji M (ed) *Study on forest ecosystems in long-term research plot*. Research Report of Grant-in-Aid for Scientific Research (B) 64-82
- Shibata, H., Ichikawa, K., Nomura, M., Sato, F., Sasa, K., Isii, Y., & Kobayashi, D. (2002)
 Elemental budgets of forest watershed at cold snowy region (in Japanese with English abstract). *Journal of Japanese Association of Hydrological Sciences* 32: 49-56
- Shibata, H., Kirikae, M., Tanaka, Y., Sakuma, T., & Hatano, R. (1998) Proton budgets of forest ecosystems on volcanogenous regosols in Hokkaido, Northern Japan. *Water, Air, & Soil Pollution* 105: 63-72
- Shinzato, T., Taba, K., Hirata, E., & Yamamori, N. (1986) Regeneration of *Castanopsis sieboldii* forest: 1. Studies on stratification and age structure of a natural stand (in Japanese with English summary). *The Science Bulletin of the Faculty of Agriculture. University of the Ryukyus* 33: 245-256
- Suzuki, S. N., Ishihara, M. I., Nakamura, M., Abe, S., Hiura, T., Homma, K., Higa, M., Hoshino, D., Hoshizaki, K., Ida, H., Ishida, K., Kawanishi, M., Kobayashi, K., Kuraji, K., Kuramoto, S., Masaki, T., Niiyama, K., Noguchi, M., Nomiya, H., Saito, S., Sakai, T., Sakimoto, M., Sakio, H., Sato, T., Shibano, H., Shibata, M., Suzuki, M., Takashima, A., Tanaka, H., Takagi, M., Tashiro, N., Tokuchi, N., Yoshida, T., & Yoshida, Y. (2012) Nation-wide litter fall data from 21 forests of the Monitoring Sites 1000 Project in Japan. *Ecological Research* 27: 989-990
- Suzuki, W. (2002) Forest vegetation in and around Ogawa Forest Reserve in relation to human impact. In: Nakashizuka T, & Matsumoto Y. (eds) *Diversity and interaction in a temperate forest community –Ogawa Forest Reserve of Japan.* Springer, Tokyo, pp 27-41
- Suzuki, W., Osumi, K., Masaki, T., Takahashi, K., Daimaru, H., & Hoshizaki, K. (2002)
 Disturbance regimes and community structures of a riparian and an adjacent terrace stand in the Kanumazawa Riparian Research Forest, northern Japan. *Forest Ecology and Management* 157: 285-301
- Takai, Y., Kanazawa, S., Asami, T., Takeshima, S., & Kawashima, N. (1976) Characteristics of soil organic matter and soil respiration in subalpine coniferous forest of Mt. Shigayama (Part 1) : On soil types and chemical properties of soil (in Japanese). *Japanese Journal of Soil Science and Plant Nutrition* 47: 33-38

- Tamai, S., & Tempo, Y. (1990) Age structure of trees in a natural cool-temperate forest (in Japanese with English summary). *Journal of the Japanese Forest Society* 72: 292-303
- Tanouchi, H., & Yamamoto, S. (1995) Structure and regeneration of canopy species in an old-growth evergreen broad-leaved forest in Aya district, southwestern Japan. *Plant Ecology* 117: 51-60
- Toda, H., Sasa, K., Sato, F., Shibata, H., Nomura, M., Ichikawa, K., Fujito, E., Takanishi, T., Seiwa, K., Tsukahara, H., Iida T., Taniguchi, N., Nakata, M., Kuwabara, S., Uchida, T., Haruta, Y., Inoue, M., Yagi, H., Tsukagoshi, T., Kuraji, K., Fukuda, M., Ono, H., Suzuki M., Imaizumi, Y., Yamaguchi, N., Takenaka, C., Yurugi, Y., Kawanabe, S., Ando, M., Nakanishi, A., Nishimura, K., Yamasaki, M., Nagayama, Y., Doi, N., Katagiri, S., Kofuji, R., Shinmura, Y., Inoue, S., Ezaki, T., Kohno, S., Fujihisa, M., Iwamatsu, I., Imayasu, K., Nakamura, S., Tsukamoto, J., Nogami, K., & Enoki, T. (2000) Stream water chemistry of university forests over Japan (in Japanese with English summary). *Journal of the Japanese Forest Society* 82: 308-312
- Tokuchi, N., Fujimaki, R., & Terai, M. (2002) Soil nitrogen dynamics of temperate conifer forest in central Japan The case study of Japanese cypress forest at Kamigamo Experimental Forest (in Japanese with English summary). *Forest research, Kyoto* 74: 47-52
- Ueda, S., Ando, M., & Kanzaki, K. (1993) Forest soil surveys of the Kyoto University Forest in Ashiu II. Soil types, grain size, and chemical and physical properties of soils. (in Japanese with English summary). *Bulletin of the Kyoto University Forests* 65: 94–112
- Ueda, S., Ando, M., & Takeuchi, M. (1994) Forest soil surveys of natural Japanese fir (*Abies firma* Sieb. et Zucc.) and hemlock (*Tsuga sieboldii* Carr.) stands and secondary broad-leaved stands in Kyoto University Forests in Wakayama (in Japanese). *Reports of the Kyoto University Forests* 26: 109–119
- University Forest in Chichibu, the University of Tokyo. (2000) *The ninth forest inventory of the University Forest in Chichibu, the University of Tokyo (in Japanese)*. University Forest in Chichibu, the University of Tokyo, Chichibu
- Watanabe, R. (1993) Forest structure of Kayanodaira beech forest of the Institute for Nature Study, Shinshu University 2: Growth of forest trees within a period (1982-1992). *Institute of Nature Education in Shiga Heights, Shinshu University* 30: 33-41

- Watanabe, R. (1994) Studies on the Fagus crenata forest in Kayanodaira 5: Forest dynamics of primary and secondary forests, 1979-1992. *Institute of Nature Education in Shiga Heights, Shinshu University* 31: 9-16
- Yamamori, N., Hirata, E., Aramoto, M., Sunakawa, S., & Asato, M. (1986) Studies on the working techniques by selection system for the broad leaved forest in the subtropics. (XII) Physical and chemical properties of soil at experimental plots (in Japanese with English summary). *The Science Bulletin of the Faculty of Agriculture. University of the Ryukyus* 33: 229-236
- Yamanaka, N., Matumoto, A., Oshima, Y., & Kawanabe, S. (1993) Stand structure of Mondori-Dani watershed, Kyoto University Forest in Ashiu (in Japanese). *Bulletin of the Kyoto University Forests* 65: 63-76
- Yoshinaga, S., Takahashi, M., & Aizawa, S. (2002) Landforms and soil characteristics in Ogawa Forest Reserve. In: Nakashizuka, T, Matsumoto, Y. (eds) *Diversity and interaction in a temperate forest community –Ogawa Forest Reserve of Japan*. Springer, Tokyo, pp 19-26