



Title	Regioselective Deaminative Allylation of Aliphatic Amines via Dual Cobalt and Organophotoredox Catalysis
Author(s)	Sekino, Tomoyuki; Sato, Shunta; Yoshino, Tatsuhiko; Kojima, Masahiro; Matsunaga, Shigeki
Citation	Organic letters, 24(11), 2120-2124 https://doi.org/10.1021/acs.orglett.2c00319
Issue Date	2022-03-09
Doc URL	http://hdl.handle.net/2115/88762
Rights	This document is the Accepted Manuscript version of a Published Work that appeared in final form in Organic Letters, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see https://pubs.acs.org/articlesonrequest/AOR-H4JCPXJAFRYRTCC68BPH .
Type	article (author version)
File Information	Main_deaminative_220304.pdf



[Instructions for use](#)


Regioselective Deaminative Allylation of Aliphatic Amines via Dual Cobalt and Organophotoredox Catalysis

Tomoyuki Sekino,[†] Shunta Sato,[†] Tatsuhiko Yoshino,^{†,‡} Masahiro Kojima,^{†,*} and Shigeki Matsunaga^{†,‡,*}

[†]Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo, Hokkaido 060-0812, Japan

[‡]Global Station for Biosurfaces and Drug Discovery, Hokkaido University, Sapporo, Hokkaido 060-0812, Japan

Supporting Information Placeholder



ABSTRACT: Despite the rapid progress in C–C-bond-forming reactions using Katritzky salts, their deaminative allylation remains a challenge. Inspired by the metallaphotoredox-catalyzed allylic substitution regime, here, we report the deaminative allylation of Katritzky salts via cobalt/organophotoredox dual catalysis. This cross-electrophile coupling enables regioselective allylation using a variety of allylic esters, overcoming the substrate limitations of reported protocols. Mechanistic studies indicate the involvement of a π -allyl cobalt complex as a radicalophile that mediates C–C bond formation.

Alkyl amines are ubiquitous structural motifs in commercially available building blocks, drug candidates, and biomolecules. Therefore, the transformation of alkyl amines into further derivatized molecules provides privileged opportunities in chemical synthesis. Since the pioneering work of Watson in 2017,¹ the reactions of Katritzky salts, which are readily synthesized in one step via the condensation of a primary amine with a pyrylium salt, have received intense attention in synthetic chemistry.^{2,3} However, in the area of C–C-bond-forming reactions,^{4–5} methods to convert amino groups into allylic substituents remain limited.⁶ Liu reported a photoredox-catalyzed desulfonylative allylation using allyl sulfones (Scheme 1A(a)),^{6a} which was later elaborated into a photocatalyst-free process by Aggarwal (Scheme 1A(b)).^{6b} By employing precisely designed homoallylic alcohols, Studer demonstrated a deaminative allylation via C–C bond cleavage (Scheme 1B).^{6c} All these allylation reactions share a common mechanism: the alkyl radical reductively generated from the Katritzky salt directly undergoes addition to the olefin, providing the stabilized radical intermediate (Scheme 1, gray inset). Thus, a substituent at the β -position of the allylating reagents is required, resulting in limitations in the substrate scope.⁷

The combination of photoredox and transition-metal catalysis, or metallaphotoredox catalysis, is a rapidly growing research area.⁸ Relevant to the present work, a radical-based approach to transition-metal-catalyzed allylic alkylation⁹ has been recognized as a complementary strategy to established protocols employing soft carbanion equivalents.


Furthermore, metallaphotoredox catalysis has been proved to be a viable strategy for cross-electrophile coupling,¹⁰ enabling selective coupling of two different electrophiles without the use of a stoichiometric metal reductant.¹¹ Inspired by these precedents, we hypothesized that π -allyl Co complexes, which are key intermediates in Co/photoredox-catalyzed allylic substitution,^{12–14} might serve as potent radicalophiles¹⁵ and realize a new deaminative allylation with a broader scope in terms of allylating reagents (Scheme 1C).

Our reaction design for the deaminative allylation is outlined in Scheme 2. Under irradiation with visible light, the photoredox catalyst (PC) in its excited state (PC*) oxidizes an electron donor (D). The reduced photocatalyst (PC^{•-}) and Co(II) undergo single electron transfer (SET), affording a Co(I) complex (Scheme 2A). Similarly, photoredox-induced SET between the reduced photocatalyst and Katritzky salts would afford an alkyl radical (Scheme 2B), which enters the dual Co and photoredox catalytic cycle (Scheme 2C). The low-valent Co(I) complex engages with an allylic acetate to form a π -allyl Co(III) intermediate. SET between the π -allyl Co(III) intermediate and the reduced photocatalyst yields a π -allyl Co(II) complex and the ground state photocatalyst. The alkyl radical from the Katritzky pyridinium salt is intercepted by the π -allyl Co(II) complex. Reductive elimination proceeds to afford the desired cross-coupled product and regenerate the Co(I) catalyst.


The results of our initial examination of the desired cross-electrophilic coupling based on this reaction proposal are summarized in Table 1. We attempted the deaminative

Scheme 1. Deaminative Allylation Reactions


A. Allylation using allyl sulfones




B. Allylation via C-C bond activation



Common mechanism



C. This work: Allylation via Co/photoredox dual catalysis



allylation of cyclohexyl Katritzky salt **2a** using (*E*)-cinnamyl acetate **1a** as the allylating reagent. After intensive investigation of the reaction conditions, the combination of 10 mol% of Co(BF₄)₂·6H₂O, 1.0 mol% of 4CzIPN as the organic PC, 4.0 equivalents of Hantzsch ester (HEH) as the electron donor, and 2.0 equivalents of 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP) as the base in the presence of MS4A in acetonitrile under blue LED irradiation provided the best result, furnishing **3aa** in 79% isolated yield with an excellent linear selectivity and *E*-selectivity (Table 1, entry 1).¹⁶ Interestingly, no additional ligands for Co were required for the reaction. Comparable results were obtained when CoBr₂ with 99.99% purity on a trace metal basis was used, clearly indicating that the reaction is mediated by cobalt and not by other metallic impurities (entry 2). The organophotocatalyst 4CzIPN proved to be more suitable for this metallaphotoredox system than iridium-centered photocatalysts (entry 3). The reaction proceeded in synthetically useful yield when the amount of HEH was reduced to 2.0 equiv (entry 4). It is noteworthy that comparable yield of **3aa** was observed when **2a** was used as a limiting reagent (entry 5). Metal reductants Mn and Zn did not promote the reaction, suggesting that photoredox conditions are uniquely effective for this type of cross-electrophilic allylation (entry 6).¹⁷

With the optimized conditions in hand, we investigated the substrate scope of this reaction with respect to allylating reagents (Table 2). In all the cases examined, the allylated products were obtained with exclusive linear

Scheme 2. Reaction Design






Table 1. Optimization of Reaction Conditions^a



entry	Deviation from above	yield (%) ^b
1	None	79 ^c
2	CoBr ₂ (99.99% purity)	63
3	Ir(ppy) ₂ (dtbbpy)PF ₆	30
4	2.0 equiv HEH	59
5	1a (2.0 equiv) and 2a as the limiting reagent	72 ^c
6	Mn or Zn (4.0 equiv) instead of 4CzIPN and HEH at 40 °C without blue LED irradiation	0



1a (0.050 mmol), **2a** (2.0 equiv), Co(BF₄)₂·6H₂O (10 mol%), 4CzIPN (1.0 mol%), HEH (4.0 equiv), DTBMP (2.0 equiv), and MS4A (200 g/mol) in MeCN (0.050 M to **1a**) at room temperature for 15 hours under blue LED irradiation unless otherwise noted. Regioselectivity and *E/Z* ratios were determined using GC-MS or ¹H NMR analysis of the crude mixture. For all cases, the linear/branched ratio was >20:1 and the *E/Z* ratio was >20:1. ^bDetermined by ¹H NMR. ^cIsolated yield in 0.15 mmol scale.

selectivity and high *E*-selectivity. From linear aromatic allyl esters, the products were obtained in good yield

irrespective of steric or electronic effects on the arene ring (**3aa–3ja**). Also, from branched aliphatic allyl acetates, linear products were obtained (**3kb–3mb**¹⁸), indicating that the regioselectivity of the products is not affected by the structure of the allylating reagents.¹⁹ An unsubstituted allylic acetate was also applicable to the deaminative reaction, delivering the terminal alkene **3nb** in 66% yield.


Table 2. Reaction Scope with Respect to Allylating Reagents^a

entry	allylating reagent 1	product 3	yield (%)	<i>E/Z</i> ^b
1			79 (84) ^c	>20:1
2		3ba	79	>20:1
3		3ca	85	>20:1
4		3da	83	>20:1
5		3ea	83	>20:1
6		3fa	60	14:1
7		3ga	73	>20:1
8		3ha	79	>20:1
9		3ia	81	>20:1
10 ^d		3ja	71	4.6:1
11		3kb	76	>20:1
12		3lb	81	>20:1
13		3mb	75	>20:1
14		3nb	66	-

^a**1** (0.15 mmol), **2** (2.0 equiv), Co(BF₄)₂·6H₂O (10 mol%), 4CzIPN (1.0 mol%), HEH (4.0 equiv), DTBMP (2.0 equiv), and MS4A (200 g/mol) in MeCN (0.050 M to **1**) at room temperature for 15 hours under blue LED irradiation. Regioselectivity was determined using GC-MS or ¹H NMR analysis of the crude mixture. For all cases, the linear/branched ratio was >20:1. Isolated yields. ^bDetermined by ¹H NMR analysis of the isolated product. ^c1.0 mmol scale. ^d**2** (1.6 equiv).

The substrate scope in terms of Katritzky pyridinium salts was also examined (Scheme 3). In essence, high regio- and geometrical selectivity was observed in all the cases examined (**3ab–3ah**). When diastereomerically pure Katritzky salt **2d** was employed as a starting material, almost no diastereoselectivity was observed in **3ad**, indicating the intermediacy of the alkyl radical.²⁰ An acyclic alkyl unit, namely, an isopropyl group, also participated in the cross-electrophilic allylation (**3ag**). The late-stage deaminative allylation of a steroid derivative successfully proceeded (**3ah**),

Scheme 3. Reaction Scope with Respect to Katritzky Salts^a




^aAs in **Table 2** (0.15 mmol scale). Regioselectivity was determined by GC-MS or ¹H NMR analysis of the crude mixture. For all cases, the linear/branched ratio was >20:1. Isolated yields. *E/Z* ratios were determined by ¹H NMR analysis of the isolated product. ^bGC yield. ^c0.10 mmol scale.

demonstrating the potential applicability of this approach in the derivatization of complex amines.²¹

To support our mechanistic proposal, the intermediacy of alkyl radicals was assessed by performing the reaction in the presence of TEMPO (Scheme 4). In this case, the TEMPO-trapped alkyl radical **4** was unequivocally obtained, and no cross-coupled product **3kb** was observed. This result, along with the loss of stereochemistry in **3ad** (Scheme 3), supports the involvement of an alkyl radical generated from the Katritzky salt.²²

Scheme 4. Intermediacy of alkyl radicals^a



^aAs in **Table 2** (0.050 mmol scale). ¹H NMR yield.

In summary, we have developed a cobalt/photoredox-catalyzed deaminative cross-electrophilic coupling of allylic acetates and Katritzky pyridinium salts. The dual catalytic system accommodated both linear and branched allylic esters irrespective of the nature of the substituents on the allyl unit, highlighting the synthetic advantages of the present system compared to known deaminative allylations. The intermediacy of a π -allyl Co complex is proposed based on preliminary mechanistic studies, which shed new light on metallaphotoredox-catalyzed allylic substitution reactions. Comprehensive investigations into the applicability of the

Co/photoredox-catalyzed allylation protocol to other radical-based bond-forming reactions, as well as further mechanistic studies including quantum chemical calculations,²³ are currently under investigation in our group.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/@@@>.

Experimental details and copies of spectroscopic data of the synthesized compounds (PDF)

AUTHOR INFORMATION

Corresponding Authors

Masahiro Kojima – Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan; orcid.org/0000-0002-4619-2621; Email: m-kojima@pharm.hokudai.ac.jp

Shigeki Matsunaga – Faculty of Pharmaceutical Sciences and Global Station for Biosurfaces and Drug Discovery, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan; orcid.org/0000-0003-4136-3548; Email: smatsuna@pharm.hokudai.ac.jp

Authors

Tomoyuki Sekino – Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan

Shunta Sato – Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan

Tatsuhiko Yoshino – Faculty of Pharmaceutical Sciences and Global Station for Biosurfaces and Drug Discovery, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, Hokkaido 060-0812, Japan; orcid.org/0000-0001-9441-9272; Email: tyoshino@pharm.hokudai.ac.jp

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported in part by JSPS KAKENHI Grants JP20H02730 (S.M.) and JP20K15946 (M.K.). This work was partly supported by Hokkaido University, Global Facility Center (GFC), Pharma Science Open Unit (PSOU), funded by MEXT under "Support Program for Implementation of New Equipment Sharing System." We thank Dr. Satoshi Maeda and Dr. Kimichi Suzuki at Hokkaido University for their support with computational studies using GRRM. Part of the calculations was performed at the Research Center for Computational Science, Okazaki, Japan. T.S. gratefully acknowledges JSPS and Nagai Memorial Research Scholarship from the Pharmaceutical Society of Japan for fellowships.

REFERENCES

(1) Basch, C. H.; Liao, J.; Xu, J.; Piane, J. J.; Watson, M. P. Harnessing Alkyl Amines as Electrophiles for Nickel-Catalyzed Cross Couplings via C–N Bond Activation. *J. Am. Chem. Soc.* **2017**, *139*, 5313–5316.
(2) Selected reviews on reactions using Katritzky pyridinium salts: (a) He, F.-S.; Ye, S.; Wu, J. Recent Advances in Pyridinium Salts as Radical Reservoirs in Organic Synthesis. *ACS Catal.* **2019**, *9*, 8943–

8960. (b) Kong, D.; Moon, P. J.; Lundgren, R. J. Radical Coupling from Alkyl Amines. *Nat. Catal.* **2019**, *2*, 473–476. (c) Correia, J. T. M.; Fernandes, V. A.; Matsuo, B. T.; Delgado, J. A. C.; de Souza, W. C.; Paixao, M. W. Photoinduced Deaminative Strategies: Katritzky Salts as Alkyl Radical Precursors. *Chem. Commun.* **2020**, *56*, 503–514. (d) Rössler, S. L.; Jelier, B. J.; Magnier, E.; Dagousset, G.; Carreira, E. M.; Togni, A. Pyridinium Salts as Redox-Active Functional Group Transfer Reagents. *Angew. Chem., Int. Ed.* **2020**, *59*, 9264–9280. (e) Li, Y.-N.; Xiao, F.; Guo, Y.; Zeng, Y.-F. Recent Developments in Deaminative Functionalization of Alkyl Amines. *Eur. J. Org. Chem.* **2021**, *2021*, 1215–1228.

(3) Tcyrulnikov, S.; Cai, Q.; Twitty, J. C.; Xu, J.; Atifi, A.; Bercher, O. P.; Yap, G. P. A.; Rosenthal, J.; Watson, M. P.; Kozłowski, M. C. Dissection of Alkylpyridinium Structures to Understand Deamination Reactions. *ACS Catal.* **2021**, *11*, 8456–8466.

(4) Selected examples of deaminative alkylation: (a) Klauck, F. J. R.; Yoon, H.; James, M. J.; Lautens, M.; Glorius, F. Visible-Light-Mediated Deaminative Three-Component Dicarbofunctionalization of Styrenes with Benzylic Radicals. *ACS Catal.* **2019**, *9*, 236–241. (b) Sun, S.-Z.; Romano, C.; Martin, R. Site-Selective Catalytic Deaminative Alkylation of Unactivated Olefins. *J. Am. Chem. Soc.* **2019**, *141*, 16197–16201. (c) Plunkett, S.; Basch, C. H.; Santana, S. O.; Watson, M. P. Harnessing Alkylpyridinium Salts as Electrophiles in Deaminative Alkyl–Alkyl Cross-Couplings. *J. Am. Chem. Soc.* **2019**, *141*, 2257–2262. (d) Wang, C.; Qi, R.; Xue, H.; Shen, Y.; Chang, M.; Chen, Y.; Wang, R.; Xu, Z. Visible-Light-Promoted C(sp³)–H Alkylation by Intermolecular Charge Transfer: Preparation of Unnatural α -Amino Acids and Late-Stage Modification of Peptides. *Angew. Chem., Int. Ed.* **2020**, *59*, 7461–7466. (e) Yang, T.; Wei, Y.; Koh, M. J. Photoinduced Nickel-Catalyzed Deaminative Cross-Electrophile Coupling for C(sp²)–C(sp³) and C(sp³)–C(sp³) Bond Formation. *ACS Catal.* **2021**, *11*, 6519–6525. (f) Sun, S.-Z.; Cai, Y.-M.; Zhang, D.-L.; Wang, J.-B.; Yao, H.-Q.; Rui, X.-Y.; Martin, R.; Shang, M. Enantioselective Deaminative Alkylation of Amino Acid Derivatives with Unactivated Olefins. *J. Am. Chem. Soc.* **2022**, *144*, 1130–1137.

(5) Selected examples of deaminative arylation: (a) Klauck, F. J. R.; James, M. J.; Glorius, F. Deaminative Strategy for the Visible-Light-Mediated Generation of Alkyl Radicals. *Angew. Chem., Int. Ed.* **2017**, *56*, 12336–12339. (b) Liao, J.; Guan, W.; Boscoe, B. P.; Tucker, J. W.; Tomlin, J. W.; Garnsey, M. R.; Watson, M. P. Transforming Benzylic Pyridinium Salts via C–N Bond Activation. *Org. Lett.* **2018**, *20*, 3030–3033. (c) Yue, H.; Zhu, C.; Shen, L.; Geng, Q.; Hock, K. J.; Yuan, T.; Cavallo, L.; Rueping, M. Nickel-Catalyzed C–N Bond Activation: Activated Primary Amines as Alkylating Reagents in Reductive Cross-Coupling. *Chem. Sci.* **2019**, *10*, 4430–4435. (d) Yi, J.; Badir, S. O.; Kammer, L. M.; Ribagorda, M.; Molander, G. A. Deaminative Reductive Arylation Enabled by Nickel/Photoredox Dual Catalysis. *Org. Lett.* **2019**, *21*, 3346–3351.

(6) Examples of deaminative allylation: (a) Zhang, M.-M.; Liu, F. Visible-Light-Mediated Allylation of Alkyl Radicals with Allylic Sulfones via a Deaminative Strategy. *Org. Chem. Front.* **2018**, *5*, 3443–3446. (b) Wu, J.; Grant, P. S.; Li, X.; Noble, A.; Aggarwal, V. K. Catalyst-Free Deaminative Functionalizations of Primary Amines by Photoinduced Single-Electron Transfer. *Angew. Chem., Int. Ed.* **2019**, *58*, 5697–5701. (c) Lübbesmeier, M.; Mackay, E. G.; Raycroft, M. A. R.; Elfert, J.; Pratt, D. A.; Studer, A. Base-Promoted C–C Bond Activation Enables Radical Allylation with Homoallylic Alcohols. *J. Am. Chem. Soc.* **2020**, *142*, 2609–2616.

(7) Similar limitations exist in γ , γ -difluoroallylations: (a) Li, Z.; Wang, K.-F.; Zhao, X.; Ti, H.; Liu, X.-G.; Wang, H. Manganese-Mediated Reductive Functionalization of Activated Aliphatic Acids and Primary Amines. *Nat. Commun.* **2020**, *11*, 5036. (b) Jin, Y.; Wu, J.; Lin, Z.; Lan, Y.; Wang, C. Merger of C–F and C–N Bond Cleavage in Cross-Electrophile Coupling for the Synthesis of *gem*-Difluoroalkenes. *Org. Lett.* **2020**, *22*, 5347–5352. (c) Liu, Y.; Tao, X.; Mao, Y.; Yuan, X.; Qiu, J.; Kong, L.; Ni, S.; Guo, K.; Wang, Y.; Pan, Y. Electrochemical C–N Bond Activation for Deaminative Reductive Coupling of Katritzky Salts. *Nat. Commun.* **2021**, *12*, 6745.

- (8) A selected review on metallaphotoredox catalysis: Chan, A. Y.; Perry, I. B.; Bissonnette, N. B.; Buksh, B. F.; Edwards, G. A.; Frye, L. I.; Garry, O. L.; Lavagnino, M. N.; Li, B. X.; Liang, Y.; Mao, E.; Millet, A.; Oakley, J. V.; Reed, N. L.; Sakai, H. A.; Seath, C. P.; MacMillan, D. W. C. Metallaphotoredox: The Merger of Photoredox and Transition Metal Catalysis. *Chem. Rev.* **2022**, *122*, 1485-1542.
- (9) Selected reviews of allylic substitutions using metallaphotoredox catalysis: (a) Huang, H.-M.; Bellotti, P.; Glorius, F. Transition metal-catalyzed allylic functionalization reactions involving radicals. *Chem. Soc. Rev.* **2020**, *49*, 6186-6197. (b) Zhang, M.-M.; Wang, Y.-N.; Lu, L.-Q.; Xiao, W.-J. Light Up the Transition Metal-Catalyzed Single-Electron Allylation. *Trends Chem.* **2020**, *2*, 764-775. Selected examples: (c) Lang, S. B.; O'Nele, K. M.; Tunge, J. A. Decarboxylative Allylation of Amino Alkanoic Acids and Esters via Dual Catalysis. *J. Am. Chem. Soc.* **2014**, *136*, 13606-13609. (d) Xuan, J.; Zeng, T.-T.; Feng, Z.-J.; Deng, Q.-H.; Chen, J.-R.; Lu, L.-Q.; Xiao, W.-J.; Alper, H. Redox-Neutral α -Allylation of Amines by Combining Palladium Catalysis and Visible-Light Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2015**, *54*, 1625-1628. (e) Matsui, K.; Gutiérrez-Bonet, Á.; Rotella, M.; Alam, R.; Gutierrez, O.; Molander, G. A. Photoredox/Nickel-Catalyzed Single-Electron Tsuji-Trost Reaction: Development and Mechanistic Insights. *Angew. Chem., Int. Ed.* **2018**, *57*, 15847-15851. (f) Zhang, H.-H.; Zhao, J.-J.; Yu, S. Enantioselective Allylic Alkylation with 4-Alkyl-1,4-dihydropyridines Enabled by Photoredox/Palladium Cocatalysis. *J. Am. Chem. Soc.* **2018**, *140*, 16914-16919. (g) Wang, Z.-J.; Zheng, S.; Romero, E.; Matsui, J. K.; Molander, G. A. Regioselective Single-Electron Tsuji-Trost Reaction of Allylic Alcohols: A Photoredox/Nickel Dual Catalytic Approach. *Org. Lett.* **2019**, *21*, 6543-6547. (h) Crisenza, G. E. M.; Faraone, A.; Gandolfo, E.; Mazarella, D.; Melchiorre, P. Catalytic Asymmetric C-C Cross-Couplings Enabled by Photoexcitation. *Nat. Chem.* **2021**, *13*, 575-580. (10) Selected reviews on cross-electrophile coupling: (a) Knappe, C. E. I.; Grupe, S.; Gärtner, D.; Corpet, M.; Gosmini, C.; Jacobi von Wangelin, A. Reductive Cross-Coupling Reactions between Two Electrophiles. *Chem. – Eur. J.* **2014**, *20*, 6828-6842. (b) Everson, D. A.; Weix, D. J. Cross-Electrophile Coupling: Principles of Reactivity and Selectivity. *J. Org. Chem.* **2014**, *79*, 4793-4798. Selected examples on cross-electrophilic allylation: (c) Qian, X.; Auffrant, A.; Felouat, A.; Gosmini, C. Cobalt-Catalyzed Reductive Allylation of Alkyl Halides with Allylic Acetates or Carbonates. *Angew. Chem., Int. Ed.* **2011**, *50*, 10402-10405. (d) Dai, Y.; Wu, F.; Zang, Z.; You, H.; Gong, H. Ni-Catalyzed Reductive Allylation of Unactivated Alkyl Halides with Allylic Carbonates. *Chem. – Eur. J.* **2012**, *18*, 808-812. (e) Anka-Lufford, L. L.; Prinsell, M. R. Weix, D. J. Selective Cross-Coupling of Organic Halides with Allylic Acetates. *J. Org. Chem.* **2012**, *77*, 9989-10000. (f) Ma, W.-Y.; Han, G.-Y.; Kang, S.; Pang, X.; Liu, X.-Y.; Shu, X.-Z. Cobalt-Catalyzed Enantiospecific Dynamic Kinetic Cross-Electrophile Vinylation of Allylic Alcohols with Vinyl Triflates. *J. Am. Chem. Soc.* **2021**, *143*, 15930. (g) Wang, L.; Wang, L.; Li, M.; Chong, Q.; Meng, F. Cobalt-Catalyzed Diastereo- and Enantioselective Reductive Allyl Additions to Aldehydes with Allylic Alcohol Derivatives via Allyl Radical Intermediates. *J. Am. Chem. Soc.* **2021**, *143*, 12755-12765. (h) Calogero, F.; Potenti, S.; Bassan, E.; Fermi, A.; Gualandi, A.; Monaldi, J.; Dereli, B.; Maity, B.; Cavallo, L.; Ceroni, P.; Cozzi, P. G. Nickel-Mediated Enantioselective Photoredox Allylation of Aldehydes with Visible Light. *Angew. Chem., Int. Ed.* **2022**, DOI: 10.1002/anie.202114981.
- (11) Selected examples of photoredox-assisted reductive coupling: (a) Zhang, P.; Le, C. C.; MacMillan, D. W. C. Silyl Radical Activation of Alkyl Halides in Metallaphotoredox Catalysis: A Unique Pathway for Cross-Electrophile Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 8084-8087. (b) Guan, H.; Zhang, Q.; Walsh, P. J.; Mao, J. Nickel/Photoredox-Catalyzed Asymmetric Reductive Cross-Coupling of Racemic α -Chloro Esters with Aryl Iodides. *Angew. Chem., Int. Ed.* **2020**, *59*, 5172-5177.
- (12) Examples of Co/photoredox-catalyzed allylic substitution: (a) Takizawa, K.; Sekino, T.; Sato, S.; Yoshino, T.; Kojima, M.; Matsunaga, S. Cobalt-Catalyzed Allylic Alkylation Enabled by Organophotoredox Catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 9199-9203. (b) Sekino, T.; Sato, S.; Kuwabara, K.; Takizawa, K.; Yoshino, T.; Kojima, M.; Matsunaga, S. Allyl 4-Chlorophenyl Sulfone as a Versatile 1,1-Synthon for Sequential α -Alkylation/Cobalt-Catalyzed Allylic Substitution. *Synthesis* **2020**, *52*, 1934-1946.
- (13) Selected examples of Co-catalyzed allylic substitution under thermal conditions: (a) Reddy, K.; Knochel, P. *Angew. Chem., Int. Ed. Engl.* New Cobalt- and Iron-Catalyzed Reactions of Organozinc Compounds. **1996**, *35*, 1700-1701. (b) Ghorai, S.; Chirke, S. S.; Xu, W.-B.; Chen, J.-F.; Li, C. Cobalt-Catalyzed Regio- and Enantioselective Allylic Amination. *J. Am. Chem. Soc.* **2019**, *141*, 11430-11434.
- (14) A selected review on cobalt/photoredox dual catalysis: (a) Kojima, M.; Matsunaga, S. The Merger of Photoredox and Cobalt Catalysis. *Trends Chem.* **2020**, *2*, 410-426. Selected examples: (b) Thullen, S. M.; Rovis, T. A Mild Hydroaminoalkylation of Conjugated Dienes Using a Unified Cobalt and Photoredox Catalytic System. *J. Am. Chem. Soc.* **2017**, *139*, 15504-15508. (c) Hu, X.; Zhang, G.; Bu, F.; Lei, A. Selective Oxidative [4+2] Imine/Alkene Annulation with H₂ Liberation Induced by Photo-Oxidation. *Angew. Chem., Int. Ed.* **2018**, *57*, 1286-1290. (d) Sun, X.; Chen, J.; Ritter, T. Catalytic Dehydrogenative Decarboxyolefination of Carboxylic Acids. *Nat. Chem.* **2018**, *10*, 1229-1233. (e) Meng, Q.-Y.; Schirmer, T. E.; Katou, K.; König, B. Controllable Isomerization of Alkenes by Dual Visible-Light-Cobalt Catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 5723-5728. (f) Cao, H.; Kuang, Y.; Shi, X.; Wong, K. L.; Tan, B. B.; Kwan, J. M. C.; Liu, X.; Wu, J. Photoinduced Site-Selective Alkenylation of Alkanes and Aldehydes with Aryl Alkenes. *Nat. Commun.* **2020**, *11*, 1956. (g) Cristófol, Á.; Limburg, B.; Kleij, A. W. Expedient Dual Co/Organophotoredox Catalyzed Stereoselective Synthesis of All-Carbon Quaternary Centers. *Angew. Chem., Int. Ed.* **2021**, *60*, 15266-15270. (h) Shi, C.; Li, F.; Chen, Y.; Lin, S.; Hao, E.; Guo, Z.; Wosqqa, U. T.; Zhang, D.; Shi, L. Photocatalytic Umpolung Synthesis of Nucleophilic π -Allylcobalt Complexes for Allylation of Aldehydes. *ACS Catal.* **2021**, *11*, 2992-2998. (i) Yasui, T.; Yamada, K.; Tatsumi, R.; Yamamoto, Y. Cobalt/Organophotoredox Dual-Catalysis-Enabled Cascade Cyclization of 1,6-Diynyl Esters via Formal 1,8-Acyloxy Migration. *ACS Catal.* **2021**, *11*, 11716-11722. (j) Zhao, H.; McMillan, A. J.; Constantin, T.; Mykura, R. C.; Juliá, F.; Leonori, D. Merging Halogen-Atom Transfer (XAT) and Cobalt Catalysis to Override E2-Selectivity in the Elimination of Alkyl Halides: A Mild Route toward *contra*-Thermodynamic Olefins. *J. Am. Chem. Soc.* **2021**, *143*, 14806-14813. (k) Occhialini, G.; Palani, V.; Wendlandt, A. E. Catalytic, *contra*-Thermodynamic Positional Alkene Isomerization. *J. Am. Chem. Soc.* **2022**, *144*, 145-152.
- (15) Capture of alkyl radicals by π -allylcobalt complexes has been proposed in previous reports in the literature. For examples, see 10c and 14b.
- (16) The reaction did not occur in the absence of cobalt salt, 4CzIPN, or light. See Table S1 in the Supporting Information for details.
- (17) See section 3 in the Supporting Information for additional investigations of the reaction conditions including the effect of other cobalt salts and ratio of **1** and **2**.
- (18) This system showed superior performance for the synthesis of **3mb** compared to the reported nickel/photoredox-catalyzed protocol. See Scheme S1 in the Supporting Information for details.
- (19) The convergence in the regioselectivity is in line with the intermediacy of π -allyl cobalt complexes. See also Scheme S2 in the Supporting Information for additional investigations.
- (20) See Scheme S3 in the Supporting Information for details.
- (21) Installation of primary alkyl group was not successful under the developed conditions. See section 4-3 of the Supporting Information for details.
- (22) The TEMPO adduct **4** was also obtained when the linear allylic ester **1a** was used instead of **1k**. See Scheme S4 in the Supporting Information for details.
- (23) See section 6-3 in the Supporting Information for preliminary investigations using DFT.

