



Title	Characterization of protease from animal periodontal pathogen <i>Porphyromonas gulae</i>
Author(s)	Alam, Urmi Saki; Inaba, Hiroaki; Yoshida, Sho; Nomura, Ryota; Nakano, Kazuhiko; Matsumoto-Nakano, Michiyo
Citation	Japanese Journal of Veterinary Research, 70(3&4), 79-90
Issue Date	2022-11
DOI	10.14943/jjvr.70.3-4.79
Doc URL	<a href="http://hdl.handle.net/2115/89308">http://hdl.handle.net/2115/89308</a>
Type	bulletin (article)
File Information	JJVR70-3-4_79-90_AlamSakiUrmi.pdf



[Instructions for use](#)

# Characterization of protease from animal periodontal pathogen *Porphyromonas gulae*

Alam Saki Urmi<sup>1)</sup>, Hiroaki Inaba<sup>1,\*</sup>, Sho Yoshida<sup>1)</sup>, Ryota Nomura<sup>2,3)</sup>, Kazuhiko Nakano<sup>3)</sup> and Michiyo Matsumoto-Nakano<sup>1)</sup>

<sup>1)</sup> Department of Pediatric Dentistry, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8556, Japan.

<sup>2)</sup> Department of Pediatric Dentistry, Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan.

<sup>3)</sup> Department of Pediatric Dentistry, Osaka University Graduate School of Dentistry, Suita-Osaka 565-0871, Japan.

Received for publication, June 23, 2022; accepted, October 7, 2022

## Abstract

*Porphyromonas gulae* is an animal-derived oral microorganisms known also to be associated with periodontal disease in humans. We previously reported that *P. gulae* proteases are possible one of virulence factors related to adhesion and invasion of gingival epithelial cells, and host cell destruction. Here we attempted to characterize bacterial proteases associated with *P. gulae*. Both alkaline phosphatase and trypsin activity were identified in all examined *P. gulae* strains and *P. gingivalis* ATCC33277. Each of the *P. gulae* strains also showed proteolytic activity in cell extract and/or culture supernatant samples. In addition, there was a significant increase in protease activity level seen in living bacterial cells, dependent on cell number, while there were no significant differences regarding proteolytic activity among the *P. gulae* strains. The present results indicate that antipain and PMSF are effective inhibitors of *P. gulae* proteases as well as *P. gingivalis*. In addition, TLCK and leupeptin significantly inhibited proteolytic activity in a dose-dependent manner. On the other hand, AEBSF, ALLN, aprotinin, bestatin, chymostatin, E64, EDTA, pepstatin, and phosphoramidon showed no inhibitory effects, while those of KYT-1 and KYT-36, *P. gingivalis* gingipain-specific inhibitors, were negligible. These results suggest that both *P. gulae* and *P. gingivalis* produce and secrete trypsin-like serine proteases, while the structure of those proteases differ between the two bacteria.

Key Words: API-ZYM, bacterial protease, *P. gulae*, protease inhibitor, proteolytic activity

## Introduction

*Porphyromonas gulae* has been isolated from the gingival sulcus of a lot of animal

species, including bear, canine, cat, coyote, marsupial, monkey, ovine, and wolf<sup>2,12,34</sup>. *P. gulae* organisms are a black-pigmented, Gram-negative coccobacillus, asaccharolytic, anaerobic, non-

\* Corresponding author : Hiroaki Inaba, DDS, PhD

Department of Pediatric Dentistry

Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences

2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan

Tel: +81-86-235-6716

Fax: +81-86-235-6719

E-mail: hinaba@okayama-u.ac.jp

doi: 10.14943/jjvr.70.3-4.79

motile, and non-spore-forming characteristics<sup>12,42</sup>, and has been observed in significant numbers of canines with periodontal diseases as compared to those with healthy gingiva<sup>42</sup>. *P. gulae* as well as human periodontal pathogen *P. gingivalis* have abilities to adhere to and invade human gingival epithelial cells, dependent on fimbria characteristics<sup>19</sup>.

Periodontal diseases, including gingivitis and chronic periodontitis, are common infection in human and dogs, left untreated, can lead to tooth loss<sup>17,35,49</sup>. The prevalence of periodontal disease in dogs reach up to 85% in individuals over the age of 4 years<sup>28</sup>. In addition, previous reports noted that *P. gulae* was detected in human gingival tissues from subjects with and without periodontitis<sup>49</sup>.

Bacterial protease, one of virulence factors, has been reported a large group of enzymes that catalyze the hydrolysis of peptide bonds and are classified into seven catalytic types; aspartyl, cysteine, serine, metallo, glutamic, threonine peptidase, and asparagine peptide lyase<sup>7,40</sup>. The roles of protease in bacterial pathogens have been reported to be degradation of host tissue components for growth and proliferation, and avoidance of the host immune defense system<sup>40</sup>. Moreover, secreted protease has been implicated in bacterial pathogenesis and shown to have several immunomodulating activities, including release of pro-inflammatory cytokines, extracellular matrix degradation, IgG cleavage, and degradation of other immunoglobulins<sup>40</sup>. Bacterial protease has been reported to cause a large number of infectious diseases, such as cholera, salmonellosis, Legionnaires' disease, bronchiectasis, cystic fibrosis, botulism, tetanus, and anthrax<sup>40</sup>.

Serine and cysteine proteases are important pathogenesis factors of the periodontal pathogens, such as *P. gingivalis*, *Treponema denticola*, *Tannerella forsythia*, and *Fusobacterium nucleatum*<sup>7,16</sup>. Protease produced by periodontal pathogens has been reported capable of degrading periodontal tissues and shown to inactivate host defense effectors, which promote pathological alterations associated with development and

progression of periodontal disease<sup>10,15</sup>. *P. gulae* protease in particular has been demonstrated to have arginine- and lysine-specific proteolytic activity<sup>30</sup>, and involved in initial attachment of that organism to human gingival epithelial cells<sup>19</sup>. Moreover, *P. gulae* protease play an important role in bacterial biology, such as hemagglutination activity, coaggregation and bacterial growth, as well as degradation of human proteins<sup>48</sup>.

Here we report that all examined strains of *P. gulae* mainly possess trypsin-like activity, and proteolytic activity was observed in both cell extracts and culture supernatants, as well as *P. gingivalis*. In addition, proteolytic activity of *P. gulae* was reduced by serine and cysteine protease inhibitors, such as antipain, phenylmethylsulfonyl fluoride (PMSF), Na-*p*-tosyl-L-lysine chloromethyl ketone (TLCK), and leupeptin. Our findings provide proteolytic characterization of *P. gulae* organisms using protease inhibitors.

## Materials and Methods

### *Bacterial culture and API-ZYM test*

*P. gingivalis* ATCC33277 and *P. gulae* ATCC51700, D040, D044, D049, D066, D077 were selected from our culture collections<sup>19,23</sup>. Bacterial cells were grown in Trypticase soy broth supplemented with yeast extract (1 mg/ml), menadione (1 µg/ml), and hemin (5 µg/ml), as described previously<sup>19</sup>. The phenotypic enzyme profiles of all strains and their supernatants were estimated using an API-ZYM test (API biomérieux, Marcy l'Etoile, France), according to the manufacturer's instructions.

### *Cell culture*

SAS cells, an oral squamous cell carcinoma cell line, were obtained from the Japanese Collection of Research Bioresources (Tokyo, Japan), and cultured in RPMI 1640 medium (Wako, Osaka, Japan) supplemented with 10% fetal bovine serum at 37°C in 5% CO<sub>2</sub>.

### Zymography

SAS cells were incubated with *P. gulae* strains ATCC 51700, D040, D044, D049, D066, and D077, and *P. gingivalis* ATCC 33277 at an MOI of 1 for 24 h. Culture supernatant samples (proMMP9) from SAS cells were collected and analyzed for proMMP9 activation using gelatin zymography, as previously described<sup>20</sup>. Briefly, samples (3 µg/lane) were mixed with SDS sample buffer without reducing reagents, then separated on 10% SDS-polyacrylamide gels containing 0.1% gelatin. The gels were incubated at 37°C with 2.5% Triton X-100 for 1 h, followed by 48 h with reaction buffer [20 mM Tris-HCl (pH 7.5), 200 mM NaCl, 5 mM CaCl<sub>2</sub>, 0.02% NaN<sub>3</sub>]. After staining with 5% Coomassie brilliant blue R-250, gelatinolytic activities were visualized as clear bands against a blue background.

### Chemicals

Eleven inhibitors were used in this study (Table 3) as follows; antipain, cysteine and serine protease inhibitor, TLCK, serine protease inhibitor; PMSF, serine protease inhibitor, were purchased from Sigma-Aldrich (St. Louis, MO). Leupeptin, cysteine and serine protease inhibitor, KYT-1, an arginine-gingipain inhibitor, and KYT-36, a lysine-gingipain inhibitor, were purchased from Peptide Institute (Osaka, Japan). A protease inhibitor set containing AEBSF, ALLN, antipain, aprotinin, bestatin, chymostatin, E-64, EDTA-Na<sub>2</sub>, leupeptin, pepstatin, phosphoramidon, and PMSF was purchased from G-Biosciences (St. Louis, MO).

### Protease activity assay

The protease activity of *P. gulae* cells and supernatants was determined using a Pierce Protease assay kit (Thermo Scientific, Rockford, IL), according to the manufacturer's instructions. Briefly, samples were mixed with a casein solution and incubation buffer, and then incubated at 37°C, after which trinitrobenzene sulfonic acid (TNBSA) was added and incubation was continued at 37°C. TNSBA reacts with exposed primary

amines to produce an orange-yellow product, which was measured on SH-1000 Lab microplate reader (Corona Electric, Ibaraki, Japan) at 450 nm. Samples were preincubated with various concentrations of each inhibitor at 37°C for 2 h before addition of the substrates. All inhibitors (1-100 µM) were dissolved in dimethyl sulfoxide.


### Statistical analyses

Data were analyzed by two-way analysis of variance (ANOVA) with Tukey multiple-comparison test and are presented as means standard deviations (SDs). Statistical significance was considered at  $P < 0.01$  and  $P < 0.05$ . The data presented are representative of at least three biological replicates.

### Results

*P. gulae* exhibits several virulence characteristics similar to those of the human periodontal pathogen *P. gingivalis*<sup>19,30,48,49</sup>. *Porphyromonas* species, isolated from companion animals, have been shown to have arginine- and lysine-specific proteolytic enzyme activities<sup>30</sup>. However, the proteolytic enzyme activities of *P. gulae* strains are not fully understood. Having previously tested the API-ZYM system, we first examined that the enzyme activities of *P. gulae* strains were assayed using bacterial cells and culture supernatants. As shown in Tables 1 and 2, all tested *P. gulae* strains as well as the *P. gingivalis* ATCC33277 consistently demonstrated alkaline phosphatase and trypsin activities, while no other enzyme activities were detected in any of the tested strains. In addition, protease activity increased in cell number dependent manner (Fig. 1A). Protease activity has been reported to differ considerably among all strains of *P. gingivalis* with type II fimbriae<sup>18</sup>. *P. gulae* protease was found to activate in both cell extracts and supernatants, with negligible differences among the strains (Fig. 1B and C).

Several protease inhibitors, including




**Fig. 1. Proteolytic activity of *P. gulae* strains.**

(A) Quantification of protease activities of *P. gulae* ATCC 51700 ( $1 \times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$  cfu), (B) ATCC 51700, D040, D044, D049, D066, and D077 ( $5 \times 10^7$  cfu), and (C) supernatants (10  $\mu$ g; the amount of protein contained in the supernatant), as compared with *P. gingivalis* ATCC33277. Protease activity was determined using a Pierce Protease assay kit, as described in Materials and Methods. Enzyme activities were recorded with a microplate reader and are expressed as arbitrary units. Data are shown as the mean  $\pm$  SD of three independent experiments. \*  $P < 0.01$  by ANOVA with Tukey multiple-comparison test.

serine and/or cysteine protease inhibitors, have been reported to inhibit the proteolytic activity of bacteria<sup>21,37</sup>. To clarify the characteristics of *P. gulae* protease, protease activity was determined using protease inhibitors, including antipain, PMSF, EBSF, ALLN, aprotinin, bestatin, chymostatin, E64, EDTA, pepstatin, phosphoramidon (Table 3). Antipain decreased *P. gulae* protease activation by  $17.54 \pm 3.83\%$ , and PMSF decreased *P. gulae* protease activation by  $38.27 \pm 3.32\%$ , whereas other inhibitors showed negligible effects. A time course analysis revealed that *P. gulae* as well as *P. gingivalis* protease activity was reduced by antipain and PMSF. In addition, both inhibitors diminished *P. gulae* protease as well as *P. gingivalis* in a concentration-dependent manner (Fig. 2). These

findings also suggest that *P. gulae* protease may possess serine protease and/or cysteine protease. Previous studies have noted that TLCK and leupeptin are inhibitors of trypsin-like and cysteine proteases<sup>14,45</sup>. In addition, TLCK and leupeptin reportedly diminishes *P. gingivalis* proteases<sup>21,37</sup>. Therefore, we examined the effects of TLCK and leupeptin on *P. gulae* protease activity using *P. gingivalis* cells as control. Both inhibitors separately and in combination significantly decreased the activities of *P. gulae* protease as well as that of *P. gingivalis* protease in time- and dose-dependent manner (Fig. 3). Together, these results suggest that *P. gulae* may produce and secrete some kinds of proteases. KYT-1 and KYT-36, specific inhibitors of *P. gingivalis* protease gingipains, have also showed significant inhibitory



**Fig. 2. Effects of antipain and PMSF inhibitors on *P. gulae* proteolytic activity.**

Quantification of protease activity of *P. gulae* ATCC 51700 ( $5 \times 10^7$  cfu) and *P. gingivalis* ATCC 33277 ( $5 \times 10^7$  cfu) with antipain or PMSF. A: Antipain. B: PMSF. Time course studies indicated for 0-4h, and dose response evaluations indicated at 100  $\mu$ M. Enzyme activities were recorded with a microplate reader and are expressed as arbitrary units. Data are shown as the mean  $\pm$  SD of three independent experiments. \*\*  $P < 0.01$  and \*  $P < 0.05$  by ANOVA with Tukey multiple-comparison test (compared with 0 h). ††  $P < 0.01$  and †  $P < 0.05$  by ANOVA with Tukey multiple-comparison test (compared with 1 h).


effects, as has been reported<sup>22)</sup>, whereas no inhibition of *P. gulae* protease activity was noted (Fig. 4). *P. gingivalis* protease has been reported to process the proenzyme to the active form of MMP9<sup>20)</sup>. Therefore, we examined the effects of selected *P. gulae* strains on proMMP9 activation. Zymography findings revealed that activation of proMMP9 was shown to be induced in vitro by all *P. gulae* strains as well as by *P. gingivalis* ATCC33277 (Fig. 5).

## Discussion

The present findings showed that all of the examined *P. gulae* strains produce serine protease with trypsin-like activity. Although details

regarding the classification and activities of *P. gulae* protease have yet to be reported, because of the enzyme characteristics similar to *P. gingivalis*, some conclusions can be drawn.

Bacterial protease activities were previously shown to have negligible differences among *P. gingivalis* strains in cell extracts and culture supernatants<sup>18)</sup>, consistent with the present findings of protease activities of *P. gulae* strains (Fig. 1, Table 2 and 3). On the other hand, *P. gingivalis* strain H66 was found to have negligible gingipain protease activities in cell extract samples, while those were remarkably increased in culture supernatant samples, in contrast to *P. gingivalis* strains ATCC33277 and W50, which showed protease activities in cell extracts<sup>38)</sup>. API ZYM systems has been




**Fig. 3. Effects of TLCK and leupeptin on *P. gulae* proteolytic activity.**

Quantification of protease activity of *P. gulae* ATCC 51700 ( $5 \times 10^7$  cfu) and *P. gingivalis* ATCC 33277 ( $5 \times 10^7$  cfu) with TLCK and/or leupeptin. A: TLCK. B: leupeptin. C: TLCK + leupeptin. Time course studies indicated for 0-4h, and dose response evaluations indicated at 100  $\mu$ M. Enzyme activities were recorded with a microplate reader and are expressed as arbitrary units. Data are shown as the mean  $\pm$  SD of three independent experiments. \*\*  $P < 0.01$  and \*  $P < 0.05$  by ANOVA with Tukey multiple-comparison test (compared with 0 h). ††  $P < 0.01$  and †  $P < 0.05$  by ANOVA with Tukey multiple-comparison test (compared with 1 h).

reportedly used to charently recognized in regard to taxonomic treatments<sup>1,24,41</sup>). Among them, three speciacterize the enzymatic profiles of the genus *Porphyromonas*. These bacteria are listed the 18 enzymes that currens, *P. gingivalis*, *P. macacae*, and *P. loveana*, have been reported to possess both alkaline phosphatase and trypsin activities<sup>8,43,46</sup>). On the other hand, other *Porphyromonas* species, such as *P. asaccharolytica*, *P. bennonis* sp, *P. canoris*, *P. cangingivalis*, *P. crevioricanis*, *P. endodontalis*, *P. leveii*, *P. somerae*, and *P. uenonis*, have alkaline phosphatase but not trypsin activities<sup>5,6,8,9,46,47</sup>). In this study, all tested *P. gulae* strains produced both alkaline phosphatase and trypsin activities (Table 1 and 2), suggesting that only four *Porphyromonas* species are phenotypically positive for trypsin and alkaline phosphatase activity. In addition, this is consistent

with reports showing that phylogenetic analysis findings presented indicating that *P. gingivalis*, *P. gulae* and *P. loveana* are relatively closely related<sup>1</sup>), though this issue requires additional study.

Protease inhibitors are known to be useful tools for identification and classification of several different bacterial proteases, including *Escherichia coli* OmpT, *Vibrio parahaemolyticus* protease A, *Yersinia pestis* YopT, and *Bacillus halodurans* keratinase<sup>26,29,39,44</sup>). Our results showed that *P. gulae* protease activity was inhibited by serine/cysteine protease inhibitors, such as antipain, PMSF, TLCK, and leupeptin, while the cysteine protease inhibitor E64 did not cause inhibition (Fig. 2 and 3, and Table 3), nor did metalloprotease, an aspartic protease inhibitor, or calpain (Table 3). Thus, those findings indicate




**Fig. 4. Effects of gingipains inhibitors on *P. gulate* proteolytic activity.**

Quantification of protease activity of *P. gulate* ATCC 51700 ( $5 \times 10^7$  cfu) with KYT-1 and KYT-36 gingipain inhibitors (100  $\mu$ M). Enzyme activities were recorded with a microplate reader and are expressed as arbitrary units. Data are shown as the mean  $\pm$  SD of three independent experiments and were analyzed with ANOVA with Tukey multiple-comparison test.

the possibility that *P. gulate* expresses trypsin-like serine proteases based on their inhibition profile, while serine proteases, such as AEBSF and aprotinin, did not work. The inhibitory effects of antipain and PMSF on proteolysis by bacteria indicate the presence of both trypsin-like serine and papain-like cysteine proteinases<sup>3,32</sup>, while AEBSF and aprotinin only inhibit the serine protease<sup>4,13</sup>. Thus, those findings suggest the possibility that *P. gulate* expresses both trypsin-like serine and papain-like cysteine proteinases based on their inhibition profile. Together, these findings suggest that both inhibitors may be unusual regarding inhibition of *P. gulate* proteolytic activity.

*P. gulate* ATCC51700 reportedly possesses *P. gingivalis* gingipain-related genes, including *rgpA*, *rgpB*, and *kgp*<sup>30</sup>. The matrix metalloproteinase 9 (MMP9) proenzyme (92KD) is activated by cleavage on the C-terminal side of arginine and lysine by trypsin, and a *P. gingivalis* gingipain, resulting in generation of active mature MMP 9 (82KD)<sup>11,20</sup>. In the present study, *P. gulate* protease exhibited arginine and lysine proteolytic activities in zymography findings, the same as *P. gingivalis*



**Fig. 5. Effects of protease inhibitors on *P. gulate* proteolytic activity.**

Following growth of SAS cells, *P. gulate* and *P. gingivalis* strains (MOI=1) were added to the culture supernatant and incubated for 24 h. proMMP9 activation was examined using gelatin zymography. No infection medium was used as a control.

(Fig. 5). These results suggest that *P. gulate* protease cleave different positions in arginine and lysine residues. Additionally, there is a possibility that Rgp and/or Kgp-like proteases of *P. gulate* are released into the extracellular environment. Those proteases have not yet been purified and their structures not observed, thus these issues require additional study. Previous study has noted that KYT-1 and KYT-36, *P. gingivalis* gingipain-specific inhibitors, were designed and synthesized using a series of small peptide analogs containing arginine at the P1 position and lysine at the P2 position<sup>22</sup>. *P. gulate* protease activity was not inhibited by KYT-1 or KYT-36, suggesting the protein structures of protease differ between *P. gulate* and *P. gingivalis*.

In conclusion, all tested *P. gulate* strains as well as *P. gingivalis* have trypsin-like protease activities. However, our results suggest that the structure of *P. gulate* protease may be different from those of *P. gingivalis* among genus Porphyromonas.

#### Acknowledgments

This research was supported by grants-in-aid for Scientific Research (20K09918 to H.I., 19K19266 to S.Y., and 20H03897 to M. M.K.) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.



**Table 1.** Enzymatic characterization of living cells of *P. gulae* strains using API ZYM® system

	<i>P. gingivalis</i> ATCC 33277	<i>P. gulae</i> ATCC 51700	<i>P. gulae</i> D040	<i>P. gulae</i> D044	<i>P. gulae</i> D049	<i>P. gulae</i> D066	<i>P. gulae</i> D077
Alkaline phosphatase	+	+	+	+	+	+	+
Esterase (C4)	-	-	-	-	-	-	-
Esterase-lipase (C8)	-	-	-	-	-	-	-
Lipase (C14)	-	-	-	-	-	-	-
Leucine arylamidase	-	-	-	-	-	-	-
Valine arylamidase	-	-	-	-	-	-	-
Cystine arylamidase	-	-	-	-	-	-	-
Trypsin	+	+	+	+	+	+	+
α-chymotrypsin	-	-	-	-	-	-	-
Acid phosphatase	-	-	-	-	-	-	-
Naphthol-AS-BI-phosphohydrolase	-	-	-	-	-	-	-
α-galactosidase	-	-	-	-	-	-	-
β-galactosidase	-	-	-	-	-	-	-
β-glucuronidase	-	-	-	-	-	-	-
α-glucosidase	-	-	-	-	-	-	-
β-glucosidase	-	-	-	-	-	-	-
N-acetyl-β-D-glucosaminidase	-	-	-	-	-	-	-
α-mannosidase	-	-	-	-	-	-	-
α-fucosidase	-	-	-	-	-	-	-
Negative control	-	-	-	-	-	-	-

**Table 2.** Enzymatic characterization of culture supernatants of *P. gulae* strains using API ZYM® system

	<i>P. gingivalis</i> ATCC 33277	<i>P. gulae</i> ATCC 51700	<i>P. gulae</i> D040	<i>P. gulae</i> D044	<i>P. gulae</i> D049	<i>P. gulae</i> D066	<i>P. gulae</i> D077
Alkaline phosphatase	+	+	+	+	+	+	+
Esterase (C4)	-	-	-	-	-	-	-
Esterase-lipase (C8)	-	-	-	-	-	-	-
Lipase (C14)	-	-	-	-	-	-	-
Leucine arylamidase	-	-	-	-	-	-	-
Valine arylamidase	-	-	-	-	-	-	-
Cystine arylamidase	-	-	-	-	-	-	-
Trypsin	+	+	+	+	+	+	+
α-chymotrypsin	-	-	-	-	-	-	-
Acid phosphatase	-	-	-	-	-	-	-
Naphthol-AS-BI-phosphohydrolase	-	-	-	-	-	-	-
α-galactosidase	-	-	-	-	-	-	-
β-galactosidase	-	-	-	-	-	-	-
β-glucuronidase	-	-	-	-	-	-	-
α-glucosidase	-	-	-	-	-	-	-
β-glucosidase	-	-	-	-	-	-	-
N-acetyl-β-D-glucosaminidase	-	-	-	-	-	-	-
α-mannosidase	-	-	-	-	-	-	-
α-fucosidase	-	-	-	-	-	-	-
Negative control	-	-	-	-	-	-	-

**Table 3.** Potency of inhibitors against *P. gulae* protease activity

Inhibitors	Target		Inhibition of Protease activity (%)	References
	Groups	Specificity		
AEBSF	Serine protease	Chymotrypsin, kallikrein, plasmin, thrombin, and trypsin	2.62 ± 6.71	4)
ALLN	Cysteine protease	Calpain	1.61 ± 2.74	37)
Antipain	Serine protease	Trypsin	17.54 ± 3.83	32)
	Cysteine protease	Papain, cathepsin A and B	1.75 ± 4.04	
Aprotinin	Serine protease	Chymotrypsin, kallikrein, plasmin, thrombin, and trypsin	0.37 ± 6.97	13)
Bestatin	Metalloprotease	Aminopeptidase N	1.53 ± 1.08	31)
Chymostatin	Serine protease	α, β, γ, δ Chymotrypsin	1.52 ± 1.04	25)
E64	Cysteine protease	Papain, cathepsin A and L	0.68 ± 3.75	3)
EDTA	Metalloprotease	Gelatinase	1.61 ± 2.74	33)
Pepstatin	Aspartic protease	Cathepsin D, chymosin, pepsin, and rennin	3.99 ± 5.10	3)
Phosphoramidon	Metalloprotease	Thermolysin, collagenase, and endothelin converting enzyme	3.09 ± 3.56	27)
PMSF	Serine protease	Chymotrypsin, thrombin, and trypsin	38.27 ± 3.32	3)
	Cysteine protease	Papain		

## References

- Bird PS, Trott DJ, Mikkelsen D, Milinovich GJ, Hillman KM, Burrell PC, Blackall LL. *Porphyromonas loveana* sp. nov., isolated from the oral cavity of Australian marsupials. *Int J Syst Evol Microbiol* 66, 3771-3778, 2016.
- Borsanelli AC, Gaetti-Jardim E Jr, Schweitzer CM, Viora L, Busin V, Riggio MP, Dutra IS. Black-pigmented anaerobic bacteria associated with ovine periodontitis. *Vet Microbiol* 203, 271-274, 2017.
- Carraro M, Zennaro C, Artero M, Candiano G, Ghiggeri GM, Musante L, Sirch C, Bruschi M, Faccini L. The effect of proteinase inhibitors on glomerular albumin permeability induced in vitro by serum from patients with idiopathic focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 19, 1969-1975, 2004.
- Citron DM, Gerardo SH, Claros MC, Abrahamian F, Talan D, Goldstein EJ. Frequency of isolation of *Porphyromonas* species from infected dog and cat bite wounds in humans and their characterization by biochemical tests and arbitrarily primed-polymerase chain reaction fingerprinting. *Clin Infect Dis* 23 Suppl 1, S78-S82, 1996.
- Citron M, Diehl TS, Capell A, Haass C, Teplow DB, Selkoe DJ. Inhibition of amyloid beta-protein production in neural cells by the serine protease inhibitor AEBSF. *Neuron* 17, 171-179, 1996.
- Collins MD, Love DN, Karjalainen J, Kanervo A, Forsblom B, Willems A, Stubbs S, Sarkiala E, Bailey GD, Wigney DI, Jousimies-Somer H. Phylogenetic analysis of members of the genus *Porphyromonas* and description of *Porphyromonas cangingivalis* sp. nov. and *Porphyromonas cansulci* sp. nov. *Int J Syst Bacteriol* 44, 674-679, 1994.
- da Silva RR. Bacterial and fungal proteolytic enzymes: production, catalysis and potential applications. *Appl Biochem Biotechnol* 183, 1-19, 2017.
- Dahlén G, Charalampakis G, Abrahamsson I, Bengtsson L, Falsen E. Predominant bacterial species in subgingival plaque in dogs. *J Periodontal Res* 47, 354-364, 2012.

- 9) Dahlén G, Gmür R, Yoshino T. Phenotypes, serotypes and antibiotic susceptibility of Swedish *Porphyromonas gingivalis* isolates from periodontitis and periodontal abscesses. *Oral Microbiol Immunol* 22, 80-86, 2007.
- 10) Doron L, Copenhagen-Glazer S, Ibrahim Y, Eini A, Naor R, Rosen G, Bachrach G. Identification and characterization of fusolisins, the *Fusobacterium nucleatum* autotransporter serine protease. *PLoS One* 9, e111329, 2014.
- 11) Duncan ME, Richardson JP, Murray GI, Melvin WT, Fothergill JE. Human matrix metalloproteinase-9: activation by limited trypsin treatment and generation of monoclonal antibodies specific for the activated form. *Eur J Biochem* 258, 37-43, 1998.
- 12) Fournier D, Mouton C, Lapierre P, Kato T, Okuda K, Ménard C. *Porphyromonas gulae* sp. nov., an anaerobic, gram-negative coccobacillus from the gingival sulcus of various animal hosts. *Int J Syst Evol Microbiol* 51, 1179-1189, 2001.
- 13) Fritz H, Wunderer G. Biochemistry and applications of aprotinin, the kallikrein inhibitor from bovine organs. *Arzneimittelforschung* 33, 479-494, 1983.
- 14) Gavioli R, Vertuani S, Masucci MG. Proteasome inhibitors reconstitute the presentation of cytotoxic T-cell epitopes in Epstein-Barr virus-associated tumors. *Int J Cancer* 101, 532-538, 2002.
- 15) Guo Y, Nguyen KA, Potempa J. Dichotomy of gingipains action as virulence factors: from cleaving substrates with the precision of a surgeon's knife to a meat chopper-like brutal degradation of proteins. *Periodontol* 2000 54, 15-44, 2010.
- 16) Holt SC, Ebersole JL. *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: the "red complex", a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol* 2000 38, 72-122, 2005.
- 17) Inaba H, Amano A. Roles of oral bacteria in cardiovascular diseases—from molecular mechanisms to clinical cases: Implication of periodontal diseases in development of systemic diseases. *J Pharmacol Sci* 113, 103-109, 2010.
- 18) Inaba H, Nakano K, Kato T, Nomura R, Kawai S, Kuboniwa M, Ishihara K, Ooshima T, Amano A. Heterogenic virulence and related factors among clinical isolates of *Porphyromonas gingivalis* with type II fimbriae. *Oral Microbiol Immunol* 23, 29-35, 2008.
- 19) Inaba H, Nomura R, Kato Y, Takeuchi H, Amano A, Asai F, Nakano K, Lamont RJ, Matsumoto-Nakano M. Adhesion and invasion of gingival epithelial cells by *Porphyromonas gulae*. *PLoS One* 14, e0213309, 2019.
- 20) Inaba H, Sugita H, Kuboniwa M, Iwai S, Hamada M, Noda T, Morisaki I, Lamont RJ, Amano A. *Porphyromonas gingivalis* promotes invasion of oral squamous cell carcinoma through induction of proMMP9 and its activation. *Cell Microbiol* 16, 131-145, 2014.
- 21) Inaba H, Tagashira M, Kanda T, Ohno T, Kawai S, Amano A. Apple- and hop-polyphenols protect periodontal ligament cells stimulated with enamel matrix derivative from *Porphyromonas gingivalis*. *J Periodontol* 76, 2223-2229, 2005.
- 22) Kadowaki T, Baba A, Abe N, Takii R, Hashimoto M, Tsukuba T, Okazaki S, Suda Y, Asao T, Yamamoto K. Suppression of pathogenicity of *Porphyromonas gingivalis* by newly developed gingipain inhibitors. *Mol Pharmacol* 66, 1599-1606, 2004.
- 23) Kato Y, Shirai M, Murakami M, Mizusawa T, Hagimoto A, Wada K, Nomura R, Nakano K, Ooshima T, Asai F. Molecular detection of human periodontal pathogens in oral swab specimens from dogs in Japan. *J Vet Dent* 28, 84-89, 2011.
- 24) Kawamura Y, Kuwabara S, Kania SA, Kato H, Hamagishi M, Fujiwara N, Sato T, Tomida J, Tanaka K, Bemis DA. *Porphyromonas pogonae* sp. nov., an anaerobic but low concentration oxygen adapted coccobacillus isolated from

- lizards (*Pogona vitticeps*) or human clinical specimens, and emended description of the genus *Porphyromonas* Shah and Collins 1988. *Syst Appl Microbiol* 38, 104-109, 2015.
- 25) Klegeris A, McGeer PL. Chymotrypsin-like proteases contribute to human monocytic THP-1 cell as well as human microglial neurotoxicity. *Glia* 51, 56-64, 2005.
- 26) Kramer RA, Dekker N, Egmond MR. Identification of active site serine and histidine residues in *Escherichia coli* outer membrane protease OmpT. *FEBS Lett* 468, 220-224, 2000.
- 27) Kumar GK. Peptidases of the peripheral chemoreceptors: biochemical, immunological, in vitro hydrolytic studies and electron microscopic analysis of neutral endopeptidase-like activity of the carotid body. *Brain Res* 748, 39-50, 1997.
- 28) Kyllar M, Witter K. Prevalence of dental disorders in pet dogs. *Vet Med (Praha)* 50, 496-505, 2005.
- 29) Lee CY, Cheng MF, Yu MS, Pan MJ. Purification and characterization of a putative virulence factor, serine protease, from *Vibrio parahaemolyticus*. *FEMS Microbiol Lett* 209, 31-37, 2002.
- 30) Lenzo JC, O'Brien-Simpson NM, Orth RK, Mitchell HL, Dashper SG, Reynolds EC. *Porphyromonas gulae* has virulence and immunological characteristics similar to those of the human periodontal pathogen *Porphyromonas gingivalis*. *Infect Immun* 84, 2575-2585, 2016.
- 31) Lkhagvaa B, Tani K, Sato K, Toyoda Y, Suzuka C, Sone S. Bestatin, an inhibitor for aminopeptidases, modulates the production of cytokines and chemokines by activated monocytes and macrophages. *Cytokine* 44, 386-391, 2008.
- 32) McGowan EB, Shafiq SA, Stracher A. Delayed degeneration of dystrophic and normal muscle cell cultures treated with pepstatin, leupeptin, and antipain. *Exp Neurol* 50, 649-657, 1976.
- 33) Métayer S, Dacheux F, Dacheux JL, Gatti JL. Comparison, characterization, and identification of proteases and protease inhibitors in epididymal fluids of domestic mammals. Matrix metalloproteinases are major fluid gelatinases. *Biol Reprod* 66, 1219-1229, 2002.
- 34) Mikkelsen D, Milinovich GJ, Burrell PC, Huynh SC, Pettett LM, Blackall LL, Trott DJ, Bird PS. Phylogenetic analysis of *Porphyromonas* species isolated from the oral cavity of Australian marsupials. *Environ Microbiol* 10, 2425-2432, 2008.
- 35) Niemiec BA. Periodontal disease. *Top Companion Anim Med.* 23, 72-80, 2008.
- 36) Patel YM, Lane MD. Mitotic clonal expansion during preadipocyte differentiation: calpain-mediated turnover of p27. *J Biol Chem* 275, 17653-17660, 2000.
- 37) Pike R, McGraw W, Potempa J, Travis J. Lysine- and arginine-specific proteinases from *Porphyromonas gingivalis*. Isolation, characterization, and evidence for the existence of complexes with hemagglutinins. *J Biol Chem* 269, 406-411, 1994.
- 38) Potempa J, Pike R, Travis J. The multiple forms of trypsin-like activity present in various strains of *Porphyromonas gingivalis* are due to the presence of either Arg-gingipain or Lys-gingipain. *Infect Immun* 63, 1176-1182, 1995.
- 39) Prakash P, Jayalakshmi SK, Sreeramulu K. Production of keratinase by free and immobilized cells of *Bacillus halodurans* strain PPKS-2: partial characterization and its application in feather degradation and dehairing of the goat skin. *Appl Biochem Biotechnol* 160, 1909-1920, 2010.
- 40) Sabotič J, Kos J. Microbial and fungal protease inhibitors—current and potential applications. *Appl Microbiol Biotechnol* 93, 1351-1375, 2012.
- 41) Sakamoto M, Li D, Shibata Y, Takeshita T, Yamashita Y, Ohkuma M. *Porphyromonas pasteri* sp. nov., isolated from human saliva. *Int J Syst Evol Microbiol* 65, 2511-2515, 2015.
- 42) Senhorinho GNA, Nakano V, Liu C, Song Y, Finegold SM, Avila-Campos MJ. Detection

- of *Porphyromonas gulae* from subgingival biofilms of dogs with and without periodontitis. *Anaerobe* 17, 257-258, 2011.
- 43) Senhorinho GNA, Nakano V, Liu C, Song Y, Finegold SM, Avila-Campos MJ. Occurrence and antimicrobial susceptibility of *Porphyromonas* spp. and *Fusobacterium* spp. in dogs with and without periodontitis. *Anaerobe* 18, 381-385, 2012.
- 44) Shao F, Merritt PM, Bao Z, Innes RW, Dixon JE. A *Yersinia* effector and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning in bacterial pathogenesis. *Cell* 109, 575-588, 2002.
- 45) Silva LB, Reis AP, Pereira EJ, Oliveira MG, Guedes RN. Partial purification and characterization of trypsin-like proteinases from insecticide-resistant and -susceptible strains of the maize weevil, *Sitophilus zeamais*. *Comp Biochem Physiol B Biochem Mol Biol* 155, 12-19, 2010.
- 46) Summanen PH, Durmaz B, Väisänen ML, Liu C, Molitoris D, Eerola E, Helander IM, Finegold SM. *Porphyromonas somerae* sp. nov., a pathogen isolated from humans and distinct from *Porphyromonas levii*. *J Clin Microbiol* 43, 4455-4459, 2005.
- 47) Summanen PH, Lawson PA, Finegold SM. *Porphyromonas bennonis* sp. nov., isolated from human clinical specimens. *Int J Syst Evol Microbiol* 59, 1727-1732, 2009.
- 48) Urmi AS, Inaba H, Nomura R, Yoshida S, Ohara N, Asai F, Nakano K, Matsumoto-Nakano M. Roles of *Porphyromonas gulae* proteases in bacterial and host cell biology. *Cell Microbiol* 23, e13312, 2021.
- 49) Yamasaki Y, Nomura R, Nakano K, Inaba H, Kuboniwa M, Hirai N, Shirai M, Kato Y, Murakami M, Naka S, Iwai S, Matsumoto-Nakano M, Ooshima T, Amano A, Asai F. Distribution and molecular characterization of *Porphyromonas gulae* carrying a new *fimA* genotype. *Vet Microbiol* 161, 196-205, 2012.