

Title	Outcomes of laparoscopic total gastrectomy in elderly patients : a propensity score matching analysis
Author(s)	Ebihara, Yuma; Kurashima, Yo; Watanabe, Yusuke; Tanaka, Kimitaka; Matsui, Aya; Nakanishi, Yoshitsugu; Asano, Toshimichi; Noji, Takehiro; Nakamura, Toru; Murakami, Soichi; Tsuchikawa, Takahiro; Okamura, Keisuke; Murakami, Yoshihiro; Murakawa, Katsuhiko; Nakamura, Fumitaka; Morita, Takayuki; Okushiba, Shunichi; Shichinohe, Toshiaki; Hirano, Satoshi
Citation	Langenbeck's archives of surgery, 407(4), 1461-1469 https://doi.org/10.1007/s00423-022-02447-2
Issue Date	2022-06-01
Doc URL	http://hdl.handle.net/2115/89786
Rights	This is a post-peer-review, pre-copyedit version of an article published in Langenbeck's archives of surgery. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00423-022-02447-2
Туре	article (author version)
File Information	LAS s00423-022-02447-2.pdf



Instructions for use

Title: Outcomes of laparoscopic total gastrectomy in elderly patients: a propensity score matching
 analysis

	h
	2
	h
٩	,

4	Yuma Ebihara, PhD ^{1,2} , Yo Kurashima, PhD ¹ , Yusuke Watanabe, PhD ¹ , Kimitaka Tanaka, PhD ¹ , Aya
5	Matsui, PhD ¹ , Yoshitsugu Nakanishi, PhD ¹ , Toshimichi Asano, PhD ¹ , Takehiro Noji, PhD ¹ , Toru
6	Nakamura, PhD ¹ , Soichi Murakami, PhD ¹ , Takahiro Tsuchikawa, PhD ¹ , Keisuke Okamura, PhD ¹ ,
7	Yoshihiro Murakami, PhD ³ , Katsuhiko Murakawa, PhD ⁴ , Fumitaka Nakamura, PhD ⁵ , Takayuki
8	Morita, PhD ⁶ , Shunichi Okushiba, PhD ⁷ , Toshiaki Shichinohe, PhD ¹ , Satoshi Hirano, PhD ¹
9	
10	¹ Department of Gastroenterological Surgery II, Faculty of Medicine, Hokkaido University, Sapporo,
11	Japan;
12	² Division of Minimally Invasive Surgery, Hokkaido University Hospital, Sapporo, Japan
13	³ Department of Surgery, Asahikawa City Hospital, Asahikawa, Japan
14	⁴ Department of Surgery, Obihiro-Kosei General Hospital, Obihiro, Japan
15	⁵ Department of Surgery, Teine Keijinkai Hospital, Sapporo, Japan
16	⁶ Department of Surgery, Hokkaido Gastroenterology Hospital, Sapporo, Japan
17	⁷ Department of Surgery, Tonan Hospital, Sapporo, Japan

1	Corresponding author: Yuma Ebihara, Department of Gastroenterological Surgery II, Hokkaido
2	University Faculty of Medicine, Kita 15 Nishi 7, Kita-ku, Sapporo City, 060-8638, Japan
3	Tel: +81 11 706 7714 Fax: +81 11 706 7158
4	Email: yuma-ebi@wc4.so-net.ne.jp
5	
6	Running head: Surgical outcomes of LTG in elderly patients aged over 80 years
7	
8	Abstract
9	Purpose
10	This study evaluated the short-term outcomes and prognosis after laparoscopic total gastrectomy
11	(LTG) in elderly patients aged \geq 80 years in a multicenter retrospective cohort study using propensity
12	score matching.
13	Methods
14	We retrospectively enrolled 440 patients who underwent curative LTG for gastric cancer at six
15	institutions between January 2004 and December 2018. Patients were categorized into an elderly
16	patient group (EG; age ≥80 years) and non-elderly patient group (non-EG; age <80 years). Patients
17	were matched using the following propensity score covariates: sex, body mass index, American
18	Society of Anesthesiologists physical status, extent of lymph node dissection, and Japanese

Classification of Gastric Carcinoma stage. Short-term outcomes and prognoses were compared.

3	We identified 37 propensity score-matched pairs. The median operative time was significantly
4	shorter and postoperative stay was longer in the EG. In terms of postoperative outcomes, the rates of
5	all complications were comparable. The median follow-up period of the EG and non-EG was 11.5 (1-
6	106.4) months and 35.7 (1-110.0) months, respectively; there were significant differences in 5-year
7	overall survival between the two groups (EG, 58.5% vs. non-EG, 91.5%; P=0.031). However, there
8	were no significant differences in 5-year disease-specific survival (EG, 62.1% vs. non-EG, 91.5%;
9	P=0.068) or 5-year disease-free survival (EG, 52.9% vs. non-EG, 60.8%; P=0.132).
10	Conclusions
11	LTG seems to be safe and feasible in elderly patients. LTG had a limited effect on morbidity, disease
12	recurrence, and survival in elderly patients. Therefore, age should not prevent elderly patients from
13	benefitting from LTG.
14	
15	Keywords: laparoscopic total gastrectomy, gastric cancer, elderly patients, multicenter retrospective
16	study
17	

18 Declarations

1	Funding:	Not app	olicable
	0		

- 2 Conflicts of interest/Competing interests: Not applicable
- 3 Availability of data and material (data transparency): The datasets during and/or analysed during the
- 4 current study available from the corresponding author on reasonable request.
- 5 Code availability: Not applicable
- 6 Ethics approval: Hokkaido University Hospital Institutional Review Board approved the data
- 7 collection and analysis (No. 016-0151).
- 8 Consent to participate (include appropriate statements): Not applicable
- 9 Consent for publication (include appropriate statements): Not applicable
- 10 Author Contributions:
- 11 Study conception and design: Yuma Ebihara and Yo Kurashima
- 12 Acquisition of data: Yuma Ebihara
- 13 Analysis and interpretation of data: Yuma Ebihara and Yo Kurashima
- 14 Drafting of manuscript: Yuma Ebihara.
- 15 Critical revision of the manuscript: Yuma Ebihara, Yo Kurashima, Yusuke Watanabe, Kimitaka Tanaka,
- 16 Aya Matsui, Yoshitsugu Nakanishi, Toshimichi Asano, Takehiro Noji, Toru Nakamura, Soichi
- 17 Murakami, Takahiro Tsuchikawa, Keisuke Okamura, Yoshihiro Murakami, Katsuhiko Murakawa,
- 18 Fumitaka Nakamura, Takayuki Morita, Shunichi Okushiba, Toshiaki Shichinohe, and Satoshi Hirano.

1 Introduction

2	The population of elderly people, especially those aged over 80 years in Japan, has been
3	dramatically increasing in number worldwide. The proportion of elderly patients with gastric cancer
4	(GC) undergoing surgery has also been increasing along with an increasing Helicobacter pylori
5	infection rate among the elderly [1]. Since the first laparoscopy-assisted distal gastrectomy for early
6	GC was performed in 1991 [2], the development of dedicated instruments and surgical techniques has
7	led to the use of laparoscopic distal gastrectomy (LDG) to treat GC [3]. However, laparoscopic total
8	gastrectomy (LTG) requires a high degree of skill in performing gastrectomy with systematic
9	lymphadenectomy, as well as postresection reconstruction; indeed, these procedures are difficult even
10	for experienced laparoscopic surgeons [4]. LTG has many advantages; however, it is associated with
11	a high incidence of postoperative complications (10-40%) [5-7]. Therefore, surgeons have some
12	difficulties in making decisions regarding the performance of LTG in elderly patients. Some authors
13	have reported the safety and effectiveness of LDG in elderly patients [8, 9]. However, few studies of
14	LTG in elderly patients, especially those aged \geq 80 years , have been reported. Therefore, it is necessary
15	to study the safety, effectiveness, and prognosis of LTG in elderly patients.
16	In this study, we aimed to clarify surgical outcomes and prognosis in the elderly patient group (EG;
17	age ≥80 years) who underwent LTG with curative intent, compared with a non-elderly patient group
18	(non-EG; age <80 years). A propensity score matching (PSM) method was used to compare the short-

term outcomes and prognosis between the EG and non-EG of LTG for GC in a multicenter setting.
 This is the first multicenter retrospective study to compare EG and non-EG in LTG using the PSM
 method.

4 Material and Methods

5 Patients

6 We retrospectively reviewed patients who underwent LTG for GC at six institutions (Hokkaido 7 University Hospital, Teine Keijinkai Hospital, Obihiro-Kosei General Hospital, Hokkaido 8 Gastroenterology Hospital, Tonan Hospital, and Asahikawa City Hospital) between January 2004 and 9 December 2018. All patients who underwent curative LTG were included in the analysis. All patients 10 were diagnosed with GC using endoscopy, computed tomography (CT), or endoscopic ultrasound. The 11 Japanese Classification of Gastric Carcinoma (JCGC) was used for tumor staging [10]. The primary 12 indication for LTG was stage I GC based on the Japanese Society of Endoscopic Surgery (JSES) 13 guidelines [11]; however, over time, we expanded the indication to include cases of advanced GC that 14 could be curatively resected. 15

16 Data collection

17 Clinicopathological data, including age, sex, body mass index (BMI), American Society of
18 Anesthesiologists physical status (ASA-PS), clinical stage, combined resection of other organs, lymph

1	node dissection, time to recurrence, and prognosis were collected. Surgical outcomes, including
2	operative time, estimated blood loss, postoperative complications, and length of postoperative hospital
3	stay were recorded.
4	Patients were categorized into an elderly patient group (EG; age ≥80 years) and non-elderly patient
5	group (non-EG; age <80 years). All patients provided informed consent, and the Hokkaido University
6	Hospital Institutional Review Board approved the data collection and analysis (No. 016-0151). This
7	study was performed in accordance with the principles of the Declaration of Helsinki.
8	
9	Surgical procedure
10	The gastric procedure type (resection and reconstruction) was determined based on the experience
11	and preference of a surgeon accredited through the endoscopic surgical skill qualification system of
12	the JSES [11]. In cases where the operating surgeon did not possess this qualification, a qualified
13	surgeon supervised the surgery. The extent of lymph node dissection was determined according to the
14	JGCA guidelines [10]. Patients who underwent D2 lymph node dissection and patients who underwent
15	D2-No.10 lymph node dissection were included in D1+. Patients were divided into three groups based
16	on the Clavien–Dindo (CD) postoperative complication classification grade [12, 13].
17	

18 Postoperative follow-up

1	All patients were observed every 3 months after surgery. Hematological analysis (including tumor
2	marker analysis for carcinoembryonic antigen and carbohydrate antigen 19-9) was performed at each
3	visit. Abdominal computed tomography scans were performed every 6 months or when clinical
4	recurrence was suspected. Gastrointestinal endoscopy was performed at 1, 3, and 5 years
5	postoperatively. Based on this surveillance, data on the 5-year overall survival (OS), disease-specific
6	survival (DSS), and disease-free survival (DFS) were collected. OS was defined as the time from
7	surgery to death for any reason or interruption of follow-up. DSS was defined as the time from surgery
8	to death from GC, including operative mortality or interruption of follow-up. DFS was defined as the
9	time from surgery to death from GC, the first recurrence of GC, or the interruption of follow-up.
10	
11	Statistical analysis
12	PSM was performed using a logistic regression model to mitigate the selection bias in the present
13	study. The parameters used for PSM were age, sex, BMI, American Society of Anesthesiologists
14	physical status, splenectomy, clinical stage, surgical method, and lymph node dissection. We matched
15	the logit of the propensity score within 0.2 standard deviations of the value based on the

- 16 recommendations by Austin [14]. Categorical variables were analyzed using the chi-squared test
- 17 before propensity score matching and the McNemar and Wilcoxon signed-rank tests after PSM.
- 18 Continuous variables were examined using the unpaired t-test before PSM and the paired t-test or

1	Wilcoxon signed-rank test after propensity score matching. Survival curves were constructed using
2	the Kaplan-Meier method and compared using the log-rank test. Statistical significance was set at
3	P<0.05. Statistical analysis was performed using the JMP [®] 15 software (SAS Institute Inc., Cary, NC,
4	USA).
5	
6	Results
7	Patients' backgrounds
8	Table 1 shows the clinical characteristics and surgical outcomes of the study population. A total of
9	440 patients were included, comprising 305 men (69.3%) and 135 women (30.7%), with a median age
10	of 67 years (range 25–88) and median BMI of 22.9 kg/m ² (range 13.6-38.9). The American Society of
11	Anesthesiologist's physical status (ASA-PS) was ≥II in 318 patients (72.3%), clinical JCGC stage was
12	≥II in 135 patients (30.7%), D2 lymphadenectomy was performed in 68 patients (15.5%), the median
13	operation time was 329 (range, 123–762) minutes, the median operative blood loss was 50 (range, 0–
14	1940) mL, postoperative complications CD ≥IIIa occurred in 66 patients (15.0%), and the median
15	postoperative hospital stay was 14 (range, 6-210) days. After applying our exclusion criteria
16	(neoadjuvant therapy, cStage IV, resection of other organs, and total number of harvested lymph nodes
17	\leq 15), 395 patients were included in the subgroup we evaluated for PSM. In this study, for accurate
18	prognostic analysis, we included more than 16 lymph nodes dissected using the exclusion criteria [15].

1	In total, 37 patients in the elderly group (EG; age \geq 80 years) were individually matched to 37 patients
2	in the non-elderly group (non-EG; age <80 years) (Fig. 1). The clinicopathological characteristics of
3	the 395 patients who underwent curative LTG and 74 propensity score-matched patients are shown in
4	Table 2. In the propensity score-matched patients, as determined by the study design, sex, BMI, ASA-
5	PS, extent of lymph node dissection, and clinical JCGC stage distributions between the EG and non-
6	EG groups were comparable.
7	Surgical outcomes
8	The surgical outcomes and postoperative complications of the 395 patients who underwent curative
9	LTG and propensity score-matched patients are shown in Table 3. In the propensity score-matched
10	group of patients, there was no significant difference in blood loss, number of harvested lymph nodes
11	and the method of esophagojejunostomy. Furthermore, there was no difference in the rate of
12	postoperative complications between the two groups (Table 4). The median operative time was
13	significantly shorter in the EG than in the non-EG (EG vs. non-EG; 260 minutes [range 171–525] vs.
14	335 minutes [range 172–553], respectively, P= 0.003). The median postoperative hospital stay was
15	significantly longer in the EG than in the non-EG (EG vs. non-EG; 16 days [range 8-82] vs. 12 days
16	[range 7–65], respectively, P=0.008).
17	Prognosis

18 The results of histological examinations of resected specimens were similar in both groups;

1	therefore, the pathological stages according to the JCGC were similar (P=0.726). The median follow-
2	up periods of the EG and non-EG groups were 11.5 (1-106.4 months) and 35.7 (1-110.0) months,
3	respectively. The number of patients who received adequate adjuvant chemotherapy according to the
4	JGCA guidelines [8] was significantly lower in the EG than in the non-EG (EG vs. non-EG; 4 patients
5	[21.1%] vs. 16 patients [84.2%], respectively, P=0.047). During the follow-up period, postoperative
6	recurrence was observed in eight patients (21.6%) in the EG and nine patients (24.3%) in the non-EG,
7	with no statistical significance (P=0.782). There was no difference in the recurrence site or length of
8	recurrence between the two groups (Table 5). The 5-year OS rates were 79.1% in all patients, were
9	58.5% and 91.5% in the EG and non-EG groups, respectively. The Kaplan-Meier analysis for OS
10	indicated that there was a significant difference between the two groups (P=0.031). The 5-year DSS
11	and DFS rates were 80.7% and 56.9% in all patients, and were 62.1% and 52.9% in the EG, and 91.5%
12	and 60.8% in the non-EG, respectively. The Kaplan-Meier analysis for DSS and DFS indicated that
13	there was no significant difference (P=0.068 and P=0.132, respectively) (Fig. 2). In the 5-year DSS,
14	the Wilcoxon test showed a significant difference between the two groups (P=0.031) because the
15	curves of the two groups overlapped at 12 months after surgery.
16	

17 Discussion

18 This is the first multicenter retrospective study comparing the surgical outcomes and prognosis of

1	elderly patients aged ≥80 years and <80 years who underwent LTG using PSM. This study aimed to
2	evaluate the short-term outcome and prognosis after LTG in elderly patients with gastric cancer after
3	LTG. After adjustment using PSM, we found no significant differences in the incidence of
4	postoperative complications and recurrence rates between the non-elderly and elderly patients.
5	The average age of patients with gastric cancer undergoing gastrectomy has been increasing in
6	recent years [8]. The natural life expectancy of elderly patients is obviously shorter than that of
7	younger patients, and elderly patients usually have various comorbidities, such as cardiovascular
8	disease [16, 17] and decreased respiratory function, leading to limited use of the procedure. In addition,
9	postoperative complications in elderly patients, such as delirium and sarcopenia, are also problems
10	[18]. In general, elderly patients often have physiological difficulties associated with aging, such as
11	decreased organ function, various complications, and mental imbalance. Surgeons are sometimes
12	hesitant to perform standard gastrectomy with lymphadenectomy in such patients because of the high
13	incidence of age-related morbidity and mortality. In addition, surgeons need to overcome perioperative
14	issues specific to elderly patients, such as postoperative delirium, sarcopenia, and frailty [18-20]. In
15	this study, many patients in the EG group did not receive adjuvant chemotherapy according to the
16	JGCA guidelines [10] due to a high preoperative complication rate. As a result, the time to relapse was
17	shorter. There was no difference in postoperative complications (CD \geq II) between the groups. In
18	addition, no deaths were observed in the hospital.

1	Laparoscopic surgery with minimal destruction of the body wall might be a beneficial approach for
2	the elderly because people with reduced organ function are more susceptible to surgical invasion.
3	Studies on long-term prognosis suggest an association between postoperative complications and long-
4	term survival in several malignancies, including breast cancer, colorectal cancer, and peripancreatic
5	cancer [21-24]. Furthermore, several reports have indicated that postoperative complications are
6	associated with the prognosis of patients with GC [25, 26]. In our previous multicenter retrospective
7	study on the long-term prognosis of laparoscopic surgery for GC, postoperative complications were
8	also shown to be associated with survival [27]. Therefore, ensuring the safety of LTG may be important
9	for the short- and long-term outcomes of patients with GC. The same was true in the present study for
10	the occurrence of severe complications of CD grade ≥III after LTG (EG vs. non-EG; 16.2 vs. 10.8%,
11	respectively, P=0.495). There was no difference in long-term prognosis (DSS and DFS) between the
12	two groups (P=0.068 and P=0.132, respectively), which may be due in part to the similarity of
13	complications. In order to keep the postoperative complication rate low in elderly patients, surgeons
14	must balance the complication rate with trefraioe oncological survival rate. The six affiliated hospitals
15	that participated in this study are high-volume centers in our prefecture, and each facility has a JSES-
16	certified laparoscopic surgeon who performed the procedure as a surgeon or teaching assistant in this
17	study. The difference in the median postoperative hospital stay between the two groups was attributed
18	to differences in the criteria for discharge among the affiliated hospitals. In addition, the median

1	postoperative hospital stay was 14 days, because the fast-track protocol was not applied in the early
2	stages of the introduction of the LTG.
3	However, the number of patients who received adequate adjuvant chemotherapy according to the
4	JGCA guidelines [10] was significantly lower in the EG than in the non-EG (P=0.047). The reasons
5	why adjuvant chemotherapy was not administered in the EG are as follows. Adjuvant chemotherapy
6	was not indicated due to poor postoperative general condition, and physicians refrained from adjuvant
7	chemotherapy at their discretion. However, postoperative recurrence was observed in eight patients
8	(21.6%) in the EG and nine patients (24.3%) in the non-EG, which was not significantly different
9	(P=0.782). There was no difference in the recurrence site or length of recurrence between the two
10	groups, and there was no effect of adjuvant chemotherapy on OS, DSS, or DFS (hazard ratio [HR] =
11	1.187, 95% confidence interval [CI]: 0.149–9.395; HR = 2.088, 95% CI: 0.242–17.963; and HR =
12	2.653, 95% CI: 0.936-7.519, respectively). Although prior randomized controlled studies indicated
13	that postoperative adjuvant treatment in patients who underwent D2 gastrectomy could improve the
14	5-year DFS and OS, the subgroup analyses showed that the survival benefits decreased with increasing
15	age. Furthermore, the ACTSGC study showed no statistically significant effects of postoperative
16	chemotherapy on DFS or OS in patients older than 70 years (DFS: $HR = 0.779, 95\%$ CI: 0.527–1.151;
17	OS: HR = 0.706, 95% CI: 0.490–1.017) [28]. Similar results for OS were observed in the CLASSIC
18	study for patients older than 65 years (HR = 0.70 , 95% CI: 0.4 – 1.12) [29]. These results might be due

1	to the considerably higher incidence of comorbidities, higher risk of complications, and shorter life
2	expectancy in elderly patients [30]. Nevertheless, when considering these conflicting results, whether
3	adjuvant chemotherapy should be administered to elderly patients with GC after D2 gastrectomy
4	remains a dilemma for physicians. In accordance with the consensus guidelines of the European
5	Society for Medical Oncology (ESMO), radical gastrectomy with free margins and an adequate
6	lymphadenectomy, and if indicated along with perioperative neoadjuvant chemotherapy, is the
7	standard of care in patients with advanced gastric cancer [31,32]. In Japan, there is still no evidence
8	of perioperative neoadjuvant therapy for gastric cancer. In the future, we look forward to the inclusion
9	of appropriate chemotherapy for elderly patients in the guidelines.
10	This study has several limitations. This was a retrospective, observational, and non-experimental
11	study. In the present study, most LTGs were performed by laparoscopic surgery experts as operators
12	or teaching assistants. In this study, 69.3% of the patients had stage I gastric cancer and only 15.5%
13	underwent D2 lymph node resection. In addition, 72.3% of all patients were ASA 1 or 2 and in
14	relatively good general condition, and there was a selection bias. The two study groups have
15	significantly different median postoperative follow-up periods (EG 11.5 months vs. non-EG 35.7
16	months). In this study, the number of cases was small because each institution was strict about the
17	indications for LTG. This may influence outcomes in term of OS, DSS, and DFS. A well-designed
18	randomized controlled trial is required to validate our findings. Selection bias was a limitation because

1	the cohort consisted of surgical patients for whom the surgeon decided the surgical indication
2	according to performance status, especially in EG. This study was conducted over a rather long period
3	between 2004 and 2018, which could have been associated with historical biases in terms of treatment
4	strategy and perioperative management, which might dictate the short-term and prognostic outcomes
5	after gastrectomy.
6	
7	Conclusions
8	LTG seems to be safe and feasible for elderly patients. LTG has had a limited effect on morbidity,
9	disease recurrence, and survival in elderly patients. Therefore, age should not prevent elderly patients
10	from benefitting from LTG. We believe that guidelines are needed to determine the appropriate extent
11	of lymph node dissection and adjuvant chemotherapy for elderly patients.
12	
13	Acknowledgments
14	The author is grateful to our colleagues (Dr. Hiroshi Kawase, Noriaki Kyogoku, and Mamoru
15	Miyasaka) and the interdisciplinary surgical team. We would like to thank Editage (www.editage.com)
16	for English language editing.
17	

1 References

2	1. Asaka M, Sugiyama T, Nobuta A, Kato M, Takeda H, Graham DY (2001) Atrophic gastritis
3	and intestinal metaplasia in Japan: results of a large multicenter study. Helicobacter 6:294–299. DOI:
4	10.1046/j.1523-5378.2001.00042.x
5	2. Kitano S, Iso Y, Moriyama M, Sugimachi K (1994) Laparoscopy-assisted Billroth I
6	gastrectomy. Surg Laparosc Endosc 4:146–148
7	3. Aiolfi A, Lombardo F, Matsushima K, Sozzi A, Cavalli M, Panizzo V, Bonitta G, Bona D
8	(2021) Systematic review and updated network meta-analysis of randomized controlled trials
9	comparing open, laparoscopic-assisted, and robotic distal gastrectomy for early and locally advanced
10	gastric cancer. Surgery. Sep;170(3):942-951. DOI: 10.1016/j.surg.2021.04.014.
11	4. Ebihara Y, Okushiba S, Kawarada Y, Kitashiro S, Katoh H (2013) Outcome of functional
12	end-to-end esophagojejunostomy in totally laparoscopic total gastrectomy. Langenbecks Arch Surg
13	398:475–479. DOI: 10.1007/s00423-013-1051-z
14	5. van der Wielen N, Straatman J, Daams F, Rosati R, Parise P, Weitz J, Reissfelder C, Diez Del
15	Val I, Loureiro C, Parada-González P, Pintos-Martínez E, Mateo Vallejo F, Medina Achirica C,
16	Sánchez-Pernaute A, Ruano Campos A, Bonavina L, Asti ELG, Alonso Poza A, Gilsanz C, Nilsson M,
17	Lindblad M, Gisbertz SS, van Berge Henegouwen MI, Fumagalli Romario U, De Pascale S, Akhtar
18	K, Jaap Bonjer H, Cuesta MA, van der Peet DL (2020) Open versus minimally invasive total

gastrectomy after neoadjuvant chemotherapy: results of a European randomized trial. Gastric Cancer
 24:258-271. DOI: 10.1007/s10120-020-01109-w.

- Lee SR, Kim HO, Son BH, Shin JH, Yoo CH (2014) Laparoscopic-assisted total gastrectomy
 versus open total gastrectomy for upper and middle gastric cancer in short-term and long-term
 outcomes. Surg Laparosc Endosc Percutan Tech 24:277–282. DOI: 10.1097/SLE.0b013e3182901290
- 6 7. Tsunoda S, Okabe H, Obama K, Tanaka E, Hisamori S, Kinjo Y, Sakai Y (2014) Short-term
- 7 outcomes of totally laparoscopic total gastrectomy: experience with the first consecutive 112 cases.
- 8 World J Surg 38:2662–2667. DOI: 10.1007/s00268-014-2611-2
- 9 8. Yoshida M, Koga S, Ishimaru K, Yamamoto Y, Matsuno Y, Akita S, Kuwabara J, Tanigawa
- 10 K, Watanabe Y (2017 Nov) Laparoscopy-assisted distal gastrectomy is feasible also for elderly patients
- aged 80 years and over: effectiveness and long-term prognosis. Surg Endosc 31:4431–4437. doi:
- 12 10.1007/s00464-017-5493-1. Epub 2017 Apr 4. PMID: 28378081
- 13 9. Tanaka R, Lee SW, Imai Y, Honda K, Matsuo K, Tashiro K, Kawai M, Uchiyama K (2021
- 14 May 21) Advantages of laparoscopic surgery for gastric cancer in elderly patients aged over 80 years:
- 15 A propensity score matching analysis. World J Surg. doi: 10.1007/s00268-021-06157-6 [Epub ahead
- 16 of print]. PMID: 34019135
- 17 10. Japanese Gastric Cancer Association (2011) Japanese gastric cancer treatment guidelines
- 18 2010 (ver 3). Gastric Cancer 14:113–123. DOI: 10.1007/s10120-011-0042-4

1	11. Japanese Society for Endoscopic Surgery (2014) Guidelines for the management of
2	endoscopic surgery. Japanese Society for Endoscopic Surgery, Tokyo
3	12. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new
4	proposal with evaluation in a cohort of 6336 patients and results of a survey. Ann Surg 240:205–213.
5	DOI: 10.1097/01.sla.0000133083.54934.ae
6	13. Clavien PA, Barkun J, de Oliveira ML, Vauthey JN, Dindo D, Schulick RD, de Santibañes
7	E, Pekolj J, Slankamenac K, Bassi C, Graf R, Vonlanthen R, Padbury R, Cameron JL, Makuuchi M
8	(2009) The Clavien-Dindo classification of surgical complications: five-year experience. Ann Surg
9	250:187–196. DOI: 10.1097/SLA.0b013e3181b13ca2
10	14. Austin PC (2011) Optimal caliper widths for propensity-score matching when estimating
11	differences in means and differences in proportions in observational studies. Pharm Stat 10:150–161.
12	DOI: 10.1002/pst.433
13	15. Amin MB, Greene FL, Edge SB, Compton CC, Gershenwald JE, Brookland RK, Meyer L,
14	Gress DM, Byrd DR, Winchester DP (2017) The Eighth Edition AJCC Cancer Staging Manual:
15	continuing to build a bridge from a population-based to a more "personalized" approach to cancer
16	staging, 8th ed CA: Cancer J Clin 67:93–99. doi: 10.3322/caac.21388
17	16. Endo S, Dousei T, Yoshikawa Y, Hatanaka N, Kamiike W, Nishijima J (2013) Prognosis of
18	gastric carcinoma patients aged 85 years or older who underwent surgery or who received best

1	supportive care only. Int J Clin Oncol 18:1014–1019. DOI: 10.1007/s10147-012-0482-9
2	17. Kim EJ, Seo KW, Yoon KY (2012) Laparoscopy-assisted distal gastrectomy for early gastric
3	cancer in the elderly. J Gastric Cancer 12:232-236. DOI: 10.5230/jgc.2012.12.4.232
4	18. Lieffers JR, Bathe OF, Fassbender K, Winget M, Baracos VE (2012) Sarcopenia is
5	associated with postoperative infection and delayed recovery from colorectal cancer resection surgery.
6	Br J Cancer 107:931–936. DOI: 10.1038/bjc.2012.350
7	19. Shen Y, Hao Q, Zhou J, Dong B (2017) The impact of frailty and sarcopenia on postoperative
8	outcomes in older patients undergoing gastrectomy surgery: a systematic review and meta-analysis.
9	BMC Geriatr 17:188. https://doi.org/10.1186/s12877-017-0569-2
10	20. Tanaka T, Suda K, Inaba K, Umeki Y, Gotoh A, Ishida Y, Uyama I (2019) Impact of frailty
11	on postoperative outcomes for laparoscopic gastrectomy in patients older than 80 years. Ann Surg
12	Oncol 26:4016–4026. DOI: 10.1245/s10434-019-07640-0
13	21. Murthy BL, Thomson CS, Dodwell D, Shenoy H, Mikeljevic JS, Forman D, Horgan K
14	(2007) Postoperative wound complications and systemic recurrence in breast cancer. Br J Cancer
15	97:1211–1217. DOI: 10.1038/sj.bjc.6604004
16	22. Kressner U, Graf W, Mahteme H, Påhlman L, Glimelius B (2002) Septic complications and
17	prognosis after surgery for rectal cancer. Dis Colon Rectum 45:316–321. DOI: 10.1007/s10350-004-
18	6174-4

1	23. Artinyan A, Orcutt ST, Anaya DA, Richardson P, Chen GJ, Berger DH (2015) Infectious
2	postoperative complications decrease long-term survival in patients undergoing curative surgery for
3	colorectal cancer: a study of 12,075 patients. Ann Surg 261:497-505. DOI:
4	10.1097/SLA.00000000000854
5	24. Cho JY, Han HS, Yoon YS, Hwang DW, Jung K, Kim YK (2013) Postoperative
6	complications influence prognosis and recurrence patterns in periampullary cancer. World J Surg
7	37:2234–2241. DOI: 10.1007/s00268-013-2106-6
8	25. Jiang N, Deng JY, Ding XW, Zhang L, Liu HG, Liang YX, Liang H (2014) Effect of
9	complication grade on survival following curative gastrectomy for carcinoma. World J Gastroenterol
10	20:8244–8252. DOI: 10.3748/wjg.v20.i25.8244
11	26. Li QG, Li P, Tang D, Chen J, Wang DR (2013) Impact of postoperative complications on
12	long-term survival after radical resection for gastric cancer. World J Gastroenterol 19:4060-4065.
13	DOI: 10.3748/wjg.v19.i25.4060
14	27. Kawase H, Ebihara Y, Shichinohe T, Nakamura F, Murakawa K, Morita T, Okushiba S,
15	Hirano S (2017) Long-term outcome after laparoscopic gastrectomy: A multicenter retrospective study.
16	Langenbecks Arch Surg 402:41–47. DOI: 10.1007/s00423-017-1559-8
17	28. Sasako M, Sakuramoto S, Katai H, Kinoshita T, Furukawa H, Yamaguchi T, Nashimoto A,
18	Fujii M, Nakajima T, Ohashi Y (2011) Five-year outcomes of a randomized phase III trial comparing

adjuvant chemotherapy with S-1 versus surgery alone in stage II or III gastric cancer. J Clin Oncol

2 29:4387–4393. DOI: 10.1200/JCO.2011.36.590)8
----------------------------------------------	----

- 3 29. Noh SH, Park SR, Yang HK, Chung HC, Chung IJ, Kim SW, Kim HH, Choi JH, Kim HK,
- 4 Yu W, Lee JI, Shin DB, Ji J, Chen JS, Lim Y, Ha S, Bang YJ, CLASSIC trial investigators (2014)
- 5 Adjuvant capecitabine plus oxaliplatin for gastric cancer after D2 gastrectomy (Classic): 5-year
- 6 follow-up of an open-label, randomised phase 3 trial. Lancet Oncol 15:1389-1396. DOI:
- 7 10.1016/S1470-2045(14)70473-5

1

- 8 30. Yoshida K, Kodera Y, Kochi M, Ichikawa W, Kakeji Y, Sano T, Nagao N, Takahashi M,
- 9 Takagane A, Watanabe T, Kaji M, Okitsu H, Nomura T, Matsui T, Yoshikawa T, Matsuyama J, Yamada
- 10 M, Ito S, Takeuchi M, Fujii M (2019) Addition of docetaxel to Oral fluoropyrimidine improves
- 11 efficacy in patients with stage III gastric Cancer: interim analysis of JACCRO GC-07, a randomized
- 12 controlled trial. J Clin Oncol 37:1296–1304. DOI: 10.1200/JCO.18.01138
- 13 31. Rausei S, Boni L, Rovera F, Dionigi G. (2013) Locally advanced gastric cancer: a new
- 14 definition to standardise. J Clin Pathol. 66(2):164-5. doi: 10.1136/jclinpath-2012-201176. Epub 2012
- 15 Dec 8. PMID: 23223567.
- 16 32. Smyth EC, Verheij M, Allum W, Cunningham D, Cervantes A, Arnold D (2016) ESMO
- 17 Guidelines Committee. Gastric cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment
- 18 and follow-up. Ann Oncol. 27(suppl 5):v38-v49. doi: 10.1093/annonc/mdw350. PMID: 27664260.

Figure legends

Fig. 1 Study enrollment. LTG: laparoscopic total gastrectomy; NAC: Neoadjuvant therapy

* In this study, for accurate prognostic analysis, we included the number of lymph nodes dissected to more than 16 in the exclusion criteria

Fig. 2 Prognosis of patients who underwent laparoscopic total gastrectomy for gastric cancer according to age. (A) Overall, (B) disease-specific, and (C) disease-free survival









Table 1. Clinical features and surgical outcomes of the study population

Variable	Overall (n=440)
Gender (M/F)	305/135
Age (year) (median, range)	67 (25-88)
BMI^{\dagger} (kg/m ²) (median, range)	22.9 (13.6-38.9)
ASA-PS* (≤II) (patients,%)	318 (72.3%)
Clinical JCGC stage** (≥II) (patients,%)	135 (30.7%)
Lymph node dissection (≥D2) (patients,%)	68 (15.5%)
Operation time (min) (median, range)	329 (123-762)
Blood loss (ml) (median, range)	50 (0-1940)
Postoperative complication (CD [§] , \geq IIIa) (patients,%)	66 (15.0%)
Postoperative hospital stays (days) (median, range)	14 (6-210)

⁺ Body mass index, *The American Society of Anesthesiologist's physical status, **According to the Japanese classification of gastric carcinoma: 3rd English edition, § Clavien-Dindo, classification

		Unmatched patients (n=395)			Propensity-matched patients (n=74)		
		EG (n=37)	non-EG (n=358)	<i>p</i> value	EG (n=37)	non-EG (n=37)	p value
		Number	Number		Number	Number	
Sex (%)				0.585			0.619
]	Male	24 (64.9)	248 (69.3)		13 (35.1)	11 (29.7)	
]	Female	13 (35.1)	110 (30.7)		24 (64.9)	26 (70.3)	
BMI ^{\dagger} (kg/m ²), median (range)		22.8 (13.6-38.9)	23.1 (15.0-30.9)	0.951	23.1 (15.6-33.4)	22.2 (14.9-34.3)	0.317
ASA-PS* (%)				0.014			0.740
	1-2	32 (86.5)	343 (95.8)		32 (86.5)	34 (91.9)	
:	3-4	5 (13.5)	15 (4.2)		5 (13.5)	3 (8.1)	
Clinical JCGC stage** (%)				0.570			0.848
]	I	22 (59.5)	251 (75.2)		22 (59.5)	22 (59.5)	
]	II	11 (29.7)	75 (16.8)		11 (29.7)	9 (24.3)	
]	III	4 (10.8)	32 (8.0)		4 (10.8)	5 (13.5)	

Table 2. Patient's characteristics who underwent LTG before and after propensity score matching

[†]Body mass index, *The American Society of Anaesthesiologist's physical status, **According to the Japanese classification of gastric carcinoma: 3rd

English edition.

	Unmatched patients (n=395)			Propensity-matched patients (n=74)			
	EG (n=37)	non-EG (n=358)	<i>p</i> value	EG (n=37)	non-EG (n=37)	<i>p</i> value	
	Number	Number		Number	Number		
Operative time (min), median (range)	260 (171-525)	330 (123-762)	< 0.001	260 (171-525)	335 (172-553)	0.003	
Blood loss (ml), median (range)	102.5 (0-909)	50 (0-1940)	0.103	10 (0-909)	40 (0-850)	0.645	
Extent of lymph node dissection			< 0.001			1.000	
D1/D1+ (%)	37 (100.0)	296 (82.7)		37 (100.0)	37 (100.0)		
D2 (%)	0	62 (17.3)		0	0		
Number of harvested lymph nodes, median (range)	29 (17-90)	41 (16-114)	0.028	29 (17-90)	37 (18-90)	0.645	
Postoperative complication (%)							
$CD \ge II$	9 (24.3)	79 (22.1)	0.756	9 (24.3)	5 (13.5)	0.233	
$CD \ge III$	6 (16.2)	55 (15.4)	0.892	6 (16.2)	4 (10.8)	0.495	
Postoperative hospital stay (day), median (range)	16 (8-82)	13 (6-210)	0.034	16 (8-82)	12 (7-65)	0.008	
Mortality (within 30 days)	0	2		0	0		

Table 3 Surgical outcomes and postoperative course in patients who underwent curative LTG before and after propensity score matching

	Unmatched patients (n=395)			Propensity-matched patients (n=74)		
	EG (n=37)	non-EG (n=358)	<i>p</i> value	EG (n=37)	non-EG (n=37)	<i>p</i> value
	Number	Number		Number	Number	
Pathological JCGC stage* (%)			0.332			0.726
Ι	17(45.9)	204(57.0)		17(45.9)	16(43.3)	
II	9(24.3)	83(23.2)		9(24.4)	7(18.9)	
III	4(10.8)	32(8.0)		11(29.7)	14(37.8)	
Adjuvant chemotherapy	4	104		4	16	
Recurrence (Y/N)	8/29	49/309	0.191	8/29	9/28	0.782
Recurrence site						
LN (Regional/Para Ao/Distant)	1/0/0	2/5/2		1/0/0	0/1/1	
Peritoneum dissemination	4	17		4	1	
Liver metastasis	3	14		3	3	
Lung metastasis	0	6		0	1	
Brain metastasis	0	1		0	1	
Bone metastasis	0	1		0	0	
Port site recurrence	0	1		0	0	
Length of recurrence (day), median (range)	249.5 (60-1480)	452 (72-2289)	0.081	249.5 (60-1480)	713 (22-3079)	0.068
*According to the Japanese classification of	gastric carcinoma: 3	rd English edition.				

Table 4.	Patient's characteristics who underwent LTG before and after propensity score matching	
----------	----------------------------------------------------------------------------------------	--