

HOKKAIDO UNIVERSITY

Title	Genital organ-associated lymphoid tissues arranged in a ring in the mucosa of cow vaginal vestibules
Author(s)	Chuluunbaatar, Tsolmon; Ichii, Osamu; Masum, Md Abdul; Namba, Takashi; Islam, Md Rashedul; Otani, Yuki; Elewa, Yaser Hosny Ali; Kon, Yasuhiro
Citation	Research in veterinary science, 145, 147-158 https://doi.org/10.1016/j.rvsc.2022.02.011
Issue Date	2022-07
Doc URL	http://hdl.handle.net/2115/90205
Rights	© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Туре	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Manuscript.pdf



Genital organ-associated lymphoid tissues arranged in a ring in the mucosa of cow vaginal vestibules

1	
2	Tsolmon Chuluunbaatar ^{1,2} , Osamu Ichii ^{1,3} , Md. Abdul Masum ⁴ , Takashi Namba ¹ , Md.
3	Rashedul Islam ^{1,5} , Yuki Otani ¹ , Yaser Hosny Ali Elewa ^{1,6} , Yasuhiro Kon ^{1*}
4	
5	^{1.} Laboratory of Anatomy, Department of Basic Veterinary Sciences, Faculty of
6	Veterinary Medicine, Hokkaido University, Hokkaido 060-0818, Japan
7	^{2.} Department of Basic Science of Veterinary Medicine, School of Veterinary Medicine,
8	Mongolian University of Life Science, Ulaanbaatar 17024, Mongolia
9	^{3.} Laboratory of Agrobiomedical Science, Faculty of Agriculture, Hokkaido University,
10	Hokkaido 060-8589, Japan
11	^{4.} Department of Anatomy, Histology, and Physiology, Faculty of Animal Science and
12	Veterinary Medicine, Sher-e-Bangla Agricultural University, Dhaka 1207, Bangladesh
13	^{5.} Department of Surgery and Theriogenology, Faculty of Animal Science and Veterinary
14	Medicine, Sher-e-Bangla Agricultural University, Dhaka 1207, Bangladesh
15	^{6.} Department of Histology, Faculty of Veterinary Medicine, Zagazig University, Zagazig
16	44519, Egypt
17	* Corresponding author: Yasuhiro Kon, DVM, PhD
18	Laboratory of Anatomy, Department of Basic Veterinary Sciences, Faculty of Veterinary
19	Medicine, Hokkaido University, Kita 18-Nishi 9, Kita-ku, 060-0818 Sapporo, JAPAN
20	Tel/Fax: +81-11-706-5188, E-mail: y-kon@vetmed.hokudai.ac.jp

21

23 Female reproductive tracts are equipped with local and mucosal immune systems; however, 24 structural information remains unclear for farm animals. In this study, the 25 mucosa-associated lymphoid tissue-like structures in cow reproductive tracts were 26 described. Vaginal vestibule (VV) and external parts of the genital organ, including the 27 clitoris and vulva, were morphologically analyzed. Whole-mount specimens revealed 28 several hematoxylin-positive spots arranged in a ring in the mucosa. Histologically, these 29 spots were aggregated immune cells and defined as genital organ-associated lymphoid 30 tissues (GOALTs). GOALTs were composed of lymphatic follicles (LFs) or diffuse 31 lymphoid tissues (DLTs) at different depths of lamina propria. LFs frequently contained 32 germinal centers. Scattered lymphocytes occupied the border area between follicles and 33 epithelium, whereas DLTs had indefinite shapes. GOALTs contained immune cells and 34 high endothelial venules. B cells were dominant both in LFs and DLTs. Abundant 35 collagenous fibers were stretched across VV lamina propria, whereas reticular fibers were 36 primarily observed in the DLT rather than LF. The epithelium covering of GOALTs was 37 partially or fully disrupted by the invasion of immune cells toward the VV lumen. These 38 findings suggest GOALTs function as a "genital lymphoid ring" as in Waldever's 39 pharyngeal ring and act as immunological gate systems in cow reproductive tracts.

40

41 Keywords

42 Cow; Genital organ associated lymphoid tissue; Lymphoid ring; Lymphatic follicle;

43 Diffuse lymphoid tissue

44

45 Introduction

46 The mucosa-associated lymphoid tissue (MALT) is a local immune system beneath the mucosal epithelium (Kenneth Murphy, 2017; Lanzkowsky et al., 2016). MALTs are located 47 48 in systemic organs that directly contact the outer environment, including alimentary tracts, 49 salivary gland ducts, conjunctiva, nasopharynx, and respiratory tracts (Forchielli and 50 Walker, 2005; Kenneth Murphy, 2017; Knop and Knop, 2011; Kracke et al., 1997; Kuper 51 et al., 1992; Moyron-Quiroz et al., 2004; Nair and Schroeder, 1986; Randall, 2010). 52 Gut-associated lymphoid tissue (GALT) is a well-known MALT and also known as Peyer's 53 patches (Forchielli and Walker, 2005). In the oropharynx, there are ring-shaped localized 54 MALTs, including the lingual, palatine, and pharyngeal areas and the tubal tonsils 55 (Hellings et al., 2000). Waldever's pharyngeal ring is the sentinel for the oral, nasal, and 56 auditory tracts toward to the pharynx and basically functions in antibody production to 57 protect against regular environmental antigens (Liebler-Tenorio et al., 2006; Manesse et al., 58 1998).

59 MALTs mainly consist of lymphoid tissues (LTs) with two types of morphology. 60 Briefly, the lamina propria of several MALT-forming organs exhibit diffused arrangement 61 of immune cells, and this type is called diffuse LT (DLT). However, in several MALTs, the 62 accumulation of immune cells is separated by connective tissue from surrounding tissues, and these are called lymphatic follicles (LFs) or lymph nodules (Wojciech and Ross, 2020).
Furthermore, the MALT is composed of B cells, CD4⁺, or CD8⁺ T-cells, antigen-presenting
cells (APCs), macrophages, and occasionally, mast cells and eosinophils. Each cell is
well-situated to initiate an immune reaction when encountering antigens passing through
the mucosal epithelium (Cesta, 2006).

68 Mammalian female reproductive tracts are also in close contact with the outer 69 environment and maintain innate and adaptive immune systems constructed by the 70 epithelial barrier, the production of antimicrobial agents, cytokines secreted by the 71 epithelial cells, and the innate immune cells (Hickey et al., 2011; Mair et al., 2014). 72 Importantly, the reproductive tracts of humans or other animal species contain MALT with 73 species- or disease-specific variations in morphology. A series of studies on LT have 74 recently been pursued in some female genital tracts, including the vaginal vestibule (VV). 75 vagina, cervix, uterus, and oviducts in non-human primates, pigs, and cows to evaluate the 76 effects of drug administration or mucosal vaccination (Blazquez et al., 1987a; Kathrine and 77 Steen, 2011; Lehner et al., 1995). Specifically, the guiana dolphin (Sotalia guianensis) 78 exhibited aggregations of LFs in the lamina propria of their uterine(Silva et al., 2016). 79 Importantly, healthy female mice did not have MALT in their reproductive tracts, but they 80 exhibited a MALT-like structure in their vaginal mucosa after receiving intravaginal 81 immunization with a specific peptide (Wang et al., 2015). Furthermore, locally organized 82 MALT in the VV observed in patients with provoked vulvodynia compared with healthy 83 women, indicated the MALT emerged because of the local alternation of VV 84 morpho-function (Liebler-Tenorio et al., 2006; Tommola et al., 2015).

85	The reproductive management of animals is crucial in veterinary medicine. In cattle,
86	gross anatomically and functionally, MALT in the alimentary and respiratory tract has been
87	well-investigated in several studies (Anderson et al., 1986; Manesse et al., 1998; Parsons et
88	al., 1989). Briefly, several types of MALTs were established according to their anatomical
89	localization, including GALTs, conjunctiva-associated LTs, and bronchus-associated LTs
90	(Anderson et al., 1986; Chodosh et al., 1998). Importantly, epithelium-associated LF
91	aggregations were suggested throughout the reproductive tracts in cattle, including the VV
92	(Blazquez et al., 1987a). The LF aggregations located in the lamina propria of the clitoris
93	and VV could play an important role in immune induction sites because of their appearance
94	during infection with cow reproductive tract pathogens (Blazquez et al., 1987b; Wang et al.,
95	2011); however, structural information on reproductive tract-associated MALTs, in
96	particular their localizations, morphological types, and cellular composition, has received
97	limited attention.

In the present study, we found a MALT-like structure arranged in a ring around the cow VV, and the stratified epithelium covering the LT was partially or completely disrupted, which permitted direct passage of immune cells and erythrocytes into the intraluminal space. We named these LTs the "genital lymphoid ring" and discussed the similarities with Waldeyer's pharyngeal ring, which acts as an immunological gate.

103

104 Materials and methods

105 Animal and sample preparation

106 Animal experimentation procedures were approved by the Institutional Animal Care and 107 Use Committee of the Faculty of Veterinary Medicine, Hokkaido University (approval no. 108 19-0097). Investigators adhered to the Guidelines for the Care and Use of Laboratory 109 Animals of Hokkaido University, Faculty of Veterinary Medicine. All animal experimental 110 protocols were approved by the Association for Assessment and Accreditation of 111 Laboratory Animal Care International. Holstein breed, normal non-pregnant cows (total n = 112 8, over 2-year-old) were sedated with xylazine (3.0 mg/kg body weight) through an 113 intramuscular route followed by general anesthesia with pentobarbital sodium (500 mg/kg 114 body weight) via the intravascular route (Supplemental table 1). The cows were sacrificed 115 by intravenous administration of a saturated potassium chloride solution of at least 0.1 116 mL/kg body weight. Then, the jugular vein was exsanguinated. After euthanasia, female genital organs were obtained, and the vagina, VV, and EPGOs, including the clitoris and 117 118 vulva were separated. They were immediately fixed in 10% neutral buffered formalin for 119 more than 7 d. These specimens were dissected using scissors at the dorsal commissure 120 along the median line of the vulva to the vagina.

121

122 Whole-mount observation

Separated genital organs were pinned flat, with the mucosa uppermost, and immersed in Mayer's hematoxylin for approximately 10 min (examining at 3–4 min intervals) to visualize LTs. LTs were then visible as small navy-blue spots. To decrease the intensity of 126 the staining background, specimens were rinsed in 70% alcohol containing 1%127 hydrochloric acid.

128 Counting of hematoxylin-positive spots was performed for the VV and EPGOs. 129 Flattened VV and EPGOs were divided into six regions (Fig. 1). Briefly, we divided the 130 VV and EPGOs into two equal regions (cranial and caudal) along the horizontal line (the 131 length of the external urethral orifice to the ventral commissure was divided into two equal 132 lengths). Then, the cranial and caudal regions were subdivided by two vertical lines into 133 three equal subregions. The external urethral orifice was located at the center of the cranial 134 median region (Fig 1, region II), and the clitoris was located at the center of the caudal 135 median region (Fig. 1, region V). Each area was measured using ImageJ (National 136 Institutes of Health; MD, Bethesda, USA) after taking photos with a DMC-FX3799 device 137 (Panasonic; Osaka, Japan). The number of hematoxylin-positive spots was counted, and 138 their density in each region was calculated.

139

140 *Histoplanimetry*

The VV of each cow was cut into more than 15 pieces according to their size. Then, whole-mount specimens with hematoxylin were embedded into paraffin and cut to a thickness of 5 μm. Sections were stained with hematoxylin and eosin (H&E) or PSR for histological analysis. Stained sections were scanned with a NanoZoomer 2.0 RS virtual slide scanner (Hamamatsu Photonics; Shizuoka, Japan). Histological images for microscopic examinations were observed using a model BZ-X710 microscope (Keyence; Osaka, Japan). The number of LFs and DLTs was counted after H&E staining. This measurement was performed for more than 120 sections of eight cows. We also characterized the LFs according to their localization (isolated, migrating, bordering) and the DLTs according to their shape (oval, c-shaped, triangle, amorphous), and their appearance % was calculated.

152

153 Immunohistochemistry (IHC)

154 IHC was performed to evaluate the presence of B-cells (CD20), T-cells (CD3), 155 macrophages (IBA1), APCs (MHCII), high endothelial venules (PNAd), and tight junctions 156 (Occludin). The staining conditions of each antibody are listed in Supplementary table 2. In 157 brief, sections were deparaffinized and incubated in 10 mM citrate buffer (pH 6.0) or 158 Tris-HCl buffer (pH 9.0) for 15 min at 110 °C. Thereafter, slides were submerged in 159 methanol containing 0.3% H_2O_2 for 20 min at 20–22 °C and blocked with normal goat 160 serum (SABPO kit, Nichirei Bioscience; Tokyo, Japan) for 1 h at room temperature. 161 Sections were incubated with primary antibody overnight at 4 °C. After rinsing in 162 phosphate-buffered saline (PBS), sections were incubated with biotinylated secondary 163 antibody (goat anti-rabbit or goat anti-rat) for 30 min and streptavidin-horseradish 164 peroxidase (Nichirei Bioscience) for 30 min at room temperature. To develop the color, 165 sections were incubated in a 3,3'-diaminobenzidine tetrahydrochloride-hydrogen peroxide 166 solution for 4 min. Finally, the sections were counterstained with hematoxylin and

- dehydrated with ascending grades of alcohols. After washing with PBS, the sections weremounted and examined using a BZ-X710 microscope.
- 169 Next, we performed the histoplanimetrical analysis of LTs using the IHC sections for CD20,
- 170 CD3, IBA1, and MHCII. More than 10 LFs and DLTs were examined. Firstly, total area of
- 171 LFs or DLTs and the positive reaction area for each marker were measured, and percentage
- 172 of the latter to the former was calculated.
- 173

174 Electron microscopy

For SEM, small pieces of VV were treated with tannic acid and post-fixed with 1% OsO4 (osmium tetroxide) for 1 h. Then, the specimens were dehydrated in ascending grades of alcohol, immersed in 3-methylbutyl acetate, and dried with the HCP-2 critical point dryer (Hitachi; Tokyo, Japan). The specimens were then observed under an S-4100 scanning electron microscope (Hitachi).

180

181 *Statistical analyses*

182 The results are expressed as the mean \pm standard error (SE). Multiple comparisons were 183 performed using the Kruskal–Wallis test followed by the Scheffé method when a 184 significant difference was observed (P < 0.01).

185

186 **Results**

187 LT found in the VV and external parts of genital organs (EPGO) mucosa of cows

188 We investigated the mucosa of the vagina and VV extending from the external urethral 189 orifice to the vulva, and the vulva along with the pudendal labia and clitoris were also 190 examined as EPGOs (Supplemental figure 1). Whole-mount specimens were visualized 191 for hematoxylin-positive spots in the examined mucosae; in particular, we focused on VV 192 and EPGOs because these areas are close to the outer environment and have contact with 193 urine and feces. All examined cows had numerous hematoxylin-positive spots in the 194 mucosal surface of the VV and EPGOs, and these spots were distributed at an 195 approximately 1 cm distance from the boundary of the un-haired pudendal labia (Fig. 1A, 196 B). Furthermore, 37.5% of examined cows showed these spots in EPGOs (three of eight 197 cows), especially in the area close to the clitoris (Fig. 1C). Then, these spots were 198 histologically identified as LT (Fig. 1D), and classified as genital organ-associated LTs 199 (GOALTs). They were further investigated in subsequent experiments.

200 We also examined the localizations of GOALTs by dividing the VV and EPGOs into 201 six regions (Fig. 1E). During the histometry of the density of GOALT spots for the six 202 examined regions, region II and V, located along the ventral median line of the VV, 203 including the external urethral orifice, clitoris, and ventral commissure of the labia, tended 204 to exhibit a higher density without significance (Fig. 1F). These results indicated that 205 GOALTs were distributed in the mucosa of the VV and EPGOs in a ring shape; therefore, 206 we denoted the GOALT distributions as a "genital lymphoid ring," based on "Waldeyer's 207 lymphoid pharyngeal ring." In subsequent analysis, we focused on VV mucosa.

209 Histology of GOALTs found in the cow VV mucosa

210 The histological observation was performed in the VV. The mucosa was covered by 211 non-keratinized and stratified squamous epithelium (Fig. 2). The GOALTs were mainly 212 localized in the lamina propria in the VV. Importantly, regarding the LT structures, two 213 different types of GOALTs were found in the VV; LF (Fig. 2A-D) and DLT (Fig. 2E-H). 214 For LF, three types were primarily observed in the VV mucosa based on their morphology. 215 First, several LFs were localized in the deep portion of the lamina propria and fully 216 surrounded by connective tissues. In this type, the VV epithelium completely maintained 217 the structure (Fig. 2A). Second, scattered lymphocytes were observed between the LF and 218 VV epithelium, and these lymphocytes were also thoroughly or partially observed within 219 the VV epithelium structure (Fig. 2B). Third, LF directly faced the VV lumen, and the VV 220 epithelium was occupied by lymphocytes (Fig. 2C). Further, most of the LFs were 221 solitarily, but paired LFs were also noticed (Fig. 2D). Additionally, several shapes of DLTs 222 were observed, such as triangle-, C-shaped-, amorphous-, and oval-types (Fig. 2E-H).

In the histometry of the VV mucosa, LF- and DLT-types accounted for 35% and 65%, respectively (**Fig. 2I**). Furthermore, as a result of morphometry according to the LF localization, migrating-, isolated-, and bordering-types accounted for 51%, 37%, and 12%, respectively (**Fig. 2J**). Finally, for the DLT shapes, amorphous-, oval-, C-shaped-, and triangle-types accounted for 57%, 30%, 9%, and 4%, respectively (**Fig. 2K**).

228

229 Cell composition of GOALTs in the cow VV mucosa

Regarding LFs of the GOALT in the VV mucosa, numerous CD20⁺ B cells were found, 230 231 and the germinal center and an area between the LF and VV epithelium were located (Fig. 232 **3A**). Further, CD3⁺T cells were diffusely distributed in LFs, including the germinal center, 233 and tended to mainly localize along the border area between the LF and VV epithelium 234 (Fig. 3B). IBA1⁺ macrophages and MHCII⁺ APCs were similar in localization and were 235 located in the germinal center and the border area between the LF and VV epithelium; the 236 former was also localized in the VV epithelium (Fig. 3C, D). Next, PNAd⁺ high endothelial 237 venules were localized around the LFs and in the border area between the LF and VV 238 epithelium (Fig. 3E). In agreement with histological observations, the percentage of B cells 239 was statistically significantly higher in LFs than the percentage of T cells, macrophages, 240 and APCs (*P* < 0.01, **Fig. 3F**).

In the DLT, B cells, T cells, macrophages, APCs, and high endothelial venules were dispersed diffusely (**Fig. 4A-E**). Similar to LFs, DLTs had a statistically significantly higher percentage of B cells than that of other examined immune cells (P < 0.01, **Fig. 4F**).

244

245 Histology of epithelial cells covering the GOALT in the cow VV mucosa

Histologically, the VV mucosa was covered by stratified epithelial cells, wherein the basal layer was smaller with narrow cytoplasm, and their nuclei were located close to one another (**Fig. 5A**). Follicular epithelial cells found in MALTs were not identified. Importantly, in several LTs, the thickness of the epithelium gradually became smaller towards the epithelium covering LTs (**Fig. 5A**). Furthermore, the epithelium covering LTs showed complete destruction of each junction of epithelial cell with numerous lymphocytes and erythrocytes, and several epithelial cells peeled off and created a large junctional gap,
which permitted direct passage of immune cells into the intraluminal space of the VV (Fig.
5B). Occludin⁺ reactions were observed in the intercellular region of the epithelial cells, but
these were unclear in the epithelium covering LTs (Fig. 5C, D).

Next, picrosirius red (PSR) staining revealed the developed connective tissue fibers in the lamina propria of the VV mucosa. Under polarized light observation, well-developed collagen fibers were visualized as bright orange to red, whereas thinner reticular fibers appeared yellow to green (**Fig. 5E**). In GOALTs, the DLT-types contained abundant collagen and reticular fibers, but they were scarce in the LF-type (**Fig. 5E**). The VV mucosa exhibited well-developed collagen and reticular fibers beneath the epithelium (**Fig. 5F**), but they were obscure in the LF (**Fig. 5G**).

263

264 Ultrastructure of surface of LT in the VV

265 Next, we observed the GOALT surface in the VV mucosa (Fig. 6). Whole-mount staining 266 with hematoxylin revealed the GOALTs as spots (Fig. 6A). Using scanning electron 267 microscopy (SEM), the epithelium covering GOALTs was found to partially lack in several 268 areas (Fig. 6B). Briefly, at these regions, several polygonal-shaped epithelial cells were 269 peeled off, and lymphocytes appeared to be directly exposed to the intraluminal space 270 through the hole (Fig. 6C, D). Figure 6E shows the large sized, dome-shaped LF covered 271 by the VV epithelium, which was approximately 350-480 µm in a diameter. Several 272 differently sized destroyed areas of the epithelium fused with each other and formed a 273 complicated cleft. From the inside of the GOALT, erythrocytes were also exposed directly

to the intraluminal space (**Fig. 6F, G**). Additionally, small protrusions occurred that appeared to be lymphocytes pushed up to the surface of the epithelium (**Fig. 6F, G**). Some connective tissue fibers were also observed inside of the GOALT and opened to the lumen (**Fig. 6G**). Summary of general quantitative data shown in **supplemental table 3**.

278

279

280 **Discussion**

281 This study clarified the morphological characteristics of GOALTs found in the VV and 282 EPGOs of the cow. In cows, the VV extends from the external urethral orifice to the ventral 283 commissure of the labia and is composed of the caudal part of the reproductive and/or 284 urinary system, localized ventral to the anus, with a length of approximately 7-11 cm 285 (Budras and E.Habel, 2003; Konig and Liebich, 2007). The LTs in the bovine VV were 286 previously suggested in cows (Blazquez et al., 1987b; Cole, 1930; Hammond, 1927), but 287 detailed morphological information was lacking. The high abundance of LTs along the VV 288 may relate to increased antigenic stimulation in this region. In the present study, the ventral median region (II and V containing EPGOs) of the VV and EPGOs tended to have a greater 289 290 number of GOALTs than other areas. We considered that the developed GOALTs in this 291 area reflected the possibility of high exposure to antigenic substances, such as urine or 292 feces, because of the anatomical localization of the VV. These relationships between 293 antigenic challenge and the localizations of LTs are compatible with statements regarding 294 the amount and type of MALTs being partially dependent on the antigenic challenge 295 (Chuluunbaatar et al., 2020; Jericho et al., 1971; Pollard and Sharon, 1970). Additionally,

the age of the cow appeared to be a factor influencing the amount of LT, and the VV of
adult cows contained a significantly higher number of LTs than 6–8-week-old calves
(Blazquez et al., 1987a).

299 GOALTs were morphologically divided into LFs and DLTs, and the latter was 300 dominant in the VV. LFs were usually dispersed solitarily in a random manner throughout 301 the mucosa of the VV or EPGOs. Most LFs had a germinal center, which initiated a series 302 of events that included lymphocyte activation and proliferation, plasma cell differentiation, 303 and antibody production. Mitotic patterns were frequently seen in the germinal center, 304 indicating that new lymphocytes were proliferating in situ (Wojciech and Ross, 2020). 305 Although our histological data showed different morphology of LFs, the dominant was the 306 migrating-type, and sectioning of LFs also could have affected their shape observed during 307 microscopy. SEM analysis revealed both an epithelium-covered area (isolated-type) and 308 non-covered area (migrating-, bordering-type) in the same LF. Furthermore, 309 three-dimensional analysis of LFs could clarify their fine structure.

310 DLTs typically do not have a distinct border with surrounding tissues, rather have an 311 amorphous shape and are tactically localized to intercept antigens and initiate an immune 312 response. In general, after contact with antigen, lymphocytes located in the LT move to 313 regional lymph nodes, where they can sustain proliferation and differentiation. Then, the 314 progenies of these cells return to the lamina propria as B and T lymphocytes with effector 315 functions. Therefore, combined localization of LFs and DLTs in the VV or EPGO mucosa 316 could act as a protector of the reproductive tract against pathogenic substances. 317 Furthermore, our histological data showed the different morphology of LFs; the dominant

amorphous-type, with variation in the morphology of DLT that might reflect the differencein the activation phase of GOALTs.

Subsequently, the percentage of the positive area for the immune cells calculated in GOALT and both DLTs and LFs was dominated by B cells in comparison with T cells, macrophages, and APCs. This was an expected finding, which is in agreement with other research data, including those for the mice spleen, Peyer's patches, isolated LFs, and cow and rabbit GALT (Hamada et al., 2002; Komban et al., 2019; Parsons et al., 1989; Schneider et al., 2001).

326 The cell components of the cow GOALT showed close similarities to those of other 327 MALTs (GALT and Waldeyer's pharyngeal ring); however, the cell distribution was 328 generally solitarily and randomly localized within the VV and EPGOs (Owen and Jones, 329 1974). One of the most important cell components in MALTs is the APC. In this study, 330 APCs were found around the DLT and LF in different regions of the mucosal epithelium, 331 such as the germinal center, mantle zone, submucosa, and middle layers of stratified 332 squamous epithelium of the VV. These strategically positioned localizations might play 333 important roles in the initiation of immune reactions and maintenance of peripheral 334 tolerance. Therefore, various types of APCs in the LTs have different ways to take up 335 antigens from the lumen, including endocytosis, phagocytosis, or transcytosis (Zhang et al., 336 2007). Although, in this study, the epithelium that covered GOALTs was thinner than areas 337 not covering GOALTs, and in some areas, the epithelium structure was completely lost, and 338 LT and the intraluminal area were directly bordered without any edge. This formation of 339 LT allows direct passage of antigens to the LT and immune cells from the intraluminal area. 340 As described in the literature (Brandtzaeg et al., 2008), MALT is divided into two specific 341 types, an organized MALT delimited by organized collagen fiber tissue and a disorganized 342 or diffuse MALT, comprising populations of lymphocytes from the lamina propria and the 343 base of the epithelial lining, which are consistent with LFs and DLTs. LFs located in the 344 conjunctiva have a lenticular shape, and occasionally around this area, the basement 345 membrane is mostly discontinuous, and the epithelium is thinner (Crespo-Moral et al., 346 2020). We classified DLTs by shape, and each shape might reflect the different LT activity 347 in the VV mucosa.

348 We studied the normal histomorphology of the GOALT, and unfortunately, the phase 349 of the estrus cycle was impossible to identify in the examined cows. Importantly, the 350 immune system in the female reproductive tract is precisely influenced and regulated by 351 sex steroid hormones, estradiol and progesterone, which are produced in cyclic variations 352 by the ovary during the estrus cycles. Throughout the estrus cycle, immune cells are present 353 in considerable numbers and inconsistently distributed in the stromal layer and the 354 epithelium of the female reproductive tract (Bulmer and Earl, 1987; Hunt, 1994). Moreover, 355 the epithelial barrier of the reproductive tract, which includes various tight junction conditions, provides immune protection by maintaining a strong physical barrier, 356 357 transferring antibodies to the mucosa, producing antibacterial composites, and recruiting 358 immune cells, all of which are influenced by hormones (Russell and Mestecky, 359 2002)(Ochiel et al., 2008). Additionally, the various shapes of GOALTs found in the VV 360 and EPGOs may be affected by different phases of the estrus cycle. A limitation of this 361 study was the small sample size available to identify whether or not the estrus cycle

influences the amount of GOALT in the VV in cows. Further study is needed to clarify theGOALT amount at every stage of the estrous cycle.

364 In summary, this study morphologically and histologically evaluated GOALTs in the 365 VV and EPGOs of cows, which constituted a "genital lymphoid ring." Regarding GOALT 366 morphology (Fig. 7), the mucosal epithelium covering DLTs and LFs was partially or 367 completely demolished, making a direct passage for immune cells to reach the luminal side. 368 The clarification of the morphology of GOALTs in cow genital tracts would contribute to 369 our understanding of its immune-associated function, the pathogenesis of diseases, and the 370 development of therapeutic strategy by applying mucosal vaccinations targeting GOALTs 371 in farm animals.

372

Declarations of competing interest

The authors declare that they have no known competing financial interest or personal relationships that could have influenced in any way the work reported in this paper.

376

377 Acknowledgments

378 This work was supported by WISE KAKENHI and JSPS KAKENHI (Grant No.:379 19H03113).

380

381 Author Contributions

382	YK and OI supervised the findings of this study. MAM, YHAE, TENA, TANA, MRI, and
383	YO provided critical feedback and helped shape the research, analysis, and manuscript. TC,
384	OI, and YK analyzed data and wrote the paper.
385	
386	References
387	Anderson, M.L., Moore, P.F., Hyde, D.M., Dungworth, D.L., 1986. Bronchus associated
388	lymphoid tissue in the lungs of cattle: relationship to age. Res. Vet. Sci. 41, 211–220.
389	Blazquez, N.B., Batten, E.H., Long, S.E., Perry, G.C., 1987a. Amount and distribution of
390	vestibular-associated lymphoid tissue in calves and adult cows. Res. Vet. Sci. 43,
391	239–242.
392	Blazquez, N.B., Batten, E.H., Long, S.E., Perry, G.C., 1987b. Histology and histochemistry
393	of the bovine reproductive tract caudal to the cervix part I. The vestibule and
394	associated glands. Br. Vet. J. 143, 328-337.
395	Brandtzaeg, P., Kiyono, H., Pabst, R., Russell, M.W., 2008. Terminology: Nomenclature of
396	mucosa-associated lymphoid tissue. Mucosal Immunol. 1, 31-37.
397	Budras, KD., E.Habel, R., 2003. Bovine Anatomy, 1st ed. Schlutersche, Hannover.
398	Bulmer, J.N., Earl, U., 1987. The expression of class II MHC gene products by fallopian
399	tube epithelium in pregnancy and throughout the menstrual cycle. Immunology 61,
400	207–213.
401	Cesta, M.F., 2006. Normal structure, function, and fistology of mucosa-associated

402 lymphoid tissue. Toxicol. Pathol. 34, 599–608.

- 403 Chodosh, J., Nordquist, R.E., Kennedy, R.C., 1998. Comparative anatomy of mammalian
- 404 conjunctival lymphoid tissue: A putative mucosal immune site. Dev. Comp. Immunol.
- 405 22, 621–630.
- 406 Chuluunbaatar, T., Ichii, O., Nakamura, T., Irie, T., Namba, T., Islam, M.R., Otani, Y.,
- 407 Masum, M.A., Okamatsu-Ogura, Y., Elewa, Y.H.A., Kon, Y., 2020. Unique running
- 408 pattern and mucosal morphology found in the colon of cotton rats. Front. Physiol. 11.
- 409 Cole, H.H., 1930. A study of the mucosa of the genital tract of the cow, with special

410 reference to the cyclic changes. Am. J. Anat. 46, 261–301.

- 411 Crespo-Moral, M., García-Posadas, L., López-García, A., Diebold, Y., 2020. Histological
- 412 and immunohistochemical characterization of the porcine ocular surface. PLoS One 15,413 e0227732.
- Forchielli, M.L., Walker, W.A., 2005. The role of gut-associated lymphoid tissues and
 mucosal defence. Br. J. Nutr. 93, S41–S48.
- 416 Hamada, H., Hiroi, T., Nishiyama, Y., Takahashi, H., Masunaga, Y., Hachimura, S.,
- 417 Kaminogawa, S., Takahashi-Iwanaga, H., Iwanaga, T., Kiyono, H., Yamamoto, H.,
- 418 Ishikawa, H., 2002. Identification of multiple isolated lymphoid follicles on the
- 419 antimesenteric wall of the mouse small intestine. J. Immunol. 168, 57–64.
- 420 Hammond, J., 1927. The physiology of reproduction in the cow. Nature 120, 37–39.
- 421 Hellings, P., Jorissen, M., Ceuppens, J., 2000. The Waldeyer's ring. Acta Otorhinolaryngol.

422 Belg. 54, 237–241.

423	Hickey, D.K.	. Patel. M. V	' Fahev	. J. V.,	Wira.	C.R.	. 2011.	Innate and ada	ptive	immunity	v at
	1110110, , D.111.	, 1 00001, 1111 1	., 1 0110 /	,,	,, , ,,	U .11	,	minute and add		111111001110	,

- mucosal surfaces of the female reproductive tract: Stratification and integration of 424
- 425 immune protection against the transmission of sexually transmitted infections. J.
- 426 Reprod. Immunol. 88, 185–194.
- 427 Hunt, J.S., 1994. Immunologically relevant cells in the uterus. Biol. Reprod. 50, 461-466.
- Jericho, K.W.F., Derbyshire, J.B., Jones, J.E.T., 1971. Intrapulmonary lymphoid tissue of 428
- 429 pigs exposed to aerosols of haemolytic streptococcus group L and porcine adenovirus. 430
- J. Comp. Pathol. 81, 1–11.
- 431 Kathrine, E., Steen, J., 2011. A review of the human vs. porcine female genital tract and 432 associated immune system in the perspective of using minipigs as a model of human 433 genital Chlamydia infection. Vet. Res. 46, 1-13.
- 434 Kenneth Murphy, C.W., 2017. Janeway's Immunobiology - 9th Edition. Garland Science,
- 435 Taylor & Francis Group, New York.
- 436 Knop, E., Knop, N., 2011. The eye-associated lymphoid tissue (EALT) - A basis of the
- 437 anatomy and immunology at the ocular surface. Acta Ophthalmol. 89, 0–0.
- 438 Komban, R.J., Strömberg, A., Biram, A., Cervin, J., Lebrero-Fernández, C., Mabbott, N.,
- 439 Yrlid, U., Shulman, Z., Bemark, M., Lycke, N., 2019. Activated Peyer's patch B cells
- 440 sample antigen directly from M cells in the subepithelial dome. Nat. Commun. 10,
- 441 1–15.

- Konig, H., Liebich, H., 2007. Veterinary anatomy of domestic mammals: Textbook and
 colour atlas, 3rd ed. Schattauer, New York.
- 444 Kracke, A., Hiller, A.S., Tschernig, H., Kasper, M., Kleemann, W.J., Troger, H.D., Pabst,
- R., 1997. Larynx-associated lymphoid tissue (LALT) in young children. Anat. Rec.
 248, 413–420.
- 447 Kuper, C.F., Koornstra, P.J., Hameleers, D.M.H., Biewenga, J., Spit, B.J., Duijvestijn,
- 448 A.M., Van Breda Vriesman, P.J.C., Sminia, T., 1992. The role of nasopharyngeal
- 449 lymphoid tissue. Immunol. Today 12, 219–224.
- 450 Lanzkowsky, P., Lipton, J.M., Fish, J.D., 2016. Lanzkowsky's Manual of Pediatric
- 451 Hematology and Oncology, Sixth Edition. Academic Press.
- 452 Lehner, T., Panagiotidi, C., Bergmeier, L.A., Tao, L., Brookes, R., Gearing, A., Adams, S.,
- 453 1995. Genital-associated lymphoid tissue in female non-human primates. Adv. Exp.
 454 Med. Biol. 371, 357–365.
- Liebler-Tenorio, E., Pabst, R., Liebler, E., Liebler-tenorio, E.M., 2006. MALT structure
 and function in farm animals. Vet. Res. 37, 257–280.
- 457 Mair, K.H., Sedlak, C., Käser, T., Pasternak, A., Levast, B., Gerner, W., Saalmüller, A.,
- 458 Summerfield, A., Gerdts, V., Wilson, H.L., Meurens, F., 2014. The porcine innate
- 459 immune system: An update. Dev. Comp. Immunol. 45, 321–343.
- 460 Manesse, M., Delverdier, M., Abella-Bourges, N., Sautet, J., Cabanié, P., Schelcher, F.,
- 461 1998. An immunohistochemical study of bovine palatine and pharyngeal tonsils at 21,

462	60 and 300 day	ys of age. Anat. Histe	ol. Embryol. 27, 179–185.
-----	----------------	------------------------	---------------------------

463	Moyron-Quiroz, J.E., Rangel-Moreno, J., Kusser, K., Hartson, L., Sprague, F., Goodrich, S.,
464	Woodland, D.L., Lund, F.E., Randall, T.D., 2004. Role of inducible bronchus
465	associated lymphoid tissue (iBALT) in respiratory immunity. Nat. Med. 10, 927–934.
466	Nair, P.N.R., Schroeder, H.E., 1986. Duct-associated lymphoid tissue (DALT) of minor
467	salivary glands and mucosal immunity. Immunology 57, 171–17180.
468	Ochiel, D., Fahey, J., Ghosh, M., Haddad, S., Wira, C., 2008. Innate immunity in the
469	female reproductive tract: Role of sex hormones in regulating uterine epithelial cell
470	protection against pathogens. Curr. Womens. Health Rev. 4, 102–117.
471	Owen, R., Jones, A., 1974. Epithelial cell specialization within human Peyer's patches: an
472	ultrastructural study of intestinal lymphoid follicles. Gastroenterology 66, 189–203.
473	Parsons, K.R., Howard, C.J., Jones, B. V, Sopp, P., 1989. Investigation of bovine gut
474	associated lymphoid tissue (GALT) using monoclonal antibodies against bovine
475	lymphocytes. Vet. Pathol. 26, 396–408.
476	Pollard, M., Sharon, N., 1970. Responses of the Peyer's patches in germ-free mice to
477	antigenic stimulation. Infect. Immun. 2, 96–100.
478	Randall, T.D., 2010. Bronchus-associated lymphoid tissue (BALT). Structure and function.
479	Adv. Immunol. 107, 187–241.
480	Russell, M.W., Mestecky, J., 2002. Humoral immune responses to microbial infections in

481 the genital tract. Microbes Infect. 4, 667–677.

482	Schneider, P., Takatsuka, H., Wilson, A., Mackay, F., Tardivel, A., Lens, S., Cachero, T.G.,
483	Finke, D., Beermann, F., Tschopp, J., 2001. Maturation of marginal zone and
484	follicular B cells requires B cell activating factor of the tumor necrosis factor family
485	and is independent of B cell maturation antigen. J. Exp. Med. 194, 1691–1697.
486	Silva, F.M.O., Guimarães, J.P., Vergara-Parente, J.E., Carvalho, V.L., Carolina, A.,
487	Meirelles, O., Marmontel, M., Oliveira, B.S.S.P., Santos, S.M., Becegato, E.Z.,
488	Evangelista, J.S.A.M., Miglino, M.A., 2016. Morphology of mucosa-associated
489	lymphoid tissue in odontocetes. Microsc. Res. Tech. 79, 845-855.
490	Tommola, P., Bützow, R., Unkila-Kallio, L., Paavonen, J., Meri, S., 2015. Activation of
491	vestibule-associated lymphoid tissue in localized provoked vulvodynia. Am. J. Obstet.
492	Gynecol. 212, 476.e1-476.e8.
493	Wang, J., Gusti, V., Saraswati, A., Lo, D.D., 2011. Convergent and divergent development
494	among M cell lineages in mouse mucosal epithelium. J. Immunol. 187, 5277–5285.
495	Wang, Y., Sui, Y., Kato, S., Hogg, A.E., Steel, J.C., Morris, J.C., Berzofsky, J.A., 2015.
496	Vaginal type-II mucosa is an inductive site for primary CD8+ T-cell mucosal
497	immunity. Nat. Commun. 6, 1–13.
498	Wojciech, P., Ross, M.H., 2020. Histology: A text and atlas: with correlated cell and
499	molecular biology, 8th ed. Wolters Kluwer, Philadelphia.
500	Zhang, J., Kuolee, R., Patel, G., Chen, W., 2007. Intestinal M cells: the fallible sentinels?
501	World J. Gastroenterol. 13, 1477–1486.

502

503 Figure legends

504 Figure 1. Distribution of GOALTs in the mucosa of the VV and EPGOs of cows.

505 (A) Gross anatomical features of the VV and EPGOs with whole-mount staining using

- 506 hematoxylin. White and red asterisks indicate the external urethral orifice and clitoris,
- 507 respectively. One scale of the ruler is 1 mm.
- 508 (B) Magnified image of the VV mucosa squared area in panel A. Scale bars = 1 cm.
- 509 (C) Magnified image of the EPGO mucosa squared area in panel A.
- 510 Black arrowheads indicate hematoxylin-positive spots. Scale bars = 1 mm.
- 511 (D) Histology of hematoxylin-positive spot squared area in panel B. Histological
- 512 observation revealed the spot is a cell aggregation, the GOALT, part of genital lymphoid
- 513 ring. H&E staining. Scale bar = $100 \mu m$.
- 514 (E) Schematic illustration of cutaway of the VV and EPGO area, including the vulva and
- 515 clitoris. For morphometry, the mucosal area is divided into six regions. Blue circles indicate
- 516 GOALTs.
- 517 (F) The density of GOALTs in the VV mucosa (number/cm²).
- 518 Values = mean \pm SE. n = 8.
- 519 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, EPGOs: external
- 520 parts of genital organs, H&E: hematoxylin & eosin
- 521
- 522 Figure 2. Histological characteristics of GOALTs in the cow VV.

- 523 (A) Isolated-type LF. LF localizes in the deep portion of the lamina propria and is fully
- 524 surrounded by connective tissues.
- 525 (**B**) Migrating-type LF. Scattered lymphocytes are observed between LF and VV 526 epithelium.
- 527 (C) Bordering-type LF. LF directly faces the VV lumen, and the VV epithelium is occupied
- 528 by lymphocytes.
- 529 (**D**) Double migrating-type LF.
- 530 Dotted lines indicate the border of LFs. H&E staining. Scale bars = $100 \mu m$.
- 531 (E) Triangle-type DLT.
- 532 (F) C-shaped-type DLT.
- 533 (G) Amorphous-type DLT. Dotted lines indicate the border of the DLT.
- 534 (H) Oval-type DLT.
- 535 Scale bars = $100 \mu m$.
- 536 (I) Appearance of the percentage of LFs and DLTs in the examined GOALTs of the VV.
- 537 n = 8.
- 538 (J) Appearance of the percentage of each type of LF in the examined LFs of the VV. n = 8.
- 539 (K) Appearance of the percentage of each type of DLT in the examined DLTs of the VV.
- 540 n = 8.
- 541 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, H&E:
- 542 hematoxylin & eosin, LF: lymphatic follicle, DLT: diffuse lymphoid tissue
- 543
- 544 Figure 3. Immune cells composing the LF-type GOALT in the cow VV.

- 545 IHC was performed in the VV mucosa, and immune cells and high endothelial venules
- 546 were found in the GOALTs.
- 547 (A) CD20 for B-cells.
- 548 **(B)** CD3 for T-cells.
- 549 (C) IBA1 for macrophages.
- 550 (**D**) MHCII for APCs.
- 551 (E) PNAd for high endothelial venules.
- 552 Squares denote the magnified area. Black arrowheads indicate positive cells for the IHC.
- 553 Scale bars = 100 and $25 \mu m$ (low and high magnifications, respectively).
- 554 (F) Percentage of each examined cell occupying the LF. Values = mean \pm SE. $n \ge 4$.
- 555 Significant differences between the immune cells indicated by *, P < 0.01;
- 556 Kruskal–Wallis test followed by the Scheffé method.
- 557 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, LF: lymphatic
- 558 follicle, IHC: immunohistochemistry, APCs: antigen-presenting cells
- 559

560 Figure 4. Immune cells composing the DLT-type GOALT in the cow VV.

- 561 IHC was performed in the VV mucosa, and immune cells and high endothelial venules
- 562 were found in the GOALTs.
- 563 (A) CD20 for B-cells.
- 564 **(B)** CD3 for T-cells.
- 565 (C) IBA1 for macrophages.
- 566 (**D**) MHCII for APCs.

- 567 (E) PNAd for high endothelial venules.
- 568 Black arrowheads indicate positive cells for the IHC. Scale bars = $100 \mu m$.
- 569 (F) Percentage of each examined cell occupying the DLT. Values = mean \pm SE. $n \ge 4$.
- 570 Significant differences between the immune cells indicated by *, P < 0.01; Kruskal–Wallis
- 571 test followed by the Scheffé method.
- 572 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, DLT: diffuse
- 573 lymphoid tissue, IHC: immunohistochemistry, APCs: antigen-presenting cells
- 574

575 Figure 5. Epithelium covering and connective tissue fibers composing the GOALTs in 576 the cow VV.

- 577 (A, B) Amorphous-type DLT. Destroyed epithelium located between DLT and intralumenal
 578 area. The square indicates the magnified area. Red arrows denote peeled-off area of
 579 epithelial cells. H&E staining. Scale bars = 100 μm.
- (C, D) Mucosal epithelium features stained using IHC for occludin. Panel C shows the
 epithelium that does not cover the LT. Panel D shows the epithelium covering the LT.
- 582 Black arrowheads indicate positive reaction for the occludin. Scale bars = $50 \mu m$.
- 583 (E) Connective tissue fibers of LF and DLT stained using PSR under polarized conditions.
- 584 White dotted line indicates the border between the lamia propria and the epithelium. Blue 585 dotted line denotes the border of the LF. Lower panel is a magnification of the squared area 586 in the upper panel.
- (F, G) Mucosal epithelium and connective tissue fibers stained using PSR. Panel F shows
 the mucosa with epithelium that does not cover the DLT. Panel G shows the mucosa with

589	epithelium covering the DLT. Upper and lower panels show optimal and polarized
590	conditions, respectively. Black and white dotted lines indicate the border between the lamia
591	propria and the epithelium.
592	Scale bars = $100 \ \mu m$.
593	GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, DLT: diffuse

- 594 lymphoid tissue, IHC: immunohistochemistry, H&E: hematoxylin & eosin, LF: lymphatic
- 595 follicle, LT: lymphoid tissue, PSR: picrosirius red

597 Figure 6. Ultrastructure of mucosal epithelium covering the GOALT of the cow VV.

- 598 (A) VV mucosa stained in whole-mount with hematoxylin. GOALTs can be observed as
- 599 hematoxylin-positive spots. Scale bars = 1 mm.
- 600 (**B**) VV mucosa examined using SEM. The same area as in panel A was examined. Scale 601 bars = $100 \mu m$.
- 602 White arrow heads indicate GOALTs.
- 603 (C, D) High-magnified areas squared in panel B. Black arrow heads indicate migrating
- 604 lymphocytes from the GOALT to the lumen of the VV. Red arrow represents small 605 protrusions that could be lymphocytes. Scale bars = $10 \mu m$.
- 606 (E) VV mucosa on large GOALTs examined using SEM. Red arrow represents small 607 protrusions that could be lymphocytes. Scale bars = $100 \mu m$.
- 608 (F) High-magnified area of panel E. High-magnified areas squared in panel E. Black
- arrowhead indicates lymphocytes. Scale bars = $10 \mu m$.
- 610 (G) High-magnified area of panel F. High-magnified areas squared in panel F. Yellow
- 611 arrowhead indicates erythrocytes. Scale bars = $2 \mu m$.
- 612 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, SEM: scanning613 electron microscopy
- 614
- Figure 7. Schematic illustration of GOALTs in the mucosa of the VV and EPGOs in
 cows.

617 GOALTs consist of various shaped LFs and DLTs that localize in different regions in the 618 lamina propria of the VV. Both LTs contain B-cells, T-cells, macrophages, and APCs. 619 B-cells are dominant. Germinal centers are frequently observed in the center of the LF. 620 High endothelial venules also appeared between LTs and surrounding connective tissues in 621 the lamina propria. Moreover, collagenous fibers are stretched across the lamina propria, whereas reticular fibers are mainly observed in DLTs rather than in LFs. Characteristically, 622 623 the epithelium covering GOALTs is partially or completely disrupted by the invasion of 624 immune cells composing LFs and DLTs. These GOALTs are arranged as ring in the VV 625 and EPGO mucosa and named the "genital lymphoid ring." 626 GOALT: genital organ-associated lymphoid tissue, VV: vaginal vestibule, DLT: diffuse 627 lymphoid tissue, LF: lymphatic follicle, LT: lymphoid tissue, APCs: antigen-presenting cells, EPGOs: external parts of genital organs 628

629

630

Highlights

- Whole-mount staining revealed lymphoid tissues on the mucosa of vaginal vestibule
- Lymphoid tissues showed two types: lymphatic nodule or diffuse lymphoid tissue
- Disrupted epithelium allowed passage for immune cells into the intraluminal space

Declarations of competing interest

The authors declare that they have no known competing financial interest or personal relationships that could have influenced in any way the work reported in this paper.











(%)

5

0

B cell

T cell

Macrophage

APC





С

D









