



Title	Simultaneous quantification of imidacloprid and its metabolites in tissues of mice upon chronic low-dose administration of imidacloprid
Author(s)	Nimako, Collins; Ikenaka, Yoshinori; Akoto, Osei; Fujioka, Kazutoshi; Taira, Kumiko; Arizono, Koji; Kato, Keisuke; Takahashi, Keisuke; Nakayama, Shouta M. M.; Ichise, Takahiro; Ishizuka, Mayumi
Citation	Journal of Chromatography A, 1652, 462350 <a href="https://doi.org/10.1016/j.chroma.2021.462350">https://doi.org/10.1016/j.chroma.2021.462350</a>
Issue Date	2021-08-30
Doc URL	<a href="http://hdl.handle.net/2115/90325">http://hdl.handle.net/2115/90325</a>
Rights	© <2021>. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Rights(URL)	<a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Highlights (7).pdf (Highlights)



[Instructions for use](#)

### ***Highlights***

- ❖ LC-MS/MS method was developed for multi-detection of imidacloprid compounds.
- ❖ Imidacloprid and its metabolites accumulated in blood, brain and testis of mice.
- ❖ IMI-olefin was the most predominant metabolite detected in mice tissues
- ❖ *N*-desnitro-IMI showed high accumulation trends in mice brain and testis
- ❖ Testis, inguinal white adipose tissue and kidney had high imidacloprid burden