



Title	Sulfuricystis multivorans gen. nov., sp. nov. and Sulfuricystis thermophila sp. nov., facultatively autotrophic sulfur-oxidizing bacteria isolated from a hot spring, and emended description of the genus Rugosibacter
Author(s)	Kojima, Hisaya; Watanabe, Miho; Miyata, Naoyuki; Fukui, Manabu
Citation	Archives of Microbiology, 204(9), 595 https://doi.org/10.1007/s00203-022-03186-0
Issue Date	2022-09-02
Doc URL	http://hdl.handle.net/2115/90337
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature 's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: https://doi.org/10.1007/s00203-022-03186-0
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Main_ver221003.pdf



[Instructions for use](#)

1

221003

2 ***Sulfuricystis multivorans* gen. nov., sp. nov. and *Sulfuricystis***
3 ***thermophila* sp. nov., facultatively autotrophic sulfur-oxidizing**
4 **bacteria isolated from a hot spring, and emended description**
5 **of the genus *Rugosibacter***

6

7 Hisaya Kojima*^{1,2}, Miho Watanabe², Naoyuki Miyata² and Manabu Fukui¹

8

9 1. The Institute of Low Temperature Science, Hokkaido University. Kita-19, Nishi-8,
10 Kita-ku, Sapporo 060-0819, Japan

11 2. Department of Biological Environment, Faculty of Bioresource Sciences, Akita
12 Prefectural University. Shimo-Shinjyo Nakano, Akita, Akita, 010-0195 Japan

13

14

15

16 *Corresponding author

17 Tel/fax number: +81 11 706 5460

18 E-mail: kojimah@lowtem.hokudai.ac.jp

19 The Institute of Low Temperature Science, Hokkaido University. Kita-19, Nishi-8,
20 Kita-ku, Sapporo 060-0819, Japan

21

22

23

24 **Abstract**

25 Strains J5B^T and M52^T are facultatively autotrophic sulfur-oxidizing bacteria isolated
26 from a microbial mat from a hot spring. They were isolated and partially
27 characterized in previous studies, as facultative anaerobes which use nitrate as
28 electron acceptor. In this study, additional characterizations were made to determine
29 their taxonomic status. In both strains, major cellular fatty acids were C_{16:1} (C_{16:1}ω7c
30 and/or C_{16:1}ω6c) and C_{16:0}. Their chemolithoautotrophic growth was supported by
31 thiosulfate and elemental sulfur. They used some organic acids as growth substrates.
32 Their 16S rRNA gene sequences indicated the highest sequence identities to species
33 in the family *Sterolibacteriaceae*, but the identities were 95% or lower. Phylogenetic
34 analysis indicated that these strains do not belong to any existing genera. Values of
35 average nucleotide identity and digital DNA–DNA hybridization between strains
36 J5B^T and M52^T were 87.93 % and 34.3 %, respectively. On the basis of phenotypic
37 and genomic characteristics, *Sulfuricystis multivorans* gen. nov. sp. nov., and
38 *Sulfuricystis thermophila* sp. nov. are proposed, with type strains of J5B^T and M52^T,
39 respectively. An emended description of the genus *Rugosibacter* is also proposed, for
40 its reclassification to the family *Sterolibacteriaceae*.

41

42

43 **Introduction**

44 The family *Sterolibacteriaceae* was proposed to encompass six genera, as a part of
45 major reconstruction of the order *Nitrosomonadales* (Boden et al., 2017). According
46 to the List of Prokaryotic Names with Standing in Nomenclature (as of 29 March 2022),
47 the family still comprises six genera, *Sterolibacterium*, *Denitratisoma*,
48 *Methyloversatilis*, *Georgfuchsia*, *Sulfuritalea* and *Sulfurisoma*. Before the
49 reclassification, some of these genera had been regarded as members of the family
50 *Rhodocyclaceae* in the order *Rhodocyclales*. Shortly before establishment of the
51 family *Sterolibacteriaceae*, a genus was proposed in the family *Rhodocyclaceae*, with
52 the name of *Rugosibacter* (Corteselli et al., 2017). In that study, phylogenetic analysis
53 of the 16S rRNA gene indicated that *Rugosibacter* belongs to a clade consisting of the
54 six genera mentioned above. The genus *Rugosibacter* was not taken into consideration
55 in the proposal of *Sterolibacteriaceae*, and it is still placed in the family
56 *Rhodocyclaceae* as noted in the original description. The close relatedness between
57 *Rugosibacter* and genera of *Sterolibacteriaceae* has also been indicated by a
58 phylogenetic analysis of concatenated ribosomal proteins (Okubo & Takami, 2021).

59 The original description of family *Sterolibacteriaceae* states that the family is
60 circumscribed on the basis of 16S rRNA gene sequences, and includes physiologically

61 diverse organisms (Boden et al., 2017). It refers to methylotrophy, autotrophy and
62 anaerobic respiration (with nitrate, ferric iron or manganic manganese), as varied
63 metabolism observed in the family. As other notable functions, anaerobic degradation
64 of aromatic compounds (Weelink et al., 2009; Sperfeld et al., 2019) and arsenate
65 respiration (Watanabe et al., 2017) have also been demonstrated in species belonging
66 to this family.

67 Besides the organisms mentioned above, some strains have been isolated and
68 characterized as members of this family. *Sterolibacteriaceae* bacterium strain J5B^T is
69 a facultatively autotrophic sulfur-oxidizing bacterium, isolated from a microbial mat
70 collected in a hot spring in Japan (Watanabe et al, 2019). As for strain J5B^T, some
71 physiological characteristics have been reported, along with genomic characteristics
72 related to sulfur oxidation (Watanabe et al, 2019). The closest relative of strain J5B^T
73 is also sulfur oxidizer, isolated from the same microbial mat. The sulfur-oxidizing
74 bacterium, strain M52^T, can oxidize arsenite anaerobically (Ospino et al., 2019). The
75 previous studies reported complete genome sequences of these two strains. The
76 genome sequences have been publicized and incorporated in the genome taxonomy
77 database (GTDB) (Parks et al., 2018). In the GTDB release 06-RS202, they are both
78 classified in a genus-level taxon (UBA2250) which has no species with validly

79 published name and regarded as representatives of independent species.

80 In this study, additional characterizations of strains J5B^T and M52^T were made
81 to determine their taxonomic status in the family *Sterolibacteriaceae*.

82

83 **Materials and methods**

84

85 Isolation and maintenance of strains

86 The strains J5B^T and M52^T were isolated from a microbial mat obtained from a hot
87 spring in Japan (42° 57' 53" N 141° 09' 47" E). They were isolated into pure cultures
88 under nitrate-reducing conditions, by using bicarbonate and thiosulfate as sole carbon
89 source and electron donor (Watanabe et al, 2019; Ospino et al., 2019). The strains were
90 maintained in the laboratory with a bicarbonate-buffered medium referred to as “S5
91 medium” (Kojima et al., 2017), supplemented with 10 mM nitrate. Purity and identity
92 of the cultures were periodically checked by microscopic observation and direct
93 sequencing of the 16S rRNA gene.

94

95 Cellular fatty acid analysis

96 For fatty acid analysis, cells of strains J5B^T and M52^T were grown at 45°C,
97 in a medium consisting of the following constituents (l⁻¹): 1 g NaNO₃, 0.5 g yeast

98 extract, 0.5 g Casamino acids, 0.5 g disodium fumarate, 0.5 g disodium succinate, and
99 50 µg cyanocobalamin. Headspace of the culturing bottles was filled with N₂ gas, and
100 pH of the medium was adjusted to 7.0. Their cellular fatty acid profiles were obtained
101 with the Sherlock Microbial Identification System (MIDI) version 6.0 (database;
102 TSBA6). As for J5B^T, cells grown in the S5 medium under oxic conditions were also
103 subjected to the same analysis.

104

105 Physiological characterizations

106 Effects of temperature on growth of strain M52^T were examined by
107 culturing at various temperatures (15, 18, 22, 25, 28, 30, 32, 35, 37, 40, 42, 45, 48, 50,
108 53, 55, 57 and 60°C), in the medium used for maintenance. The other tests for
109 phenotypic characterization were all conducted at 45°C.

110 Chemolithoautotrophic growth under nitrate-reducing conditions was tested
111 in the medium used for the maintenance, by replacing thiosulfate with one of electron
112 donors listed below; elemental sulfur (0.5 g l⁻¹), sulfide (2 mM), tetrathionate (10 mM)
113 and hydrogen gas (H₂/N₂/CO₂ 50:40:10 v/v/v; 200 kPa in total pressure). Utilization
114 of organic substrates for heterotrophic growth was tested in 10 mM MOPS-NaOH
115 buffer (pH 7.0), supplemented with NaNO₃ (1 g l⁻¹), yeast extract (0.1 g l⁻¹), Casamino
116 acids (0.1 g l⁻¹) and cyanocobalamin (50 µg l⁻¹). The medium was dispensed in closed

117 culture bottles and the headspace was filled with N₂ gas. One of the following organic
118 substrates were added to the medium (mM); pyruvate (5), lactate (5), acetate (5),
119 propionate (2.5), succinate (2.5), fumarate (2.5), malate (2.5), butyrate (2.5), benzoate
120 (2.5), isobutyrate (2.5), methanol (5), ethanol (2.5), formate (5), citrate (5), glucose
121 (2.5), xylose (2.5), phenol (2), *o*-cresol (1), *m*-cresol (1).

122 Effect of pH on growth of strain M52^T was tested with pH-buffered media
123 modified from the medium used for the fatty acid analysis. The media were buffered
124 with 10 mM of MES, MOPS or Tricine, and adjusted to varying pH (with 0.1- or 0.2-
125 unit intervals) by adding NaOH. The tested pH were as follows: 5.4–7.3 with MES;
126 6.4–8.0 with MOPS; 7.0–9.0 with Tricine.

127

128 Genomic characterization

129 The complete genome sequences of strains J5B^T and M52^T were obtained in
130 the previous studies (Watanabe et al, 2019; Ospino et al., 2019). As overall genome
131 relatedness indices between them, values of average nucleotide identity (ANI) and
132 digital DNA–DNA hybridization (dDDH) were calculated by using online tools as
133 follows. The ANI was computed by ANI calculator available in EzBioCloud, based on
134 the OrthoANIu algorithm (Yoon et al., 2017). The dDDH value was calculated using
135 Genome-to-Genome Distance Calculator (GGDC) provided by DSMZ (Meier-

136 Kolthoff et al., 2013), applying the formula 2. To evaluate genomic similarity with
137 related bacteria, values of average amino acid identity (AAI) were calculated with
138 EzAAI version 1.0 (Kim et al. 2021).

139 The genome sequences were annotated with DFAST (Tanizawa et al., 2018),
140 to identify protein coding sequences and RNA genes. Genes for sulfur oxidation were
141 identified as described previously (Watanabe et al, 2019).

142 The full sequences of the 16S rRNA gene identified in the genomes were
143 subjected to the blastn search at NCBI against the nucleotide collection (nr/nt) database.
144 Phylogenetic analyses were conducted using the program MEGA version 11 (Tamura
145 et al., 2021). The 16S rRNA gene sequences were aligned with reference sequences,
146 using the MUSCLE algorithm. The references sequences were those of type strains of
147 species with validly published names in the order *Nitrosomonadales* and the genus
148 *Rugosibacter*. The best model for calculation of genetic distances was selected by
149 using the model selection tool in MEGA, as the model with the lowest score of the
150 Bayesian Information Criterion (BIC).

151 Genome-based phylogenetic analysis of family *Sterolibacteriaceae* was
152 conducted by using ezTree pipeline (Wu, 2018). The genome sequences of strains J5B^T
153 and M52^T were input into the pipeline, along with those of the type strains in the family

154 and *Rugosibacter aromaticivorans* Ca6^T. *Thiobacillus thioparus* DSM 505^T was also
155 included in the analysis as an outgroup. In the pipeline with default settings, single-
156 copy marker genes were automatically identified, and their coding sequences were
157 separately aligned. The resulting alignments were concatenated to generate a
158 maximum-likelihood tree.

159

160 **Results and Discussion**

161

162 Physiological and chemotaxonomic characteristics

163 The fundamental characteristics of strains J5B^T and M52^T are presented in
164 the species descriptions, and some of them are summarized in Table 1. Their cells
165 grown in the maintenance medium were both motile rods, with width of 0.4–0.5 μm.
166 The cell lengths of strain J5B^T and M52^T were 0.8–2.0 μm and 1.8–3.2 μm,
167 respectively. The strain M52^T grew at 18–55°C, with optimum growth at 50°C. Its
168 pH range for growth was 5.5–8.6, and optimum pH was 6.7–6.9. These
169 characteristics of strain J5B^T have been reported previously.

170 The strain M52^T was isolated and maintained under thiosulfate-oxidizing
171 conditions, without organic carbon source. Chemolithoautotrophic growth of the
172 strain was also supported by electron donors of tetrathionate and elemental sulfur. On

173 the other hand, hydrogen gas and sulfide did not support growth strain M52^T. As
174 reported previous, strain M52^T can grow on acetate and lactate. The culturing
175 experiments in this study revealed that the following organic acids are also used as
176 growth substrates; pyruvate, propionate, succinate, fumarate, malate, and butyrate
177 and isobutyrate. None of tested sugars, alcohols or cresols supported growth of strain
178 M52^T.

179 The cellular fatty acid profiles of strains J5B^T and M52^T are shown in Table
180 S1. For strain J5B^T, effects of growth conditions on fatty acid composition was
181 investigated by analyzing cells grown autotrophically and heterotrophically. Under
182 both conditions, C_{16:0} was the most abundant fatty acid, followed by summed feature
183 3 (C_{16:1} ω 7*c* and/or C_{16:1} ω 6*c*). These two major components accounted for 78–80% of
184 total, irrespective of the culture conditions. The third most abundant fatty acid
185 differed depending on the growth conditions. That was cyclo-C_{17:0} under the
186 heterotrophic condition, whereas that under the autotrophic condition was summed
187 feature 8 (C_{18:1} ω 7*c* and/or C_{18:1} ω 6*c*). The fatty acid profile of strain M52^T was
188 similar to that of J5B^T grown under the same conditions, sharing the major fatty
189 acids (Table S1).

190

191 Genomic characteristics

192 Complete genome sequences of strains J5B^T and M52^T were independently
193 obtained in the previous studies. The values of ANI and dDDH between them were
194 calculated to be 87.93 % and 34.3 %, respectively. These values are lower than
195 thresholds for species delineation (Richter & Rosselló-Móra, 2009; Meier-Kolthoff et
196 al., 2013), suggesting that strains J5B^T and M52^T are representatives of different
197 species.

198 As reported previously, strains J5B^T and M52^T are chemolithoautotrophs
199 growing on thiosulfate. According to these physiological properties, genes for sulfur
200 oxidation and carbon fixation were identified in their genomes. As those for sulfur
201 oxidation, genes encoding proteins involved in the Sox-Dsr-Soe pathway were
202 identified in the genome of strain M52^T. The Sox-Dsr-Soe pathway is one of the
203 three core pathways of sulfur oxidation, and shared by sulfur oxidizers in the family
204 *Sterolibacteriaceae*, i.e., *Sulfuritalea hydrogenivorans* sk43H^T, *Sulfurisoma*
205 *sediminicola* BSN1^T and strain J5B^T (Watanabe et al. 2019). These sulfur oxidizers
206 seem to fix inorganic carbon via the Calvin–Benson–Bascham cycle. They have
207 genes encoding key enzyme of the cycle, ribulose-1,5-bisphosphate
208 carboxylase/oxygenase (RuBisCO). They all have the *cbbM* gene which encodes

209 form II RuBisCO, whereas the *cbbLS* genes encoding form I RuBisCO were
210 identified only in genomes of strain M52^T and *Sulfurisoma sediminicola* BSN1^T.
211 *Sulfuritalea hydrogenivorans* sk43H^T is known to have key genes for anaerobic
212 aromatics degradation and arsenate respiration (Sperfeld et al., 2019; Watanabe et al.
213 2017). These genes were not identified in the genomes of strains J5B^T and M52^T.

214

215 Taxonomic assignment

216 The strains J5B^T and M52^T have two copies of the 16S rRNA gene in
217 their genomes, respectively. The copies of strain J5B^T are both 1544 nt in length, and
218 there are five mismatches between their sequences. Those of strain M52^T are
219 different in length, 1544 nt and 1534 nt respectively. These sequences indicated the
220 highest sequence identities to species belonging to family *Sterolibacteriaceae*, but
221 the identities were 95% or lower. The sequence identities between strains J5B^T and
222 M52^T ranged 98.3–98.8%. The sequence variations between the copies of each strain
223 did not affect the phylogenetic analysis described below, because they were located
224 at positions with gaps and thus excluded from the calculation.

225 Phylogenetic positions of strains J5B^T and M52^T within the order
226 *Nitrosomonadales* are shown in Fig. 1, as the maximum likelihood tree of the 16S

227 rRNA gene (full version is shown in Figure S1). The tree indicated that the strains
228 belong to the family *Sterolibacteriaceae* but not to any existing genera. This means
229 that a novel genus should be created to accommodate strains J5B^T and M52^T. The
230 tree also indicate that the genus *Rugosibacter* should be reclassified to the family
231 *Sterolibacteriaceae*, at least at present. These conclusions did not conflict with the
232 phylogenetic analysis based on 92 marker genes identified in the genomes (Fig. S2).

233 As an attempt to evaluate reasonability of genus-level classification, AAI
234 values were calculated for strains J5B^T and M52^T, against all type strains within the
235 family *Sterolibacteriaceae* (Table 1). The calculated values ranged from 63.2 to
236 71.5%. These results do not provide solid support for creation of a new genus,
237 because AAI values between different species in a same genus are 60–80% in many
238 cases (Luo et al., 2014). In the family *Sterolibacteriaceae*, AAI may not work as a
239 key criterion for genus-level classification. The AAI values were also calculated in
240 all combinations of the type strains in this family (Table S2). The values between
241 *Sterolibacteriaceae* strains from different genera were within the range of 60–80%,
242 in all combinations.

243

244 **Conclusion**

245 On the basis of the results of the previous and present studies, strain J5B^T is
246 proposed as the type strain of a novel species of a new genus, with the name
247 *Sulfuricystis multivorans* gen. nov., sp. nov. In addition, strain M52^T is proposed as
248 the type strain of another species of the novel genus, with the name *Sulfuricystis*
249 *thermophila* sp. nov. Further, emended description of the genus *Rugosibacter* is also
250 proposed to indicate its affiliation to the family *Sterolibacteriaceae*.

251

252 **Description of *Sulfuricystis* gen. nov.**

253 *Sulfuricystis* (Sul.fu.ri.cys'tis. L. neut. n. *sulfur*, sulfur; Gr. fem. n. *kystis*, a bag; N.L.
254 fem. n. *Sulfuricystis*, sulfur-oxidizing bag).

255 Grows by oxidation of sulfur compounds. Facultatively anaerobic and neutrophilic.
256 Gram-stain-negative. Major cellular fatty acids are C_{16:0} and C_{16:1} (C_{16:1}ω7c and/or
257 C_{16:1}ω6c). Phylogenetically, belongs the family *Sterolibacteriaceae*. The type species
258 is *Sulfuricystis multivorans*.

259

260 **Description of *Sulfuricystis multivorans* sp. nov.**

261 *Sulfuricystis multivorans* (mul.ti.vo'rans. L. masc. adj. *multus*, many; L. pres. part.

262 vorans, devouring, eating; N.L. part. adj. *multivorans*, devouring various substrates).

263 In addition to properties listed in the genus description, cells are rod-shaped, 0.8–2.0

264 µm long and 0.4–0.5 µm wide. Uses oxygen and nitrate as electron acceptor. Under

265 nitrate-reducing conditions, grows chemolithoautotrophically on thiosulfate and

266 elemental sulfur, but not on sulfide, tetrathionate, or hydrogen gas. Grows

267 heterotrophically on pyruvate, lactate, acetate, propionate, succinate, fumarate, malate,

268 and butyrate. Does not grow on benzoate, isobutyrate, methanol, ethanol, formate,

269 citrate, glucose, xylose, phenol, *o*-cresol, and *m*-cresol. Temperature range for growth

270 is 28–55°C, with an optimum of 45–50°C. Growth occurs at pH 5.8–8.7, with an

271 optimum of pH 6.7–7.4. G + C content of genomic DNA of the type strain is 62.4

272 mol%.

273 The type strain J5B^T (= BCRC 81316^T = DSM 104688^T = NBRC 112605^T) was

274 isolated from a microbial mat of a hot spring in Japan.

275 The GenBank/EMBL/DDBJ accession numbers for the chromosome and two

276 plasmids of type strain are AP018718 and AP018719-AP018720, respectively.

277

278 **Description of *Sulfuricystis thermophila* sp. nov.**

279 *Sulfuricystis thermophila* (ther.mo'phi.la. Gr. masc. adj. *thermos*, hot; Gr. masc. adj.
280 *philos*, loving; N.L. fem. adj. *thermophila*, heat-loving).

281 In addition to properties listed in the genus description, cells are rod-shaped, 1.8–3.2
282 μm long and 0.4–0.5 μm wide. Uses oxygen and nitrate as electron acceptor. Under
283 nitrate-reducing conditions, grows chemolithoautotrophically on thiosulfate,
284 tetrathionate and elemental sulfur, but not on sulfide or hydrogen gas. Grows
285 heterotrophically on pyruvate, lactate, acetate, propionate, succinate, fumarate, malate,
286 and butyrate and isobutyrate. Does not grow on benzoate, methanol, ethanol, formate,
287 citrate, glucose, xylose, phenol, *o*-cresol, and *m*-cresol. Temperature range for growth
288 is 18–55°C, with an optimum of 50°C. Growth occurs at pH 5.5–8.6, with an optimum
289 of pH 6.6–6.9. G+C content of genomic DNA of the type strain is 63.6 mol%.

290 Type strain M52^T (= BCRC 81317^T = NBRC 114016^T) was isolated from a
291 microbial mat of a hot spring in Japan.

292 The GenBank/EMBL/DDBJ accession number for the complete genome of
293 the type strain is AP019373.

294

295 **Emended description of *Rugosibacter* (Corteselli et al., 2017)**

296 *Rugosibacter* (Ru.go.si.bac'ter. L. adj. *rugosus* wrinkled; N. L. masc. n. *bacter* a rod;
297 N. L. masc. n. *Rugosibacter* a wrinkled rod).

298 Cells are Gram-stain-negative and non-motile. Aerobic. Catalase-negative and
299 oxidase-positive. Heterotrophic growth occurs on organic acids. Predominant fatty
300 acids are summed feature 3 (C_{16:1}ω7c and/or C_{16:1}ω7c) and C_{16:0}. The major polar
301 lipids are phosphatidylethanolamine, phosphatidylglycerol, diphosphatidylglycerol
302 and phospholipid. The major respiratory quinone is ubiquinone-8. Phylogenetically,
303 belongs the family *Sterolibacteriaceae*. The type species is *Rugosibacter*
304 *aromaticivorans*.

305

306 **Acknowledgments**

307 Authors thank Y. Kato and A. Shinohara for technical assistance.

308

309 **Reference**

310 Boden R, Hutt LP, Rae AW (2017) Reclassification of *Thiobacillus aquaesulis* (Wood
311 & Kelly, 1995) as *Annwoodia aquaesulis* gen. nov., comb. nov., transfer of
312 *Thiobacillus* (Beijerinck, 1904) from the *Hydrogenophilales* to the

313 *Nitrosomonadales*, proposal of *Hydrogenophilalia* class. nov. within the
314 'Proteobacteria', and four new families within the orders *Nitrosomonadales* and
315 *Rhodocyclales*. Int J Syst Evol Microbiol 67:1191–1205.
316 <https://doi.org/10.1099/ijsem.0.001927>

317 Corteselli EM, Aitken MD, Singleton DR (2017) *Rugosibacter aromaticivorans* gen.
318 nov., sp. nov., a bacterium within the family *Rhodocyclaceae*, isolated from
319 contaminated soil, capable of degrading aromatic compounds. Int J Syst Evol
320 Microbiol 67: 311–318. <https://doi.org/10.1099/ijsem.0.001622>

321 Doronina NV, Kaparullina EN, Trotsenko, YA (2014) *Methyloversatilis*
322 *thermotolerans* sp. nov., a novel thermotolerant facultative methylotroph isolated
323 from a hot spring. Int J Syst Evol Microbiol 64:158–164
324 <https://doi.org/10.1099/ijms.0.055046-0>

325 Fahrbach M, Kuever J, Meinke R, Kämpfer, P, Hollender J (2006) *Denitratisoma*
326 *oestradiolicum* gen. nov., sp. nov., a 17 β -estradiol-degrading, denitrifying
327 betaproteobacterium. Int J Syst Evol Microbiol 56:1547–1552.
328 <https://doi.org/10.1099/ijms.0.63672-0>

329 Kalyuzhnaya MG, De Marco P, Bowerman S, Pacheco CC, Lara JC, Lidstrom ME,
330 Chistoserdova L (2006) *Methyloversatilis universalis* gen. nov., sp. nov., a novel

331 taxon within the Betaproteobacteria represented by three methylophilic isolates.
332 Int J Syst Evol Microbiol 56:2517–2522. <https://doi.org/10.1099/ijs.0.64422-0>

333 Kim D, Park S, Chun J (2021) Introducing EzAAI: a pipeline for high throughput
334 calculations of prokaryotic average amino acid identity. J Microbiol 59:476–480.
335 <https://doi.org/10.1007/s12275-021-1154-0>

336 Kojima H, Fukui M (2011) *Sulfuritalea hydrogenivorans* gen. nov., sp. nov., a
337 facultative autotroph isolated from a freshwater lake. Int J Syst Evol Microbiol
338 61:1651–1655. <https://doi.org/10.1099/ijs.0.024968-0>

339 Kojima H, Fukui M (2014) *Sulfurisoma sedimicola* gen. nov., sp. nov., a facultative
340 autotroph isolated from a freshwater lake. Int J Syst Evol Microbiol 64:1587–1592.
341 <https://doi.org/10.1099/ijs.0.057281-0>

342 Kojima H, Watanabe M, Fukui M (2017) *Sulfurivermis fontis* gen. nov., sp. nov., a
343 sulfur-oxidizing autotroph, and proposal of *Thioprofundaceae* fam. nov. Int J Syst
344 Evol Microbiol 67:3458–3461. <https://doi.org/10.1099/ijsem.0.002137>

345 Luo C, Rodriguez-r LM, Konstantinidis KT (2014) MyTaxa: an advanced taxonomic
346 classifier for genomic and metagenomic sequences. Nucl Acids Res 42:e73–e73
347 <https://doi.org/10.1093/nar/gku169>

348 Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M (2013) Genome sequence-based
349 species delimitation with confidence intervals and improved distance functions.
350 BMC Bioinformatics 14:60–14. <https://doi.org/10.1186/1471-2105-14-60>

351 Okubo T, Takami H (2021) Metabolic potential of the imperfect denitrifier *Candidatus*
352 *Desulfobacillus denitrificans* in an anammox bioreactor *MicrobiologyOpen* 10:
353 e1227 <https://doi.org/10.1002/mbo3.1227>

354 Ospino MC, Kojima H, Fukui M (2019) Arsenite oxidation by a newly isolated
355 betaproteobacterium possessing *arx* genes and diversity of the *arx* gene cluster in
356 bacterial genomes. *Front Microbiol* 12:10.
357 <https://doi.org/10.3389/fmicb.2019.01210>

358 Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA,
359 Hugenholtz P (2018) A standardized bacterial taxonomy based on genome
360 phylogeny substantially revises the tree of life. *Nat Biotechnol* 36:996–1004.
361 <https://doi.org/10.1038/nbt.4229>

362 Richter M, Rosselló-Móra R (2009) Shifting the genomic gold standard for the
363 prokaryotic species definition. *Proc Natl Acad Sci* 106:19126–19131.
364 <https://doi.org/10.1073/pnas.0906412106>

365 Smalley NE, Taipale S, De Marco P, Doronina NV, Kyrpides N, Shapiro N, Woyke T,
366 Kalyuzhnaya MG (2015) Functional and genomic diversity of methylotrophic
367 *Rhodocyclaceae*: description of *Methyloversatilis discipulorum* sp. nov. Int J Syst
368 Evol Microbiol 65:2227–2233. <https://doi.org/10.1099/ijs.0.000190>

369 Sperfeld M, Diekert G, Studenik S (2019) Anaerobic aromatic compound degradation
370 in *Sulfuritalea hydrogenivorans* sk43H. FEMS Microbiol Ecol 95:fiy199. doi:
371 [10.1093/femsec/fiy199](https://doi.org/10.1093/femsec/fiy199)

372 Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics
373 Analysis version 11. Mol Biol Evol 38:3022–3027.
374 <https://doi.org/10.1093/molbev/msab120>.

375 Tanizawa Y, Fujisawa T, Nakamura Y (2018) DFAST: a flexible prokaryotic genome
376 annotation pipeline for faster genome publication. Bioinformatics 34:1037–1039.
377 <https://doi.org/10.1093/bioinformatics/btx713>

378 Tarlera S, Denner EB (2003) *Sterolibacterium denitrificans* gen. nov., sp. nov., a novel
379 cholesterol-oxidizing, denitrifying member of the β -Proteobacteria. Int J Syst Evol
380 Microbiol 53:1085–1091. <https://doi.org/10.1099/ijs.0.02039-0>

381 Watanabe T, Miura A, Iwata T, Kojima H, Fukui M (2017) Dominance of *Sulfuritalea*

382 species in nitrate-depleted water of a stratified freshwater lake and arsenate
383 respiration ability within the genus. *Env Microbiol Rep* 9:522–527.
384 <https://doi.org/10.1111/1758-2229.12557>

385 Watanabe T, Kojima H, Umezawa K, Hori C, Takasuka ET, Kato Y, Fukui M (2019)
386 Genomes of neutrophilic sulfur-oxidizing chemolithoautotrophs representing 9
387 proteobacterial species from 8 genera *Front Microbiol* 10:316. [https://doi.org/](https://doi.org/10.3389/fmicb.2019.00316)
388 [10.3389/fmicb.2019.00316](https://doi.org/10.3389/fmicb.2019.00316)

389 Weelink SA, van Doesburg W, Saia FT, Rijpstra WI, Röling WF, Smidt H, Stams AJ
390 (2009) A strictly anaerobic betaproteobacterium *Georgfuchsia toluolica* gen. nov.,
391 sp. nov. degrades aromatic compounds with Fe(III), Mn(IV) or nitrate as an
392 electron acceptor. *FEMS Microbiol Ecol* 70:575–585.
393 <https://doi.org/10.1111/j.1574-6941.2009.00778.x>

394 Yoon S-H, Ha S-M, Lim J, Kwon S, Chun J (2017) A large-scale evaluation of
395 algorithms to calculate average nucleotide identity. *Antonie van Leeuwenhoek*
396 110:1281–1286. <https://doi.org/10.1007/s10482-017-0844-4>

397 Wu YW (2018) ezTree: an automated pipeline for identifying phylogenetic marker
398 genes and inferring evolutionary relationships among uncultivated prokaryotic
399 draft genomes. *BMC genomics* 19:921. <https://doi.org/10.1186/s12864-017-4327->

400 9

401

402 **Statements and Declarations**

403 This study was and supported by JSPS KAKENHI, Grant Number 20K12215.

404 The authors have no relevant financial or non-financial interests to disclose.

405

406 **Figure legends**

407 Fig. 1. Maximum likelihood tree of the 16S rRNA gene sequences, showing
408 phylogenetic positions of strains J5B^T and M52^T within the order *Nitrosomonadales*.
409 The Kimura 2-parameter model with a discrete gamma distribution and invariable
410 sites was used as the best model with the lowest BIC. All positions containing gaps
411 and missing data were eliminated and there were 1,279 positions in the final dataset.
412 Numbers on nodes represent percentage values of 1,000 bootstrap resampling (values
413 greater than 50 are shown). Names of all strains included in the analysis are shown in
414 full version of the tree provided as Figure S1.

Table 1. Differential characteristics of strains J5B^T, M52^T and type strains of species in the family *Sterolibacteriaceae*. Strains: 1, J5B^T (Watanabe et al., 2019); 2, M52^T (Ospino et al., 2019; this study); 3, *Sterolibacterium denitrificans* Chol-1S^T (Tarlera & Denner, 2003); 4, *Denitratisona oestradiolicum* AcBE2-1^T (Fahrbach et al., 2006); 5, *Methyloversatilis universalis* FAM5^T (Kalyuzhnaya et al., 2006); 6, *M. thermotolerans* 3t^T (Doronina et al., 2014); 7, *M. discipulorum* FAM1^T (Smalley et al., 2015); 8, *Georgfuchsia toluolica* G5G6^T (Weelink et al., 2009); 9, *Sulfuritalea hydrogenivorans* sk43H^T (Kojima & Fukui, 2011); 10, *Sulfurisoma sedimnicola* BSN1^T (Kojima & Fukui, 2014); 11, *Rugosibacter aromaticivorans* Ca6^T (Corteselli et al., 2017). Data were retrieved from respective references, except for AAI values calculated in this study.

	1	2	3	4	5	6	7	8	9	10	11
Optimum temp.	45–50	50	30-32	28-30	37	30-37	25–37	25-30	25	30-32	30-34
Temp. range	28–55	18–55	15-35	4-38	10-42	10–45	7-37	20-37	8-32	8-34	20-35
Optimum pH	6.7–7.4	6.6–6.9	7.0	7.0-7.2	8.0	7.0-7.5	6.6-8.0	7.3	6.7-6.9	7.8-8.1	6.5
pH range	5.8–8.7	5.5–8.6	5.8-8.0	6.4-8.5	6.5-9.0	6.5-8.5	8.8-8.0	6.6-9.0	6.4-7.6	6.8–8.8	6.5-7.5
Electron acceptor											
oxygen	+	+	+	+	+	+	+	-	+	+	+
nitrate	+	+	+	+	-	+	+	+	+	+	-
Electron donor											
methanol	-	-	ND	-	+	+	+	-	-	-	ND
ethanol	-	-	-	-	+	+	+	-	-	-	ND
thiosulfate	+	+	ND	-	ND	ND	ND		+	+	ND
hydrogen	-	-	ND	ND	ND	ND	ND	-	+	+	ND
benzoate	-	-	-	-	ND	ND	ND	-	+	-	ND
phenol	-	-	-	-	ND	ND	+	+	-	-	ND
m-cresol	-	-	ND	-	ND	ND	ND	+	ND	-	ND
isobutyrate	-	+	-	+	ND	ND	ND	ND	+	-	ND
pyruvate	+	+	-	+	-	ND	+	-	+	+	ND
lactate	+	+	-	+	ND	ND	ND	-	+	+	-
acetate	+	+	+	+	-	+	+	-	+	+	-
propionate	-	+	+	+	ND	ND	ND	-	+	+	-
succinate	+	+	-	+	+	+	+	-	+	+	-
fumarate	+	+	-	+	ND	+	ND	-	+	+	ND
malate	+	+	-	ND	+	ND	+	-	+	+	ND
butyrate	+	+	+	+	ND	ND	ND	-	+	-	ND
glucose	-	-	-	-	+	+	+	-	-	-	ND
xylose	-	-	-	-	ND	ND	ND	-	-	-	ND
citrate	-	-	-	-	+	-	ND	-	-	-	-
formate	-	-	-	-	+	+	+	ND	-	-	+
AAI to J5B ^T (%)	100	89.9	66.4	67.4	63.5	63.2	63.4	67.0	69.1	71.1	67.0
AAI to M52 ^T (%)	89.9	100	66.2	68.1	63.6	63.6	63.5	67.3	69.6	71.5	67.8

