

# HOKKAIDO UNIVERSITY

Title	Catchment geology preconditions spatio-temporal heterogeneity of ecosystem functioning in forested headwater streams
Author(s)	Tolod, Janine Rodulfo; Negishi, Junjiro N.; Ishiyama, Nobuo; Alam, Md Khorshed; Rahman, Mirza A. T. M. Tanvir; Pongsivapai, Pongpet; Gao, Yiyang; Sueyoshi, Masanao; Nakamura, Futoshi
Citation	Hydrobiologia, 849, 4307-4324 https://doi.org/10.1007/s10750-022-04992-9
Issue Date	2022-09-07
Doc URL	http://hdl.handle.net/2115/90344
Rights	This is a post-peer-review, pre-copyedit version of an article published in Hydrobiologia. The final authenticated version is available online at:https://doi.org/10.1007/s10750-022-04992-9
Туре	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Tolod et al. MAIN TEXT_final revision.pdf



1	A manuscript prepared as a submission to <u>Hydrobiologia</u>
2	
3	Title: Catchment geology preconditions spatio-temporal heterogeneity of ecosystem functioning in forested
4	headwater streams
5	
6	Janine Rodulfo Tolod1*, Junjiro N. Negishi2*, Nobuo Ishiyama3, Md. Khorshed Alam1, Mirza A.T.M.
7	Tanvir Rahman <sup>1,4</sup> , Pongpet Pongsivapai <sup>1,7</sup> , Gao Yiyang <sup>1</sup> , Masanao Sueyoshi <sup>5</sup> , Futoshi Nakamura <sup>6</sup>
8	<sup>1</sup> Graduate School of Environmental Science, Hokkaido University, Kita 10 Nishi 5 Sapporo, Hokkaido
9	060-0810, Japan
10	<sup>2</sup> Faculty of Environmental Earth Science, Hokkaido University, Kita 10 Nishi 5 Sapporo, Hokkaido 060-
11	0810, Japan
12	<sup>3</sup> Hokkaido Research Organization, Koshunaicho, Bibai, Hokkaido 079-0198, Japan
13	<sup>4</sup> Department of Environmental Sciences, Jahangirnagar University, Savar, Dhaka 1342, Bangladesh
14	<sup>5</sup> National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba City, Ibaraki 3305-8506, Japan
15	<sup>6</sup> Faculty of Agriculture, Hokkaido University, Kita 9 Nishi 9 Sapporo, Hokkaido 060-8589, Japan
16 17 18	<sup>7</sup> Department of Soil Science, Faculty of Agriculture at Kamphaeng Saen, Kasetsart University, Kanphaeng Saen Campus, Nakhon Pathom, 73140, Thailand
19	
20	corresponding authors:
21	Janine R. Tolod; e-mail: jrtolod@gmail.com,
22	Junjiro N. Negishi; e-mail: negishi@ees.hokudai.ac.jp
23	
24	Abstract
25	Catchment geology can affect water chemistry and groundwater influence, eventually affecting
26	macroinvertebrate communities, but its effects on stream functions such as leaf decomposition have been
27	scarcely investigated. To understand the effects of geology on leaf decomposition, we conducted leaf litter

28 experiments in streams with volcanic and non-volcanic substrata using fine and coarse mesh bags. Volcanic 29 spring-fed streams showed lower temperature in summer and higher temperature in winter (with 30 temperature difference being more pronounced later in incubation) than non-volcanic streams. 31 Macroinvertebrate communities captured inside coarse litter bags differed in the two stream types in both 32 seasons, mainly because of shredder communities. Shredder abundance and biomass were higher in 33 volcanic streams in both seasons. Geology-dependent temperature influenced microbe-mediated 34 decomposition in both seasons, with total phosphorus as an additional driver in winter. Summer temperature 35 was associated with an overall positive effect on the abundance of shredders, which affected invertebrate-36 mediated decomposition, but this was not evident in winter. Shredder activity in volcanic streams 37 compensated for temperature-dependent microbial activity resulting in an overall balance in leaf 38 decomposition. Spring-fed systems are valuable ecosystems, particularly for cold-adapted species. Thus, 39 understanding these understudied ecosystems will significantly aid in their appropriate conservation.

40 Keywords: leaf decomposition, shredders, spring-fed streams, temperature, invertebrates

## 41 Statements and Declarations

42 The authors have no competing interests to declare that are relevant to the contents of this research article.

#### 43 Acknowledgements

We are grateful to the staff and management of the University of Tokyo Hokkaido Forest (UTHF) for allowing us to conduct our research within the UTHF and for their assistance in field activities, to Dr. Kazuhiro Toyoda for his assistance in the laboratory analyses of water samples, to Ms. Aiko Agui for her assistance in laboratory work, and Mr. Hokuto Izumi and Mr. Kunihito Mihara for their help on field activities. Lastly, we are very grateful for our two expert reviewers who devoted valuable time and effort to help improve this manuscript.

## 50 Funding

51 This study was possible through the financial support from the research fund for the Ishikari and Tokachi

52 Rivers provided by the Ministry of Land, Infrastructure, Transport, and Tourism of Japan.

#### 54 Data Availability Statement

55 Data are present in the Supplementary Information and available from the corresponding author upon
 56 reasonable request.

57 Declarations

#### 58 **Conflict of interest**

- 59 The authors have no competing interests to declare that are relevant to the contents of this research article.
- 60

## 61 Introduction

62 Aquatic communities and ecosystem functions are affected by abiotic factors such as water chemistry, 63 temperature, and nutrient availability. Studies assessing the direct effects of physical and chemical 64 characteristics of the river on different aquatic organisms and processes are extensive (see Boyero et al., 65 2011; Dallas & Ross-Gillespie, 2015; Chase et al., 2017), but few have considered the possible contribution 66 of regional factors such as underlying geology (i.e., lithology of the substratum), which could indirectly 67 affect aquatic processes through their influence on water quality. Underlying geology can dictate 68 groundwater influence. Rivers with highly permeable underlying rocks would mean a stronger influence of 69 groundwater to surface water (i.e., in spring-fed streams) than those with less permeable substratum (i.e., 70 in run-off streams). A noticeable influence of groundwater to surface water is demonstrated in the relatively 71 stable temperature regime of spring-fed streams, mainly due to the buffering effect of groundwater 72 temperature to surface water temperature (Kaandorp et al., 2019).

Studies that relate geology and limnology are primarily in the field of hydrology, hydrogeology, and geomorphology (Younger, 1995; Mosher et al., 2010), and those that involve ecological components are mostly done on spring-fed rivers draining chalk and limestone aquifers (Berrie, 1992; Lusardi et al., 2021), with very few comparing rivers with different groundwater influence (Sear et al., 1999). Existing literature showed consistency in assessing spring-fed streams as relatively stable physical and chemical environments in terms of flow, temperature regime, and food availability compared to run-off streams, with variations in water quality resulting in different adaptations of invertebrate communities (see Death & Winterbourn, 1995; Jin & Ward, 2007; Lusardi et al., 2016). In addition, few studies that investigated the effect of geology on aquatic components considered structural characteristics of macroinvertebrate communities such as invertebrate diversity and abundance (i.e., Barquin & Death, 2006; Olson, 2012; Carrie et al., 2015; Lusardi et al., 2016) but did not go further as to the effects of their observed differences on stream functions (but see Griffiths & Tiegs, 2016) such as ecosystem metabolism and leaf litter decomposition. The latter is the focus of this study.

86 In headwater streams, allochthonous organic matter in the form of plant litter fuels the aquatic 87 food web (Vannote et al., 1980). Leaves are the main components of litterfall, amounting to as high as 73% 88 of total annual litterfall during the autumn season (Abelho, 2001). Upon reaching the aquatic system, leaf 89 litter undergoes decomposition and is either directly incorporated into the biomass of detritivores, 90 metabolized to carbon dioxide, or transformed into other products such as dissolved organic matter and fine 91 particulate organic matter (Wallace et al., 1995). Disentangling organic matter processing pathways and 92 the major factors influencing them, including those of anthropogenic and natural origins, is essential for 93 both aquatic biodiversity conservation and global carbon cycling. Organic matter exported in various forms 94 is provided to downstream reaches to sustain productivity (Wipfli et al., 2007), and the emitted carbon 95 dioxide can contribute substantial amounts to global carbon fluxes (Raymond et al., 2013). Decomposition 96 processes integrate a myriad of biological, physical, and chemical processes in rivers, providing an 97 important metric of river health assessment with a focus on functions (Young et al., 2008; Niyogi et al., 98 2013). Organisms (mainly fungi, bacteria, and macroinvertebrates) are the driving forces in aquatic 99 decomposition; however, these organisms depend on the substrate and surrounding environment (Graça, 100 2001). Thus, intrinsic factors (i.e., leaf chemistry, macroinvertebrate preference) and extrinsic 101 environmental factors (i.e., water quality, temperature, nutrient availability, seasonality) can play important 102 roles in decomposition (Boyero et al., 2021a, b). Geology is a primary driver of variation in both factors 103 via controls on water chemistry, nutrient content, channel geomorphology, and watershed hydrology, which 104 elicit different responses from aquatic organisms, including benthic macroinvertebrates through adaptations 105 in osmoregulation, feeding, habitat, and temperature regimes (Olson, 2012; Ishiyama et al., 2021; Lusardi 106 et al., 2021).

107

This study examined if geology affects water quality and macroinvertebrate detritivores, indirectly

108 resulting in spatial and temporal variations in leaf litter decomposition in forested headwater streams. To 109 represent geological variation, we used two types of streams with different underlying geology: volcanic 110 spring-fed streams (hereafter volcanic streams) and non-volcanic run-off streams (hereafter non-volcanic 111 streams). We hypothesized that there would be a variation in surface water characteristics between the two 112 stream types, with temperature being the most obvious difference (i.e., higher in non-volcanic streams 113 during summer but lower during winter), resulting in variations in decomposition rates by affecting both 114 microbial decomposition (leaf conditioning) and macroinvertebrate-mediated decomposition (biological 115 shredding). Using fine mesh and coarse mesh leaf litter bags, we aimed to elucidate how microbe- and 116 macroinvertebrate-mediated decomposition differ between stream types and determine the role of geology 117 in the interplay of the factors affecting decomposition in two seasons (summer and winter). Specific 118 predictions in both seasons were: 1) shredding invertebrates would be more abundant in volcanic streams 119 because of the benign temperature that is optimal for the growth and development of shredders; 2) overall 120 decomposition rate would be higher in volcanic streams because of the overriding importance of biological 121 shredding relative to temperature-dependent microbial decomposition.

122

# 123 Materials and Methods

124 Study site

125 The study was conducted within 100-m long sections of ten forested streams in Furano, Central Hokkaido, 126 Japan, encompassing an area of approximately 600 km<sup>2</sup> (Figure 1). All study sites are tributaries of the 127 Sorachi River, which belongs to the Ishikari River drainage system, and are drained by similar-sized 128 catchments. During the Quaternary Period, eruptions of the Daisetsu Mountain system covered the area 129 from the northeast with pyroclastic flow deposits and formed surficial geology dominated by volcano-130 derived rock (Ikeda & Mukoyama, 1983; Nakamura, 2016). The study streams were chosen such that they 131 differed in geological characteristics and were categorized into two types; volcanic and non-volcanic (five 132 sites/streams each) according to the dominant underlying rock type based on the Seamless Digital 133 Geological Map of Japan (Geological Survey of Japan, National Institute of Advanced Industrial Science 134 and Technology). Volcanic stream types have underlying geology composed of Early Pleistocene nonalkaline pyroclastic flow volcanic rocks. In contrast, non-volcanic stream types have Early to Late Cretaceous accretionary complexes, metamorphic rocks, marine muddy turbidite, and late Pleistocene to Holocene marine and non-marine sediments in their respective watersheds. The mean ( $\pm$ standard deviation) percentage of the volcanic geological layer for each catchment (volcanic proportion) was 99.9 ( $\pm$  2.0) for volcanic streams and 6.4 ( $\pm$  12.80) for non-volcanic streams (Table S1.1).

140 According to the weather record at the Furano weather station of the Japan Meteorological Agency, 141 approximately 20 km northwest of the study area, the mean annual temperature and annual precipitation 142 ( $\pm$ standard deviation) for the period 2010 – 2021 were 6.98 ( $\pm$  0.32) °C and 1060.09 ( $\pm$  208.86) mm, 143 respectively; mean monthly air temperature was highest in July (21.31 °C) and lowest in January (-8.25 °C). 144 Higher resolution (1km x 1km) of mean annual precipitation and air temperature data obtained for each 145 catchment using 30-year data (1980 - 2010) showed similar climatic characteristics between the two stream 146 types (P=0.475 for mean annual precipitation and P=0.174 for mean annual temperature) (adjusted alpha = 147 0.008, see Statistical Analyses). Volcanic and non-volcanic streams recorded 1063.71 - 1292.58 mm and 148 1088.85 - 1580.80 mm mean annual precipitation, and 2.66 - 4.13 °C and 3.11 - 4.82 °C mean annual air 149 temperature, respectively (Table S1.1). Selected volcanic and non-volcanic streams are similar in elevation 150 (P=0.019) (Table S1.1), slope (P=0.034) (adjusted alpha = 0.008, see Statistical Analyses), and catchment 151 size (P=0.897) (Table S1.2). High-resolution land-use and land cover maps used to calculate the proportion 152 of forest type in the catchment using ArcGIS Pro (ESRI, version 2.4.0) showed that stream catchments 153 were mainly composed of broadleaf forest and small percentages of needleleaf forest (Table S1.3). Overall, 154 forest cover was high (92% in volcanic and 99% in non-volcanic streams), but there was a slight difference 155 in forest cover between the two stream types due to the relatively lower broadleaf forest cover in one of the 156 volcanic sites (P4) (Table S1.3).

157

158

159 *Leaf decomposition experiment* 

160

161 Leaf litter packs were mimicked using  $3.0 \pm 0.1$  g (mean dry weight  $\pm$  standard errors) of *Alnus japonica* 162 (Thunb.) Steud. (Japanese Alder) placed inside leaf litter bags (hereafter litter bags) with fine and coarse 163 mesh nets (see Appendix S1). Fine mesh (257 x 266 µm) bags allow only microorganisms (bacteria and 164 fungal conidia) to enter, while coarse mesh bags enable access to both microorganisms and 165 macroinvertebrates. The coarse mesh bag is a hybrid of two net sizes (black net: 3.9 x 3.9 mm; green net: 166 1 x 1 cm), with the bottom portion (touching the riverbed) having black net only and the top portion having 167 half green and half black net. The green part (facing upstream) accommodates larger shredders observed in 168 the study sites during reconnaissance, and the black part (facing downstream) prevents the loss of 169 fragmented alder leaves. Gholz et al. (2000) applied a similar principle of using two mesh sizes for leaf 170 decomposition experiments. Two types of leaves were used to capture heterogeneous litter decomposition 171 associated with leaf conditions: green and old senescent leaves during summer and old senescent and new 172 senescent leaves during winter. Green leaves represent fresh leaves falling in summer due to wind or rain 173 action. Old senescent leaves represent lateral input of remaining litter from the previous fall season while 174 new senescent leaves represent both vertical and lateral input during the current fall season (see Appendix 175 S2 for litter bag preparation).

176 Overall, 100 pairs of coarse and fine mesh bags were incubated each season (50 pairs in volcanic 177 streams and 50 pairs in non-volcanic streams). Five replicates of four litter bags (factorial combinations of 178 two mesh types and leaf types) were used for every stream, with each replicate tethered to a metal lever 179 driven into the riverbed (20 bags in each stream per season). The distance between replicates within each 180 stream was 10 to 25 m apart. Incubation of leaf litter was done in 2018 in two seasons, from 11th July to 3rd 181 August during summer and 2<sup>nd</sup> November to 4<sup>th</sup> December during winter. December was chosen as the 182 winter sampling period because this was the third coldest month (mean monthly air temperature for the 183 period 2010 – 2021 at the Furano station: -4.84 °C) of the year after January and February (-7.15 °C) and 184 the coldest snow-free month with road access to all the sites. The collection was done twice for each season, 185 after approximately two- and three weeks of incubation during summer and after two- and four weeks 186 during winter. Two replicates from each stream were retrieved during the first collection, and the remaining 187 three replicates were retrieved during the second collection. Because this was the first time to perform a 188 litter decomposition experiment on the sites, the speed of decomposition was only roughly estimated, and 189 conditions such as lower water levels later in summer or heavy snow later in winter were anticipated. Thus, 190 more replicates (three pairs) were left for the second collection to have a higher possibility of retrieving at 191 least two pairs of litter bags.

#### 192

# 193 Collection and processing of litter bags

194 Litter bag collection was done using laundry nets (0.4 mm  $\times$  0.4 mm mesh size) to avoid loss of both 195 macroinvertebrates and remaining leaf litter which may be washed away upon moving the litter bags. Each 196 pair of litter bags was enclosed in one laundry net while still underwater and raised slowly from the water, 197 maintaining as little disturbance as possible. Some replicates were found slightly buried in deposited fine 198 sediments. To quantify the extent of sediment burial, coarse litter bags were collected with the deposited 199 sediment. Sediment did not penetrate in substantial amounts through the fine mesh and thus, was not 200 quantified for the fine mesh bags. Collected litter bags were immediately placed inside a clean Ziploc bag 201 and transported to the laboratory.

202 The remaining leaf litter was processed immediately upon reaching the laboratory. Contents of 203 litter bags were washed and sorted, and the recovered A. japonica leaf litter was oven-dried at 60°C for 204 seven days, weighed, combusted at 500 °C for 4 hours (FO310, Yamato Scientific Co., Ltd., Japan), and 205 weighed again to obtain ash-free dry mass (AFDM). Macroinvertebrates were also recovered from the 206 coarse litter bags and, upon reaching the laboratory, were immediately separated from remaining leaf litter 207 and preserved in 75% Ethanol. All invertebrates, except for Nematoda, were counted and identified to 208 family level using existing keys (Merritt & Cummins, 1996; Kawai & Tanida, 2005). Potential shredder 209 families were determined according to the classification of Merritt & Cummins (1996) as "generally 210 shredders". Some invertebrates, including all potential shredder families, were collected from the study 211 sites and subjected to C and N stable isotope analyses along with potential primary food resources such as 212 leaf litter (Figure S1) to confirm the invertebrate shredder families. Consequently, six families (Capniidae, 213 Nemouridae, Lepidostomatidae, Limnephilidae, Tipulidae, Gammaridae) were considered shredders and 214 examined in detail in relation to leaf litter decomposition. For all taxa, individuals were grouped into size 215 classes by visual identification, and representatives were taken from each size class to measure the dry 216 biomass of individuals from the respective season (Table S2). Average biomass per family per season was 217 obtained and was multiplied by each individual count to obtain approximate total shredder biomass and 218 total invertebrate biomass in each litter bag.

219

220

221

In coarse litter bags, the fine sediment removed from *A. japonica* and from other organic components (leaves and twigs), as well as sediments attached to invertebrates (i.e., in Limnephilidae cases), were collected, oven-dried at 60°C until fully dried and weighed with an analytical balance.

222

# 223 Estimation of physical and chemical parameters

224 For each site, one temperature logger (Hobo pendant temperature/light logger, UA-002-08, Onset Computer 225 Corp., Massachusetts, USA) recording every 15 minutes was set up from the start of the litter bag incubation 226 period until the collection of the last set of litter bags. Two types of temperature data were used for 227 subsequent analyses. For comparing stream types (in Table 1), the average temperature was obtained by 228 getting the mean value of all sites per day for each stream type. For structural equation modeling (SEM) 229 analyses (see Statistical analyses), the average temperature used was the mean temperature for each site for 230 every incubation period. Electrical conductivity (EC) and pH were recorded in each stream during set-up 231 and every litter bag collection using a handheld EC/pH meter (WM 32EP, DKK-TOA Co., Tokyo, Japan). 232 Water samples were collected using 250-mL acid-washed polyethylene bottles once from each site for each 233 season. The collected water was filtered through a 0.5µm filter paper (GC-50 Advantec, Tokyo, Japan) and 234 subjected to an ion analyzer (IA-300, DKK-TOA Corp., Tokyo, Japan) to determine chloride (Cl<sup>-</sup>), 235 phosphate ( $PO_4^{3-}$ ), sulfate ( $SO_4^{2-}$ ), nitrite ( $NO_2^{-}$ ) and nitrate ( $NO_3^{-}$ ) concentration. Nitrite and phosphate 236 were below the detection limit (0.05 mg/L) in all cases and thus were no longer reported hereafter. An 237 unfiltered portion was subjected to digestion using the colorimetric method and eventually to 238 spectrophotometry (UV-1280, Shimadzu Corp., Japan) to obtain total nitrogen (TN) and total phosphorus 239 (TP) concentration using a method described in Rahman et al. (2021). Winter water samples were obtained 240 in November 2018, while for the summer data, water samples were obtained in August 2019.

During litter bag collection, water depth was measured thrice for every replicate, with the 1<sup>st</sup> measurement immediate to the left side of the 1<sup>st</sup> bag, the 2<sup>nd</sup> measurement in-between the 2<sup>nd</sup> and 3<sup>rd</sup> bags, and the 3<sup>rd</sup> measurement immediate to the right side of the 4<sup>th</sup> bag. At the same time, flow velocity (Propeller-based Velocity Meter, CR-11, Cosmo-Riken Inc., Tokyo, Japan) on top of litter bags was also recorded by taking three points (in between every bag) for each replicate. Flow velocity measurements were consistently taken at approximately 60% of the depth relative to the water surface.

247

# 248 Estimation of decomposition rates

249 Decomposition rates were estimated for every litter bag collected at the second collection according to their

250 treatment types (n=12 for each stream per season) by employing the exponential decay model (Fig. S2):

$$251 \qquad \qquad AFDM_r = e^{-kt}$$

where:

253

252

254	$AFDM_r$ (remaining AFDM) = final AFDM (g) / initial AFDM (g)
255	t = incubation duration (days)
256	k = breakdown rate coefficient
257	

To fit the model,  $AFDM_r$  obtained for leaf bags at the first collection was processed to provide mean values that corresponded to each combination of mesh type and leaf type (mean  $AFDM_r$  at the first collection). The initial  $AFDM_r$ , which was assumed to be 100, the mean  $AFDM_r$  at the first collection, and  $AFDM_r$  at the second collection were used in non-linear least square regression (nls function in R software; see Statistical analyses below) with the corresponding *t* to obtain a total of 12 *k* values (six each for fine *k* and coarse *k*) for each stream. Fine *k* represents largely microbial decomposition process (hereafter microbe *k*), whereas coarse *k* represents microbial and macroinvertebrate shredder activities (hereafter overall *k*).

Decomposition rate due to shredder activity only (invertebrate *k*) was also approximated. To obtain the approximate value of final AFDM if only detritivore consumption happened (i.e., excluding the contribution of microbes), the lost AFDM due to microorganism (fine mesh) decomposition was added to the remaining AFDM obtained for coarse mesh samples. In instances where approximate final AFDM due to macroinvertebrates only exceeds the initial AFDM (fine mesh decomposition was greater compared to that in the coarse-mesh bags because of slight differences in leaf mass at the beginning), the value of initial AFDM was adopted as the final AFDM, so that contribution due to macroinvertebrates is approximated as 272 zero and is considered irrelevant. In total, 18 k values (six replicates for each type of k) were estimated for 273 each stream in each season.

274

#### 275 *Statistical analyses*

276 General characteristics of the watershed (elevation, forest cover, slope, catchment area, annual mean 277temperature, annual mean precipitation) and physical and chemical characteristics of streams (temperature, 278 ions, TN, TP, EC, pH, discharge) were compared between stream types per season using Generalized Linear 279 Mixed Models (GLMMs) with the main effect of stream type and random factor of river identity (or river 280 identity and sampling occasion/incubation day for EC, pH, and temperature) and with Gaussian error 281 distribution. Because multiple comparisons (disjunction testing) (Rubin 2021) were done within these two 282 data sets, alpha was adjusted using the Holm's Sequential Bonferroni Procedure to control for Type I error 283 (Abdi, 2010). Adjusted alpha for general watershed characteristics and physicochemical data are P=0.008284and *P*=0.006, respectively.

285 The taxonomic richness and Shannon-Weiner diversity index of the entire community and 286 abundance and biomass of shredders were compared between stream types and seasons. GLMMs were 287 developed with the main effects of stream types and seasons and their interaction, and random factors of 288 leaf type, river identity, and sampling occasion and with the following error distributions: Poisson for 289 taxonomic richness, negative binomial for shredder abundance, and Gaussian for diversity index and 290 shredder biomass). The same analyses were conducted for overall k, microbe k, and invertebrate k with 291 Gaussian error distribution. Permutational multivariate analysis of variance (PERMANOVA) was used to 292 test the compositional differences of invertebrates between stream types. For this analysis,  $\log_{10^{-1}}$ 293 transformed abundance and raw biomass of family-level macroinvertebrates were used. Moreover, 294 similarity percentages analysis (SIMPER) (Clarke, 1993) was used to determine the percentage contribution 295 of shredder taxa to the average dissimilarity between communities in the two stream types.

To identify the drivers through which geological differences in catchments affect decomposition rates directly and indirectly, structural equation modeling (SEM) was adopted, with microbe k and invertebrate k as response variables. The modeling was performed in three cases in each season: one for 299 microbe k and two for invertebrate k, which used either shredder abundance or biomass as one of the factors. 300 There were 60 data points for each type of k in each season. Physical and chemical data available only at 301 the stream scale (e.g., volcanic proportion) were assigned to all data points in corresponding streams. In the 302 analyses for microbe k and invertebrate k, volcanic proportion, temperature, and other local environmental 303 factors (local flow velocity for microbe k, and both local flow velocity and sediment deposited on coarse 304 litter bags for invertebrate k) that can affect decomposition rates were included as direct or indirect factors. 305 Nitrate and TN, which are known factors that affect leaf decomposition, were not included in the SEM 306 analyses due to the lack of significant difference in nitrate and TN between the stream types for both seasons 307 (as stated in the results). On the other hand, discharge and TP were included in the SEM models even 308 though they did not show significant differences between the stream types because of their relatively higher 309 values in one of the stream types in a certain season. The a priori path structures were decided using existing 310 knowledge about invertebrates and physical and chemical environments (Allan et al., 2021). For the 311 invertebrate k analyses, we initially did not include direct pathways from volcanic proportion to the 312 invertebrate abundance or biomass and invertebrate k as unknown effects in our hypothetical SEMs. 313 However, these relationships appeared as important latent relationships in all cases without which the model 314 did not become statistically significant. This was because there were other potentially important factors, 315 such as temporal variability of flow and water quality in the study periods. Furthermore, factors directly 316 related to geology, such as rock type, erodibility of underlying rocks and substrate lithology, can also 317 influence invertebrates (see Shearer & Young, 2011 and Nitzu et al., 2018). Thus, direct pathways from 318 volcanic proportion to invertebrate abundance/biomass were eventually included to represent these possible 319 direct effects. The Piecewise SEM framework was adopted, and each relationship was constructed using 320 generalized linear models with Gaussian error distribution (Lefcheck, 2016; Miller et al., 2018). Variables 321 were log<sub>10</sub>-transformed and scaled when appropriate. Type of leaf and location of subsite were included as 322 random factors.

323 All statistical analyses were performed using R version 4.0.1 (R Core Team, 2020) using 324 'glmmTMB', 'lme4', 'vegan', 'ADONIS', and 'piecewiseSEM' packages. The statistical significance level 325 for analyses was set at 0.05. For piecewise SEM, Fisher's C statistics (p>0.05) was used to determine the 326 model's goodness of fit.

## 328 Results

329 Stream-type difference reflected through water temperature was more evident in summer than in winter, 330 with volcanic and non-volcanic streams ranging from 9.10 - 10.86 °C and 13.63 - 16.14 °C, respectively, 331 during summer incubation and 4.28 - 6.06 °C and 3.63 - 4.87 °C, respectively, during winter incubation 332 (Table S3.1). In summer, volcanic streams recorded significantly lower water temperature while in winter, 333 temperature was significantly higher in volcanic than non-volcanic streams (Figure 2, Table 1). Based on 334 this result, water temperature was considered as a possible factor affecting k in the SEM analyses later. 335 Although not statistically significant, discharge in summer was relatively higher in non-volcanic streams 336  $(0.432 \pm 0.198 \text{ m}^3/\text{s})$  than volcanic streams  $(0.209 \pm 0.048 \text{ m}^3/\text{s})$ , and TP concentration in winter was 337 relatively higher in volcanic  $(0.025 \pm 0.009 \text{ mg/L})$  than non-volcanic streams  $(0.015 \pm 0.004 \text{ mg/L})$  (Table 338 S3.1 and S3.2). The pattern observed in TP was due to a decrease in TP concentration in non-volcanic 339 streams during winter while the concentration in volcanic streams remained similar in the two seasons.

340 A total of 18,938 macroinvertebrate individuals (7,013 in summer and 11,925 in winter) were 341 collected from the coarse litter bags. In eight fine litter bags, an average of four early instars Chironomidae 342 (17 individuals at the most) per bag were found, which were 0.57-7.69% of the Chironomidae abundance 343 in their paired coarse bag. Because of the small body size and lower abundance compared to those found 344 in the coarse litter bags, the presence of Chironomidae larvae in the fine mesh bags was assumed to have 345 no substantial contribution to decomposition and was not included in the analyses. Community differences 346 between river types and seasons were observed in diversity indices and shredder community structure. 347 Taxonomic richness and Shannon-Weiner index were higher in volcanic streams in summer (Table 2 a, b; 348 Figure 3 a, b). Detritivores comprised 68.9 and 80.8% of the abundance and biomass of all samples, 349 respectively, with some variations between seasons and river types (Figure S3). Shredder abundance was 350 higher in volcanic streams in summer (Table 2 c, Figure 3 c). Shredder biomass was higher in winter 351 compared to summer, and volcanic streams exhibited higher values in both seasons (Table 2 d, Figure 3 d). 352 The results of SIMPER indicated that shredder taxa were among the most critical groups in distinguishing 353 river-type communities (Table 3).

354

Decomposition rate exhibited various responses to seasons and river types depending on the type

355 of k, with all cases having statistically significant interactions between two factors (Table 4, Figure 4). 356 Overall, k seasonally fluctuated in nonvolcanic streams, whereas volcanic streams showed relatively stable 357 values. Microbe k was higher in summer and decreased in winter in non-volcanic streams, whereas the 358 decreasing trend in volcanic streams was statistically insignificant. These seasonal trends in microbe k were 359 consistent with those in overall k, although the degree of the change was lesser in the latter. Damped 360 seasonal contrasts in overall k were explained by the trends in invertebrate k, which offset the seasonal 361 changes in microbe k. The invertebrate k was lowest in non-volcanic streams in summer and was relatively 362 higher in volcanic streams in both seasons.

363 All SEM exhibited acceptable levels of goodness of fit (Fisher's C statistics, p > 0.09 in all cases). 364 Temperature was the most important environmental factor by which surface geology affected microbe k in 365 both seasons (summer: standard coefficient (coef) = 0.48,  $r^2 = 0.24$ ; winter: coef = 0.47,  $r^2 = 0.48$ ) (Figure 366 5 a, d). These pathways were directionally contrasting, with colder water in volcanic streams (coef = -0.96, 367  $r^2 = 0.93$ ) limiting the microbe k in summer and warmer water in volcanic streams (coef = 0.67,  $r^2 = 0.45$ ) 368 promoting the microbe k in winter (Figure 5 a, d). In addition to temperature, relatively higher TP levels 369 associated with volcanic geology (coef = 0.68,  $r^2 = 0.45$ ) also contributed to an increased microbe k in 370 winter (coef = 0.26,  $r^2 = 0.48$ ). Indirect effects of surface geology affecting invertebrate k via shredding 371 invertebrates were apparent in three out of four cases (except for biomass in summer) (Figure 5 b, e, f). In 372 summer, the colder temperature in volcanic streams was associated with positive overall effects on shredder 373 abundance (coef = -1.45,  $r^2 = 0.37$ ) and thus invertebrate k (coef = 0.28,  $r^2 = 0.28$ ) (Figure 5 b); unknown 374 direct pathways were also detected for geology-invertebrate k relations (coef = 0.29 for abundance and 0.34 375 for biomass). In winter, both abundance (coef = 0.29) and biomass (coef = 0.36) adequately explained the 376 variation in invertebrate k ( $r^2 = 0.45$  for abundance and 0.46 for biomass), although temperature-mediated 377 effects of surface geology on invertebrate k were undetected. The detected effects of surface geology on 378 invertebrate k via measured environmental factors included indirect effects of flow velocity through its 379 direct effect on abundance (coef = -0.38,  $r^2 = 0.21$ ) and biomass (coef = -0.28,  $r^2 = 0.33$ ) in winter; unknown 380 direct pathways were also detected for geology-shredder relations (abundance: coef = 0.68,  $r^2 = 0.21$ ; 381 biomass:  $coef = 0.79, r^2 = 0.33$ ).

## 383 Discussion

384 Spatial heterogeneity in surface geological characteristics at the landscape level and its effects on physical 385 properties of riverine ecosystems such as catchment hydrology have been well recognized (McDonnell et 386 al., 2007; Olson, 2012; Kaandorp et al. 2019). Despite the known effects of flow regime and catchment 387 hydrology on biota (Leland & Porter, 2000; Newton et al., 2008; Wenger et al., 2011; Lusardi et al., 2021), 388 our understanding of catchment-geology controls on ecosystem functioning remained limited. By using leaf 389 litter decomposition as a measure of key ecosystem functioning in forested headwater streams, our findings 390 supported the hypothesis that catchment geology mediates decomposer community structures, leading to 391 variations in ecosystem functioning. However, our specific prediction was partially unsupported because 392 the overall decomposition rate in coarse litter bags was not significantly higher in shredder-abundant 393 volcanic streams but was more seasonally stable than in non-volcanic streams.

394 The observed physical environmental gradients across study streams can be attributed to 395 geological characteristics of the surficial deposits because other factors that could dictate water temperature, 396 such as climatic characteristics, were shown to be similar between the two stream types. Although there 397 was a slight difference in overall forest cover due to one volcanic site (P4) with a lower cover, this did not 398 affect the observed temperature regime characteristic of the volcanic sites. Specifically, although P4 399 showed a slightly higher temperature than other volcanic streams during summer, it was still lower than the 400 non-volcanic streams. This could also highlight the importance of underlying geology in relation to the 401 effects of forest cover on water temperature in headwaters. Moreover, the thermal regimes of the two stream 402 types are similar to those observed in other studies (particularly in the middle of summer and beginning of 403 winter) (i.e., Jin & Ward, 2007; Lusardi et al., 2021) which reported difference in temperature pattern of 404 streams due to variation in substrata. Geology was found to be a strong predictor of stream water 405 temperature. Pyroclastic flow volcanic rocks found in the underlying geology of volcanic streams were 406 mainly pumice deposits characterized by high porosity, with large pores connected by small channels 407 (Bernard et al., 2007). High groundwater input allows a stable temperature regime due to the strong 408 interaction of groundwater with surface water (Kaandorp et al., 2019). In volcanic streams, the stability of 409 groundwater temperature would result in colder summer temperatures and warmer winter temperatures than 410 non-volcanic streams (Crisp & Le Cren, 1970). As observed in our study sites, the temperature is colder in

411 volcanic sites than in non-volcanic sites during summer and the expected shifting of the pattern was also 412 observed during winter, with volcanic streams having warmer water than non-volcanic streams especially 413 in the latter part of incubation. This contrast may be more pronounced later in winter because water 414 temperature in non-volcanic streams would have been more strongly affected by colder temperatures in the 415 middle of winter (i.e., January).

416 The structure of macroinvertebrate communities was distinctive in relation to the geological 417 characteristics of the contributing catchments. Variations in macroinvertebrate community composition 418 between streams having different groundwater influences have been reported by several studies on benthic 419 invertebrates (i.e., Barquin & Death, 2006; Lusardi et al., 2016). It has been noted that the relative stability 420 of the spring-fed system enables the provision of good quality and quantity of food resources. Interestingly, 421 invertebrate shredders appear to be particularly responsive to geological differences. Volcanic spring-fed 422 rivers were observed to have higher retention of large woody debris (Barquin & Death, 2004; Lusardi et al., 423 2021), which, in addition to stable flow, could contribute to the possible retention of leaf litter, thereby 424 increasing the food availability for shredders.

425 An important limiting factor of community structure highlighted in this study is water 426 temperature in summer. In aquatic insects, the temperature regime can be the most important factor 427 influencing growth in areas where food is not limited. Because of this, the emergence of terrestrial form 428 can be significantly affected (Watanabe et al., 1999; Bayoh & Lindsay, 2003; Jin & Ward, 2007). High 429 temperatures hasten physiological development and might shorten the available time for organisms to 430 accumulate the required nutrients needed for complex processes such as eclosion and emergence to the 431 adult stage of some species (Lassiter et al., 1994). These direct limiting effects of temperature can be 432 detrimental for shredders. The majority of the shredders collected in the study area are from the orders 433 Plecoptera and Trichoptera, which are evolutionarily adapted to cold waters (Balian et al., 2008) - the same 434 reason that there are more shredders in temperate areas compared to the tropics (Irons et al., 1994; Boyero 435 et al., 2011). Thus, higher water temperature in non-volcanic streams in summer could limit the abundance 436 and diversity of invertebrate communities.

437

It was also possible that the physical and chemical environment influenced by geology indirectly

438 affected invertebrates via mediation of food quality through leaf litter conditioning. This pathway was 439 visualized by dividing k into several constituents together with SEM analyses. It is noticeable that microbe-440 mediated decomposition follows the temperature pattern. This trend was expected since the positive effect 441 of temperature on microbial activity has already been established in both laboratory and field settings. 442 Temperature has a positive effect on microbial mass and enzymatic activity (Ferreira & Chauvet, 2011; 443 Ferreira et al., 2015), explaining about 40% of variations in microbe-mediated decomposition (Boyero et 444 al., 2011). During decomposition, microorganisms use cellulases (e.g.,  $\beta$ -glucosidase) to degrade cellulose 445 fibers through cellulolysis. Temperature was reported to have the strongest interaction with substrate 446 chemistry and cellulolytic activities compared to other water quality parameters and was shown to cause 447 higher decomposition efficiency and cellulase activity, giving it a leading role as a predictor of 448 decomposition rates (Fenoy et al., 2016). Ultimately, temperature can positively affect the leaf conditioning 449 process of microorganisms, which is crucial for the palatability of leaf litter for shredder invertebrates. 450 However, in this study, summer temperature did not probably lead to positive effects on shredder 451 invertebrates through litter conditioning, as evidenced by the lower abundance and biomass of shredders in 452 the warmer non-volcanic streams. In contrast, temperature positively affected microbe k in winter, 453 suggesting possible positive effects of volcanic geology on shredders via increased food resource quality 454 in this season. Furthermore, microbial processes and production in winter might have been stimulated by 455 the relatively higher concentration of phosphorus in volcanic streams, as observed in the positive pathway 456 from geology to TP and microbe k in the SEM plot (Fig. 5). This illustrates the effect of phosphorus-457 stimulated food quality on leaf decomposition (Gulis et al., 2006; Ardon et al., 2020). The TP concentration 458 in volcanic streams was similar in the two seasons and was relatively higher than in non-volcanic streams 459 during winter, probably due to the different sources of TP for each stream type. In contrast with volcanic 460 streams, which are highly influenced by groundwater, non-volcanic streams are highly affected by 461 contributions from surface runoff, which are lower during winter. Volcanic streams can have high 462 phosphorus levels because groundwater can move through the young volcanic rocks rich in phosphorus. 463 This may also be brought about by the weathering of rock materials associated with volcanic deposits (e.g., 464 Timperley & Vigor-Brown, 1985). Substantial contribution of volcanic geology to phosphorus load has 465 also been documented elsewhere (Felitsyn, 2002; Domagalski & Saleh, 2015; Records et al., 2016). 466 Although there was no significant difference in TP between stream types, the noticeable decrease in TP

467 concentration in non-volcanic streams during winter showed substantial contribution to the microbe-468 mediated decomposition SEM, highlighting the importance of stability of TP concentration in the leaf 469 decomposition in volcanic streams.

470

471 The significant contribution of invertebrates to macroinvertebrate-mediated decomposition, 472 particularly in volcanic areas, can be attributed to the presence of more shredders in the relatively 473 temperature-stable volcanic streams. However, this causal pathway was not reflected as the overarching 474 effect of geology on shredders and invertebrate k when examined separately for each season in the SEM 475 plot. First, temperature-mediated effects of geology on invertebrate k were not detected in winter. Second, 476 the TP-mediated effect of geology on invertebrate k, which was expected from the phosphorus-stimulated 477 food quality hypothesis above, was not observed in winter. Also, direct pathways from geology to 478 invertebrate abundance and biomass unexplained by the measured environmental variables were rather 479 substantial as seen in the values of  $R^2$  and standard coefficients. A plausible explanation for the above-480 mentioned discrepancies in seasonal patterns and the unexpected extent of the direct effects of geology on 481 invertebrate-related response variables would be the temporal mismatch between the recorded biota 482 response and the environmental parameters used in the analyses. Most shredder taxa are considered to have 483 univoltine to semivoltine life cycles (personal observations based on body size frequency distribution and 484 emergent adults in the study sites, JNN), being potentially susceptible to any seasonal environmental filters. 485 Moreover, there is inherent complexity in the response of different shredder invertebrate species to 486 environmental factors according to their life history strategies (i.e., some species undergo aestivation, some 487 undergo winter diapause, while others migrate to warmer areas in the river to prepare for winter) which 488 might be entirely different with the other species under the same family. In our case, as mentioned earlier, 489 the summer water temperature may strongly limit macroinvertebrate communities, including shredders, 490 while community structure could have little association with the same environmental variable measured in 491 different seasons. For the same reason, positive effects of phosphorus-stimulated food quality may affect 492 other properties of the community, such as individual growth rate or survival rates, which may not be 493 detected as strong associations between TP and shredders in winter.

494 Recognizing the effects of geology on the structural and functional characteristics of streams is 495 essential due to the observed influence of geology on both biotic and abiotic components of the stream. 496 Leaf decomposition provides the primary energy source for the aquatic food web, which is further linked 497 with the terrestrial food web in headwater streams, making it an important component of complex aquatic-498 terrestrial relationships. This highlights the importance of quantifying and understanding factors that could 499 affect the decomposition process. Although the results of this study were able to pinpoint temperature as 500 one of the vital factors in the geology-leaf decomposition connection, the robustness of data and accuracy 501 of the conclusion can still be improved by adding more environmental factors to the analyses and 502 conducting manipulative experiments to elucidate causal relationships. Because summer conditions or 503 winter conditions each by themselves or in combination across two seasons can create bottlenecks in the 504 invertebrate community (i.e., temperature and velocity limiting shredder abundance and biomass) as well 505 as associated ecosystem functioning (i.e., temperature and/or TP influencing microbe decomposition), 506 future studies would need to have a design to empirically disentangle them.

507

# 508 Conclusions

509 This study established a strong conclusion regarding the differences in volcanic and non-volcanic systems 510 in terms of macroinvertebrate community structure and functional attributes such as leaf litter 511 decomposition, which is vital in continuing aquatic ecosystem processes. Importantly, temporally stable 512 functioning was maintained by shredding invertebrate activity that balanced well with temperature-513 dependent microbial activity in volcanic streams. Implications of temperature as a strong factor influencing 514 decomposition call attention to the prospected increase in global atmospheric temperature. There is a 515 growing threat to the survival of cold-adapted aquatic organisms as increasing temperature and broader 516 temperature fluctuations limit their distribution and survival. As thermally and functionally stable systems, 517 volcanic spring-fed systems can be valuable, and identification and conservation of these systems, including 518 invertebrate diversity, is important in this age of increasing global atmospheric temperature.

519

520 **References** 

- Abelho, M. 2001. From Litterfall to Breakdown in Streams: A Review. The Scientific World Journal 1:
  656-680. DOI:10.1100/tsw.2001.103
- 523 Abdi, H. 2010. Holm's Sequential Bonferroni Procedure. In Neil Salkind (Ed), Encyclopedia of Research
- 524 Design, Thousand Oaks, CA: Sage. DOI:10.4135/9781412961288.n178
- Allan JD, Castillo MM, Capps KA. 2021. Stream ecology: structure and function of running waters (3<sup>rd</sup> ed).
  Springer, Cham.
- Ardón M, Zeglin LH, Utz RM, Cooper SD, Dodds WK, Bixby RJ, ... & Ward A. 2020. Experimental
  nitrogen and phosphorus enrichment stimulates multiple trophic levels of algal and detrital-based food
  webs: a global meta-analysis from streams and rivers. Biological Reviews 000-000. DOI:
  10.1111/brv.12673
- Balian EV, Segers H, Lévèque C, Martens K. 2008. The Freshwater Animal Diversity Assessment: an
  overview of the results. Hydrobiologia 595: 627–637. DOI:10.1007/s10750-007-9246-3
- Barquín J & RG Death. 2004. Patterns of invertebrate diversity in streams and freshwater springs in
  Northern Spain. Fundamental and Applied Limnology 161: 329-349. DOI:10.1127/0003-9136/2004/01610329
- Barquín J & RG Death. 2006. Spatial patterns of macroinvertebrate diversity in New Zealand springbrooks
  and rhithral streams. Journal of the North American Benthological Society 25: 768-786.
  DOI:10.1899/0887-3593(2006)025[0768:SPOMDI]2.0.CO;2
- 539 Bayoh MN & SW Lindsay. 2003. Effect of temperature on the development of the aquatic stages of
- 540 Anopheles gambiae sensu stricto (Diptera: Culicidae). Bulletin of Entomological Research 93: 375-381.
- 541 DOI:10.1079/ber2003259
- 542 Bernard ML, Zamora M, Géraud Y, Boudon G. 2007. Transport properties of pyroclastic rocks from
- 543 Montagne Pelée volcano (Martinique, Lesser Antilles). Journal of Geophysical Research: Solid Earth 112:
- 544 B5. DOI:10.1029/2006JB004385
- 545 Berrie AD. 1992. The chalk-stream environment. Hydrobiologia 248: 3-9. DOI: 10.1007/BF00008881

- 546 Boyero L, Pearson RG, Gessner MO, Barmuta LA, Ferreira V, Graca MAS, ... & West DC. 2011. A global
- 547 experiment suggests climate warming will not accelerate litter decomposition in streams but might reduce
- 548 carbon sequestration. Ecology Letters 14: 289-94. DOI: 10.1111/j.1461-0248.2010.01578.x
- 549 Boyero L, Pérez J, López-Rojo N, Tonin AM, Correa-Araneda F, Pearson RG, ... & Yule CM. 2021a.
- 550 Latitude dictates plant diversity effects on instream decomposition. Science Advances 7: eabe7860. DOI:
- 551 10.1126/sciadv.abe7860
- 552 Boyero L, López-Rojo N, Tonin AM, Perez J, Correa-Araneda F, Pearson RG, ... & Yule CM. 2021b.
- Impacts of detritivore diversity loss on instream decomposition are greatest in the tropics. Nature
  Communications 12: 3700. DOI:10.1038/s41467-021-23930-2
- Carrie R, Dobson M, Barlow J. 2015. The influence of geology and season on macroinvertebrates in
  Belizean streams: implications for tropical bioassessment. Freshwater Science 34: 648-662.
  DOI:10.1086/681541
- 558 Chase JW, Benoy AG, Culp JM. 2017. Combined effects of nutrient enrichment and inorganic
- sedimentation on benthic biota in an experimental stream system. Water Quality Research Journal 52: 151-
- 560 165. DOI:10.2166/wqrj.2017.038
- 561 Clarke KR. 1993. Non-parametric multivariate analyses of changes in community structure. Australian
- 562 Journal of Ecology 18: 117-143. DOI: 10.1111/j.1442-9993.1993.tb00438.x
- 563 Crisp DT & ED Le Cren. 1970. The temperature of three different small streams in northwest England.
- 564 Hydrobiologia 35: 305-323. DOI:10.1007/BF00181734
- 565 Dallas HF & V Ross-Gillespie. 2015. Review: Sublethal effects of temperature on freshwater organisms,
- 566 with special reference to aquatic insects.Water SA 41: 712-726. DOI:10.4314/was.v41i5.15
- 567 Death RG & MJ Winterbourn. 1995. Diversity patterns in stream benthic invertebrate communities: the
- 568 influence of habitat stability. Ecology 76: 1446-1460. DOI:10.2307/1938147
- 569 Domagalski JL & D Saleh. 2015. Sources and transport of phosphorus to rivers in California and adjacent
- 570 states, U.S., as determined by SPARROW modelling. Journal of the American Water Resources

571 Association 6: 1463-1486. DOI: 10.1111/1752-1688.12326

Felitsyn SB. 2002. The redistribution of phosphorus in basic volcanic rocks. Lithology and Mineral
Resources 37: 94-96. DOI: 10.1023/A:1013695708085

Fenoy E, Casas JJ, Díaz-López M, Rubio J, Guil-Guerrero JL, Moyano-López FJ. 2016. Temperature and
substrate chemistry as major drivers of interregional variability of leaf microbial decomposition and
cellulolytic activity in headwater streams. FEMS Microbiology Ecology 92: fiw169.
DOI:10.1093/femsec/fiw169

- Ferreira V & E Chauvet. 2011. Synergistic effects of water temperature and dissolved nutrients on litter
  decomposition and associated fungi. Global Change Biology 17: 551-564. DOI:10.1111/j.13652486.2010.02185.x
- 581 Ferreira V, Chauvet E, Canhoto C. 2015. Effects of experimental warming, litter species, and presence of
- 582 macroinvertebrates on litter decomposition and associated decomposers in a temperature mountain stream.
- 583 Canadian Journal of Fisheries and Aquatic Sciences 72: 206-215. DOI:10.1139/cjfas-2014-0119
- 584 Gholz HL, Wedin DA, Smitherman SA, Harmon ME, Parton WJ. 2000. Long-term dynamics of pine 585 hardwood litter in contrasting environments: toward a global model of decomposition. Global Change 586 Biology 6: 751-765. DOI: 10.1046/j.1365-2486.2000.00349.x
- 587 Graça MA. 2001. The role of invertebrates on leaf litter decomposition in streams–a review. International
- 588 Review of Hydrobiology 86: 383-393. DOI:10.1002/1522-2632(200107)86:4/5<383::AID-589 IROH383>3.0.CO;2-D
- 590 Griffiths NA & SD Tiegs. 2016. Organic-matter decomposition along a temperature gradient in a forested
- headwater stream. Freshwater Science 35: 518-533. DOI:10.1086/685657
- 592 Gulis V, Ferreira V, Graca MAS. 2006. Stimulation of leaf litter decomposition and associated fungi and
- 593 invertebrates by moderate eutrophication: implications for stream assessment. Freshwater Biology 51:
- 594 1655-1669. DOI: 10.1111/j.1365-2427.2006.01615.x
- 595 Ikeda Y & S Mukoyama. 1983. Stratigraphy and correlation of the pyroclastic flow deposits in the Furano-

- Asahikawa area, central Hokkaido, Japan. Journal of the Geological Society of Japan 89: 163-172. DOI:
  10.5575/geosoc.89.163
- 598 Irons JG, Oswood MW, Stout J, Pringle CM. 1994. Latitudinal patterns in leaf litter breakdown: is
- temperature really important? Freshwater Biology 32: 401-411. DOI:10.1111/j.1365-2427.1994.tb01135.x
- 600 Ishiyama N, Miura K, Inoue T, Sueyoshi M, Nakamura F. 2021. Geology-dependent impacts of forest
- 601 conversion on stream fish diversity. Conservation Biology 35: 884-896. DOI:10.1111/cobi.13655
- 502 Jin H & M Ward. 2007. Life history and secondary production of Glossosoma nigrior Banks (Trichoptera:
- 603 Glossosomatidae) in two Alabama streams with different geology. Hydrobiologia 572: 245-258. DOI:
- 604 10.1007/s10750-006-0370-2
- 605 Kaandorp VP, Doornenbal PJ, Kooi H, Broers HP, de Louw PGB. 2019. Temperature buffering by
- 606 groundwater in ecologically valuable lowland streams under current and future climate conditions. Journal
- 607 of Hydrology X 3: 100031. DOI: 10.1016/j.hydroa.2019.100031
- Kawai T & K Tanida. 2005. Aquatic insects of Japan: manual with keys and illustrations. Tokai University,
  Kanagawa, Japan, 1-1342.
- 610 Lassiter MT, Apperson CS, Crawford CL, Roe RM. 1994. Juvenile hormone metabolism during adult
- 611 development of Culex quinquefasciatus (Diptera: Culicidae). Journal of Medical Entomology 31: 586-593.
- 612 DOI:10.1093/jmedent/31.4.586
- 613 Lefcheck JS. 2016. piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution,
- and systematics. Methods in Ecology and Evolution 7: 573-579. DOI:10.1111/2041-210X.12512
- 615 Leland HV & SD Porter. 2000. Distribution of benthic algae in the upper Illinois River basin in relation to
- 616 geology and land use. Freshwater Biology 44: 279-301. DOI:10.1046/j.1365-2427.2000.00536.x
- 617 Lusardi RA, Bogan MT, Moyle PB, Dahlgren RA. 2016. Environment shapes invertebrate assemblage
- 618 structure differences between volcanic spring-fed and runoff rivers in northern California. Freshwater
- 619 Science 35: 1010-1022. DOI:10.1086/687114
- 620 Lusardi RA, Nichols AL, Willis AD, Jeffres CA, Kiers AH, Van Nieuwenhuyse EE, Dahlgren RA. 2021.

- Not all rivers are created equal: The importance of spring-fed rivers under a changing climate. Water 13:
  1652. DOI:10.3390/w13121652
- 623 McDonnell JJ, Sivapalan M, Vaché K, Dunn S, Grant G, Haggerty R, ... & Weiler M. 2007. Moving beyond
- heterogeneity and process complexity: A new vision for watershed hydrology. Water Resources Research
- 625 43: W07301. DOI:10.1029/2006WR005467
- Merritt RW & Cummins KW. 1996. An introduction to the aquatic insects of North America, 3rd edition.
  Kendall/Hunt Publishing Co., Iowa
- 628 Miller RJ, Lafferty KD, Lamy T, Kui L, Rassweiler A, Reed DC. 2018. Giant kelp, Macrocystis pyrifera,
- 629 increases faunal diversity through physical engineering. Proceedings of the Royal Society B: Biological
- 630 Sciences 285: 1874. DOI: 10.1098/rspb.2017.2571
- Mosher JJ, Klein GC, Marshall AG, Findlay RH. 2010. Influence of bedrock geology on dissolved organic
  matter quality in stream water. Organic Geochemistry 41: 1177-1188.
  DOI:10.1016/j.orggeochem.2010.08.004
- 634 Nakamura Y. 2016. Stratigraphy, distribution, and petrographic properties of Holocene tephras in Hokkaido,
- 635 northern Japan. Quaternary International 397: 52-62. DOI:10.1016/j.quaint.2015.07.056
- Newton TJ, Woolnough DA, Strayer DL. 2008. Using landscape ecology to understand and manage
  freshwater mussel populations. Journal of the North American Benthological Society 27: 424-439.
  DOI:10.1899/07-076.
- 639 Nitzu E, Dorobat ML, Popa I, Giurginca A, Baba S. 2018. The influence of geological substrate on the
- 640 faunal structure of the superficial subterranean habitats. Carpathian Journal of Earth and Environmental
- 641 Sciences 13: 383-393. DOI:10.26471/cjees/2018/013/033
- 642 Niyogi DK, Harding JS, Simon KS. 2013. Organic matter breakdown as a measure of stream health in New
- 643 Zealand streams affected by acid mine drainage. Ecological Indicator 24: 510–517.
  644 DOI:10.1016/j.ecolind.2012.08.003
- Olson JR. 2012. The influence of geology and other environmental factors on stream water chemistry and

- 646 benthic invertebrate assemblages. All Graduate Theses and Dissertations, Utah State University. Retrieved
- 647 from https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=2340&context=etd
- 648 Rahman MATMT, Negishi JN, Alam, MdK, Yiyang G, Tolod JR, Pongsivapai P. 2021. Lateral and
- 649 longitudinal flight dispersals of a stonefly, Alloperla ishikariana (Plecoptera, Chloroperlidae), from the
- hyporheic zone in a gravel-bed river in Japan. Limnologica 89: 125886. DOI:10.1016/j.limno.2021.125886
- 651 R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical
- 652 Computing, Vienna, Austria. URL https://www.R-project.org/
- 653 Raymond PA, Hartmann J, Lauerwald R, Sobek S, McDonald C, Hoover M, ... & Guth P. 2013. Global
- carbon dioxide emissions from inland waters. Nature 503: 355-359. DOI:10.1038/nature12760
- 655 Records RM, Wohl E, Arabi M. 2016. Phosphorus in the river corridor. Earth Science Reviews 158: 65-88.
- 656 DOI:10.1016/j.earscirev.2016.04.010
- 657 Rubin, M. 2021. When to adjust alpha during multiple testing: A consideration of disjunction, conjunction,
- 658 and individual testing. Synthese 199: 10969-1100. DOI:10.1007/s11229-021-03276-4
- 659 Sear DA, Armitage PD, Dawson FH. 1999. Groundwater dominated rivers. Hydrological Processes 13:
- 660 255-276. DOI:10.1002/(SICI)1099-1085(19990228)13:3<255::AID-HYP737>3.0.CO;2-Y
- 661 Shearer KA & RG Young. 2011. Influences of geology and land use on macroinvertebrate communities
- across Motueka River catchment, New Zealand. New Zealand Journal of Marine Freshwater Research 45:
- 663 437-454. DOI:10.1080/00288330.2011.587823
- Timperley MH & RJ Vigor-Brown. 1985. Weathering of pumice in the sediments as a possible source of
- major ions for the waters of Lake Taupo, New Zealand. Chemical Geology 49: 43-52. DOI:10.1016/0009-
- 666 2541(85)90146-9
- 667 Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE. 1980. The river continuum concept.
- 668 Canadian Journal of Fisheries and Aquatic Sciences 37: 130-37. DOI:10.1139/f80-017
- 669 Wallace JB, Whiles MR, Eggert S, Cuffney TF, Lugthart GJ, Chung K. 1995. Long-Term Dynamics of
- 670 Coarse Particulate Organic Matter in Three Appalachian Mountain Streams. Journal of the North American

## 671 Benthological Society 14: 217-232. DOI:10.2307/1467775

- 672 Watanabe NC, Mori C, Yoshitaka I. 1999. Effect of water temperature on the mass emergence of the mayfly,
- 673 Ephron shigae, in a Japanese river (Ephemeroptera: Polymitarcyidae). Freshwater Biology 41: 537-541.
- Wenger SJ, Isaak DJ, Luce CH, Neville HM, Fausch KD, Dunham JB, ... & Williams JE. 2011. Flow
- regime, temperature, and biotic interactions drive differential declines of trout species under climate change.
- 676 Proceedings of the National Academy of Sciences 108: 14175-14180. DOI:10.1073/pnas.1103097108
- 677 Wipfli MS, Richardson JS, Naiman RJ. 2007. Ecological linkages between headwaters and downstream
- 678 ecosystems: Transport of organic matter, invertebrates, and wood down headwater channels 1. Journal of
- 679 the American Water Resources Association 43: 72-85. DOI:10.1111/j.1752-1688.2007.00007.x
- 680 Young RG, Matthaei CD, Townsend CR. 2008. Organic matter breakdown and ecosystem metabolism:
- 681 functional indicators for assessing river ecosystem health. Journal of the North American Benthological
- 682 Society 27:605-25. DOI:10.1899/07-121.1
- 683 Younger PL. 1995. Modelling river-aquifer interactions. British Hydrological Society Occassional Paper
- No. 6. BHS, p. 165. In Sear, D.A., P.D. Armitage, & F.H. Dawson, 1999. Groundwater dominated rivers.
- 685 Hydrological Processes 13: 255-276. DOI:10.1002/(SICI)1099-1085(19990228)13:3<255::AID-</li>
  686 HYP737>3.0.CO;2-Y

687

## 688 Tables

- 689 Table 1 Results of GLMMs testing for the effect of stream type on physical and chemical variables. p-
- 690 values shown for the two seasons were produced for the Wald Z-statistic. The significance level determined
- by Holm-Bonferroni Sequential Procedure requires a *p*-value of 0.006 after alpha adjustment. Bold letters
- 692 denote statistical significance. The sample size for each analysis is n=10 except for EC (summer=40,

693 winter=60), pH (summer=30, winter=60), and temperature (summer=44, winter=58).

Water	EC		Cl-	NO <sub>3</sub> -	SO4 <sup>2-</sup>	ТР	TN	Discharge	Temperature
Quality	(mS/m)	рН	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(m <sup>3</sup> /s)	(°C)

Parameter									
Summer	0.153	0.568	0.243	0.327	0.953	0.493	0.354	0.014	p<0.006
Winter	0.145	0.745	0.0234	0.234	0.279	0.019	0.264	0.204	p<0.006

**Table 2** Results of GLMMs testing of the effects of stream types related to geological characteristics (G),

696 season (S), and their interactions on the taxonomic richness (a), Shannon-Wiener diversity index (b),

697 shredder abundance (c), and shredder biomass (d) in coarse litter bags. P-values were produced in GLMM

698 for the Wald Z-statistic. Bold letters indicate statistical significance. SD represents standard deviations.

(a) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	1.583	0.108	<0.001
G (volcanic relative to non-volcanic)	0.530	0.147	<0.001
S (winter relative to summer)	0.480	0.081	<0.001
G x S	-0.215	0.104	0.038

(b) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	1.280	0.161	<0.001
G (volcanic relative to non-volcanic)	0.878	0.192	<0.001
S (winter relative to summer)	0.643	0.170	<0.001
G x S	-0.483	0.167	0.004

(c) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	1.824	0.231	<0.001
G (volcanic relative to non-volcanic)	0.853	0.320	0.008
S (winter relative to summer)	1.397	0.199	<0.001
G x S	-0.356	0.265	0.180

(d) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	1.021	0.283	<0.001
G (volcanic relative to non-volcanic)	1.501	0.400	<0.001
S (winter relative to summer)	1.224	0.400	0.002
G x S	-0.647	0.565	0.252

<sup>702</sup> 

703 **Table 3** Results of SIMPER analyses with abundance (a) and biomass (b) showing overall dissimilarity

between communities of the two stream types, average contribution of ten most important taxa to overall

705 dissimilarity, and their cumulative contribution.

(a) Summer			Winter			
Overall	62.8%		Overall	51 4%		
dissimilarity	02.070		dissimilarity	51.770		
	Average	Cumulative		Average	Cumulative	
Contributory taxa	contribution	contribution	Contributory taxa	contribution	contribution	
	(%)	(%)		(%)	(%)	
Leptophlebiidae	7.9	12.6	Ephemerellidae	5.6	10.8	
Chironomidae	7.8	25.0	Lepidostomatidae <sup>†</sup>	4.6	19.7	
Nemouridae <sup>†</sup>	6.9	35.9	Leptophlebiidae	4.3	27.9	
Perlodidae	6.0	45.5	Capniidae <sup>†</sup>	4.2	36.1	
Ephemerellidae	5.3	53.9	Chironomidae	4.0	43.9	
Tipulidae <sup>†</sup>	3.7	59.8	Nemouridae <sup>†</sup>	3.8	51.3	
Lepidostomatidae <sup>†</sup>	3.5	65.4	Perlodidae	3.3	57.7	
Chloroperlidae	3.4	70.8	Chloroperlidae	2.8	63.2	
Limnephilidae <sup>†</sup>	2.9	75.4	Limnephilidae <sup>†</sup>	2.5	68.0	
Uenoidae	2.2	78.8	Baetidae	2.1	72.2	

<sup>†</sup>shredder invertebrates

(b) Summer			Winter		
Overall	68 23%		Overall	77 70%	
dissimilarity	00.2370		dissimilarity	//./0/0	
	Average	Cumulative		Average	Cumulative
Contributory taxa	contribution	contribution	Contributory taxa	contribution	contribution
	(%)	(%)		(%)	(%)
Tipulidae†	10.8	15.8	Ephemerellidae	16.0	20.6
Limnephilidae†	7.8	27.2	Lepidostomatidae†	13.1	37.5
Chironomidae	7.6	38.4	Limnephilidae†	12.8	53.9
Uenoidae	6.1	47.4	Perlodidae	12.0	69.3
Rhyacophilidae	5.5	55.5	Capniidae†	5.0	75.7
Ephemerellidae	4.9	62.7	Tipulidae†	5.0	82.1
Nemouridae†	4.0	68.6	Chironomidae	1.8	84.5
Perlodidae	3.9	74.3	Nemouridae†	1.8	86.7
Lepidostomatidae†	3.6	79.5	Chloroperlidae	1.6	88.7
Leptophlebiidae	3.4	84.5	Gammaridae†	1.5	90.7

<sup>†</sup>shredder invertebrates

707

708 Tab	le 4 Results of (	GLMMs testing	of the effects o	of stream typ	pes related to	geological	characteristics (	(G),
---------	-------------------	---------------	------------------	---------------	----------------	------------	-------------------	------

season (S), and their interactions on overall (a), microbial (b), and invertebrate (c) decomposition rates (k)

710 of leaf litter. *p*-values were produced in GLMM for the Wald Z-statistic. Bold letters indicate statistical

<i>i</i> i significance. SD represents standard deviations	71	]	l sig	nificance.	SD :	represents	standard	deviations
--	----	---	-------	------------	------	------------	----------	------------

(a) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	-3.687	0.2017	<0.001
G (volcanic relative to non-volcanic)	0.0483	0.1722	0.779
S (winter relative to summer)	-0.6046	0.2466	0.014
G×S	0.285	0.135	0.035

(b) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	-3.6691	0.1485	<0.001
G (volcanic relative to non-volcanic)	-0.3713	0.0775	<0.001
S (winter relative to summer)	-1.0229	0.2022	<0.001
$G \times S$	0.5336	0.0748	<0.001

713

(c) Coefficients	Estimates	SD	<i>p</i> -value
Intercept	-6.756	0.407	<0.001
G (volcanic relative to non-volcanic)	1.371	0.407	<0.001
S (winter relative to summer)	1.201	0.494	0.015
G×S	-1.086	0.398	0.006

714

715

# 716 Figure Captions

Fig. 1 Study sites in headwater watersheds of the Sorachi River, a tributary of the Ishikari River in Furano,
Central Hokkaido, Japan. Symbols denote the site locations and their catchments are indicated by solid
lines. The five volcanic sites are represented by open circles with site names accompanied by asterisks,
while open triangles represent the five non-volcanic sites. Map colors correspond to different rock types in
the area.

**Fig. 2** Daily average temperature of the two stream types during summer (top) and winter (bottom) obtained from temperature loggers measuring continuously at 15-minute intervals. Data shown are only for time periods where all temperature loggers were able to record at the same time. Mean value was obtained from simultaneous temperature readings of different sites for each stream type (n=4 for summer volcanic, one site was not included due to logger malfunction; n=5 for summer non-volcanic; n=5 for winter volcanic; n=5 for winter non-volcanic). Solid lines represent volcanic stream type and dashed lines represent nonvolcanic stream type. 729 Fig. 3 Boxplots showing taxonomic richness (a) and Shannon-Weiner diversity index (b) of the entire 730 invertebrate community and the abundance (c) and biomass (d) of shredder invertebrates in two seasons 731 (summer and winter) in the two stream types (non-volcanic and volcanic). Alphabetical letters indicate the 732 results of multiple comparisons among groups after GLMMs testing of effects of season, stream type, and 733 their interactions. For taxonomic richness, diversity index, and abundance, four groups were compared; 734 groups accompanied by the same letter are statistically indistinguishable. For biomass, group comparisons 735 were conducted for two main factors (season and stream type); unbracketed capital letters indicate 736 statistically different groups between seasons, while bracketed capital letters indicate different groups 737 between stream types across two seasons.

Fig. 4 Boxplots showing decomposition rate (*k*) for overall decomposition (a), microbe-mediated decomposition (b), and invertebrate-mediated decomposition (c) observed within each litter bag in two seasons (summer and winter) in the two stream types (non-volcanic and volcanic). Alphabetical letters indicate the results of multiple comparisons among groups after GLMMs testing for effects of season, stream type, and their interactions. In all cases, four groups were compared with each other; groups accompanied by the same letters are statistically indistinguishable.

744 Fig. 5 Graphical summary of results of SEM analyses in two seasons (summer and winter) for effects of 745 geology on microbe k via environmental variables (a, d), effects of geology on invertebrate k via 746 environmental variables, and shredder invertebrate abundance (b, e), and effects of geology on invertebrate 747 k via environmental variables and shredder invertebrate biomass (c, f). Only statistically significant 748 pathways are shown as arrows with positive effects in red and negative effects in blue. Standardized 749 coefficients of the effects are listed in boldface beside corresponding arrows. R-squared values (in normal 750 typeface) are shown for significantly explained response variables. Abbreviations: geo for volcanic 751 proportion; temp for temperature; disc for flow discharge; vel for local flow velocity; sed for the total 752amount of sediment on coarse litter bag; abun for abundance; mass for biomass; and TP for total phosphorus.

753

754

756		
757		
758		



Fig. 1 Tolod et al. 2022







Summer



Winter



Fig. 5 Tolod et al. 2022