



Title	Prominence of NUDT15 genetic variation associated with 6-mercaptopurine tolerance in a genome-wide association study of Japanese children with acute lymphoblastic leukaemia
Author(s)	Tanaka, Yoichi; Urayama, Kevin Y.; Mori, Makiko; Arakawa, Yuki; Hasegawa, Daisuke; Noguchi, Yasushi; Yanagimachi, Masakatsu; Keino, Dai; Ota, Setsuo; Akahane, Koshi; Inukai, Takeshi; Hangai, Mayumi; Kawaguchi, Takahisa; Takagi, Masatoshi; Koh, Katsuyoshi; Matsuda, Fumihiko; Manabe, Atsushi
Citation	British journal of haematology, 199(2), 260-269 https://doi.org/10.1111/bjh.18405
Issue Date	2022-10-01
Doc URL	http://hdl.handle.net/2115/90588
Rights	This is the peer reviewed version of the following article: Tanaka, Y, Urayama, KY, Mori, M, Arakawa, Y, Hasegawa, D & Noguchi, Y, et al. Prominence of NUDT15 genetic variation associated with 6-mercaptopurine tolerance in a genome-wide association study of Japanese children with acute lymphoblastic leukaemia. Br J Haematol. 2022; 199(2): 260– 269, which has been published in final form at 10.1111/bjh.18405. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley ' s version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.
Type	article (author version)
File Information	Br J Haematol_bjh.18405.pdf



[Instructions for use](#)

Prominence of *NUDT15* genetic variation associated with 6-mercaptopurine tolerance in a genome-wide association study of Japanese children with acute lymphoblastic leukemia

Yoichi Tanaka¹, PhD; Kevin Y. Urayama^{2,3}, PhD, MPH; Makiko Mori⁴, MD; Yuki Arakawa⁴, MD; Daisuke Hasegawa⁵, MD, PhD; Yasushi Noguchi⁶, MD; Masakatsu Yanagimachi^{7,8}, MD, PhD; Dai Keino^{7,9}, MD, PhD; Setsuo Ota¹⁰, MD, PhD; Koshi Akahane¹¹, MD, PhD; Takeshi Inukai¹¹, MD, PhD; Mayumi Hangai³, MD; Takahisa Kawaguchi¹², PhD; Masatoshi Takagi¹³, MD, PhD; Katsuyoshi Koh⁴, MD; Fumihiko Matsuda¹², PhD; Atsushi Manabe¹⁴, MD, PhD

¹Division of Medical Safety Sciences, National Institute of Health Sciences, Kanagawa, Japan

²Graduate School of Public Health, St. Luke's International University, Tokyo, Japan

³Department of Social Medicine, National Center for Child Health and Development, Tokyo, Japan

⁴Department of Hematology/Oncology, Saitama Children's Medical Center, Saitama, Japan

⁵Department of Pediatrics, St. Luke's International Hospital, Tokyo, Japan

⁶Department of Pediatrics, Japanese Red Cross Narita Hospital, Chiba, Japan

⁷Department of Hematology/Oncology, Children's Cancer Center, Kanagawa Children's Medical Center, Kanagawa, Japan

⁸Department of Pediatrics, Yokohama City University Hospital, Kanagawa, Japan

⁹Department of Pediatrics, St. Marianna University, Kanagawa, Japan

¹⁰Department of Pediatrics, Teikyo University Chiba Medical Center, Chiba, Japan

¹¹Department of Pediatrics, University of Yamanashi, Yamanashi, Japan

¹²Center for Genomic Medicine, Kyoto University, Kyoto, Japan

¹³ Department of Pediatrics, Tokyo Medical and Dental University, Tokyo, Japan

¹⁴Department of Pediatrics, Hokkaido University, Hokkaido, Japan

Correspondence: Yoichi Tanaka

Division of Medical Safety Sciences, National Institute of Health Sciences

3-25-26, Tonomachi, Kawasaki-ku, Kawasaki-shi, Kanagawa 210-9501 JAPAN

Tel: +81-44-270-6626; FAX: +81-44-270-6627

E-mail: tanakayoichi@nihs.go.jp

Running Title: Genetic risk factors for 6-MP tolerance in Japanese ALL

Keywords: 6-mercaptopurine, tolerance, genetic variant, genome-wide association study, Japanese

Counts: Abstract- 193 words; Text- 3,106 words; Tables- 3; Figure- 2; References- 37

Summary

Inherited genetic variation is associated with 6-mercaptopurine (6-MP) dose reduction and frequent 6-MP induced toxicities. However, tolerable dose for 6-MP is not completely predicted by the known variation in *NUDT15* and *TPMT* among Asian children with acute lymphoblastic leukemia (ALL). We performed a genome-wide association study (GWAS) related to 6-MP dose among Japanese children with ALL. This GWAS comprised 224 patients previously enrolled in Tokyo Children's Cancer Study Group clinical studies with replication attempted in 55 patients. Genome-wide single nucleotide polymorphism (SNP) genotypes were evaluated for association with 6-MP average dose during initial 168 days of maintenance therapy. Possible associations were observed across 5 gene coding regions, among which only variants at 13q14.2 were genome-wide significant and replicated (rs116855232, *NUDT15*, $\beta=-10.99$, $P=3.7\times 10^{-13}$). Notable findings were observed for variants in *AFF3* (rs75364948, $P=2.05\times 10^{-6}$) and *CHST11* (rs1148407, $P=2.09\times 10^{-6}$), but were not replicated possibly due to small numbers. A previously reported candidate SNP in *MTHFR* was associated with higher 6-MP average dose (rs1801133, $P=0.045$), and *FOLH1* (rs12574928) was associated in a candidate region evaluation ($P_{adjust}=0.013$). This study provides strong evidence that rs116855232 in *NUDT15* is the prominent genetic factor associated with 6-MP tolerable dose in Japanese.

Introduction

A survival probability of 80-90% in children with acute lymphoblastic leukemia (ALL) has been achieved due largely in part to advances in combination chemotherapy¹. 6-mercaptopurine (6-MP) is a main component for improving therapeutic outcomes, but tolerability is different in each patient. Failure to minimize incidents of therapy interruption may affect prognostic outcomes for childhood ALL patients². Response to 6-MP has been associated with variants in genes participating in the 6-MP metabolism pathway³. It is well known that genetic variation of the *TPMT* gene is associated with 6-MP intolerance³. However, the *TPMT* variant was not shown to be associated with thiopurine induced toxicities in Japanese⁴, and the frequencies of *TPMT* poor metabolizer was lower in East Asians compared with other races. In 2014, the *NUDT15* rs116855232 variant was reported to influence 6-MP intolerance identified through a genome-wide association study (GWAS), and the frequency of that variant is higher in Asians compared to populations of European and African ancestries⁵. Subsequently, *NUDT15* genotype was shown to affect 6-MP tolerability among Asians undergoing childhood ALL therapy⁴.

To date, several candidate gene studies have shown that genetic variation in *NUDT15* is associated with 6-MP dose reduction and frequent 6-MP-induced toxicities in childhood ALL^{4,6-13}. *NUDT15* enzyme dephosphorylates thio-guanosine triphosphate (GTP) and deoxy thio-GTP to thio-guanosine monophosphate (GMP) and deoxy thio-GMP, respectively. Lower *NUDT15* enzyme leads to increased thioguanosine incorporation ratio to 6-MP dose and 6-MP related toxicities. In some instances, severe toxicities requiring significant 6-MP dose reduction have been observed in patients who do not carry the known genetic risk factors for 6-MP tolerability. Thus, tolerable dose for 6-MP is not completely predicted by the variation in *NUDT15* and *TPMT*.

The aim of this study was to perform a GWAS to examine the role of inherited genetic variation related to 6-MP dose, with objectives to both identify newly associated variation, as well as to

characterize the variants previously reported in diverse populations, among Japanese childhood ALL patients. Moreover, we pursued a targeted examination of genetic variation across the candidate gene regions to identify additional associations with 6-MP dose.

Methods

Patients and sample collection

This GWAS comprised patients previously enrolled in a Tokyo Children's Cancer Study Group (TCCSG) clinical study including L89-12^{14,15}, L95-14^{15,16}, L99-15¹⁷ and L04-16¹⁸ from 23 clinical centers collaborating on this genomic study. Patients were recruited at the time of routine outpatient follow-up visit between 2013 and 2015, as previously described¹⁹. Briefly, patients were considered eligible if they were aged 19 years or younger at the time of ALL diagnosis and self-identified as Japanese. Saliva samples were collected at the time of follow-up visit after remission. DNA was extracted using the Oragene prepIT DNA extraction kit (DNA Genotek, Ottawa, Canada), and stored at -80 °C. The study protocol was approved by the institutional review boards of all collaborating institute and hospital involved in patient recruitment. Written informed consent was obtained from the parents of each participant together with a written informed consent or assent by the child depend on age.

Genotyping and quality control

Whole genome single nucleotide polymorphism (SNP) microarray genotyping was performed using the Illumina HumanCoreExome-12 v1.1 BeadChip (San Diego, CA) and has been described previously¹⁹. Briefly, quality control (QC) procedures comprised excluding samples if the genotype call-rate was below 95% and samples that exhibited relatedness based on an identity-by-descent analysis. SNPs were excluded if genotype call-rate was less than 99%, genotype distribution deviated

from that expected based on Hardy-Weinberg equilibrium ($P > 1 \times 10^{-6}$), or the minor allele frequency (MAF) was less than 0.01. Additionally, principal components (PC) analysis based on a subset of the post-QC genome-wide SNPs in low linkage disequilibrium (LD) was performed using EIGENSTRAT 2.0 software together with HapMap data from Japanese, and sample outliers were excluded based on a plot of the leading PCs.

Genome-wide SNP imputation was performed using ShapeIT2 and Minimac4 with reference population from the 1000 Genomes Project Phase III Version 5. SNP imputation QC comprised excluding poorly imputed SNPs defined by an R^2 of less than 0.5, resulting in a total of 6,236,137 SNPs available for analysis.

Clinical data

The clinical data and details of the administration of 6-MP doses during maintenance therapy were available for 289 Japanese patients. In TCCSG protocols, maintenance therapy is initiated with 40 mg/m²/day of 6-MP and 25 mg/m²/week of oral methotrexate. These dosages were adjusted to maintain the target leucocyte count at 1,500 – 3,000/mm³. We excluded patients who started at more than 125% or less than 75% of protocol 6-MP dose, as therapeutic dose for 6-MP and MTX were adjusted for drug induced toxicities before initiation of maintenance therapy. In total, we included 224 ALL patients (discovery cohort) who started maintenance therapy using the normal protocol dose (30–50 mg/m²/day). The outcome variable for this GWAS was defined as 6-MP average dose for initial 168 days of maintenance therapy.

Replication series

The replication series included 55 patients who started maintenance therapy by normal protocol recruited previously through TCCSG as part of a separate study⁴. DNA were extracted from peripheral

blood obtained at remission, and SNP genotyping was performed using Taqman real-time PCR assays (Applied Biosystems, Waltham, MA).

Statistical analysis

We performed genome-wide association analyses of SNPs in relation to 6-MP average dose for initial 168 days of maintenance therapy among the discovery cohort using linear regression assuming a log-additive genetic model of inheritance and adjusting for age at diagnosis. Association analysis assuming dominant and recessive genetic inheritance models were also performed. Results showing a $P < 5 \times 10^{-8}$ were considered statistically significance at the genome-wide level, and a $P < 1 \times 10^{-5}$ was considered as showing a suggestive association. In the replication series, association analysis was conducted similarly using linear regression assuming a log-additive genetic model and adjusting for age at diagnosis. We defined a Bonferroni corrected $P < 0.05$ as statistically significant in the replication. Additionally, among the top SNPs, the difference of 6-MP average dose across the three possible genotypes was evaluated using the Kruskal-Wallis test. Normality test of 6-MP distribution was evaluated using the D'Agostino-Pearson test.

The association of specific candidate SNPs reported previously, as well as the variants in coding regions of those candidate genes were examined. Association results with a nominal p-value of less than 0.05 was considered statistically significant for specific candidate SNPs. For regional examination of association results across candidate genes, SNPs with a nominal p-value of less than 0.05 were considered noteworthy, and a condensed list of SNPs pruned on LD ($r^2 > 0.50$) within each gene were adjusted for multiple testing based on 10,000 permutation of the data on the 6-MP dose outcome in which a p-value below a family-wise type I error rate threshold of 0.05 was considered statistically significant. Statistical analyses were performed using PLINK version 1.9, R software (version 3.6.1) and Prism 9 (GraphPad Software, San Diego, CA).

Results

Genome-wide association analysis of 6-MP average dose during 168 days of maintenance therapy

A total of 224 patients were included in the genome-wide association analysis. No patients needed to be excluded due to outlying genetic ancestry based on PC analysis evaluations. The characteristics of patients were shown in Table 1. Age at diagnosis was significantly correlated with 6-MP average dose ($\beta = -0.88$, $P = 6.96 \times 10^{-5}$). The median 6-MP average doses for the initial 168 days were 41.1 (interquartile range: IQR 25% – 75%, 32.8 – 48.6), and did not appear to deviate from a normal distribution ($P > 0.05$).

In the genome-wide analysis of the discovery cohort, linear regression of the 6-MP average dose adjusted for age at diagnosis showed minimal evidence of genomic inflation ($\lambda = 1.01$) (Supplementary Figure S1). The Manhattan plot of the results showed potential association of variants representing 5 genetic coding regions ($P < 1 \times 10^{-5}$) with 6-MP average dose (Figure 1, Table 2 and Supplementary Table S1). Genome-wide significant associations with 6-MP average dose were observed for variants at the chromosome 13q14.2 region in which the leading SNP was rs116855232 located in *NUDT15* SNP ($\beta = -11.45$, $P_{additive} = 8.5 \times 10^{-10}$, $P_{dominant} = 2.27 \times 10^{-9}$). Other variants within this cluster of associated SNPs were in LD with this SNP (Supplementary Figure S2, $r^2 > 0.8$). Suggestive associations were observed at chromosome 2q11.2 (rs75364948, *AFF3*), 2p21 (rs14452634, *THADA*), 12q23.3 (rs1148407, *CHST11*) and 16q23.2-3 (rs12934986 and rs10153053, *CMIP*) (Table 2). In a sensitivity analysis, the results excluding outlier values of 6-MP (defined as the values of more than the third IQR + $1.5 \times$ IQR or less than the first IQR + $1.5 \times$ IQR) showed a persistent association with the chromosome 13q14.2 region, and slight attenuation of association for the other loci (Supplementary Figure S3).

To identify associations in the absence of the effect of the well-known loci for 6-MP intolerance,

we excluded patients who were carriers of the *TPMT* rs1142345 and *NUDT15* rs116855232 variants. In this cohort, 7 and 41 patients were carriers of the *TPMT* and *NUDT15* variants, respectively. Only one patient carried both the *TPMT* and *NUDT15* variants. Among the remaining 178 patients, the genome-wide analysis of 6-MP average dose showed potential associations with variants representing 13 genetic coding regions ($P < 1 \times 10^{-5}$) (Supplementary Table S2). The results of the suggestive variants for *CHST11*, *CMIP* and *THADA* were similar to that of the full analysis.

Replication analysis

For the replication series, we selected three potentially associated SNPs that reside within genes that have been previously implicated in 6-MP tolerability or progression of hematological malignancy (Table 2). The association with *NUDT15* rs116855232 strongly replicated in this series ($P = 5.43 \times 10^{-5}$); however, while the direction of association was consistent, rs75364948 (*AFF3*) and rs1148407 (*CHST11*) were not significantly associated with 6-MP average dose (Table 2). Combining the estimates from the discovery and replication cohorts, a strong genome-wide significant association between rs116855232 (*NUDT15*) and 6-MP average dose was observed ($\beta = -10.99$, $P = 3.66 \times 10^{-13}$).

Variant rs116855232 in the *NUDT15* gene represents the prominent association with 6-MP average dose within our Japanese study. Among the combined discovery and replication cohorts, 51 patients were heterozygous for rs116855232 (C/T), 5 patients were homozygous for the variant (T/T). The median of the 6-MP average dose for patients with CT and TT were 31.6 and 13.1 mg/m²/day, respectively; these 6-MP doses were significantly lower compared to that of patients with the CC genotype (41.9 mg/m²/day, $P = 3.74 \times 10^{-11}$, Figure 2A).

Previously reported candidate genes and 6-MP tolerable dose

As a secondary aim, we selected candidate loci based on review of the literature and the

PharmGKB database (<https://www.pharmgkb.org>) and performed targeted examination of these candidate regions. SNPs rs1142345 (*TPMT*)⁵, rs3765534 (*ABCC4*)²⁰, rs1142345 (*ITPA*)^{21,22}, rs4149056 (*SLCO1B1*)²³, rs1801133 (*MTHFR*)²⁴, rs61886492 (*FOLH1*)²⁵, rs12199316 (*NHLRC1*)²⁶ and rs72846714 (*NT5C2*)²⁷ have been studied previously as candidate loci for 6-MP tolerable dose and/or toxicities. Among these, *MTHFR* rs1801133 showed a significant association with 6-MP average dose in our study ($\beta = 2.26$, $P = 0.045$). However, the other reported genetic variants were not related with 6-MP dose in this analysis (Table 3). In addition, other variants in LD with specific candidate SNPs were not associated. A range of genetic variation across these candidate genes were targeted to examine whether other SNPs may be associated with 6-MP average dose in Japanese. SNP associations showing nominal p-values of less than 0.05 were observed for *MTHFR*, *FOLH1*, *ABCC4*, and *ITPA* (Table 3). Among these, rs12574928 in *FOLH1* showed a statistically significant association with 6-MP average dose after adjusting for multiple testing ($\beta = -8.02$, $P = 0.013$).

Discussion

We observed strong evidence showing that genetic variation at the chromosome 13q14.2 region is significantly associated with 6-MP average dose for initial 168 days of maintenance therapy. Among the cluster of association signals, *NUDT15* rs116855232 was the leading SNP, and the other variants spanning the *SUCLA2* and *MED4* genes were in strong LD with this SNP. *NUDT15* genetic variation has been shown to effect thiopurine tolerance in our previous studies, as well as others from Asian countries^{4,12,28}. In this study, 19 of 279 patients experienced a reduction of average 6-MP dose during the 168 days to a level less than 20 mg/m²/day. Five of 19 patients (26%) who needed a dose reduction were carriers of the rs116855232 variant. Lower *NUDT15* enzyme leads to increased thioguanosine incorporation ratio to 6-MP dose and 6-MP related toxicities. Based on the characteristics of the

current study comprising an ethnically homogeneous group of Japanese children with ALL, *NUDT15* was observed to have the strongest effect on 6-MP tolerance, and the associated variant rs116855232 is more frequent in Asians than other races/ethnicities.

The *TPMT* variant is a well-known genetic risk factor for 6-MP tolerance³. In this population, only 10 patients (1.8%) carried the *TPMT* rs1142345 variant, and their 6-MP average dose was 40.0 (19.1 – 57.4) mg/m²/day. The MAF of rs1142345 in this Japanese cohort was lower than populations of European and African ancestry (MAF; 4 – 6%), and there were no patients who were homozygous carriers. While not significant possibly due to limitations in statistical power, the effect size of this variant was also weaker compared to reports from other races/ethnicities.

Interestingly, 2 patients with *TPMT* rs1142345 requiring 6-MP average dose of less than 20 mg/m², were carriers of the *NUDT15* rs116855232 variant. Yang et al. reported that patients with heterozygous genotypes for both *TPMT* rs1142345 and *NUDT15* rs116855232 had 30-60% of 6-MP standard dose during 6-months of maintenance therapy among the multi-ethnic patient cohort⁵. Choi et al. reported *NUDT15* rs116855232 was significantly associated with high 6-thioguanine nucleotide levels in erythrocytes, 6-MP dose ratio, and frequency of leukopenia, but *TPMT* and other variants were not associated among Korean children with ALL⁹. In our population, the average 6-MP dose for heterozygous carriers of *TPMT* rs1142345 was similar to wild-type patients, but the average dose for patients with both *TPMT* rs1142345 and *NUDT15* rs116855232 was 50% of tolerable dose of the wild-type patients. Moreover, the standard dose of 6-MP on protocol for childhood ALL in Asian countries²⁹ is set at 70% of the dose (40-60 mg/m²) applied to protocols in Europe and US (75 mg/m²). Although there does appear to be an effect of *TPMT* rs1142345 on 6-MP tolerability in Asians in the presence of *NUDT15* variation, these effects may be small due to low prevalence of the variant and a low standard dose. Results suggest that heterozygous variants of both *NUDT15* and *TPMT* may induce less tolerability for 6-MP than *NUDT15* heterozygous genotype alone.

Other genetic variants identified through candidate gene approaches have been reported to be associated with thiopurine tolerable dose. In our study, *MTHFR* rs1801133 was associated with 6-MP dose, but the direction of association is in contrast to other reports²⁴. *MTHFR* is involved in folate metabolism and rs1801133 has been associated with hepatotoxicity. However, our study showed rs1801133 to be associated with high 6-MP average dose. Other previously reported candidate genetic variants were not associated with 6-MP average dose in our study.

In maintenance therapy, 6-MP dose and therapeutic duration is different by country, and allele frequencies of candidate variants may be different. This circumstance makes validation of SNP associations and consistency a challenge. The candidate missense SNP rs61886492 in *FOLH1* could not be evaluated in the current study due to its absence in the Japanese population. This variant was previously reported to be associated with 6-MP mediated toxicity²⁵. Interestingly, regional evaluation of genetic variation of this gene in our study identified a significant association signal led by SNP rs12574928 located in an intronic region. This SNP has been described as an expression quantitative trait locus (eQTL) for *FOLH1* in multiple tissues as documented in the Genotype-Tissue Expression (GTEx) Portal. Due to LD broadly spread across this relatively short gene, localizing a defined causal region will require additional studies.

We showed that *NUDT15* rs116855232 is the most prominently associated genetic loci for 6-MP tolerance in Japanese children with ALL. However, 6 of 20 patients who required significant 6-MP dose reduction to a level below 20 mg/m²/day did not carry the known risk variants for 6-MP tolerance. Moreover, 17 patients needed to increase 6-MP dose to a level greater than 60 mg/m²/day. Five previously unreported variants located in four genes showed suggestive associations with 6-MP tolerance, including loci residing in *AFF3*, *CHST11*, *THADA* and *CMIP*. However, these suggestive variants are located in intronic regions, and their functional significance has not been reported. The associated *AFF3* rs75364948 and surrounding SNPs in strong LD showed to be potential eQTL for

AFF3 and *KIAA1211L*. *AFF3* encodes a tissue-restricted nuclear transcriptional activator, and *AFF3* upregulation was reported to be high in tamoxifen resistant breast cancer³⁰. While the function of *KIAA1211L* is not clear, a previous study of molecular markers among childhood leukemia patients showed this gene to be differentially expressed between patients with and without central nervous system involvement³¹. *CHSTII* was reported previously in relation to the efficacy of methotrexate treatment in rheumatoid arthritis³². *THADA* and *CMIP* genetic variants were identified in GWAS as a susceptibility locus for type 2 diabetes mellitus^{33,34}, and other diseases^{35,36}. These suggestive findings could not be replicated in our small patient series in the present study. Future activities will comprise securing additional patient numbers for a statistically robust opportunity to confirm these findings in further replication attempts.

In this study, 6-MP average dose for initial 168 days of maintenance therapy was negatively correlated with age at diagnosis. Therefore, the association of 6-MP and genetic variant was adjusted by age. Although there have been no reports on the relationship between 6-MP tolerable dose and age, some reports have shown that age was negatively correlated with TPMT activity³⁷.

There are some limitations of this study to acknowledge. Due the retrospective nature of this study, we decided to exclude patients who started at more than 125% or less than 75% of protocol 6-MP dose. These patients were not suitable for evaluation for 6-MP response to standard therapy in maintenance therapy because therapeutic dose for 6-MP and MTX were adjusted for drug induced toxicities before maintenance therapy. Therefore, it is possible that our study cohort underrepresented patients who deviated from the normal 6-MP tolerance. Despite this limitation, our study is meaningful in describing the responsible genetic factors of 6-MP tolerance among patients starting with standard protocol dose.

Conclusion

We provide strong evidence that *NUDT15* rs116855232 is the prominent genetic factor associated with 6-MP tolerable dose, and it currently represents the most meaningful clinically predictive marker in Japanese. These findings indicated that *NUDT15* genotyping should be considered prior to initiation of 6-MP treatment. Prospects for the identification of additional loci of 6-MP tolerance are high after further validation of suggestive loci observed in this Japanese study.

Acknowledgements

This work was supported by funding from St. Luke's Life Science Institute (Tokyo, Japan), the Japan Society for the Promotion of Science KAKENHI grant numbers (15K18932, 26253041), the Children's Cancer Association of Japan, and the Japan Leukemia Research Fund.

Conflict of interest

There are no conflicts of interest to declare.

Authorship contributions

Y.T., K.Y.U., M.T. and A.T. designed the experiments. Y.T., K.Y.U. and M.H. analyzed data. M.M., Y.A., D.H., Y.N., M.Y., D.K., S.O., K.A., T.I., M.T. K.K., and A.M. collected the clinical data. T.K. and F.M. contributed to perform genotyping. Y.T. and K.Y.U. wrote the paper. All authors approved the manuscript and the interpretation of the data.

References

1. Kato M, Manabe A. Treatment and biology of pediatric acute lymphoblastic leukemia. *Pediatr Int*. 2018;60(1):4–12.
2. Relling M V., Hancock ML, Boyett JM, Pui CH, Evans WE. Prognostic importance of 6-mercaptopurine dose intensity in acute lymphoblastic leukemia. *Blood*. 1999;93(9):2817–23.

- 326 3. Relling M V, Schwab M, Whirl-Carrillo M, Suarez-Kurtz G, Pui CH, Stein CM, et al. Clinical
327 Pharmacogenetics Implementation Consortium Guideline for Thiopurine Dosing Based on TPMT
328 and NUDT 15 Genotypes: 2018 Update. *Clin Pharmacol Ther.* 2019;105(5):1095–105.
- 329 4. Tanaka Y, Kato M, Hasegawa D, Urayama KY, Nakadate H, Kondoh K, et al. Susceptibility to 6-
330 MP toxicity conferred by a NUDT15 variant in Japanese children with acute lymphoblastic
331 leukaemia. *Br J Haematol.* 2015;171(1):109–15.
- 332 5. Yang JJ, Landier W, Yang W, Liu C, Hageman L, Cheng C, et al. Inherited NUDT15 Variant Is a
333 Genetic Determinant of Mercaptopurine Intolerance in Children With Acute Lymphoblastic
334 Leukemia. *J Clin Oncol.* 2015;33(11):1235–42.
- 335 6. Wang DS, Yu CH, Lin CY, Chang YH, Lin KH, Lin DT, et al. Childhood acute lymphoblastic
336 leukemia mercaptopurine intolerance is associated with NUDT15 variants. *Pediatr Res.*
337 2021;89(1):217–22.
- 338 7. Wahlund M, Nilsson A, Kahlin AZ, Broliden K, Myrberg IH, Appell ML, et al. The Role of
339 TPMT, ITPA, and NUDT15 Variants during Mercaptopurine Treatment of Swedish Pediatric
340 Patients with Acute Lymphoblastic Leukemia. *J Pediatr.* 2020;216:150-157.e1.
- 341 8. Buaboonnam J, Sripatanatadasakul P, Treesucon A, Glomglao W, Siraprapapat P, Narkbunnam
342 N, et al. Effect of NUDT15 on incidence of neutropenia in children with acute lymphoblastic
343 leukemia. *Pediatr Int.* 2019;61(8):754–8.
- 344 9. Choi R, Sohn I, Kim MJ, Woo HI, Lee JW, Ma Y, et al. Pathway genes and metabolites in
345 thiopurine therapy in Korean children with acute lymphoblastic leukaemia. *Br J Clin Pharmacol.*
346 2019;85(7):1585–97.
- 347 10. Khera S, Trehan A, Bhatia P, Singh M, Bansal D, Varma N. Prevalence of TPMT, ITPA and
348 NUDT 15 genetic polymorphisms and their relation to 6MP toxicity in north Indian children with
349 acute lymphoblastic leukemia. *Cancer Chemother Pharmacol.* 2019;83(2):341–8.

- 350 11. Soler AM, Olano N, Méndez Y, Lopes A, Silveira A, Dabezies A, et al. TPMT and NUDT15
351 genes are both related to mercaptopurine intolerance in acute lymphoblastic leukaemia patients
352 from Uruguay. *Br J Haematol.* 2018;181(2):252–5.
- 353 12. Liang DC, Yang CP, Liu HC, Jaing TH, Chen SH, Hung IJ, et al. NUDT15 gene polymorphism
354 related to mercaptopurine intolerance in Taiwan Chinese children with acute lymphoblastic
355 leukemia. *Pharmacogenomics J.* 2016;16(6):536–9.
- 356 13. Chiengthong K, Ittiwut C, Muensri S, Sophonphan J, Sosothikul D, Seksan P, et al. NUDT15
357 c.415C>T increases risk of 6-mercaptopurine induced myelosuppression during maintenance
358 therapy in children with acute lymphoblastic leukemia. *Haematologica.* 2016;101(1):e24–6.
- 359 14. Manabe A, Tsuchida M, Hanada R, Ikuta K, Toyoda Y, Okimoto Y, et al. Delay of the diagnostic
360 lumbar puncture and intrathecal chemotherapy in children with acute lymphoblastic leukemia
361 who undergo routine corticosteroid testing: Tokyo Children’s Cancer Study Group Study L89-12.
362 *J Clin Oncol.* 2001;19(13):3182–7.
- 363 15. Tsuchida M, Ohara A, Manabe A, Kumagai M, Shimada H, Kikuchi A, et al. Long-term results
364 of Tokyo children’s cancer study group trials for childhood acute lymphoblastic leukemia, 1984-
365 1999. *Leukemia.* 2010;24(2):383–96.
- 366 16. Igarashi S, Manabe A, Ohara A, Kumagai M, Saito T, Okimoto Y, et al. No advantage of
367 dexamethasone over prednisolone for the outcome of standard- and intermediate-risk childhood
368 acute lymphoblastic leukemia in the Tokyo Children’s Cancer Study Group L95-14 protocol. *J*
369 *Clin Oncol.* 2005;23(27):6489–98.
- 370 17. Manabe A, Ohara A, Hasegawa D, Koh K, Saito T, Kiyokawa N, et al. Significance of the
371 complete clearance of peripheral blasts after 7 days of prednisolone treatment in children with
372 acute lymphoblastic leukemia: The Tokyo Children’s Cancer Study Group Study L99-15.
373 *Haematologica.* 2008;93(8):1155–60.

- 374 18. Takahashi H, Kajiwara R, Kato M, Hasegawa D, Tomizawa D, Noguchi Y, et al. Treatment
375 outcome of children with acute lymphoblastic leukemia: the Tokyo Children's Cancer Study
376 Group (TCCSG) Study L04-16. *Int J Hematol*. 2018 Jul;108(1):98–108.
- 377 19. Urayama KY, Takagi M, Kawaguchi T, Matsuo K, Tanaka Y, Ayukawa Y, et al. Regional
378 evaluation of childhood acute lymphoblastic leukemia genetic susceptibility loci among Japanese.
379 *Sci Rep*. 2018;8(1):789.
- 380 20. Tanaka Y, Nakadate H, Kondoh K, Nakamura K, Koh K, Manabe A. Interaction between
381 NUDT15 and ABCC4 variants enhances intolerance of 6-mercaptopurine in Japanese patients
382 with childhood acute lymphoblastic leukemia. *Pharmacogenomics J*. 2018;18(2):275–80.
- 383 21. Stocco G, Cheok M, Crews K, Dervieux T, French D, Pei D, et al. Genetic Polymorphism of
384 Inosine Triphosphate Pyrophosphatase Is a Determinant of Mercaptopurine Metabolism and
385 Toxicity During Treatment for Acute Lymphoblastic Leukemia. *Clin Pharmacol Ther*.
386 2009;85(2):164–72.
- 387 22. Hareedy MS, El Desoky ES, Woillard JB, Thabet RH, Ali AM, Marquet P, et al. Genetic variants
388 in 6-mercaptopurine pathway as potential factors of hematological toxicity in acute lymphoblastic
389 leukemia patients. *Pharmacogenomics*. 2015;16(10):1119–34.
- 390 23. Eldem I, Yavuz D, Cumaogullari Ö, Ileri T, Ünal Ince E, Ertem M, et al. SLC01B1
391 polymorphisms are associated with drug intolerance in childhood leukemia maintenance therapy.
392 *J Pediatr Hematol Oncol*. 2018;40(5):e289–94.
- 393 24. Shimasaki N, Mori T, Torii C, Sato R, Shimada H, Tanigawara Y, et al. Influence of MTHFR and
394 RFC1 Polymorphisms on Toxicities During Maintenance Chemotherapy for Childhood Acute
395 Lymphoblastic Leukemia or Lymphoma. *J Pediatr Hematol Oncol*. 2008;30(5):347–52.
- 396 25. Dorababu P, Naushad SM, Linga VG, Gundeti S, Nagesh N, Kutala VK, et al. Genetic variants of
397 thiopurine and folate metabolic pathways determine 6-MP-mediated hematological toxicity in

childhood ALL. *Pharmacogenomics*. 2012;13(9):1001–8.

26. Moriyama T, Yang W, Smith C, Pui CH, Evans WE, Relling M V., et al. Comprehensive characterization of pharmacogenetic variants in TPMT and NUDT15 in children with acute lymphoblastic leukemia. *Pharmacogenet Genomics*. 2022;32(2):60–6.

27. Jiang C, Yang W, Moriyama T, Liu C, Smith C, Yang W, et al. Effects of NT5C2 Germline Variants on 6-Mercaptopurine Metabolism in Children With Acute Lymphoblastic Leukemia. *Clin Pharmacol Ther*. 2021;109(6):1538–45.

28. Chiengthong K, Ittiwut C, Muensri S, Sophonphan J, Sosothikul D, Seksan P, et al. NUDT15 c.415C>T increases risk of 6-mercaptopurine induced myelosuppression during maintenance therapy in children with acute lymphoblastic leukemia. *Haematologica*. 2016;101(1):e24-6.

29. Tanaka Y, Yeoh AEJ, Moriyama T, Li CK, Kudo K, Arakawa Y, et al. An international retrospective study for tolerability of 6-mercaptopurine on NUDT15 bi-allelic variants in children with acute lymphoblastic leukemia. *Haematologica*. 2021;106(7):2026–9.

30. Shi Y, Zhao Y, Zhang Y, AiErken N, Shao N, Ye R, et al. AFB3 upregulation mediates tamoxifen resistance in breast cancers. *J Exp Clin Cancer Res*. 2018;37(1):254.

31. Hicks C, Sitthi-Amorn J, Douglas J, Ramani R, Miele L, Vijayakumar V, et al. Molecular Analysis of Central Nervous System Disease Spectrum in Childhood Acute Lymphoblastic Leukemia. *Clin Med Insights Oncol*. 2016;10:5–15.

32. Aslibekyan S, Brown EE, Reynolds RJ, Redden DT, Morgan S, Baggott JE, et al. Genetic variants associated with methotrexate efficacy and toxicity in early rheumatoid arthritis: Results from the treatment of early aggressive rheumatoid arthritis trial. *Pharmacogenomics J*. 2014;14(1):48–53.

33. Vujkovic M, Keaton JM, Lynch JA, Miller DR, Zhou J, Tcheandjieu C, et al. Discovery of 318 new risk loci for type 2 diabetes and related vascular outcomes among 1.4 million participants in

a multi-ancestry meta-analysis. Nat Genet. 2020;52(7):680–91.

34. Keaton JM, Gao C, Guan M, Hellwege JN, Palmer ND, Pankow JS, et al. Genome-wide interaction with the insulin secretion locus MTNR1B reveals CMIP as a novel type 2 diabetes susceptibility gene in African Americans. Genet Epidemiol. 2018;42(6):559–70.
35. Liu JZ, van Sommeren S, Huang H, Ng SC, Alberts R, Takahashi A, et al. Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. Nat Genet. 2015;47(9):979–86.
36. Conti D V., Darst BF, Moss LC, Saunders EJ, Sheng X, Chou A, et al. Trans-ancestry genome-wide association meta-analysis of prostate cancer identifies new susceptibility loci and informs genetic risk prediction. Nat Genet. 2021;53(1):65–75.
37. Liu C, Yang W, Pei D, Cheng C, Smith C, Landier W, et al. Genomewide Approach Validates Thiopurine Methyltransferase Activity Is a Monogenic Pharmacogenomic Trait. Clin Pharmacol Ther. 2017;101(3):373–81.

Figure legends

Figure 1. Manhattan plot of results of the GWAS using linear regression adjusting for age at diagnosis.

Figure 2. Average 6-MP dose for 168 days from initial maintenance therapy in the combined Discovery and replication cohort (N = 279). Difference between genotypes were analyzed by the Kruskal-Wallis test. (A) *NUDT15* rs116855232; (B) *AFF3* rs75364948; (C) *CHST11* rs1148407.

Table 1. Demographic and clinical characteristics of patients

	Discovery	Replication
Patient number	224	55
Gender		
Female	99 (44.2%)	24 (43.6%)
Male	125 (55.8%)	31 (56.4%)
Age at diagnosis (years)		
0-1	14 (6.3%)	3 (5.5%)
2-4	104 (46.4%)	19 (34.5%)
5-10	79 (35.3%)	26 (47.3%)
11-19	27 (12.0%)	7 (12.7%)
Median (range)	4.7 (0.9 – 15.7)	5.0 (1 – 17)
6-mercaptopurine initial dose (mg/m ² , median)	39.9 (30.2 - 50.0)	40.1 (30.8 – 49.5)
6-mercaptopurine average dose for 168 days (mg/m ² , median)	41.1 (13.3 – 78.5)	37.2 (5.5 – 60.0)
methotrexate dose (mg/m ² /week, median)	24.5 (0 – 31.5)	24.7 (0 – 30)

Table 2. Results of the genome-wide association analysis of 6-mercaptopurine dose and replication (age adjusted)

Chr	Position (GRCh37)	Gene	SNP	MAF in discovery	Discovery (N = 224)		Replication (N = 55)		Combined (N = 279)	
					β (95% CI)	P	β (95% CI)	P	β (95% CI)	P
13	48619855	<i>NUDT15</i>	rs116855232	0.094	-11.45 (-14.95, -7.95)	8.46×10^{-10}	-9.13 (-13.29, -4.96)	5.43×10^{-5}	-10.99 (-13.82, -8.16)	3.66×10^{-13}
2	100399866	<i>AFF3</i>	rs75364948	0.203	6.49 (3.88, 9.10)	2.05×10^{-6}	0.84 (-4.76, 6.44)	0.766	5.83 (3.42, 8.24)	3.12×10^{-6}
12	105058593	<i>CHST11</i>	rs1148407	0.138	8.08 (4.83, 11.3)	2.09×10^{-6}	2.22 (-3.28, 7.74)	0.421	6.52 (3.56, 9.49)	2.03×10^{-5}
2	43501170	<i>THADA</i>	rs144526347	0.015	21.37 (12.74, 30.00)	2.28×10^{-6}	--	--	--	--
16	81553070	<i>CMIP</i>	rs12934986	0.277	5.63 (3.24, 8.01)	6.43×10^{-6}	--	--	--	--
16	81659083	<i>CMIP</i>	rs10153053	0.234	-5.91 (-8.40, -3.14)	5.79×10^{-6}	--	--	--	--

Abbreviations; Chr, chromosome; SNP, single nucleotide polymorphism; MAF, minor allele frequency.

Table 3. Results of the relationship between previously reported candidate genes and average 6-mercaptopurine dose for 168 days (age adjusted)

Chr	Position (GRCh37)	Gene	SNP	Function	Alleles	MAF	Discovery (N = 224)			Reference
							β (95% CI)	P (nominal)	P (adjusted)	
Previous reported SNPs										
1	11856378	<i>MTHFR</i>	rs1801133	Missense	G/A	0.375	2.26 (0.06, 4.46)	0.0454	--	24
6	18123502	<i>NHLRC1</i>	rs12199316	Upstream	C/G	0.394	-0.69 (-3.06, 1.69)	0.553	--	26
6	18130918	<i>TPMT</i>	rs1142345	Missense	T/C	0.0156	-2.38 (-11.45, 6.70)	0.608	--	2
10	104878454	<i>NT5C2</i>	rs72846714	Intron	G/A	0.024	3.75 (-6.00, 13.5)	0.452	--	27
11	49186274	<i>FOLH1</i>	rs61886492	Missense	G/A	--	--	--	--	25
12	21331549	<i>SLCO1B1</i>	rs4149056	Missense	T/C	0.147	-2.45 (-5.47, 0.57)	0.113	--	23
13	95815415	<i>ABCC4</i>	rs3765534	Missense	C/T	0.143	-1.66 (-4.88, 1.55)	0.310	--	20
20	3193842	<i>ITPA</i>	rs1127354	Missense	C/A	0.152	1.01 (-2.05, 4.07)	0.520	--	21, 22
Genetic variants across previously reported candidate gene regions (nominal P-value<0.05)										
11	49213504	<i>FOLH1</i>	rs12574928	Intron	C/T	0.056	-8.02 (-12.9, -3.21)	1.59 × 10 ⁻³	0.013	25
13	95707834	<i>ABCC4</i>	rs9561773	Intron	C/T	0.228	3.69 (1.02, 6.37)	7.35 × 10 ⁻³	0.577	20
13	95953517	<i>ABCC4</i>	rs11568681	Missense	G/T	0.018	10.91 (2.54, 19.29)	0.011	0.707	20
20	3200778	<i>ITPA</i>	rs959815466	Intron	C/T	0.013	-11.55 (-21.2, -1.90)	0.019	0.126	21,22

Abbreviations; Chr, chromosome; SNP, single nucleotide polymorphism; MAF, minor allele frequency.



