



Title	Heterologous complementation systems verify the mosaic distribution of three distinct protoporphyrinogen IX oxidase in the cyanobacterial phylum
Author(s)	Kohata, Ryoya; Lim, HyunSeok; Kanamoto, Yuki; Murakami, Akio; Fujita, Yuichi; Tanaka, Ayumi; Swingley, Wesley; Ito, Hisashi; Tanaka, Ryouichi
Citation	Journal of plant research, 136, 107-115 https://doi.org/10.1007/s10265-022-01423-7
Issue Date	2022-11-10
Doc URL	http://hdl.handle.net/2115/90680
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature 's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s10265-022-01423-7
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	PPOX-supplementary-figures-final.pdf (Supplementary Information)



[Instructions for use](#)

Electronic supplementary materials

Title:

Heterologous complementation systems verify the mosaic distribution of three distinct protoporphyrinogen IX oxidase in the cyanobacterium phylum.

Journal:

Journal of Plant Research

Author:

Ryoya Kohata, HyunSeok Lim, Yuki Kanamoto, Akio Murakami, Yuichi Fujita, Ayumi Tanaka, Wesley Swingley, Hisashi Ito and Ryouichi Tanaka

Corresponding author:


Ryouichi Tanaka

Institute of Low Temperature Science, Hokkaido University


Kita-ku, N19W8, Sapporo 060-0819, JAPAN

rtanaka@lowtem.hokudai.ac.jp


a Expression cassette for a PPOX homologue



b *hemJ* knock-out cassette



c Verification of *psbA2-Gv_hemY/ΔhemJ* transformant



d Verification of *psbA2-Te_hemY/ΔhemJ* transformant




Fig. S1. Disruption of the *hemJ* gene in the *psbA2-Gv_hemY/ΔhemJ* and *psbA2-Te_hemY/ΔhemJ* transformants. First, a cassette comprising the *psbA2* promoter, a PPOX homologue, the kanamycin resistance gene and the *psbA2* terminator (a) were introduced into the *Synechocystis* genome. Subsequently, the endogenous *HemJ* gene of these transformants were disrupted by inserting the chloramphenicol marker using the cassette shown in (b). Approximate positions of the primers used in panels (c) and (d) are shown with arrows in (b), which span the disrupted region of the *hemJ* sequence. Verification of segregation of the disrupted *hemJ* gene was assessed in the *psbA2-Gv_hemY/ΔhemJ* transformant (c) or the *psbA2-Te_hemY/ΔhemJ* transformant (d) by PCR using the primer set shown in B. The primer set gave a 250-bp product which amplified a part of the intact *hemJ* gene (lane WT of panels c and d). In contrast, the same primer set gave a 1.6-kbp product when the chloramphenicol marker was inserted into the *hemJ* locus (lane T of panels c and d). M: 1-kb DNA ladder from Watson BRG, WT: wild-type *Synechocystis* cells. N: no template control, T: *psbA2-Gv_hemY/ΔhemJ* transformant cells (c) or *psbA2-Te_hemY/ΔhemJ* transformant cells (d). It should be noted that a single gel image was cropped to show only relevant lanes in panels c and d. The original images are shown at the right side of cropped images to indicate no signs of inappropriate image manipulations.




Fig. S2. Verification of the *hemJ* replacement with the cassette of PPOX homologue and the kanamycin (Kn) resistance marker. (a) Verification of the replacement of *HemJ* with *Gv_hemG*. M: Marker (lambda DNA/Styl), W: wild type DNA, P: *Gv_hemG-Km* plasmid (positive control), N: No template (negative control), T: Synechocystis Kn resistant transformant. The original picture which is shown on the right side contained extra lanes which are not relevant to the analysis mentioned here. Therefore, only relevant lanes are shown in this panel. (b) Verification of the replacement of *hemJ* with *Pm_hemG*. M: Marker (lambda DNA/Styl), W: wild type DNA, P: *Pm_hemG-Km* plasmid (positive control), N: No template (negative control), T: Synechocystis Kn resistant transformant.




Fig. S3. Effects of acifluorfen on the growth of *psbA2-Gv_hemY/ΔhemJ*.

The cultures of WT and *psbA2-Gv_hemY/ΔhemJ* cells were divided into four groups and diluted to $OD_{750} = 0.1$. Acifluorfen dissolved in dimethylformamide was added to WT and *psbA2-Gv_hemY/ΔhemJ* at the final concentration of 1 mM, which groups were denoted as WT+A and *Gv_hemY+A*, respectively. As a control, dimethylformamide was added to the other groups, which were denoted as WT and *Gv_hemY*, respectively. $n = 3$. Error bars indicate standard deviation.