



Title	Measurement of serum procalcitonin concentrations in calves with bovine respiratory disease
Author(s)	Koshiishi, Tomoko; Shibutani, Shusaku; Chuma, Takehisa; Iwata, Hiroyuki
Citation	Japanese Journal of Veterinary Research, 71(2), 65-71 https://doi.org/10.57494/jjvr.71.2_65
Issue Date	2023-11-27
Doc URL	http://hdl.handle.net/2115/90821
Type	bulletin (article)
File Information	JJVR71-2_65-71_TomokoKoshiishi.pdf



[Instructions for use](#)

SHORT COMMUNICATION

Experimental Research

Measurement of serum procalcitonin concentrations in calves with bovine respiratory disease

Tomoko Koshiishi¹⁾, Shusaku Shibutani^{1,*}, Takehisa Chuma²⁾
and Hiroyuki Iwata^{1,*}

¹⁾ Laboratory of Veterinary Hygiene, The United Graduate School of Veterinary Science Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8515, Japan.

²⁾ Laboratory of Veterinary Public Health, Joint Faculty of Veterinary Medicine, Kagoshima University, 1-21-24 Korimoto, Kagoshima, 890-0065 Japan

Received for publication, January 17, 2023; accepted, May 1, 2023

Abstract

Bovine respiratory disease (BRD) is an important and complex disease caused by various pathogens and stressors. Although BRD is often caused by viruses, secondary bacterial infections can worsen symptoms and prognosis. Due to this viral-bacterial synergy, the evaluation of bacterial infections in BRD is important for the diagnosis and prognosis of BRD. In this study, we analyzed the serum concentrations of procalcitonin (PCT), a biomarker of bacterial infection, in 41 calves with BRD. We demonstrate that when calves are categorized into PCT-positive and PCT-negative groups based on serum PCT concentrations, the PCT-positive group shows more persistently high BRD scores. These findings suggest that the measurement of serum PCT is potentially useful for the diagnosis and prognosis of calves with BRD.

Key Words: bovine respiratory disease, procalcitonin, respiratory score

Bovine respiratory disease (BRD) is a complex disease caused by various pathogens, including viruses and bacteria. BRD is one of the most important diseases affecting beef and dairy calves because it causes retarded growth and affects industrial productivity, resulting in significant economic losses^{19,25)}. Therefore, effective tools are required for early diagnosis and prognosis.

Various biomarkers have been evaluated for the diagnosis of human community-acquired pneumonia, including procalcitonin (PCT), C-reactive protein (CRP), tumor necrosis factor-related apoptosis ligand, interferon- γ -induced protein 10, erythrocyte sedimentation rate, white blood cell (WBC) count, and neutrophil percentage. Among these, PCT has shown

* Corresponding author:

Shusaku Shibutani

Laboratory of Veterinary Hygiene, The United Graduate School of Veterinary Science Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8515, Japan

FAX: 083-933-5900

E-mail: shushibu@yamaguchi-u.ac.jp

Hiroyuki Iwata

Laboratory of Veterinary Hygiene, The United Graduate School of Veterinary Science Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8515, Japan

FAX: 083-933-5899

E-mail: hiwata@yamaguchi-u.ac.jp

doi: 10.57494/jjvr.71.2_65

Table 1. Clinical signs and blood chemistry results on day 1 in relation to PCT status

	PCT+ group	PCT- group	Normal value ^{a)}
Age (days)	218.5 ± 25.7 ^{b)}	172.0 ± 18.6	
Temperature (°C)	40.2 ± 0.3	40.4 ± 0.1	38-39
Heart rate (/min)	109.5 ± 6.2	119.3 ± 5.4	40-80
Respiration (/min)	72.0 ± 12.7	63.1 ± 3.6	12-36
BRD score	9.1 ± 1.4	9.4 ± 0.6	
WBC (/μl)	8870 ± 621	10028 ± 590	4000-12000
Neu (/μl)	2105 ± 427	2021 ± 319	600-4000
Band (/μl)	163 ± 55	121 ± 56	0-120
Lym (/μl)	6388 ± 406*	7839 ± 453	2500-7500
Mon (/μl)	250 ± 81	162 ± 39	25-840
Eos (/μl)	155 ± 63	81 ± 24	0-2400
RBC (×10 ⁴ /l)	803 ± 10	801 ± 24	500-1000
Ht (%)	32.8 ± 0.8	33.9 ± 1.0	29-39
Hb (g/dl)	10.7 ± 0.1	11.3 ± 0.7	8-15
AST (IU/l)	67.9 ± 4.6*	52.7 ± 2.2	43-127
TP (g/dl)	6.5 ± 0.3	6.4 ± 0.1	6.7-7.5
Alb (g/dl)	2.4 ± 0.1	2.8 ± 0.2	3-3.6
Tchol (mg/dl)	88.8 ± 9.4	100.0 ± 7.7	80-120
Glu (mg/dl)	77.3 ± 5.7	78.9 ± 3.9	45-75
BUN (mg/dl)	11.7 ± 1.0	10.8 ± 1.0	20-30
NEFA (μEq/l)	242.1 ± 49.7	224.0 ± 25.2	144.6 ± 4.3

a) Normal value cited from the references [5, 14, 15, 22]

b) Data represent mean ± SE (*: $P < 0.05$ vs PCT- group; Welch's t test)**Table 2.** Clinical symptoms of the 41 calves diagnosed with BRD on initial examination

	n	(%)
Auscultation of abnormal lung sounds		
Discontinuous sounds		
Coarse crackles	2	(4.9)
Fine crackles	0	(0)
Continuous sounds		
Wheezes	8	(19.5)
Rhonchi	3	(7.3)
Pleural friction rub	1	(2.4)
Bronchial sounds in the lung fields	11	(26.8)
Clinical signs		
Abdominal respiration	7	(17.1)
Oral respiration	1	(2.4)
Cough	41	(100)
Purulent nasal discharge	2	(4.9)
Dyspnea	22	(53.7)
Polypnea ^{a)}	36	(87.8)
Low auricular reflex	19	(46.3)
Cyanosis	0	(0)

a) Normal respiratory rates of adult and young (< 30 days of age) are 12-36 and 30-60, respectively.

excellent diagnostic and prognostic performance for severe bacterial infections^{6,16,20,23)}. PCT is a precursor of the calcium-regulating hormone calcitonin (CT). CT is produced intracellularly by proteolytic splicing of PCT and is normally secreted by parafollicular cells of the thyroid gland in response to hypercalcemia. In contrast, PCT does not enter the bloodstream of healthy animals. However, in response to bacterial infection, PCT is induced directly by lipopolysaccharide and other bacterial toxins and indirectly by proinflammatory cytokines, such as interleukin (IL)-1 β , tumor necrosis factor (TNF)- α , and IL-6, which are secreted in response to infection¹⁸⁾. Prominent increases in serum PCT concentrations were observed in systemic bacterial infections, but not in noninfectious inflammatory responses or viral infections^{7,11)}. During viral infection, PCT expression is suppressed by interferon- γ , which plays a central role in antiviral immunity¹²⁾. Bacterial infection can worsen the severity of BRD even if the primary cause of the disease is a viral infection. This interaction is called viral-bacterial synergy¹⁰⁾. Therefore, the severity of BRD may be

related to biomarkers specific to bacterial infections such as PCT.

In addition to humans, elevated serum PCT has been detected in mice^{3,26)}, hamsters¹⁷⁾, pigs²⁴⁾, horses²⁾, and calves^{4,9)} in response to bacterial infection. Similarly, blood PCT concentrations increase in calves with sepsis^{4,9)}. A previous study has shown that serum concentrations of PCT, along with other biological markers, are increased in BRD-suspected calves with any of the BRD symptoms when compared with healthy calves⁸⁾. However, whether PCT concentrations of BRD are associated with prognosis remains unclear. Therefore, in this study, we investigated the relationship between serum PCT positivity on the first medical day (day 1) and the subsequent severity of BRD using a scoring system that quantifies the severity of BRD.

This study included 41 calves (30 Japanese Black, 1 Holstein, and 10 first filial (F1) generations between Japanese Black and Holstein) fed commercial feedlots in Ehime prefecture, Japan, which were diagnosed with BRD. These calves were examined for California BRD scores,




Fig. 1. BRD scores in relation to PCT status

(A) Distribution of BRD scores in the BRD calves of the PCT+ (upper panels, $n = 10$) and PCT- (lower panels, $n = 31$) groups.

(B) Time course of BRD scores in the BRD calves of the PCT+ and PCT- groups. Data represent mean \pm SE. The BRD scores of the PCT+ group are significantly higher than those of the PCT- group on days 3 and 7 (**; $P < 0.01$; Mann-Whitney's U test)

which were determined by the presence/absence of an eye discharge (score of 0 or 2), nasal discharge (score of 0 or 4), ear droop/head tilt (score of 0 or 5), cough (score of 0 or 2), forced breathing (score of 0 or 2), and rectal temperature (score of 0 or 2; $\geq 39.2^\circ\text{C}$)¹⁾. BRD scores of ≥ 5 are proposed to be “BRD positive”¹³⁾. Most (37 of 41) calves exhibited BRD scores of ≥ 5 on day 1. The BRD score was also evaluated on days 2, 3, and 7. The ages of the calves ranged from 38 to 399 days. None of the patients died during the study. More detailed clinical findings, including blood cell counts, serum biochemistry, and clinical symptoms, are summarized in Tables 1 and 2. BRD cases showed low serum albumin, increased band neutrophils, slightly increased lymphocytes, decreased blood

urea nitrogen (BUN), and increased levels of non-esterified fatty acids (Table 1). Auscultation revealed coarse crackles in two cases (4.9%), wheezing in eight cases (19.5%), rhonchi in three cases (7.3%), pleural friction rub in one case (2.4%) and bronchial sounds in the lung fields in 11 cases (26.8%) (Table 2). In addition to BRD calves, five healthy F1 calves were used as controls.

Whole blood and serum samples were collected from the jugular vein of the calves on days 1, 2, 3, and 7 using collection tubes with and without an anticoagulant (heparin). Whole blood samples were immediately analyzed using a multi-parameter automated blood cell counter (KX-21NV; Sysmex, Kobe, Japan) and tested for differential WBC counts. Blood samples were centrifuged at $2500\times g$




Fig. 2. Time course of lymphocyte counts, monocyte counts, and AST concentrations in relation to PCT status

(A) Time course of lymphocyte counts in BRD calves of the PCT+ and PCT- groups.

(B) Time course of monocyte counts in the BRD calves of the PCT+ and PCT- groups.

(C) Time course of AST concentrations in the BRD calves of the PCT+ and PCT- groups.

Data represent mean \pm SE. The asterisks indicate significant differences between the groups (*: $P < 0.05$; Welch's *t* test).

for 20 min at 4°C and stored at -20°C until analysis of blood biochemistry using a chemistry analyzer (AU400; Beckman Coulter, Brea, CA, USA). Serum PCT concentration was measured using a bovine procalcitonin ELISA kit (Cusabio Biotech, Houston, TX, USA). The detection range of the ELISA kit is 40–2500 pg/mL. In all control calves, serum PCT concentrations were below the detection limit. In contrast, elevated serum PCT

concentrations were detected in 10 of the 41 calves with BRD on day 1 or day 2. The calves were categorized into the PCT+ group. The PCT+ calves continuously showed high PCT concentrations during the test period (mean \pm SE values of 411 ± 160 , 358 ± 156 , 569 ± 231 , and 530 ± 254 pg/mL on days 1, 2, 3, and 7, respectively). The remaining 31 BRD calves, in which serum PCT concentrations were lower than the detection limit (40 pg/mL) on day 1, were categorized into the PCT- group.

The relationship between BRD scores (days 1, 2, 3, and 7) and PCT positive/negative status (day 1) is shown in Fig. 1. Although the distribution of the BRD scores was not significantly different on day 1 (Fig. 1A), the BRD scores of the PCT+ group were significantly higher than those of the PCT- group on days 3 and 7 (Fig. 1B; $P = 0.003$ and 0.048 , respectively; Mann-Whitney's *U* test).

The clinical signs and serum biochemistry were then analyzed in relation to the positive/negative status of PCT on day 1. Table 1 lists the measurements on day 1. Among the measured parameters, the lymphocyte count and aspartate aminotransferase (AST) concentration showed significant differences on day 1. Lymphocyte counts in the PCT+ group were significantly lower than those of the PCT- group, and AST concentrations in the PCT+ group were significantly higher than those in the PCT- group (Table 1). The other parameters did not show statistical significance between the PCT+ and PCT- groups on day 1. The time course of lymphocyte counts showed that lymphocyte numbers were higher in the PCT- group on days 1 and 2 (Fig. 2A). In contrast, we observed that monocyte counts on day 2 were significantly higher in the PCT+ group than in the PCT- group (Fig. 2B). Monocytes are increased by acute stress and in the healing phase of acute and chronic infections²¹⁾, suggesting that elevated PCT concentrations might reflect these activities. AST concentrations were also higher in the PCT+ group than in the PCT- group on day 1 (Fig. 2C). Elevated AST concentrations may have resulted from more severe tissue damage in the PCT+ group. Although lymphocytes, monocytes, and AST showed significant differences between the

PCT+ and PCT− groups, these values were within the normal range in both groups, and thus, using them alone as biomarkers will be difficult. In contrast, serum PCT is not usually detected in uninfected healthy calves, making it particularly specific for infection by bacteria and useful as a biomarker.

Bacteriological tests for two major bacterial species that cause BRD, *Pasteurella multocida* and *Mannheimia haemolytica*⁸⁾, were performed on 11 cases (4 PCT+ and 7 PCT−). On Day 1, samples were taken from the right and left nasal cavities of affected calves using sterile swabs, applied to 5% sheep blood agar plates (Nissui Pharmaceutical, Tokyo, Japan), and incubated aerobically at 37°C for 24 h. *Pasteurella multocida* and *Mannheimia haemolytica* were attempted to be identified by microscopic observation after Gram staining and using ID Test HN-20 Rapid (Nissui Pharmaceutical) according to the analysis profile of the kit. *Pasteurella multocida* was isolated in all cases except for one PCT− case. In contrast, *Mannheimia haemolytica* was not detected in any case. Thus, at least for *Pasteurella multocida* and *Mannheimia haemolytica*, no clear association was noted between bacterial infection and PCT positive/negative status.

BRD is often diagnosed based on clinical signs (e.g., coughing, breathing difficulty, and fever), physical examination, auscultation, and laboratory tests. The BRD scoring system used in this study provides a simple and convenient method for quantifying multiple clinical signs. While it is useful, the distribution of the BRD scores was not apparently different between the PCT+ and PCT− groups (Fig. 1). However, our data indicated that PCT status on day 1 correlated with BRD scores on days 3 and 7 (Fig. 1B). These findings suggest that measurement of serum PCT concentrations can provide an additional layer of information for the diagnosis and prognosis of BRD; the PCT-positive status on day 1 is associated with subsequent severity of BRD. The specificity of PCT for bacterial infection is particularly important for the evaluation of the state of viral-bacterial synergy that exacerbates symptoms. In summary, this study showed the

potential diagnostic and prognostic utility of PCT measurement in BRD. Further investigation of the relationship between changes in serum PCT concentration and the various bacterial species underlying the pathogenesis of BRD, as well as changes in PCT in animals with more chronic or severe BRD, will provide a more accurate and extensive understanding of the PCT response to BRD.

Acknowledgments

The authors thank the members of the Laboratory of Veterinary Hygiene for their helpful discussions. This study was supported by JSPS KAKENHI (Grant Number 15K07745) provided to H. I.

Conflict of interest

The authors declare that they have no conflict of interest regarding the content of this manuscript.

References

- 1) Aly SS, Love WJ, Williams DR, Lehenbauer TW, Van Eenennaam A, Drake C, Kass PH, Farver TB. Agreement between bovine respiratory disease scoring systems for pre-weaned dairy calves. *Anim Health Res Rev* 15, 148–150, 2014. doi: 10.1017/S1466252314000164.
- 2) Barton AK, Pelli A, Rieger M, Gehlen H. Procalcitonin as a biomarker in equine chronic pneumopathies. *BMC Vet Res* 12, 281, 2016. doi: 10.1186/s12917-016-0912-4.
- 3) Biju PG, Garg S, Wang W, Choudhry MA, Kovacs EJ, Fink LM, Hauer-Jensen M. Procalcitonin as a predictive biomarker for total body irradiation-induced bacterial load and lethality in mice. *Shock* 38, 170–176, 2012. doi: 10.1097/SHK.0b013e31825b2db3.
- 4) Bonelli F, Meucci V, Divers TJ, Boccardo A,

- Pravettoni D, Meylan M, Belloli AG, Sgorbini M. Plasma procalcitonin concentration in healthy calves and those with septic systemic inflammatory response syndrome. *Vet J* 234,61–65, 2018. doi: 10.1016/j.tvjl.2018.02.003.
- 5) Carlson GP. Clinical chemistry tests. In: Large animal internal medicine, 4th ed. Smith BP. ed. Elsevier, St. Louis. pp. 375–397, 2009.
 - 6) De Azevedo JRA, Torres OJM, Czczeko NG, Tuon FF, Nassif PAN, De Souza GD. Procalcitonin as a prognostic biomarker of severe sepsis and septic shock. *Rev Col Bras Cir* 39, 456–461, 2012. doi: 10.1590/s0100-69912012000600003.
 - 7) Eberhard OK, Haubitz M, Brunkhorst FM, Kliem V, Koch KM, Brunkhorst R. Usefulness of procalcitonin for differentiation between activity of systemic autoimmune disease (systemic lupus erythematosus/systemic antineutrophil cytoplasmic antibody-associated vasculitis) and invasive bacterial infection. *Arthritis Rheum* 40, 1250–1256, 1997. doi: 10.1002/art.1780400709.
 - 8) El-Deeb W, Elsohaby I, Fayez M, Mkrtchyan HV, El-Etriby D, ElGioushy M. Use of procalcitonin, neopterin, haptoglobin, serum amyloid A and proinflammatory cytokines in diagnosis and prognosis of bovine respiratory disease in feedlot calves under field conditions. *Acta Trop* 204, 105336, 2020. doi: 10.1016/j.actatropica.2020.105336.
 - 9) Ercan N, Tuzcu N, Başbug O, Tuzcu M, Alim A. Diagnostic value of serum procalcitonin, neopterin, and gamma interferon in neonatal calves with septicemic colibacillosis. *J Vet Diagn Invest* 28, 180–183, 2016. doi: 10.1177/1040638715626488.
 - 10) Hodgson PD, Aich P, Manuja A, Hokamp K, Roche FM, Brinkman FSL, Potter A, Babiuk LA, Griebel PJ. Effect of stress on viral-bacterial synergy in bovine respiratory disease: novel mechanisms to regulate inflammation. *Comp Funct Genomics* 6, 244–250, 2005. doi: 10.1002/cfg.474.
 - 11) Jarešová M, Stríz I, Čermáková J, Lácha J, Sedláček J, Mudra K, Hána I, Vítko Š. Serum procalcitonin concentrations in transplant patients with acute rejection and bacterial infections. *Immunol Lett* 69, 355–358, 1999. doi: 10.1016/s0165-2478(99)00120-0.
 - 12) Linscheid P, Seboek D, Nylen ES, Langer I, Schlatter M, Becker KL, Keller U, Müller B. In vitro and in vivo calcitonin I gene expression in parenchymal cells: a novel product of human adipose tissue. *Endocrinology* 144, 5578–5584, 2003. doi: 10.1210/en.2003-0854.
 - 13) Love WJ, Lehenbauer TW, Kass PH, Van Eenennaam AL, Aly SS. Development of a novel clinical scoring system for on-farm diagnosis of bovine respiratory disease in pre-weaned dairy calves. *PeerJ* 2, e238, 2014. doi: 10.7717/peerj.238.
 - 14) Morris DD. Alterations in the erythron. In: Large animal internal medicine, 4th ed. Smith BP. ed. Elsevier, St. Louis. pp. 400–404, 2009.
 - 15) Morris DD. Alterations in the leukogram. In: Large animal internal medicine, 4th ed. Smith BP. ed. Elsevier, St. Louis. pp. 405–410, 2009.
 - 16) Nobre V, Borges I; Núcleo Interdisciplinar de Investigação em Medicina Intensiva (NIIMI). Prognostic value of procalcitonin in hospitalized patients with lower respiratory tract infections. *Rev Bras Ter Intensiva* 28, 179–189, 2016. doi: 10.5935/0103-507X.20160019.
 - 17) Nylen ES, Whang KT, Snider RH, Steinwald PM, White JC, Becker KL. Mortality is increased by procalcitonin and decreased by an antiserum reactive to procalcitonin in experimental sepsis. *Crit Care Med* 26, 1001–1006, 1998. doi: 10.1097/00003246-199806000-00015.
 - 18) Oberhoffer M, Stonans I, Russwurm S, Stonane E, Vogelsang H, Junker U, Jäger L, Reinhart K. Procalcitonin expression in human peripheral blood mononuclear cells and its modulation by lipopolysaccharides and sepsis-related cytokines in vitro. *J Lab Clin Med* 134, 49–55, 1999. doi: 10.1016/s0022-2143(99)90053-7.
 - 19) Pardon B, Hostens M, Duchateau L, Dewulf J, De Bleecker K, Deprez P. Impact of respiratory disease, diarrhea, otitis and

- arthritis on mortality and carcass traits in white veal calves. *BMC Vet Res* 9, 79, 2013. doi: 10.1186/1746-6148-9-79.
- 20) Principi N, Esposito S. Biomarkers in pediatric community-acquired pneumonia. *Int J Mol Sci* 18, 447, 2017. doi: 10.3390/ijms18020447.
- 21) Roland L, Drillich M, Iwersen M. Hematology as a diagnostic tool in bovine medicine. *J Vet Diagn Invest* 26, 592–598, 2014. doi: 10.1177/1040638714546490.
- 22) Terra RL. Ruminant history, physical examination, and records. In: Large animal internal medicine, 4th ed. Smith BP. ed. Elsevier, St. Louis. pp. 3–14, 2009.
- 23) Wacker C, Prkno A, Brunkhorst FM, Schlattmann P. Procalcitonin as a diagnostic marker for sepsis: a systematic review and meta-analysis. *Lancet Infect Dis* 13, 426–435, 2013. doi: 10.1016/S1473-3099(12)70323-7.
- 24) Wagner KE, Martinez JM, Vath SD, Snider RH, Nylén ES, Becker KL, Müller B, White JC. Early immunoneutralization of calcitonin precursors attenuates the adverse physiologic response to sepsis in pigs. *Crit Care Med* 30, 2313–2321, 2002. doi: 10.1097/00003246-200210000-00021.
- 25) Windeyer MC, Leslie KE, Godden SM, Hodgins DC, Lissemore KD, LeBlanc SJ. Factors associated with morbidity, mortality, and growth of dairy heifer calves up to 3 months of age. *Prev Vet Med* 113, 231–240, 2014. doi: 10.1016/j.prevetmed.2013.10.019.
- 26) Wu D, Zhou S, Hu S, Liu B. Inflammatory responses and histopathological changes in a mouse model of *Staphylococcus aureus*-induced bloodstream infections. *J Infect Dev Ctries* 11, 294–305, 2017. doi: 10.3855/jidc.7800.