



Title	MILNOR K-THEORY, F-ISOCRYSTALS AND SYNTOMIC REGULATORS
Author(s)	Asakura, Masanori; Miyatani, Kazuaki
Citation	Journal of the Institute of Mathematics of Jussieu, 23, 1-59 https://doi.org/10.1017/S1474748023000208
Issue Date	2024
Doc URL	http://hdl.handle.net/2115/91149
Rights	This article has been published in a revised form in [Journal of the Institute of Mathematics of Jussieu] [https://doi.org/10.1017/S1474748023000208]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © copyright holder.
Type	article (author version)
File Information	F-Isoc.HG.main.revised.pdf



[Instructions for use](#)

MILNOR K -THEORY, F -ISOCRYSTALS AND SYNTOMIC REGULATORS

MASANORI ASAKURA AND KAZUAKI MIYATANI

ABSTRACT. We introduce a category of filtered F -isocrystals and construct a symbol map from Milnor K -theory to the group of 1-extensions of filtered F -isocrystals. We show that our symbol map is compatible with the syntomic symbol map to the log syntomic cohomology by Kato and Tsuji. These are fundamental materials in our computations of syntomic regulators which we work in other papers.

1. INTRODUCTION

The purpose of this paper is to provide fundamental materials for computing the syntomic regulators on Milnor K -theory, which is based on the theory of F -isocrystals.

Let V be a complete discrete valuation ring such that the residue field k is a perfect field of characteristic $p > 0$ and the fractional field K is of characteristic 0. For a smooth affine scheme S over V , we introduce a *category of filtered F -isocrystals*, which is denoted by $\mathrm{Fil}\text{-}F\text{-MIC}(S)$ (see §?? for the details). Roughly speaking, an isocrystal is a crystalline sheaf which corresponds to a smooth \mathbb{Q}_l -sheaf, and “ F ” means Frobenius action. We refer the book [?] for the general terminology of F -isocrystals. As is well-known, it is equivalent to a notion of an integrable connection with Frobenius action. According to this, we shall define the category $\mathrm{Fil}\text{-}F\text{-MIC}(S)$ without the terminology of F -isocrystals. Namely we define it to be a category of coherent modules endowed with Hodge filtration, integrable connection and Frobenius action, so that the objects are described by familiar and elementary notation. However, the theory of F -isocrystals plays an essential role in verifying several functorial properties. The purpose of this paper is to introduce a symbol map on the Milnor K -group to the group of 1-extensions of filtered F -isocrystals. To be precise, let S be a smooth affine scheme over V and $U \rightarrow S$ a smooth V -morphism having a good compactification, which means that $U \rightarrow S$ extends to a projective smooth morphism $X \rightarrow S$ such that $X \setminus U$ is a relative simple normal crossing divisor (*abbreviated to* NCD) over S . Suppose that the comparison isomorphism

$$\mathcal{O}(S)_K^\dagger \otimes_{\mathcal{O}(S)_K} H_{\mathrm{dR}}^i(U_K/S_K) \xrightarrow{\cong} H_{\mathrm{rig}}^i(U_k/S_k)$$

holds for each $i \geq 0$ where $U_K = U \times_V \mathrm{Spec} K$ and $U_k = U \times_V \mathrm{Spec} k$ etc. Then, for an integer $n \geq 0$ such that $\mathrm{Fil}^{n+1} H_{\mathrm{dR}}^{n+1}(U_K/S_K) = 0$ where Fil^\bullet is the Hodge filtration, we construct a homomorphism

$$[-]_{U/S} : K_{n+1}^M(\mathcal{O}(U)) \longrightarrow \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-MIC}(S)}^1(\mathcal{O}_S, H^n(U/S)(n+1))$$

from the Milnor K -group of the affine ring $\mathcal{O}(U)$ to the group of 1-extensions in the category of filtered F -isocrystals (Theorem ??). We call this the *symbol map for U/S* . We provide an

explicit formula of our symbol map (Theorem ??). Moreover we shall give the comparison of our symbol map with the syntomic symbol map

$$[-]_{\text{syn}} : K_{n+1}^M(\mathcal{O}(U)) \longrightarrow H_{\text{syn}}^{n+1}(U, \mathbb{Z}_p(n+1))$$

to the syntomic cohomology of Fontaine-Messing, or more generally the log syntomic cohomology (cf. [?, Chapter I §3], [?, §2.2]). See Theorem ?? for the details.

Thanks to the recent work by Nekovář-Nizioł [?], there are the syntomic regulator maps

$$\text{reg}_{\text{syn}}^{i,j} : K_i(X) \otimes \mathbb{Q} \longrightarrow H_{\text{syn}}^{2j-i}(X, \mathbb{Q}_p(j))$$

in very general setting, which includes the syntomic symbol maps (up to torsion) and the rigid syntomic regulator maps by Besser [?]. They play the central role in the Bloch-Kato conjecture [?], and in the p -adic Beilinson conjecture by Perrin-Riou [?, 4.2.2] (see also [?, Conjecture 2.7]). On the other hand, the authors do not know how to construct $\text{reg}_{\text{syn}}^{i,j}$ without “ $\otimes \mathbb{Q}$ ”. We focus on the log syntomic cohomology with \mathbb{Z}_p -coefficients since the integral structure is important in our ongoing applications (e.g. [?]), namely a *deformation method* for computing syntomic regulators.

It is a notorious fact that, it is never easy to compute the syntomic regulator maps. Indeed it is non-trivial even for showing non-vanishing of $\text{reg}_{\text{syn}}^{i,j}$ in a general situation. The deformation method is a method to employ differential equations, which is motivated by Lauder [?] who provided the method for computing the Frobenius eigenvalues of a smooth projective variety over a finite field. The overview is as follows. Suppose that a variety X extends to a projective smooth family $f : Y \rightarrow S$ with $X = f^{-1}(a)$ and suppose that an element $\xi_X \in K_i(X)$ extends to an element $\xi \in K_i(Y)$. We deduce a differential equation such that a “function” $F(t) = \text{reg}_{\text{syn}}(\xi|_{f^{-1}(t)})$ is a solution. Solve the differential equation. Then we get the $\text{reg}_{\text{syn}}(\xi_X)$ by evaluating $F(t)$ at the point $a \in S$. Of course, this method works only in a good situation, for example, it is powerless if f is a constant family. However once it works, it has a big advantage in explicit computation of the syntomic regulators. We demonstrate it by a particular example, namely an elliptic curve with 3-torsion points.

Theorem 1.1 (Corollary ??). *Let $p \geq 5$ be a prime. Let $W = W(\overline{\mathbb{F}}_p)$ be the Witt ring, and $K := \text{Frac}(W)$. Let $a \in W$ satisfy $a \not\equiv 0, 1 \pmod{p}$. Let E_a be the elliptic curve over W defined by a Weierstrass equation $y^2 = x^3 + (3x + 4 - 4a)^2$. Let*

$$\xi_a = \{h_1, h_2\} = \left\{ \frac{y - 3x - 4 + 4a}{-8(1-a)}, \frac{y + 3x + 4 - 4a}{8(1-a)} \right\} \in K_2(E_a),$$

where we note that the divisors $\text{div}(h_i)$ have supports in 3-torsion points. Then there are overconvergent functions $\varepsilon_1(t), \varepsilon_2(t) \in K \otimes W[t, (1-t)^{-1}]^\dagger$ which are explicitly given as in Theorem ?? together with (??) and (??), and we have

$$\text{reg}_{\text{syn}}(\xi_a) = \varepsilon_1(a) \frac{dx}{y} + \varepsilon_2(a) \frac{xdx}{y} \in H_{\text{syn}}^2(E_a, \mathbb{Q}_p(2)) \cong H_{\text{dR}}^1(E_a/K).$$

We note that the function $\varepsilon_i(t)$ are defined in terms of the hypergeometric series

$${}_2F_1\left(\begin{matrix} \frac{1}{3}, \frac{2}{3} \\ 1 \end{matrix}; t\right) = \sum_{n=0}^{\infty} \frac{(\frac{1}{3})_n (\frac{2}{3})_n}{n!} t^n, \quad (\alpha)_n := \alpha(\alpha+1) \cdots (\alpha+n-1).$$

Concerning hypergeometric functions and regulators, the first author obtains more examples in [?]. There he introduces certain convergent functions which satisfy Dwork type congruence relations [?] to describe the syntomic regulators. Besides, in the joint paper [?], Chida and the first author discuss K_2 of elliptic curves in more general situation, and obtain a number of numerical verifications of the p -adic Beilinson conjecture. In both works, our category $\text{Fil-}F\text{-MIC}(S)$ and symbol maps play as fundamental materials.

Finally we comment on the category of syntomic coefficients by Bannai [?, 1.8]. His category is close to our $\text{Fil-}F\text{-MIC}(S)$. The difference is that, Bannai takes account into the boundary condition at $\bar{S} \setminus S$ on the Hodge filtration, while we do not. In this sense our category is less polish than his. On the other hand, he did not work on the symbol maps or regulator maps. Our main interest is the syntomic regulators, especially the deformation method. For this ours is sufficient.

Notation. For a integral domain V and a V -algebra R (resp. V -scheme X), let R_K (resp. X_K) denote the tensoring $R \otimes_V K$ (resp. $X \times_V K$) with the fractional field K .

Suppose that V is a complete valuation ring V endowed with a non-archimedian valuation $|\cdot|$. For a V -algebra B of finite type, let B^\dagger denote the weak completion of B . Namely if $B = V[T_1, \dots, T_n]/I$, then $B^\dagger = V[T_1, \dots, T_n]^\dagger/I$ where $V[T_1, \dots, T_n]^\dagger$ is the ring of power series $\sum a_\alpha T^\alpha$ such that for some $r > 1$, $|a_\alpha| r^{|\alpha|} \rightarrow 0$ as $|\alpha| \rightarrow \infty$. We simply write $B_K^\dagger = K \otimes_V B^\dagger$.

Acknowledgements. The authors are grateful to Atsushi Shiho for kindly answering their questions and telling us about his previous results.

2. FILTERED F -ISOCRYSTALS AND MILNOR K -THEORY

In this section, we work over a complete discrete valuation ring V of characteristic 0 such that the residue field k is a perfect field of characteristic $p > 0$. We suppose that V has a p -th Frobenius F_V , namely an endomorphism on V such that $F_V(x) \equiv x^p \pmod{pV}$, and fix it throughout this section. Let $K = \text{Frac}(V)$ be the fractional field. The extension of F_V to K is also denoted by F_V .

A scheme means a separated scheme which is of finite type over V unless otherwise specified. If X is a V -scheme (separated and of finite type), then \hat{X}_K will denote Raynaud's generic fiber of the formal completion \hat{X} , X_K^{an} will denote the analytification of the K -scheme X_K , and $j_X: \hat{X}_K \hookrightarrow X_K^{\text{an}}$ will denote the canonical immersion [?, (0.3.5)].

2.1. The category of Filtered F -isocrystals. Let $S = \text{Spec}(B)$ be an affine smooth variety over V . Let $\sigma: B^\dagger \rightarrow B^\dagger$ be a p -th Frobenius compatible with F_V on V , which means that σ is F_V -linear and satisfies $\sigma(x) \equiv x^p \pmod{pB}$. The induced endomorphism $\sigma \otimes_{\mathbb{Z}} \mathbb{Q}: B_K^\dagger \rightarrow B_K^\dagger$ is also denoted by σ . We define the category $\text{Fil-}F\text{-MIC}(S, \sigma)$ (which we call the category of *filtered F -isocrystals* on S) as follows.

Definition 2.1. An object of $\text{Fil-}F\text{-MIC}(S, \sigma)$ is a datum $H = (H_{\text{dR}}, H_{\text{rig}}, c, \Phi, \nabla, \text{Fil}^\bullet)$, where

- H_{dR} is a coherent B_K -module,
- H_{rig} is a coherent B_K^\dagger -module,

- $c: H_{\text{dR}} \otimes_{B_K} B_K^\dagger \xrightarrow{\cong} H_{\text{rig}}$ is a B_K^\dagger -linear isomorphism,
- $\Phi: \sigma^* H_{\text{rig}} \xrightarrow{\cong} H_{\text{rig}}$ is an isomorphism of B_K^\dagger -algebra,
- $\nabla: H_{\text{dR}} \rightarrow \Omega_{B_K}^1 \otimes H_{\text{dR}}$ is an (algebraic) integrable connection and
- Fil^\bullet is a finite descending filtration on H_{dR} of locally free B_K -module (i.e. each graded piece is locally free),

that satisfies $\nabla(\text{Fil}^i) \subset \Omega_{B_K}^1 \otimes \text{Fil}^{i-1}$ and the compatibility of Φ and ∇ in the following sense. Note first that ∇ induces an integrable connection $\nabla_{\text{rig}}: H_{\text{rig}} \rightarrow \Omega_{B_K^\dagger}^1 \otimes H_{\text{rig}}$, where $\Omega_{B_K^\dagger}^1$ denotes the sheaf of continuous differentials. In fact, firstly regard H_{dR} as a coherent \mathcal{O}_{S_K} -module. Then, by (rigid) analytification, we get an integrable connection ∇^{an} on the coherent $\mathcal{O}_{S_K^{\text{an}}}$ -module $(H_{\text{dR}})^{\text{an}}$. Then, apply the functor j_S^\dagger to ∇^{an} to obtain an integrable connection on $\Gamma(S_K^{\text{an}}, j_S^\dagger((H_{\text{dR}})^{\text{an}})) = H_{\text{dR}} \otimes_{B_K} B_K^\dagger$. This gives an integrable connection ∇_{rig} on H_{rig} via the isomorphism c . Then the compatibility of Φ and ∇ means that Φ is horizontal with respect to ∇_{rig} , namely $\Phi \nabla_{\text{rig}} = \nabla_{\text{rig}} \Phi$. We usually write $\nabla_{\text{rig}} = \nabla$ to simplify the notation.

A morphism $H' \rightarrow H$ in $\text{Fil-}F\text{-MIC}(S, \sigma)$ is a pair of homomorphisms $(h_{\text{dR}}: H'_{\text{dR}} \rightarrow H_{\text{dR}}, h_{\text{rig}}: H'_{\text{rig}} \rightarrow H_{\text{rig}})$, such that h_{rig} is compatible with Φ 's, h_{dR} is compatible with ∇ 's and Fil^\bullet 's, and moreover they agree under the isomorphism c .

Remark 2.2. (1) The category $\text{Fil-}F\text{-MIC}(S, \sigma)$ can also be described by using simpler categories as follows. Let $\text{Fil-MIC}(S_K)$ denote the category of filtered S_K -modules with integrable connection, that is, the category of data $(M_{\text{dR}}, \nabla, \text{Fil}^\bullet)$ with M_{dR} a coherent B_K -module, ∇ an integrable connection on M_{dR} , and Fil^\bullet a finite descending filtration on M_{dR} of locally free B_K -module. Let $\text{MIC}(B_K^\dagger)$ denote the category of coherent B_K^\dagger -modules with integrable connections (M_{rig}, ∇) on B_K^\dagger , and let $F\text{-MIC}(B_K^\dagger, \sigma)$ denote the category of the coherent B_K^\dagger -modules with integrable connections $(M_{\text{rig}}, \nabla, \Phi)$ with σ -linear Frobenius endomorphisms. Then $\text{Fil-}F\text{-MIC}(S, \sigma)$ is identified with the fiber product

$$F\text{-MIC}(B_K^\dagger, \sigma) \times_{\text{MIC}(B_K^\dagger)} \text{Fil-MIC}(S_K).$$

(2) Let $F\text{-Isoc}^\dagger(B_k)$ denote the category of overconvergent F -isocrystals on S_k . Then there is the equivalence of categories ([?, Theorem 8.3.10])

$$F\text{-Isoc}^\dagger(B_k) \xrightarrow{\cong} F\text{-MIC}(B_K^\dagger, \sigma).$$

Therefore, by combining with the description in (1), we see that our category $\text{Fil-}F\text{-MIC}(S, \sigma)$ does not depend on σ , which means that there is the natural equivalence $\text{Fil-}F\text{-MIC}(S, \sigma) \cong \text{Fil-}F\text{-MIC}(S, \sigma')$ (see also Lemma ?? in Appendix). By virtue of this fact, we often drop “ σ ” in the notation.

For two objects H, H' in the category $\text{Fil-}F\text{-MIC}(S, \sigma)$, we have a direct sum $H \oplus H'$ and the tensor product $H \otimes H'$ in a customary manner. The unit object for the tensor product, denoted by B or \mathcal{O}_S , is $(B_K, B_K^\dagger, c, \sigma_B, d, \text{Fil}^\bullet)$, where c is the natural isomorphism, d is the usual differential and Fil^\bullet is defined by $\text{Fil}^0 B_K = B_K$ and $\text{Fil}^1 B_K = 0$. The category $\text{Fil-}F\text{-MIC}(S, \sigma)$ forms a tensor category with this tensor product and the unit object B or \mathcal{O}_S .

The unit object can also be described as $B = B(0)$ or $\mathcal{O}_S = \mathcal{O}_S(0)$ by using the following notion of *Tate object*.

Definition 2.3. Let n be an integer.

(1) The *Tate object* in $\text{Fil-}F\text{-MIC}(S)$, which we denote by $B(n)$ or $\mathcal{O}_S(n)$, is defined to be $(B_K, B_K^\dagger, c, p^{-n}\sigma_B, d, \text{Fil}^\bullet)$, where c is the natural isomorphism, $d : B_K \rightarrow \Omega_{B_K}^1$ is the usual differential, and Fil^\bullet is defined by $\text{Fil}^{-n}B_K = B_K$ and $\text{Fil}^{-n+1}B_K = 0$.

(2) For an object H of $\text{Fil-}F\text{-MIC}(S)$, we write $H(n) := H \otimes B(n)$.

Now, we discuss the Yoneda extension groups in the category $\text{Fil-}F\text{-MIC}(S)$. A sequence

$$H_1 \longrightarrow H_2 \longrightarrow H_3$$

in $\text{Fil-}F\text{-MIC}(S)$ (or in $\text{Fil-MIC}(S_K)$) is called *exact* if

$$\text{Fil}^i H_{1,\text{dR}} \longrightarrow \text{Fil}^i H_{2,\text{dR}} \longrightarrow \text{Fil}^i H_{3,\text{dR}}$$

are exact for all i . Then the category $\text{Fil-}F\text{-MIC}(S)$ forms an exact category which has kernel and cokernel objects for any morphisms. Thus, the Yoneda extension groups

$$\text{Ext}^\bullet(H, H') = \text{Ext}_{\text{Fil-}F\text{-MIC}(S)}^\bullet(H, H')$$

in $\text{Fil-}F\text{-MIC}(S)$ are defined in the canonical way (or one can further define the derived category of $\text{Fil-}F\text{-MIC}(S)$ [?, 1.1]). An element of $\text{Ext}^j(H, H')$ is represented by an exact sequence

$$0 \longrightarrow H' \longrightarrow M_1 \longrightarrow \cdots \longrightarrow M_j \longrightarrow H \longrightarrow 0$$

and is subject to the equivalence relation generated by commutative diagrams

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H' & \longrightarrow & M_1 & \longrightarrow & \cdots & \longrightarrow & M_j & \longrightarrow & H & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & H' & \longrightarrow & N_1 & \longrightarrow & \cdots & \longrightarrow & N_j & \longrightarrow & H & \longrightarrow & 0. \end{array}$$

Note that $\text{Ext}^\bullet(H, H')$ is uniquely divisible (i.e. a \mathbb{Q} -module) as the multiplication by $m \in \mathbb{Z}_{>0}$ on H or H' is bijective.

Nextly, we discuss the functoriality of the category $\text{Fil-}F\text{-MIC}(S)$ with respect to S . Let $S' = \text{Spec}(B')$ be another affine smooth variety and $i : S' \rightarrow S$ a morphism of V -schemes. Then, i induces a pull-back functor

$$(2.1) \quad i^* : \text{Fil-}F\text{-MIC}(S) \rightarrow \text{Fil-}F\text{-MIC}(S')$$

in a natural way. In fact, if $H = (H_{\text{dR}}, H_{\text{rig}}, c, \Phi, \nabla, \text{Fil}^\bullet)$ is an object of $\text{Fil-}F\text{-MIC}(S)$, then we define $i^*H = (H_{\text{dR}} \otimes_{B_K} B'_K, H_{\text{rig}} \otimes_{B_K^\dagger} B'^{\dagger}_K, c \otimes_{B_K^\dagger} B'^{\dagger}_K, \Phi', \nabla', \text{Fil}^\bullet)$, where ∇' and Fil^\bullet are natural pull-backs of ∇ and Fil^\bullet respectively, and where Φ' is the natural Frobenius structure obtained as follows. We may regard $(H_{\text{rig}}, \nabla_{\text{rig}}, \Phi)$ as an overconvergent F -isocrystal on S via the equivalence $F\text{-MIC}(B_K^\dagger) \simeq F\text{-Isoc}^\dagger(B_K)$ [?, Theorem 8.3.10]. Then, its pull-back along $i_k : S'_k \rightarrow S_k$ is an overconvergent F -isocrystal on S'_k , which is again identified with an object of $F\text{-MIC}(B'^{\dagger}_K)$. Thus it is of the form $(H'_{\text{rig}}, \nabla'_{\text{rig}}, \Phi')$, and H'_{rig} is naturally isomorphic to $H_{\text{rig}} \otimes_{B_K^\dagger} B'^{\dagger}_K$ ([?, Prop 8.1.15]). Now, Φ' gives the desired Frobenius structure.

2.2. The complex $\mathcal{S}(M)$. In this subsection, we introduce a complex $\mathcal{S}(M)$ for each object M in $\text{Fil-}F\text{-MIC}(S, \sigma)$ which is, in the case where $M = \mathcal{O}_S(r)$, close to the syntomic complex $\mathcal{S}_n(R)_{S, \sigma}$ of Fontaine–Messing.

Before the definition, we prepare a morphism attached to each object of $\text{Fil-MIC}(S_K)$. Let $H = (H_{\text{dR}}, \nabla, \text{Fil}^\bullet)$ be an object of $\text{Fil-MIC}(S_K)$. Let $\Omega_{B_K}^\bullet \otimes \text{Fil}^{i-\bullet} H_{\text{dR}}$ denote the de Rham complex

$$\text{Fil}^i H_{\text{dR}} \rightarrow \Omega_{B_K}^1 \otimes \text{Fil}^{i-1} H_{\text{dR}} \rightarrow \cdots \rightarrow \Omega_{B_K}^n \otimes \text{Fil}^{i-n} H_{\text{dR}} \rightarrow \cdots,$$

where \otimes denotes \otimes_{B_K} and the differentials are given by

$$(2.2) \quad \omega \otimes x \mapsto d\omega^j \otimes x + (-1)^j \omega \wedge \nabla(x), \quad (\omega \otimes x \in \Omega_{B_K}^j \otimes H_{\text{dR}}).$$

Now, we define a natural map

$$(2.3) \quad \text{Ext}_{\text{Fil-MIC}(S_K)}^i(\mathcal{O}_{S_K}, H) \longrightarrow H^i(\Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} H_{\text{dR}})$$

in the following way. Let

$$(2.4) \quad 0 \longrightarrow H \longrightarrow M_i \longrightarrow M_{i-1} \longrightarrow \cdots \longrightarrow M_0 \longrightarrow \mathcal{O}_{S_K} \longrightarrow 0$$

be an exact sequence in $\text{Fil-MIC}(S_K)$. This induces an exact sequence

$$0 \longrightarrow \Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} H \longrightarrow \Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} M_i \longrightarrow \cdots \longrightarrow \Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} M_0 \longrightarrow \Omega_{B_K}^\bullet \longrightarrow 0$$

of complexes, and hence a connecting homomorphism $\delta: H^0(\Omega_{B_K}^\bullet) \rightarrow H^i(\Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} H)$. Then the map (??) is defined by sending the sequence (??) to $\delta(1)$.

By the forgetful functor $\text{Fil-}F\text{-MIC}(S, \sigma) \rightarrow \text{Fil-MIC}(S)$, the morphism (??) clearly induces a canonical morphism

$$(2.5) \quad \text{Ext}_{\text{Fil-}F\text{-MIC}(S, \sigma)}^i(\mathcal{O}_S, H) \longrightarrow H^i(\Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} H_{\text{dR}}).$$

Let $F\text{-mod} = F\text{-MIC}(\text{Spec } K)$, namely the category of K -modules endowed with F_V -linear endomorphisms. Then, we have a functor

$$F\text{-MIC}(B_K^\dagger, \sigma) \longrightarrow D^b(F\text{-mod}), \quad (M_{\text{rig}}, \Phi) \longmapsto (\Omega_{B_K^\dagger}^\bullet \otimes M_{\text{rig}}, \Phi)$$

to the derived category of complexes in $F\text{-mod}$, where Φ in the target is defined to be $\sigma \otimes \Phi$ (we use the same notation because we always extend the Frobenius action on the de Rham complex $\Omega_{B_K^\dagger}^\bullet \otimes M_{\text{rig}}$ by this rule). We note that the above functor does not depend on σ . Indeed, the composition

$$F\text{-Isoc}^\dagger(B_K) \xrightarrow{\cong} F\text{-MIC}(B_K^\dagger, \sigma) \longrightarrow D^b(F\text{-mod})$$

is the functor $(E, \Phi) \mapsto (R\Gamma_{\text{rig}}(S_K, E), \Phi_{\text{rig}})$ where Φ_{rig} denotes the Frobenius action on the rigid cohomology ([?, Proposition 8.3.12]), and this does not depend on σ .

Definition 2.4. For an object $M \in \text{Fil-}F\text{-MIC}(S, \sigma)$, we define $\mathcal{S}(M)$ to be the mapping fiber of the morphism

$$1 - \Phi : \Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} M_{\text{dR}} \longrightarrow \Omega_{B_K^\dagger}^\bullet \otimes M_{\text{rig}},$$

where 1 denotes the inclusion $\Omega_{B_K}^\bullet \otimes \text{Fil}^{-\bullet} M_{\text{dR}} \hookrightarrow \Omega_{B_K^\dagger}^\bullet \otimes M_{\text{rig}}$ via the comparison and Φ is the composition of it with Φ on $\Omega_{B_K^\dagger}^\bullet \otimes M_{\text{rig}}$.

Note that, in more down-to-earth manner, each term of $\mathcal{S}(M)$ is given by

$$(2.6) \quad \mathcal{S}(M)^i = \Omega_{B_K}^i \otimes \mathrm{Fil}^{-i} M_{\mathrm{dR}} \oplus \Omega_{B_K^\dagger}^{i-1} \otimes M_{\mathrm{rig}}$$

and the differential $\mathcal{S}(M)^i \rightarrow \mathcal{S}(M)^{i+1}$ is given by

$$(\omega, \xi) \mapsto (d_M \omega, (1 - \Phi)\omega - d_M \xi), \quad (\omega \in \Omega_{B_K}^i \otimes \mathrm{Fil}^{-i} M_{\mathrm{dR}}, \xi \in \Omega_{B_K^\dagger}^{i-1} \otimes M_{\mathrm{rig}})$$

where d_M is the differential (??).

An exact sequence

$$0 \longrightarrow H \longrightarrow M_i \longrightarrow M_{i-1} \longrightarrow \cdots \longrightarrow M_0 \longrightarrow \mathcal{O}_S \longrightarrow 0$$

in $\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)$ gives rise to an exact sequence

$$0 \longrightarrow \mathcal{S}(H) \longrightarrow \mathcal{S}(M_i) \longrightarrow \mathcal{S}(M_{i-1}) \longrightarrow \cdots \longrightarrow \mathcal{S}(M_0) \longrightarrow \mathcal{S}(\mathcal{O}_S) \longrightarrow 0$$

of complexes of \mathbb{Q}_p -modules. Let $\delta: \mathbb{Q}_p \cong H^0(\mathcal{S}(\mathcal{O}_S)) \rightarrow H^i(\mathcal{S}(H))$ be the connecting homomorphism. We define a homomorphism

$$(2.7) \quad u: \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)}^i(\mathcal{O}_S, H) \longrightarrow H^i(\mathcal{S}(H))$$

by associating $\delta(1)$ to the above extension. The composition of u with the natural map $H^i(\mathcal{S}(H)) \rightarrow H^i(\Omega_{B_K}^\bullet \otimes \mathrm{Fil}^{-\bullet} H_{\mathrm{dR}})$ agrees with (??).

The complex $\mathcal{S}(\mathcal{O}_S(r))$ is close to the syntomic complex of Fontaine-Messing. More precisely, let $S_n := S \times_W \mathrm{Spec} W/p^n W$ and $B_n := B/p^n B$. The syntomic complex $\mathcal{S}_n(r)_{S, \sigma}$ is the mapping fiber of the morphism

$$1 - p^{-r} \sigma: \Omega_{B_n}^{\bullet \geq r} \longrightarrow \Omega_{B_n}^\bullet$$

of complexes where we note that $p^{-r} \sigma$ is well-defined (see [?, p.410–411]). Hence there is the natural map

$$(2.8) \quad H^i(\mathcal{S}(\mathcal{O}_S(r))) \longrightarrow \mathbb{Q} \otimes \varprojlim_n H_{\mathrm{zar}}^i(S_1, \mathcal{S}_n(r)_{S, \sigma}) =: H_{\mathrm{syn}}^i(S, \mathbb{Q}_p(r)).$$

Let

$$(2.9) \quad \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)}^i(\mathcal{O}_S, \mathcal{O}_S(r)) \longrightarrow H_{\mathrm{syn}}^i(S, \mathbb{Q}_p(r))$$

be the composition morphism. Apparently, the both side of (??) depend on σ . However, if we replace σ with σ' , there is the natural transformation between $\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)$ and $\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma')$ thanks to the theory of F -isocrystals, and there is also a natural transformation on the syntomic cohomology. The map (??) is compatible under these transformations. In this sense, (??) does not depend on the Frobenius σ .

Lemma 2.5. *Suppose $\mathrm{Fil}^0 H_{\mathrm{dR}} = 0$. Then the map u in (??) is injective when $i = 1$. Moreover, the map (??) is injective when $i = 1$ and $r \geq 0$.*

Proof. Let

$$0 \longrightarrow H \longrightarrow M \longrightarrow \mathcal{O}_S \longrightarrow 0$$

be an exact sequence in $\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)$. Since $\mathrm{Fil}^0 H_{\mathrm{dR}} = 0$, there is the unique lifting $e \in \mathrm{Fil}^0 M_{\mathrm{rig}}$. Then

$$(2.10) \quad u(M) = (\nabla(e), (1 - \Phi)e) \in H^1(\mathcal{S}(H)) \subset \Omega_{B_K^\dagger}^1 \otimes \mathrm{Fil}^{-1} H_{\mathrm{rig}} \oplus H_{\mathrm{rig}}$$

by definition of u . If $u(M) = 0$, then the datum $(B_K e, B_K^\dagger e, c, \Phi, \nabla, \text{Fil}^\bullet)$ forms a subobject of M which is isomorphic to the unit object \mathcal{O}_S . This gives a splitting of the above exact sequence. The latter follows from the fact that

$$H_{\text{syn}}^1(S, \mathbb{Q}_p(1)) \subset \mathbb{Q} \otimes \varprojlim_n (\Omega_{B_n}^1 \oplus B_n), \quad H_{\text{syn}}^1(S, \mathbb{Q}_p(r)) \subset \mathbb{Q} \otimes \varprojlim_n (\{0\} \oplus B_n), \quad (r \geq 2)$$

and

$$H^1(\mathcal{S}(\mathcal{O}_S(1))) \subset \Omega_{B_K^\dagger}^1 \oplus B_K^\dagger, \quad H^1(\mathcal{S}(\mathcal{O}_S(r))) \subset \{0\} \oplus B_K^\dagger, \quad (r \geq 2).$$

□

2.3. Log objects. In this subsection, we introduce the “log object” in $\text{Fil-}F\text{-MIC}(S)$ concerning a p -adic logarithmic function. In the next subsection, it will be generalized to a notion of “polylog object”.

For $f \in B^\times$ let

$$\log^{(\sigma)}(f) := p^{-1} \log \left(\frac{f^p}{f^\sigma} \right) = -p^{-1} \sum_{n=1}^{\infty} \frac{(1 - f^p/f^\sigma)^n}{n} \in B^\dagger.$$

An elementary computation yields $\log^{(\sigma)}(f) + \log^{(\sigma)}(g) = \log^{(\sigma)}(fg)$ for $f, g \in B^\times$.

Definition 2.6. For $f \in B^\times$, we define the *log object* $\mathcal{L}og(f)$ in $\text{Fil-}F\text{-MIC}(S, \sigma)$ as follows.

- $\mathcal{L}og(f)_{\text{dR}}$ is a free B_K -module of rank two; $\mathcal{L}og(f)_{\text{dR}} = B_K e_{-2} \oplus B_K e_0$.
- $\mathcal{L}og(f)_{\text{rig}} = B_K^\dagger e_{-2} \oplus B_K^\dagger e_0$.
- c is the natural isomorphism.
- Φ is the σ -linear morphism defined by

$$\Phi(e_{-2}) = p^{-1} e_{-2}, \quad \Phi(e_0) = e_0 - \log^{(\sigma)}(f) e_{-2}.$$

- ∇ is the connection defined by $\nabla(e_{-2}) = 0$ and $\nabla(e_0) = \frac{df}{f} e_{-2}$.
- Fil^\bullet is defined by

$$\text{Fil}^i \mathcal{L}og(f)_{\text{dR}} = \begin{cases} \mathcal{L}og(f)_{\text{dR}} & \text{if } i \leq -1, \\ B_K e_0 & \text{if } i = 0, \\ 0 & \text{if } i > 0. \end{cases}$$

This is fit into the exact sequence

$$(2.11) \quad 0 \longrightarrow \mathcal{O}_S(1) \xrightarrow{\epsilon} \mathcal{L}og(f) \xrightarrow{\pi} \mathcal{O}_S \longrightarrow 0$$

in $\text{Fil-}F\text{-MIC}(S)$ where the two arrows are defined by $\epsilon(1) = e_{-2}$ and $\pi(e_0) = 1$. This defines a class in $\text{Ext}_{\text{Fil-}F\text{-MIC}(S, \sigma)}^1(\mathcal{O}_S, \mathcal{O}_S(1))$, which we write by $[f]_S$. It is easy to see that $f \mapsto [f]_S$ is additive. We call the group homomorphism

$$(2.12) \quad [-]_S : \mathcal{O}(S)^\times \longrightarrow \text{Ext}_{\text{Fil-}F\text{-MIC}(S, \sigma)}^1(\mathcal{O}_S, \mathcal{O}_S(1))$$

the symbol map.

Lemma 2.7. *The composition*

$$\mathcal{O}(S)^\times \xrightarrow{[-]_S} \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S,\sigma)}^1(\mathcal{O}_S, \mathcal{O}_S(1)) \longrightarrow H_{\mathrm{syn}}^1(S, \mathbb{Q}_p(1))$$

agrees with the symbol map by Kato [?] where the second arrow is the map (??). Namely, it is explicitly described as follows,

$$f \longmapsto \left(\frac{df}{f}, \log^{(\sigma)}(f) \right) \in H_{\mathrm{syn}}^1(S, \mathbb{Q}_p(1)) \subset \Omega_{\widehat{B}_K}^1 \oplus \widehat{B}_K$$

where $\widehat{B} := \varprojlim_n B/p^n B$ and $\widehat{B}_K := \mathbb{Q} \otimes \widehat{B}$.

Proof. By definition of ρ in (??), one has

$$\rho(\mathcal{L}og(f)) = (\nabla(e_0), (1 - \Phi)e_0) = \left(\frac{df}{f}, \log^{(\sigma)}(f) \right)$$

as desired. \square

Lemma 2.8. *The symbol map (??) is functorial with respect to the pull-back. Namely, for a morphism $i: S' \rightarrow S$, the diagram*

$$\begin{array}{ccc} \mathcal{O}(S)^\times & \xrightarrow{[-]_S} & \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S,\sigma)}^1(\mathcal{O}_S, \mathcal{O}_S(1)) \\ i \downarrow & & \downarrow i^* \\ \mathcal{O}(S')^\times & \xrightarrow{[-]_{S'}} & \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S',\sigma)}^1(\mathcal{O}_{S'}, \mathcal{O}_{S'}(1)) \end{array}$$

is commutative where i^* denotes the map induced from the pull-back functor (??).

Proof. Because of the injectivity of (??) (Lemma ??), the assertion can be reduced to the compatibility of Kato's symbol maps by Lemma ?? \square

2.4. Polylog objects. In this subsection, we generalize the log object to polylog objects. To define the polylog objects, we need the p -adic polylog function.

For an integer r , we denote the p -adic polylog function by

$$(2.13) \quad \ln_r^{(p)}(z) := \sum_{n \geq 1, p \nmid n} \frac{z^n}{n^r} = \lim_{s \rightarrow \infty} \frac{1}{1 - z^{p^s}} \sum_{1 \leq n < p^s, p \nmid n} \frac{z^n}{n^r} \in \mathbb{Z}_p \left[z, \frac{1}{1 - z} \right]^\wedge$$

where A^\wedge denotes the p -adic completion of a ring A . As is easily seen, one has

$$\ln_r^{(p)}(z) = (-1)^{r+1} \ln_r^{(p)}(z^{-1}), \quad z \frac{d}{dz} \ln_{r+1}^{(p)}(z) = \ln_r^{(p)}(z).$$

If $r \leq 0$, this is a rational function. Indeed

$$\ln_0^{(p)}(z) = \frac{1}{1 - z} - \frac{1}{1 - z^p}, \quad \ln_{-r}^{(p)}(z) = \left(z \frac{d}{dz} \right)^r \ln_0^{(p)}(z).$$

If $r \geq 1$, it is no longer a rational function but an overconvergent function.

Proposition 2.9. *Let $r \geq 1$. Put $x := (1 - z)^{-1}$. Then $\ln_r^{(p)}(z) \in (x - x^2)\mathbb{Z}_p[x]^\dagger$.*

Proof. Since $\ln_r^{(p)}(z)$ has \mathbb{Z}_p -coefficients, it is enough to check $\ln_r^{(p)}(z) \in (x - x^2)\mathbb{Q}_p[x]^\dagger$. We first note

$$(2.14) \quad (x^2 - x) \frac{d}{dx} \ln_{k+1}^{(p)}(z) = \ln_k^{(p)}(z).$$

The limit in (??) can be rewritten as

$$\lim_{s \rightarrow \infty} \frac{1}{x^{p^s} - (x-1)^{p^s}} \sum_{1 \leq n < p^s, p \nmid n} \frac{x^{p^s}(1-x^{-1})^n}{n^r}.$$

This shows that $\ln_r^{(p)}(z)$ is divided by $x - x^2$. Let $w(x) \in \mathbb{Z}_p[x]$ be defined by

$$\frac{1 - z^p}{(1 - z)^p} = x^p - (x - 1)^p = 1 - pw(x).$$

Then

$$\ln_1^{(p)}(z) = p^{-1} \log \left(\frac{1 - z^p}{(1 - z)^p} \right) = -p^{-1} \sum_{n=1}^{\infty} \frac{p^n w(x)^n}{n} \in \mathbb{Z}_p[x]^\dagger.$$

This shows $\ln_1^{(p)}(z) \in (x - x^2)\mathbb{Q}_p[x]^\dagger$, as required in case $r = 1$. Let

$$-(x - x^2)^{-1} \ln_1^{(p)}(z) = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n + \cdots \in \mathbb{Z}_p[x]^\dagger.$$

By (??) one has

$$\ln_2^{(p)}(z) = c + x + \frac{a_1}{2} x^2 + \cdots + \frac{a_n}{n+1} x^{n+1} + \cdots \in \mathbb{Q}_p[x]^\dagger$$

and hence $\ln_2^{(p)}(z) \in (x - x^2)\mathbb{Q}_p[x]^\dagger$, as required in case $r = 2$. Continuing the same argument, we obtain $\ln_r^{(p)}(z) \in (x - x^2)\mathbb{Q}_p[x]^\dagger$ for every r . \square

Remark 2.10. The proof shows that $\ln_r^{(p)}(z)$ converges on an open disk $|x| < |1 - \zeta_p|$.

Definition 2.11. Let $C = V[T, T^{-1}, (1-T)^{-1}]$, and σ_C a p -th Frobenius such that $\sigma_C(T) = T^p$. Let $n \geq 1$ be an integer. We define the n -th polylog object $\mathcal{P}ol_n(T)$ of Fil- F -MIC(Spec C) as follows.

- $\mathcal{P}ol_n(T)_{\text{dR}}$ is a free C_K -module of rank $n + 1$; $\mathcal{P}ol_n(T)_{\text{dR}} = \bigoplus_{j=0}^n C_K e_{-2j}$.
- $\mathcal{P}ol_n(T)_{\text{rig}} := \mathcal{P}ol_n(T)_{\text{dR}} \otimes_{C_K} C_K^\dagger$.
- c is the natural isomorphism.
- Φ is the C_K^\dagger -linear morphism defined by

$$\Phi(e_0) = e_0 - \sum_{j=1}^n (-1)^j \ln_j^{(p)}(T) e_{-2j}, \quad \Phi(e_{-2j}) = p^{-j} e_0, \quad (j \geq 1).$$

- ∇ is the connection defined by

$$\nabla(e_0) = \frac{dT}{T-1} e_{-2}, \quad \nabla(e_{-2j}) = \frac{dT}{T} e_{-2j-2}, \quad (j \geq 1)$$

where $e_{-2n-2} := 0$.

- Fil $^\bullet$ is defined by $\text{Fil}^m \mathcal{P}ol_n(T)_{\text{dR}} = \bigoplus_{0 \leq j \leq -m} C_K e_{-2j}$. In particular, $\text{Fil}^m \mathcal{P}ol_n(T)_{\text{dR}} = \mathcal{P}ol_n(T)_{\text{dR}}$ if $m \leq -n$ and $= 0$ if $m \geq 1$.

When $n = 2$, we also write $\mathcal{D}ilog(T) = \mathcal{P}ol_2(T)$, and call it the *dilog object*.

For a general $S = \text{Spec}(B)$ and $f \in B$ satisfying $f, 1 - f \in B^\times$, we define the polylog object $\mathcal{P}ol_n(f)$ to be the pull-back $u^* \mathcal{P}ol_n(T)$ where $u: \text{Spec}(B) \rightarrow \text{Spec } V[T, T^{-1}, (1 - T)^{-1}]$ is given by $u(T) = f$. When $n = 1$, $\mathcal{P}ol_1(T)$ coincides with the log object $\mathcal{L}og(1 - T)$ in $\text{Fil-}F\text{-MIC}(C)$.

2.5. Relative cohomologies. Let $S = \text{Spec}(B)$ be a smooth affine V -scheme and let σ be a p -th Frobenius on B^\dagger . In this subsection, we discuss objects in $\text{Fil-}F\text{-MIC}(S)$ arising as a relative cohomology of smooth morphisms.

Let $u: U \rightarrow S$ be a quasi-projective smooth morphism. We firstly describe a datum which we discuss in this subsection.

Definition 2.12. We define a datum

$$H^i(U/S) = (H_{\text{dR}}^i(U_K/S_K), H_{\text{rig}}^i(U_k/S_k), c, \nabla, \Phi, \text{Fil}^\bullet)$$

as follows.

- $H_{\text{dR}}^i(U_K/S_K)$ is the i -th relative algebraic de Rham cohomology of u_K , namely the module of global sections of the i -th cohomology sheaf $R^i(u_K)_* \Omega_{U_K/S_K}^\bullet$, and ∇ (resp. Fil^\bullet) is the Gauss–Manin connection (resp. the Hodge filtration) on $H_{\text{dR}}^i(U_K/S_K)$.
- $(H_{\text{rig}}^i(U_k/S_k), \Phi)$ is the B_K^\dagger -module with σ -linear Frobenius structure obtained as the i -th relative rigid cohomology of u_k . In particular,

$$H_{\text{rig}}^i(U_k/S_k) = \Gamma(S_K^{\text{an}}, R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}),$$

where $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}} := R^i(u_K^{\text{an}})_* j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet$ is the i -th relative rigid cohomology sheaf. (We justify this notation and definition in Remark ?? below.)

- $c: H_{\text{dR}}^i(U_K/S_K) \otimes_{B_K} B_K^\dagger \rightarrow H_{\text{rig}}^i(U_k/S_k)$ is the natural morphism between the algebraic de Rham cohomology and the rigid cohomology.

Let us give a construction of the comparison morphism c in this datum. We basically follow the construction in [?, 5.8.2]. Let $\iota: S_K^{\text{an}} \rightarrow S_K$ be the natural morphism of ringed topoi [?, 0.3]. Then, by adjunction, we have a natural morphism $R^i u_{K,*} \Omega_{U_K/S_K}^\bullet \rightarrow \iota_* (R^i u_{K,*} \Omega_{U_K/S_K}^\bullet)^{\text{an}}$. Now, together with the natural morphisms

$$(R^i u_{K,*} \Omega_{U_K/S_K}^\bullet)^{\text{an}} \rightarrow R^i(u_K^{\text{an}})_* \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet \rightarrow R^i(u_K^{\text{an}})_* j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet,$$

we get a morphism $R^i u_{K,*} \Omega_{U_K/S_K}^\bullet \rightarrow \iota_* R^i(u_K^{\text{an}})_* j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet$. By taking the module of global sections, we get a morphism $H_{\text{dR}}^i(U_K/S_K) \rightarrow H_{\text{rig}}^i(U_k/S_k)$, and therefore the desired morphism $c: H_{\text{dR}}^i(U_K/S_K) \otimes_{B_K} B_K^\dagger \rightarrow H_{\text{rig}}^i(U_k/S_k)$ because $H_{\text{rig}}^i(U_k/S_k)$ is a B_K^\dagger -module.

Remark 2.13. (1) Let us justify our description of the rigid cohomology $H_{\text{rig}}^i(U_k/S_k)$ in Definition ??.

We begin by recalling a usual definition of the rigid cohomology of $u_k: U_k \rightarrow S_k$ over a frame $(S_k, \overline{S}_k, \widehat{S})$, where \overline{S} is a closure of S in a projective space over V and \widehat{S} is its completion. Let \overline{X} be the closure of U in a projective space over \overline{S} and let $\overline{f}: \overline{X} \rightarrow \overline{S}$ be the extension of u . (We choose this notation because, in this article, X usually denotes $\overline{f}^{-1}(S)$ and f denotes $\overline{f}|_X$.) Then, by definition, the i -th relative rigid cohomology sheaf is

$R^i(\bar{f}_K^{\text{an}})_* \bar{j}_U^\dagger \Omega_{\bar{X}_K^{\text{an}}/\bar{S}_K^{\text{an}}}^\bullet$, where $\bar{j}_U: \widehat{U}_K \hookrightarrow \bar{X}_K^{\text{an}}$ is the natural inclusion, and $H_{\text{rig}}^i(U_k/S_k)$ is the module of global sections of this sheaf on \bar{S}_K .

However, by using the fact that $u: U \rightarrow S$ is a lift of $u_k: U_k \rightarrow S_k$ to a smooth morphism of smooth *algebraic* V -schemes, we may obtain the same module without referring to compactifications. In fact, firstly, since S_K^{an} is a strict neighborhood of \widehat{S}_K [?, (1.2.4)(ii)], $H_{\text{rig}}^i(U_k/S_k)$ is also the module of global sections of $R^i(\bar{f}_K^{\text{an}})_* \bar{j}_U^\dagger \Omega_{\bar{X}_K^{\text{an}}/\bar{S}_K^{\text{an}}}^\bullet$ on S_K^{an} by overconvergence. Moreover, the restriction of this sheaf on S_K^{an} is isomorphic to $R^i(u_K^{\text{an}})_* j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet$ by [?, 6.2.2] because again U_K^{an} is a strict neighborhood of \widehat{U}_K . Our definition of $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ and the description of $H_{\text{rig}}^i(U_k/S_k)$ in Definition ?? are thus justified.

(2) We will use the datum $H^i(U/S)$ only in the case where the rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ is known to be a coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -module for all $i \geq 0$. In this case, we also have

$$H_{\text{rig}}^i(U_k/S_k) = \Gamma(S_K^{\text{an}}, R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}})$$

by the vanishing of higher sheaf cohomologies for coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -modules. (This vanishing is perhaps well-known, but we included it as Lemma ?? at the end of this subsection because we could not find an appropriate reference.)

Now, we have defined the datum $H^i(U/S)$. This, however, does not immediately mean that it is an object of $\text{Fil-}F\text{-MIC}(S)$. For this datum to be an object in $\text{Fil-}F\text{-MIC}(S)$, we need the i -th relative cohomology $H_{\text{rig}}^i(U_k/S_k)$ to be a coherent B_K^\dagger -module and we need the morphism c to be an isomorphism. In the rest of this subsection, we discuss two settings under which these conditions hold. Briefly said, these two settings are: the case of proper smooth morphisms (Setting ??) and the case of smooth families of general dimension with “good compactification” of both the source and the target (Setting ??).

We start with the first setting.

Setting 2.14. $u: U \rightarrow S$ is a projective smooth morphism of smooth V -schemes with $S = \text{Spec}(B)$.

Proposition 2.15. *Under Setting ??, the relative rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ is a coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -module for each $i \geq 0$. Consequently, $H_{\text{rig}}^i(U_k/S_k)$ is a coherent B_K^\dagger -module for each $i \geq 0$. Moreover, the comparison morphism*

$$(2.15) \quad c: H_{\text{dR}}^i(U_K/S_K) \otimes_{B_K} B_K^\dagger \rightarrow H_{\text{rig}}^i(U_k/S_k)$$

is bijective for each $i \geq 0$.

In particular, the datum

$$H^i(U/S) = (H_{\text{dR}}^i(U_K/S_K), H_{\text{rig}}^i(U_k/S_k), c, \nabla, \Phi, \text{Fil}^\bullet)$$

is an object of $\text{Fil-}F\text{-MIC}(S)$.

Proof. Firstly, the coherence of the rigid cohomology follows from [?, Théorème 5]. Now, since c is a morphism of coherent modules over the noetherian ring B_K^\dagger , it suffices to prove

that it is an isomorphism on the reduction by each maximal ideal of B_K^\dagger , which is the extension of a maximal ideal of B_K . Therefore we may assume that $S = \bar{S} = \text{Spec}(k)$ (after a possible extension of k), and then the claim follows from comparison of the (absolute) algebraic de Rham cohomology and the rigid (or, in this case, crystalline) cohomology (e.g. [?, 4.2], [?, (7)]). \square

The second sufficient condition for $H^i(U/S)$ to be an object of $\text{Fil-}F\text{-MIC}(S)$ is, briefly said, that u has a “good compactification”.

Setting 2.16. Let $S = \text{Spec}(B)$ be a smooth affine V -scheme, let \bar{S} be a projective smooth V -scheme with an open immersion $S \hookrightarrow \bar{S}$ such that the complement T is a relative simple NCD on \bar{S} over V . Let \bar{X} be a projective smooth V -scheme, and let $\bar{f}: \bar{X} \rightarrow \bar{S}$ be a projective morphism. Let \bar{D}^h be a relative NCD on \bar{X} over V , and put $\bar{D}^v = \bar{f}^{-1}(T)$ and $\bar{D} = \bar{D}^h \cup \bar{D}^v$. We put $X = \bar{X} \setminus \bar{D}^v$, $f = \bar{f}|_X$ and $U = \bar{X} \setminus \bar{D}$. We then assume that the following conditions hold:

- (1) $\bar{D} = \bar{D}^h \cup \bar{D}^v$ is also a relative NCD over V .
- (2) $D := \bar{D}^h \cap X \hookrightarrow X$ is a relative NCD over S .
- (3) The morphism $\bar{f}: (\bar{X}, \bar{D}) \rightarrow (\bar{S}, T)$ is log smooth and integral, and (\bar{S}, T) is of Zariski type.

The notation in Setting ?? can be summarized by the diagram

$$\begin{array}{ccccc} X & \xhookrightarrow{\bar{D}^v} & \bar{X} & \xhookleftarrow{\bar{D}} & U \\ f \downarrow & & \downarrow \bar{f} & \swarrow u & \\ S & \xhookrightarrow{T} & \bar{S} & & \end{array}$$

where the notation above \hookrightarrow shows the complement of the subscheme.

Proposition 2.17. *Under Setting ??, the relative rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ is a coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -module for each $i \geq 0$. Consequently, $H_{\text{rig}}^i(U_k/S_k)$ is a coherent B_K^\dagger -module for each $i \geq 0$. Moreover, the comparison morphism*

$$c: H_{\text{dR}}^i(U_K/S_K) \otimes_{B_K} B_K^\dagger \rightarrow H_{\text{rig}}^i(U_k/S_k)$$

is bijective for each $i \geq 0$.

In particular, the datum

$$H^i(U/S) = (H_{\text{dR}}^i(U_K/S_K), H_{\text{rig}}^i(U_k/S_k), c, \nabla, \Phi, \text{Fil}^\bullet)$$

is an object of $\text{Fil-}F\text{-MIC}(S)$.

Proof. Our setting assures us that we are in the situation of [?, Section 2], i.e. the assumptions before [?, Theorem 2.1] are satisfied. Moreover, since f is integral, the assumption (\star) in [?, Theorem 2.1] is also satisfied [?, Corollary 4.7]. Therefore, the coherence of the rigid cohomology sheaves follows from [?, Theorem 2.2].

Now that the coherence is proved for all i , the proof reduces to the absolute case as in the proof of Proposition ??. Then, since the algebraic de Rham cohomology (resp. the rigid cohomology) is isomorphic to the algebraic log de Rham cohomology (resp. log rigid

cohomology by e.g. [?, Theorem 3.5.1]), the claim follows from the comparison theorem between algebraic log de Rham cohomology and log rigid cohomology [?, Corollary 2.6]. \square

We also have a Gysin exact sequence in $\text{Fil-}F\text{-MIC}(S)$ for curves under this setting.

Proposition 2.18 (Gysin exact sequence). *Let $U = \text{Spec}(A)$ and $S = \text{Spec}(B)$ be smooth affine V -schemes and let $u: U \rightarrow S$ be a smooth morphism of relative dimension one with connected fibers. Assume that there exists a projective smooth curve $f: X \rightarrow S$ with an open immersion $U \hookrightarrow X$ such that $f|_U = u$ and that the complementary divisor $D := X \setminus U$ is finite étale over S . Moreover, assume that u satisfies the conclusions of Proposition ?? (namely, the coherence of the rigid cohomology and the bijectivity of the comparison morphism), e.g. that we are in Setting ??.*

Then, we have an exact sequence

$$0 \rightarrow H^1(X/S) \rightarrow H^1(U/S) \rightarrow H^0(D/S)(-1) \rightarrow H^2(X/S) \rightarrow 0.$$

in $\text{Fil-}F\text{-MIC}(S)$.

Proof. Firstly, $H^i(X/S)$ and $H^i(D/S)$ are objects of $\text{Fil-}F\text{-MIC}(S)$ by Proposition ??, and so is $H^1(U/S)$ by assumption. Next, it is a standard fact about de Rham cohomology that we have an exact sequence

$$0 \rightarrow H_{\text{dR}}^1(X_K/S_K) \rightarrow H_{\text{dR}}^1(U_K/S_K) \rightarrow H_{\text{dR}}^0(D_K/S_K)(-1) \rightarrow H_{\text{dR}}^2(X_K/S_K) \rightarrow 0$$

whose morphisms are horizontal and compatible with respect to Fil . Therefore, by the comparison isomorphism on each term and by the flatness of B_K^\dagger over B_K , we get a corresponding exact sequence

$$0 \rightarrow H_{\text{rig}}^1(X_k/S_k) \rightarrow H_{\text{rig}}^1(U_k/S_k) \rightarrow H_{\text{rig}}^0(D_k/S_k)(-1) \rightarrow H_{\text{rig}}^2(X_k/S_k) \rightarrow 0$$

for rigid cohomologies. The compatibility of this sequence with Frobenius structures on each term can be checked on each closed point of \widehat{S}_K , therefore reduced to the absolute case [?, Theorem 2.19]. \square

The following lemma is the promised statement in Remark ?? (2).

Lemma 2.19. *Let X be a smooth V -scheme and let \mathcal{M} be a coherent $j_X^\dagger \mathcal{O}_{X_K^{\text{an}}}$ -module. Then, for any $j \geq 1$, we have $H^j(X_K^{\text{an}}, \mathcal{M}) = 0$.*

Proof. This lemma is essentially given in the proof of [?, 6.2.12]. We recall the argument for the convenience for the reader. Choose a closed immersion $X \hookrightarrow \mathbb{A}_V^N$ to an affine space and let Y be the closure of X in $X \hookrightarrow \mathbb{A}_V^N \hookrightarrow \mathbb{P}_V^N$. Then, $V_\rho := X_K^{\text{an}} \cap \mathbb{B}^N(0, \rho^+)$ for $\rho > 1$ form a cofinal family of strict neighborhoods of \widehat{X}_K .

By the coherence of \mathcal{M} , we can take a coherent $\mathcal{O}_{V_{\rho_0}}$ -module M for some $\rho_0 > 1$ such that $\mathcal{M}|_{V_{\rho_0}} = j_X^\dagger M$ [?, 5.4.4]. Then, if $j_\rho: V_\rho \hookrightarrow V_{\rho_0}$ denotes the inclusion for $1 < \rho < \rho_0$, we have isomorphisms

$$R\Gamma(X_K^{\text{an}}, \mathcal{M}) = R\Gamma(V_{\rho_0}, \mathcal{M}|_{V_{\rho_0}}) = \varinjlim_{\rho} R\Gamma(V_{\rho_0}, (j_\rho)_* M) = \varinjlim_{\rho} R\Gamma(V_\rho, M|_{V_\rho}).$$

In fact, the second identification follows from quasi-compactness and separatedness of V_{ρ_0} , and the third one holds because j_ρ is affinoid. Now, the claim follows because each V_ρ is affinoid and $M|_{V_\rho}$ is coherent. \square

2.6. Extensions associated to Milnor symbols. In this subsection, we discuss how we associate an extension in $\text{Fil-}F\text{-MIC}(U)$ to a Milnor symbol.

Let $u: U = \text{Spec}(A) \rightarrow S = \text{Spec}(B)$ be a smooth morphism of smooth affine V -scheme. We assume that u satisfies the consequences of Proposition ??, namely:

Assumption 2.20. For each $i \geq 0$, the i -th relative rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ is a coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -module and the natural morphism

$$(2.16) \quad H_{\text{dR}}^i(U_K/S_K) \otimes_{B_K} B_K^\dagger \longrightarrow H_{\text{rig}}^i(U_k/S_k)$$

is bijective.

Remark 2.21. (1) As we have discussed in Proposition ??, Assumption ?? is satisfied if we are in Setting ??.

(2) If there is a projective smooth morphism $f: X \rightarrow S$ with an open immersion $U \hookrightarrow X$ such that $X \setminus U$ is a relative simple normal crossing divisor over S , then the bijectivity of the morphism (??) is deduced from the former half of the Assumption ?? (the coherence of the i -th relative rigid cohomology sheaf for each $i \geq 0$) as in the proof of Proposition ??.

(3) Note that (a part of) Assumption ?? allows us to interpret the relative rigid cohomology as a cohomology of Monsky–Washnitzer type.

More precisely, for a smooth morphism $u: U = \text{Spec}(A) \rightarrow S = \text{Spec}(B)$ of affine smooth V -schemes, assume that, for each $i \geq 0$, the i -th rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ satisfies $H^j(S_K^{\text{an}}, R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}) = 0$ for all $j \geq 1$ (e.g. if all $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ are coherent). Then, the B_K^\dagger -module $H_{\text{rig}}^i(U_k/S_k) = \Gamma(S_K^{\text{an}}, R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{S_K^{\text{an}}})$ is isomorphic to the cohomology $H_{\text{MW}}^i(U_k/A_k) = H^i(\Omega_{A_K^\dagger/B_K^\dagger}^\bullet)$ of the complex of continuous differentials $\Omega_{A_K^\dagger/B_K^\dagger}^\bullet = \Gamma(U_K^{\text{an}}, j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet)$. This follows from our assumption and the vanishing of the cohomologies $H^j(U_K^{\text{an}}, j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^k) = 0$ for all $j \geq 1$ and $k \geq 0$ (which also follows from Lemma ??).

Recall that, for a commutative ring R , the r -th Milnor K -group $K_r^M(R)$ is defined to be the quotient of $(R^\times)^{\otimes r}$ by the subgroup generated by

$$a_1 \otimes \cdots \otimes b \otimes \cdots \otimes (-b) \otimes \cdots \otimes a_r, \quad a_1 \otimes \cdots \otimes b \otimes \cdots \otimes (1-b) \otimes \cdots \otimes a_r.$$

Recall from §?? the log object $\mathcal{L}og(f)$ in $\text{Fil-}F\text{-MIC}(U)$ for $f \in \mathcal{O}(U)^\times$, and the extension (??) which represents the class

$$[f]_U \in \text{Ext}_{\text{Fil-}F\text{-MIC}(U)}^1(\mathcal{O}_U, \mathcal{O}_U(1)).$$

For $h_0, h_1, \dots, h_n \in \mathcal{O}(U)^\times$, we associate an $(n+1)$ -extension

$$0 \longrightarrow \mathcal{O}_U(n+1) \longrightarrow \mathcal{L}og(h_n)(n) \longrightarrow \cdots \longrightarrow \mathcal{L}og(h_1)(1) \longrightarrow \mathcal{L}og(h_0) \longrightarrow \mathcal{O}_U \longrightarrow 0$$

which represents the class

$$[h_0]_U \cup \cdots \cup [h_n]_U \in \text{Ext}_{\text{Fil-}F\text{-MIC}(U)}^{n+1}(\mathcal{O}_U, \mathcal{O}_U(n+1)).$$

It is a standard argument to show that the above cup-product is additive with respect to each h_i , so that we have an additive map

$$(\mathcal{O}(U)^\times)^{\otimes n+1} \longrightarrow \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(U)}^{n+1}(\mathcal{O}_U, \mathcal{O}_U(n+1)).$$

Proposition 2.22. $[f]_U \cup [f]_U = 0$ for $f \in \mathcal{O}(U)^\times$ and $[f]_U \cup [1-f]_U = 0$ for $f \in \mathcal{O}(U)^\times$ such that $1-f \in \mathcal{O}(U)^\times$. Hence the homomorphism

$$K_{n+1}^M(\mathcal{O}(U)) \longrightarrow \mathrm{Ext}_{\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(U)}^{n+1}(\mathcal{O}_U, \mathcal{O}_U(n+1)), \quad \{h_0, \dots, h_n\} \mapsto [h_0]_U \cup \dots \cup [h_n]_U$$

is well-defined (note $\mathcal{L}og(-f) = \mathcal{L}og(f)$ by definition).

Proof. To prove this, it follows from Lemma ?? that we may assume $U = \mathrm{Spec} V[T, T^{-1}]$ and $f = T$ for the vanishing $[f]_U \cup [f]_U = 0$ and $U = \mathrm{Spec} V[T, (T - T^2)^{-1}]$ and $f = T$ for the vanishing $[f]_U \cup [1-f]_U = 0$.

Here, we show the latter vanishing. Let $C = V[T, (T - T^2)^{-1}]$ and $U = \mathrm{Spec}(C)$. Recall the dilog object $D := \mathcal{D}ilog(T)$ which has a unique increasing filtration W_\bullet (as an object of $\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(U)$) that satisfies

$$W_j D_{\mathrm{dR}} = \begin{cases} 0 & \text{if } j \leq -5, \\ C_K e_{-4} & \text{if } j = -4, -3, \\ C_K e_{-4} \oplus C_K e_{-2} & \text{if } j = -2, -1, \\ \mathcal{D}ilog(T)_{\mathrm{dR}} & \text{if } j \geq 0 \end{cases}$$

and the filtration Fil^\bullet on $W_j D_{\mathrm{dR}}$ is given to be $\mathrm{Fil}^i W_j D_{\mathrm{dR}} = W_j D_{\mathrm{dR}} \cap \mathrm{Fil}^i D_{\mathrm{dR}}$. Then, it is straightforward to check that $W_{-4} \cong \mathcal{O}_U(2)$, $W_{-2} \cong \mathcal{L}og(T)(1)$, $W_0/W_{-4} \cong \mathcal{L}og(1-T)$ and $W_0/W_{-2} \cong \mathcal{O}_U$. Thus, $[T]_U \cup [1-T]_U$ is realized by the extension

$$0 \longrightarrow W_{-4} \longrightarrow W_{-2} \longrightarrow W_0/W_{-4} \longrightarrow W_0/W_{-2} \longrightarrow 0.$$

Consider a commutative diagram

$$(2.17) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_U(2) & \xrightarrow{\mathrm{id}} & \mathcal{O}_U(2) & \xrightarrow{0} & \mathcal{O}_U & \xrightarrow{\mathrm{id}} & \mathcal{O}_U & \longrightarrow & 0 \\ & & \parallel & & \uparrow \pi_1 & & \uparrow \pi_3 & & \parallel & & \\ 0 & \longrightarrow & \mathcal{O}_U(2) & \xrightarrow{\iota} & \mathcal{O}_U(2) \oplus W_{-2} & \xrightarrow{\pi_2} & W_0 & \longrightarrow & \mathcal{O}_U & \longrightarrow & 0 \\ & & \parallel & & \downarrow \mathrm{add} & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & W_{-4} & \longrightarrow & W_{-2} & \longrightarrow & W_0/W_{-4} & \longrightarrow & \mathcal{O}_U & \longrightarrow & 0, \end{array}$$

where ι is the first inclusion, π_1 is the first projection, π_2 is the composition with the second projection and the inclusion $W_{-2} \hookrightarrow W_0$, π_3 is the quotient $W_0 \rightarrow W_0/W_{-2} \cong \mathcal{O}_U$, and $\mathrm{add}: (x, y) \mapsto x + y$. The above diagram shows the vanishing $[T]_U \cup [1-T]_U = 0$.

The proof of the vanishing $[T]_U \cup [T]_U = 0$ goes in a similar way by replacing $\mathcal{D}ilog(T)$ with $\mathrm{Sym}^2 \mathcal{L}og(T)$. \square

Let n be a non-negative integer, let $h_0, \dots, h_n \in \mathcal{O}(U)^\times$ and suppose that $\mathrm{Fil}^{n+1} H^{n+1}(U/S) = 0$. Under this setting, we define an object

$$M(U/S)_{h_0, \dots, h_n}$$

in $\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(U)$ in the following way.

Let $\mathcal{M}_{h_0, \dots, h_n}$ be the complex

$$\mathcal{L}og(h_n)(n) \longrightarrow \cdots \longrightarrow \mathcal{L}og(h_1)(1) \longrightarrow \mathcal{L}og(h_0)$$

in $\text{Fil-}F\text{-MIC}(U)$ where the first term is placed in degree 0, which fits into a distinguished triangle

$$(2.18) \quad 0 \longrightarrow \mathcal{O}_U(n) \longrightarrow \mathcal{M}_{h_0, \dots, h_n} \longrightarrow \mathcal{O}_U[-n] \longrightarrow 0$$

in the derived category of $\text{Fil-}F\text{-MIC}(U)$.

Firstly, we define the de Rham part of $M(U/S)$. Let $\mathcal{M}_{h_0, \dots, h_n, \text{dR}}$ denote the de Rham realization of $\mathcal{M}_{h_0, \dots, h_n}$. This can be seen as a complex of mixed Hodge modules by M. Saito [?]. Therefore the de Rham cohomology

$$M(U/S)_{h_0, \dots, h_n, \text{dR}} := H_{\text{dR}}^n(U_K/S_K, \mathcal{M}_{h_0, \dots, h_n, \text{dR}}) = H^n(U_K, \Omega_{U_K/S_K}^\bullet \otimes \mathcal{M}_{h_0, \dots, h_n, \text{dR}})$$

carries the Hodge filtration which we write by Fil^\bullet . Now, consider the exact sequence

$$0 \longrightarrow H_{\text{dR}}^n(U_K/S_K) \longrightarrow M(U/S)_{h_0, \dots, h_n, \text{dR}} \longrightarrow \mathcal{O}_{S_K} \xrightarrow{\delta} H_{\text{dR}}^{n+1}(U_K/S_K).$$

It follows from the Hodge theory that $\text{Im}(\delta) \subset \text{Fil}^{n+1} H_{\text{dR}}^{n+1}(U_K/S_K) = 0$. Hence we have an exact sequence

$$0 \longrightarrow H_{\text{dR}}^n(U_K/S_K) \longrightarrow M(U/S)_{h_0, \dots, h_n, \text{dR}} \longrightarrow \mathcal{O}_{S_K} \longrightarrow 0.$$

In particular, $M(U/S)_{h_0, \dots, h_n, \text{dR}}$ is locally free. Therefore, together with the connection ∇ as relative de Rham cohomology, $(M(U/S)_{h_0, \dots, h_n, \text{dR}}, \nabla, \text{Fil}^\bullet)$ is an object of $\text{Fil-MIC}(S_K)$. Here, The strictness with respect to Fil^\bullet follows from the fact that the above can be seen as an exact sequence of variations of mixed Hodge structures, again by the theory of Hodge modules [?], [?].

Let $\mathcal{M}_{h_0, \dots, h_n, \text{rig}}$ be the corresponding complex in $F\text{-MIC}(A_K^\dagger)$ to $\mathcal{M}_{h_0, \dots, h_n}$ which can be seen as a complex of overconvergent F -isocrystals. In particular, this can also be seen as a complex of coherent $j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ -modules with Frobenius structure. We define

$$M(U/S)_{h_0, \dots, h_n, \text{rig}} := H_{\text{rig}}^n(U_k/S_k, \mathcal{M}_{h_0, \dots, h_n, \text{rig}}) := \Gamma(S_K^{\text{an}}, R^n u_{\text{rig}} \mathcal{M}_{h_0, \dots, h_n, \text{rig}}),$$

where $R^n u_{\text{rig}} \mathcal{M}_{h_0, \dots, h_n, \text{rig}} = R^n(u_K^{\text{an}})_* \left(j_U^\dagger \Omega_{U_K^{\text{an}}/S_K^{\text{an}}}^\bullet \otimes \mathcal{M}_{h_0, \dots, h_n, \text{rig}} \right)$.

Now, we explain how we get a comparison isomorphism

$$c: M(U/S)_{h_0, \dots, h_n, \text{dR}} \otimes_{B_K} B_K^\dagger \xrightarrow{\cong} M(U/S)_{h_0, \dots, h_n, \text{rig}}.$$

Firstly, we have a canonical homomorphism

$$H_{\text{dR}}^i(U_K/S_K, \mathcal{M}_{h_0, \dots, h_n, \text{dR}}) \otimes_{B_K} B_K^\dagger \longrightarrow H_{\text{rig}}^i(U_k/S_k, \mathcal{M}_{h_0, \dots, h_n, \text{rig}})$$

for each i . (The construction is the same as in the case of trivial coefficients in the beginning of the previous subsection.) To prove that this is an isomorphism, as in the de Rham part, note that we have an exact sequence

$$0 \longrightarrow H_{\text{rig}}^n(U_k/S_k) \longrightarrow M(U/S)_{h_0, \dots, h_n, \text{rig}} \longrightarrow B_K^\dagger \longrightarrow H_{\text{rig}}^{n+1}(U_k/S_k)$$

because $H^1(S_K^{\text{an}}, R^n u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}/S_K^{\text{an}}}) = 0$ by Assumption ???. Thus, the comparison homomorphism is an isomorphism by the flatness of B_K^\dagger over B_K , by Assumption ??? and by five lemma.

Now, we have constructed an object $M_{h_0, \dots, h_n}(U/S)$ in $\text{Fil-}F\text{-MIC}(S)$ which fits into the exact sequence

$$(2.19) \quad 0 \longrightarrow H^n(U/S)(n+1) \longrightarrow M_{h_0, \dots, h_n}(U/S) \longrightarrow \mathcal{O}_S \longrightarrow 0.$$

The extension class (??) is additive with respect to each h_i (one can show this in the same way as the proof of bi-additivity of $[h_0]_U \cup \dots \cup [h_n]_U$, but based on the theory of mixed Hodge modules concerning the strictness of the filtrations; for the rigid part, we use the functoriality of the relative rigid cohomology and the comparison to algebraic de Rham cohomology), so that we have a homomorphism

$$(2.20) \quad (\mathcal{O}(U)^\times)^{\otimes n+1} \longrightarrow \text{Ext}_{\text{Fil-}F\text{-MIC}(S)}^1(\mathcal{O}_S, H^n(U/S)(n+1)).$$

Theorem 2.23. *Suppose $\text{Fil}^{n+1}H_{\text{dR}}^{n+1}(U_K/S_K) = 0$. Then, under the assumption ??, the homomorphism (??) factors through the Milnor K -group, so that we have a map*

$$(2.21) \quad [-]_{U/S} : K_{n+1}^M(\mathcal{O}(U)) \longrightarrow \text{Ext}_{\text{Fil-}F\text{-MIC}(S)}^1(\mathcal{O}_S, H^n(U/S)(n+1))$$

which we call the symbol map for U/S .

Remark 2.24. If it were possible to define a natural projection

$$\text{Ext}^{n+1}(\mathcal{O}_U, \mathcal{O}_U(n+1)) \longrightarrow \text{Ext}^1(\mathcal{O}_S, H^n(U/S)(n+1))$$

under the assumption $H^{n+1}(U/S) = 0$, then the object $M_{h_0, \dots, h_n}(U/S)$ should correspond to the class $[h_0]_U \cup \dots \cup [h_n]_U$, and hence Theorem ?? would be immediate from Proposition ??. However this is impossible since we do not take into consideration the boundary conditions such as admissibility etc. for constructing our category. We need to prove Theorem ?? independently while almost same argument works as well.

Proof. Write the homomorphism (??) by $\tilde{\rho}$. It is enough to show the following.

$$\tilde{\rho}(h_0 \otimes \dots \otimes f \otimes f \otimes \dots \otimes h_n) = 0, \quad \tilde{\rho}(h_0 \otimes \dots \otimes f \otimes (1-f) \otimes \dots \otimes h_n) = 0.$$

We show the latter. Let $f \in \mathcal{O}(U)^\times$ such that $1-f \in \mathcal{O}(U)^\times$. Let $u : U \rightarrow \text{Spec } V[T, (T-T^2)^{-1}]$ be the morphism given by $u^*T = f$. Recall the diagram (??), and take the pull-back by u . It follows that

$$0 \rightarrow \mathcal{O}_U(n+1) \rightarrow \overbrace{\text{Log}(h_n) \rightarrow \dots \rightarrow \text{Log}(f)(i+1) \rightarrow \text{Log}(f)(i) \rightarrow \dots \rightarrow \text{Log}(h_0)}^{\mathcal{M}} \rightarrow \mathcal{O}_U \rightarrow 0$$

is equivalent to

$$0 \rightarrow \mathcal{O}_U(n+1) \rightarrow \overbrace{\text{Log}(h_n) \rightarrow \dots \rightarrow \mathcal{O}_U(i+2)}^{\mathcal{M}'} \xrightarrow{0} \underbrace{\mathcal{O}_U(i) \rightarrow \dots \rightarrow \text{Log}(h_0)}_{\mathcal{N}} \rightarrow \mathcal{O}_U \rightarrow 0$$

as $(n+1)$ -extension in $\text{Fil-}F\text{-MIC}(U)$. This implies that \mathcal{M} is quasi-isomorphic to \mathcal{M}' as mixed Hodge modules or F -isocrystals. Hence \mathcal{N} gives rise to a splitting of

$$0 \longrightarrow H^n(U/S)(n+1) \longrightarrow M_{h_0, \dots, f, 1-f, \dots, h_n}(U/S) \longrightarrow \mathcal{O}_S \longrightarrow 0.$$

This completes the proof of $\tilde{\rho}(h_0 \otimes \dots \otimes f \otimes (1-f) \otimes \dots \otimes h_n) = 0$.

To see $\tilde{\rho}(h_0 \otimes \dots \otimes f \otimes f \otimes \dots \otimes h_n) = 0$, we consider the similar diagram to (??) obtained from $\text{Sym}^2 \text{Log}(T)$. Then the rest is the same. \square

2.7. Explicit formula. In this subsection, we continue to use the setting of the previous subsection. In particular, $U = \operatorname{Spec}(A) \rightarrow S = \operatorname{Spec}(B)$ is a smooth morphism of smooth V -schemes that satisfies Assumption ???. We fix a p -th Frobenius endomorphism φ (resp. σ) on A^\dagger (resp. B^\dagger).

Let $n \geq 0$ be an integer. Suppose that $\operatorname{Fil}^{n+1} H_{\operatorname{dR}}^{n+1}(U_K/S_K) = 0$. Recall the maps

$$K_{n+1}^M(A) \xrightarrow{[-]_{U/S}} \operatorname{Ext}_{\operatorname{Fil-F-MIC}(S)}^1(\mathcal{O}_S, H^n(U/S)(n+1)) \xrightarrow{(?)} H^1(\mathcal{S}(H^n(U/S)(n+1))).$$

Note that the last cohomology group is

$$\left\{ (\omega, \xi) \in (\Omega_{B_K}^1 \otimes \operatorname{Fil}^n H_{\operatorname{dR}}^n(U_K/S_K)) \oplus (B_K^\dagger \otimes H_{\operatorname{dR}}^n(U_K/S_K)) \mid (1 - \varphi_{n+1})\omega = d\xi, d(\omega) = 0 \right\}$$

where $\varphi_i := p^{-i}\varphi$ and d is the differential map induced from the Gauss-Manin connection as in (?). Let

(2.22)

$$\begin{array}{ccc} & \operatorname{Ext}_{\operatorname{Fil-F-MIC}(S)}^1(\mathcal{O}_S, H^n(U/S)(n+1)) & \\ \delta \swarrow & & \searrow R_\sigma \\ \Omega_{B_K}^1 \otimes \operatorname{Fil}^n H_{\operatorname{dR}}^n(U_K/S_K) & & B_K^\dagger \otimes H_{\operatorname{dR}}^n(U_K/S_K) \end{array}$$

be the compositions of (?) and the projections $(\omega, \xi) \mapsto \omega$ and $(\omega, \xi) \mapsto \xi$ respectively. The map δ agrees with the map (?), and hence it does not depend on σ , while R_σ does. We put

$$(2.23) \quad D_{U/S} := \delta \circ [-]_{U/S}, \quad \operatorname{reg}_{U/S}^{(\sigma)} := R_\sigma \circ [-]_{U/S}.$$

The purpose of this section is to give an explicit description of these maps.

Theorem 2.25. Suppose $\operatorname{Fil}^{n+1} H_{\operatorname{dR}}^{n+1}(U_K/S_K) = 0$. Let $\xi = \{h_0, \dots, h_n\} \in K_{n+1}^M(A)$. Then

$$(2.24) \quad D_{U/S}(\xi) = (-1)^n \frac{dh_0}{h_0} \wedge \frac{dh_1}{h_1} \wedge \dots \wedge \frac{dh_n}{h_n},$$

(2.25)

$$\operatorname{reg}_{U/S}^{(\sigma)}(\xi) = (-1)^n \sum_{i=0}^n (-1)^i p^{-1} \log \left(\frac{h_i^p}{h_i^\varphi} \right) \left(\frac{dh_0}{h_0} \right)^{\varphi_1} \wedge \dots \wedge \left(\frac{dh_{i-1}}{h_{i-1}} \right)^{\varphi_1} \wedge \frac{dh_{i+1}}{h_{i+1}} \wedge \dots \wedge \frac{dh_n}{h_n}.$$

Here one can think of (?) as an element of

$$\Omega_{B_K}^1 \otimes \operatorname{Fil}^n H_{\operatorname{dR}}^n(U_K/S_K) = \Omega_{B_K}^1 \otimes \Gamma(X_K, \Omega_{X_K/S_K}^n(\log D_K))$$

in the following way. Let $\tilde{\Omega}_{X_K}^\bullet(\log D_K) := \Omega_{X_K}^\bullet(\log D_K) / \operatorname{Im}(\Omega_{S_K}^2 \otimes \Omega_{X_K}^{\bullet-2}(\log D_K))$ which fits into the exact sequence

$$0 \longrightarrow \Omega_{S_K}^1 \otimes \Omega_{X_K/S_K}^{i-1}(\log D_K) \longrightarrow \tilde{\Omega}_{X_K}^i(\log D_K) \longrightarrow \Omega_{X_K/S_K}^i(\log D_K) \longrightarrow 0.$$

We think (?) of being an element of $\Gamma(X_K, \tilde{\Omega}_{X_K}^{n+1}(\log D_K))$. However, since $\operatorname{Fil}^{n+1} H_{\operatorname{dR}}^{n+1}(U_K/S_K) = 0$ by the assumption, it turns out to be an element of

$$\Gamma(X_K, \Omega_{S_K}^1 \otimes \Omega_{X_K/S_K}^n(\log D_K)) = \Omega_{B_K}^1 \otimes \Gamma(X_K, \Omega_{X_K/S_K}^n(\log D_K)).$$

Proof. In this proof, we omit to write the symbol “ \wedge ”.

Firstly, we describe the extension $[\xi]_{U/S}$. Let

$$0 \longrightarrow H_{\mathrm{dR}}^n(U/S)(n+1) \xrightarrow{\iota} M_\xi(U/S) \longrightarrow \mathcal{O}_S \longrightarrow 0$$

be the extension $[\xi]_{U/S}$ in $\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S, \sigma)$. Let $e_\xi \in \mathrm{Fil}^0 M_\xi(U/S)_{\mathrm{dR}}$ be the unique lifting of $1 \in \mathcal{O}(S_K)$. Then

$$(2.26) \quad \iota(D_{U/S}(\xi)) = \nabla(e_\xi), \quad \iota(\mathrm{reg}_{U/S}^{(\sigma)}(\xi)) = (1 - \Phi)e_\xi$$

by definition (see also (??)) where ∇ and Φ are the data in $M_\xi(U/S) \in \mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S, \sigma)$.

We first write down the term $M_\xi(U/S)$ explicitly. Write $l_i := p^{-1} \log(h_i^p/h_i^\varphi)$. Let $\{e_{i,0}, e_{i,-2}\}$ be the basis of $\mathcal{L}og(h_i)(i)$ such that $\mathrm{Fil}^{-i} \mathcal{L}og(h_i)(i)_{\mathrm{dR}} = A_K e_{i,0}$ and

$$\nabla(e_{i,0}) = \frac{dh_i}{h_i} e_{i,-2}, \quad \Phi(e_{i,0}) = p^{-i} e_{i,0} - p^{-i} l_i e_{i,-2}, \quad \Phi(e_{i,-2}) = p^{-i-1} e_{i,-2}.$$

Recall the $(n+1)$ -extension

$$0 \rightarrow \mathcal{O}_U(n+1) \rightarrow \mathcal{L}og(h_n)(n) \rightarrow \cdots \rightarrow \mathcal{L}og(h_1)(1) \rightarrow \mathcal{L}og(h_0) \rightarrow \mathcal{O}_U \rightarrow 0.$$

Let $(T_{A_K/B_K}^\bullet, D)$ be the total complex of the double complex $\Omega_{A_K/B_K}^\bullet \otimes \mathcal{L}og(h_\star)_{\mathrm{dR}}$. In more down-to-earth manner, we have

$$T_{A_K/B_K}^q := \bigoplus_{i=0}^d \Omega_{A_K/B_K}^i \otimes \mathcal{L}og(h_{i-q})(i-q)_{\mathrm{dR}}, \quad q \in \mathbb{Z}$$

where we denote $\mathcal{L}og(h_j)(j) := 0$ if $j < 0$ or $j > n$, and the differential $D : T^q \rightarrow T^{q+1}$ is defined by

$$D(\omega^i \otimes x_j) = d\omega^i \otimes x_j + (-1)^i \omega^i \wedge \nabla(x_j) + (-1)^i \omega^i \otimes \pi(x_j)$$

for $\omega^i \otimes x_j \in \Omega_{A_K/B_K}^i \otimes \mathcal{L}og(h_j)(j)_{\mathrm{dR}}$, where $\pi : \mathcal{L}og(h_i)(i) \rightarrow A_K e_{i,0}$ is the projection. We have the exact sequence

$$(2.27) \quad 0 \longrightarrow \Omega_{A_K/B_K}^{\bullet+n} \longrightarrow T_{A_K/B_K}^\bullet \longrightarrow \Omega_{A_K/B_K}^\bullet \longrightarrow 0$$

where the first arrow is induced from $\mathcal{O}_{U_K} \cong \mathcal{O}_{U_K} e_{d,-2} \hookrightarrow \mathcal{L}og(h_d)_{\mathrm{dR}}$, the second arrow induced from the projection $\mathcal{L}og(h_0)_{\mathrm{dR}} \rightarrow \mathcal{O}_{U_K} e_{0,0} \cong \mathcal{O}_{U_K}$, and the differential on $\Omega_{A_K/B_K}^{\bullet+n}$ is the usual differential operator d (not $(-1)^n d$). For the rigid part, let $(T_{A_K^\dagger/B_K^\dagger}^\bullet, D)$ be defined in the same way by replacing Ω_{A_K/B_K}^\bullet with $\Omega_{A_K^\dagger/B_K^\dagger}^\bullet$ (see Remark ?? (3)). Then, we also have an exact sequence corresponding to (??). Now, we have a description

$$M_\xi(U/S)_{\mathrm{dR}} = H^0(T_{A_K/B_K}^\bullet), \quad M_\xi(U/S)_{\mathrm{rig}} = H^0(T_{A_K^\dagger/B_K^\dagger}^\bullet)$$

(the description of the rigid part follows from the exact sequence (??) on two sides with Remark ?? (3)), and we have an exact sequence

$$0 \longrightarrow H^n(U/S)(n+1) \longrightarrow M_\xi(U/S) \longrightarrow \mathcal{O}_S \longrightarrow 0$$

in $\mathrm{Fil}\text{-}F\text{-}\mathrm{MIC}(S, \sigma)$.

Before going to the proof of Theorem ??, we give an explicit description of e_ξ in (?). Put

$$\omega^0 := 1, \quad \omega^i := \frac{dh_0}{h_0} \frac{dh_1}{h_1} \cdots \frac{dh_{i-1}}{h_{i-1}} \in \Omega_{A_K/B_K}^i \quad (1 \leq i \leq n+1).$$

However we note that $\omega^{n+1} = \omega^n \wedge \frac{dh_n}{h_n} = 0$ as $\omega^n \in \text{Fil}^{n+1} H_{\text{dR}}^{n+1}(U_K/S_K) = 0$. We claim

$$(2.28) \quad e_\xi = \sum_{i=0}^n \omega^i e_{i,0} \in T_{A_K/B_K}^0.$$

Indeed it is a direct computation to show $D(e_\xi) = 0$. We see $e_\xi \in \text{Fil}^0 M_\xi(U/S)_{\text{dR}}$ in the following way. Let $j : U_K \hookrightarrow X_K$. We think T_{A_K/B_K}^\bullet of being a complex of \mathcal{O}_{U_K} -modules. Let $\text{Fil}^0 \mathcal{L}og(h_i)_{X_K} = \mathcal{O}_{X_K} e_{i,0}$, $\text{Fil}^k \mathcal{L}og(h_i)_{X_K} = \mathcal{O}_{X_K} e_{i,0} + \mathcal{O}_{X_K} e_{i,-2}$ for $k < 0$, $\text{Fil}^k \mathcal{L}og(h_i)_{X_K} = 0$ for $k > 0$, and put

$$\text{Fil}^k \mathcal{T}_{U/S}^q = \bigoplus_{i=0}^d \Omega_{X_K/S_K}^i (\log D_K) \otimes \text{Fil}^{k-q} \mathcal{L}og(h_{i-q})_{X_K} \subset j_* T_{A_K/B_K}^q$$

locally free \mathcal{O}_{X_K} -modules. Then

$$F^k M_\xi(U/S)_{\text{dR}} = \Gamma(X_K, \text{Fil}^k \mathcal{T}_{U/S}^\bullet).$$

Therefore $e_\xi \in \text{Fil}^0 M_\xi(U/S)_{\text{dR}}$.

We prove (?). Apply $1 - \Phi$ on (?). We have

$$(2.29) \quad (1 - \Phi)e_\xi = \sum_{i=0}^n \omega^i e_{i,0} - \varphi_i(\omega^i)(e_{i,0} - l_i e_{i,-2}) = \sum_{i=0}^n (1 - \varphi_i)(\omega^i) e_{i,0} + l_i \varphi_i(\omega^i) e_{i,-2}.$$

Put

$$Q_0 := l_0, \quad Q_k := \sum_{i=0}^k (-1)^i l_i \left(\frac{dh_0}{h_0} \right)^{\varphi_1} \cdots \left(\frac{dh_{i-1}}{h_{i-1}} \right)^{\varphi_1} \frac{dh_{i+1}}{h_{i+1}} \cdots \frac{dh_k}{h_k} \quad (1 \leq k \leq n).$$

A direct calculation yields

$$dQ_k = (1 - \varphi_{k+1}) \left(\frac{dh_0}{h_0} \cdots \frac{dh_k}{h_k} \right) = (1 - \varphi_{k+1})(\omega^{k+1}).$$

It follows

$$\begin{aligned} D(Q_k e_{k+1,0}) &= (1 - \varphi_{k+1})(\omega^{k+1}) e_{k+1,0} + (-1)^k Q_k \frac{dh_{k+1}}{h_{k+1}} e_{k+1,-2} + (-1)^k Q_k e_{k,-2} \\ &= (1 - \varphi_{k+1})(\omega^{k+1}) e_{k+1,0} + (-1)^k (Q_{k+1} - (-1)^{k+1} l_{k+1} \varphi_{k+1}(\omega^{k+1})) e_{k+1,-2} + (-1)^k Q_k e_{k,-2} \\ &= [(1 - \varphi_{k+1})(\omega^{k+1}) e_{k+1,0} + l_{k+1} \varphi_{k+1}(\omega^{k+1}) e_{k+1,-2}] + (-1)^k Q_k e_{k,-2} - (-1)^{k+1} Q_{k+1} e_{k+1,-2} \end{aligned}$$

for $0 \leq k \leq n-1$. Hence

$$\begin{aligned} D \left(\sum_{k=0}^{n-1} Q_k e_{k+1,0} \right) &= (1 - \Phi)(e_\xi) - l_0 e_{0,-2} + (-1)^n (1 - \varphi_{n+1})(\eta) e_{n,-2} + Q_0 e_{0,-2} - (-1)^n Q_n e_{n,-2} \\ &= (1 - \Phi)(e_\xi) - (-1)^n Q_n e_{n,-2} \end{aligned}$$

by (??). This shows (cf. (??))

$$\mathrm{reg}_{U/S}^{(\sigma)}(\xi) = (-1)^n Q_n = (-1)^n \sum_{i=0}^n (-1)^i l_i \left(\frac{dh_0}{h_0} \right)^{\varphi_1} \cdots \left(\frac{dh_{i-1}}{h_{i-1}} \right)^{\varphi_1} \frac{dh_{i+1}}{h_{i+1}} \cdots \frac{dh_n}{h_n}$$

as required.

We prove (??). Note that $D_{U/S}(\xi)$ is characterized by $D_{U/S}(\xi)e_{n,-2} = \nabla(e_\xi)$ where ∇ is the Gauss-Manin connection on $M_\xi(U/S)_{\mathrm{dR}}$ (cf. (??)). Let $T_{A_K}^\bullet$ be the complex defined in the same way as T_{A_K/B_K}^\bullet by replacing Ω_{A_K/B_K}^\bullet with $\Omega_{A_K}^\bullet$. Let

$$\tilde{\omega}^0 := 1, \quad \tilde{\omega}^i := \frac{dh_0}{h_0} \frac{dh_1}{h_1} \cdots \frac{dh_{i-1}}{h_{i-1}} \in \Omega_{A_K}^i \quad (1 \leq i \leq n+1),$$

and $\tilde{e}_\xi := \sum_{i=0}^n \tilde{\omega}^i e_{i,0} \in T_{A_K}^0$. Then

$$D(\tilde{e}_\xi) = \sum_{i=0}^n (-1)^i \left(\frac{dh_0}{h_0} \cdots \frac{dh_i}{h_i} e_{i,-2} + \frac{dh_0}{h_0} \cdots \frac{dh_{i-1}}{h_{i-1}} e_{i-1,-2} \right) = (-1)^n \frac{dh_0}{h_0} \cdots \frac{dh_n}{h_n} e_{n,-2}$$

and this shows (cf. (??)),

$$(2.30) \quad \nabla(e_\xi) = (-1)^n \frac{dh_0}{h_0} \cdots \frac{dh_n}{h_n} e_{n,-2} \in \Omega_{B_K}^1 \otimes \Gamma(X_K, \Omega_{X_K/S_K}^n(\log D_K)) e_{n,-2}$$

as required. \square

3. COMPARISON WITH THE SYNTOMIC SYMBOL MAPS

In this section we compare the symbol map $[-]_{U/S}$ introduced in §?? with the symbol maps to the log syntomic cohomology. We refer to Kato's article [?] for the formulation and terminology of log schemes. Throughout this section, we write $(X_n, M_n) := (X, M) \otimes \mathbb{Z}/p^n \mathbb{Z}$ for a log scheme (X, M) .

3.1. Log syntomic cohomology. We work over a fine log scheme (S, L) , flat over $\mathbb{Z}_{(p)}$. We endow the DP-structure γ on $I = p\mathcal{O}_S$ compatible with the canonical DP-structure on $p\mathbb{Z}_{(p)}$. The log de Rham complex for a morphism $f : (X, M_X) \rightarrow (S, L)$ of log schemes is denoted by $\omega_{X/S}^\bullet$ ([?, (1.7)]).

Proposition 3.1 ([?, Corollary 1.11]). *Let (Y_n, M_n) be a fine log scheme over (S_n, L_n) . Let $(Y_n, M_n) \hookrightarrow (Z_n, N_n)$ be an (S_n, L_n) -closed immersion. Assume that (Z_n, N_n) has p -bases over (S_n, L_n) locally in the sense of [?, Definition 1.4]. Let (D_n, M_{D_n}) be the DP-envelope of (Y_n, M_n) in (Z_n, N_n) . Let $J_{D_n}^{[r]} \subset \mathcal{O}_{Z_n}$ be the r -th DP-ideal of D_n . Then the complex*

$$J_{D_n}^{[r-\bullet]} \otimes \omega_{Z_n/S_n}^\bullet$$

of sheaves on $(D_n)_{\mathrm{\acute{e}t}} = (Y_n)_{\mathrm{\acute{e}t}}$ does not depend on the embedding $(Y_n, M_n) \hookrightarrow (Z_n, N_n)$. In particular if (Y_n, M_n) has p -bases over (S_n, L_n) locally, then the natural morphism

$$(3.1) \quad J_{D_n}^{[r-\bullet]} \otimes \omega_{Z_n/S_n}^\bullet \longrightarrow \omega_{Y_n/S_n}^{\bullet \geq r}$$

is a quasi-isomorphism.

Concerning p -bases of log schemes, the following result is sufficient in most cases.

Lemma 3.2 ([?, Lemma 1.5]). *If $f : (X, M_X) \rightarrow (Y, M_Y)$ is a smooth morphism of fine log schemes over $\mathbb{Z}/p^n\mathbb{Z}$, then f has p -bases locally.*

Following [?, §2], we say a collection $\{(X_n, M_{X,n}, i_n)\}_n$ (abbreviated to $\{(X_n, M_{X,n})\}_n$) an adic inductive system of fine log schemes, where $(X_n, M_{X,n})$ are fine log schemes over $\mathbb{Z}/p^n\mathbb{Z}$, and $i_n : (X_n, M_{X,n}) \rightarrow (X_{n+1}, M_{X,n+1}) \otimes \mathbb{Z}/p^n\mathbb{Z}$ are isomorphisms.

Let $\{(Y_n, M_n)\}_n \hookrightarrow \{(Z_n, N_n)\}_n$ be an exact closed immersion of adic inductive systems of fine log schemes over $\{(S_n, L_n)\}_n$. We consider the following condition.

Condition 3.3. (1) For each $n \geq 1$, (Z_n, N_n) has p -bases over (S_n, L_n) locally,
 (2) Let $(D_n, M_{D_n}) \rightarrow (Z_n, N_n)$ be the DP-envelope of (Y_n, M_n) compatible with the DP-structure on (S_n, L_n) , and let $J_{D_n}^{[r]} \subset \mathcal{O}_{D_n}$ be the r -th DP-ideal. Then $J_{D_n}^{[r]}$ is flat over $\mathbb{Z}/p^n\mathbb{Z}$ and $J_{D_{n+1}}^{[r]} \otimes \mathbb{Z}/p^n\mathbb{Z} \cong J_{D_n}^{[r]}$.

Suppose that there is a compatible system $\sigma = \{\sigma_n : (S_n, L_n) \rightarrow (S_n, L_n)\}_n$ of p -th Frobenius endomorphisms.

Condition 3.4. There is a hypercovering $\{Y_n^*\}_n \rightarrow \{Y_n\}_n$ in the étale topology which admits an embedding $\{(Y_n^*, M_n^*)\}_n \hookrightarrow \{(Z_n^*, N_n^*)\}_n$ into an adic inductive system of simplicial log schemes over $\{(S_n, L_n)\}_n$, such that each $\{(Y_n^\nu, M_n^\nu)\}_n \hookrightarrow \{(Z_n^\nu, N_n^\nu)\}_n$ satisfies Condition ??, and there is a p -th Frobenius $\{\varphi_{Z_n}^\nu : (Z_n^\nu, N_n^\nu) \rightarrow (Z_n^\nu, N_n^\nu)\}_n$ compatible with σ .

Let $(Y, M) = \{(Y_n, M_n)\}_n \rightarrow \{(S_n, L_n)\}_n$ together with the p -th Frobenius σ on (S, L) satisfy Condition ??. We define the log syntomic complexes according to the construction in [?, p.539–541]¹. Let $r \geq 0$ be an integer. Let $(D_n^\nu, M_{D_n^\nu}) \rightarrow (Z_n^\nu, N_n^\nu)$ be the DP-envelopes of (Y_n^ν, M_n^ν) compatible with the DP-structure on (S_n, L_n) , and $J_{D_n^\nu}^{[r]} \subset \mathcal{O}_{D_n^\nu}$ be the r -th DP-ideal. Let

$$\mathbf{J}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^{[r], \bullet} := J_{D_n^\nu}^{[r-\bullet]} \otimes \omega_{Z_n^\nu/S_n}^\bullet$$

be the complex of sheaves on $(D_1^\nu)_{\text{ét}} = (Y_1^\nu)_{\text{ét}}$. We also write

$$\mathbf{O}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^\bullet = \mathbf{J}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^{[0], \bullet}.$$

If $r < p$, then there is the well-defined morphism

$$\varphi_r^\nu : \mathbf{J}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^{[r], \bullet} \longrightarrow \mathbf{O}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^\bullet$$

satisfying $p^r \varphi_r^\nu = (\varphi_{Z_n}^\nu)^*$ (cf. [?, p.540]). It follows from Proposition ?? that one can “glue” the complexes $\mathbf{J}_{n, (Y^\nu, M^\nu), (Z^\nu, N^\nu)/(S, L)}^{[r], \bullet}$ so that we have a complex

$$(3.2) \quad \mathbf{J}_{n, (Y, M), (Z^*, N^*)/(S, L)}^{[r], \bullet}$$

in the derived category of sheaves on $(Y_1)_{\text{ét}}$ ([?, p.541], see also [?, Remark(1.8)]). Moreover the Frobenius φ_r^ν are also glued as σ is fixed (this can be shown by the same argument as in [?, p.212]), we have

$$\varphi_r : \mathbf{J}_{n, (Y, M), (Z^*, N^*)/(S, L)}^{[r], \bullet} \longrightarrow \mathbf{O}_{n, (Y, M), (Z^*, N^*)/(S, L)}^\bullet.$$

¹Tsuji [?] defined the log syntomic complexes only in case $S = \text{Spec } \mathbb{Z}_p$ with trivial log structure. However the same construction works in general as long as σ is fixed.

Then we define the log syntomic complex $\mathcal{S}_n(r)_{(Y,M),(Z^*,N^*)/(S,L,\sigma)}$ to be the mapping fiber of

$$1 - \varphi_r : \mathbf{J}_{n,(Y,M),(Z^*,N^*)/(S,L)}^{[r],\bullet} \longrightarrow \mathbf{O}_{n,(Y,M),(Z^*,N^*)/(S,L)}^\bullet.$$

In more down-to-earth manner, the degree q -component of $\mathcal{S}_n(r)_{(Y,M),(Z^*,N^*)/(S,L,\sigma)}$ is

$$\mathbf{J}_{n,(Y,M),(Z^*,N^*)/(S,L)}^{[r],q} \oplus \mathbf{O}_{n,(Y,M),(Z^*,N^*)/(S,L)}^{q-1}$$

and the differential maps are given by $(\alpha, \beta) \mapsto (d\alpha, (1 - \varphi_{d+1})\alpha - d\beta)$. We define the log syntomic cohomology groups by

$$H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}/p^n(r)) := H_{\text{ét}}^i(Y_1, \mathcal{S}_n(r)_{(Y,M),(Z^*,N^*)/(S,L,\sigma)}),$$

$$H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}_p(r)) := \varprojlim_n H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}/p^n(r))$$

and $H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Q}_p(r)) := H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}_p(r)) \otimes \mathbb{Q}$. Note that they depend on the choice of σ .

Let $(Y', M') \rightarrow (S', L', \sigma')$ satisfy Condition ??, and let

$$\begin{array}{ccc} (Y', M') & \xrightarrow{f} & (Y, M) \\ \downarrow & & \downarrow \\ (S', L') & \longrightarrow & (S, L) \end{array}$$

be a commutative diagram of fine log schemes in which σ and σ' are compatible. Then there is the pull-back

$$f^* : H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}/p^n(r)) \longrightarrow H_{\text{syn}}^i((Y', M')/(S', L', \sigma'), \mathbb{Z}/p^n(r)).$$

Proposition 3.5. *Let $0 \leq r < p$. Suppose that $(Y, M) \rightarrow (S, L)$ is smooth. Let $\Phi_r : H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^{\bullet \geq r}) \rightarrow H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^\bullet)$ be the σ -linear map induced from φ_r . Then there is an exact sequence*

$$\begin{aligned} \cdots \longrightarrow & H_{\text{zar}}^{i-1}(Y_n, \omega_{Y_n/S_n}^{\bullet \geq r}) \xrightarrow{1-\Phi_r} H_{\text{zar}}^{i-1}(Y_n, \omega_{Y_n/S_n}^\bullet) \longrightarrow \\ & H_{\text{syn}}^i((Y, M)/(S, L, \sigma), \mathbb{Z}/p^n(r)) \longrightarrow H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^{\bullet \geq r}) \xrightarrow{1-\Phi_r} H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^\bullet) \longrightarrow \cdots \end{aligned}$$

Proof. Since $(Y, M) \rightarrow (S, L)$ is smooth, it has p -bases locally by Lemma ?. Therefore the natural morphism (??) is a quasi-isomorphism, and the exact sequence follows. \square

If $S = \text{Spec } V$ with V a p -adically complete discrete valuation ring and $Y \rightarrow S$ is proper, the exact sequence in Proposition ?? remains true after taking the projective limit with respect to n . Indeed, each term is a $V/p^n V$ -module of finite length, so that the Mittag-Leffler condition holds. The author does not know whether it is true for general (S, L) .

We attach the following lemma which we shall often use.

Lemma 3.6. *For a ring A , let $A^\wedge := \varprojlim A/p^n A$ denote the p -adic completion. Let (S, L) be a fine log scheme flat over $\mathbb{Z}_{(p)}$ such that S is affine and noetherian. Let $(Y, M) \rightarrow (S, L)$*

be a smooth morphism of fine log schemes such that $Y \rightarrow S$ is proper. Let $(Y_n, M_n) = (Y, M) \otimes \mathbb{Z}/p^n\mathbb{Z}$ and $(S_n, L_n) = (S, L) \otimes \mathbb{Z}/p^n\mathbb{Z}$. Then

$$\mathcal{O}(S)^\wedge \otimes_{\mathcal{O}(S)} H_{\text{zar}}^i(Y, \omega_{Y/S}^{\bullet \geq k}) \xrightarrow{\cong} \varprojlim_n H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^{\bullet \geq k})$$

for any k . Note that $\mathcal{O}(S)^\wedge$ is a noetherian ring ([?, Theorem 10.26]).

Proof. For an abelian group M and an integer n , we denote by $M[n]$ the kernel of the multiplication by n . An exact sequence

$$0 \longrightarrow \omega_{Y/S}^{\bullet \geq k} \xrightarrow{p^n} \omega_{Y/S}^{\bullet \geq k} \longrightarrow \omega_{Y_n/S_n}^{\bullet \geq k} \longrightarrow 0$$

gives rise to an exact sequence

$$0 \longrightarrow H_{\text{zar}}^i(Y, \omega_{Y/S}^{\bullet \geq k})/p^n \longrightarrow H_{\text{zar}}^i(Y_n, \omega_{Y_n/S_n}^{\bullet \geq k}) \longrightarrow H_{\text{zar}}^{i+1}(Y, \omega_{Y/S}^{\bullet \geq k})[p^n] \longrightarrow 0$$

of finitely generated $\mathcal{O}(S)$ -modules as $Y \rightarrow S$ is proper. Therefore the assertion is reduced to show that, for any p -torsion free noetherian ring A and any finitely generated A -module M ,

$$(3.3) \quad A^\wedge \otimes_A M \xrightarrow{\cong} \varprojlim_n M/p^n M$$

where the transition map $M/p^{n+1} \rightarrow M/p^n$ is the natural surjection, and

$$(3.4) \quad \varprojlim_n M[p^n] = 0$$

where the transition map $M[p^{n+1}] \rightarrow M[p^n]$ is multiplication by p . The isomorphism (??) is well-known ([?, Prop. 10.13]). We show (??). Let

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be any exact sequence of finitely generated A -modules. Then

$$0 \rightarrow M_1[p^n] \rightarrow M_2[p^n] \rightarrow M_3[p^n] \rightarrow M_1/p^n \rightarrow M_2/p^n \rightarrow M_3/p^n \rightarrow 0$$

is exact. Suppose that $M_2[p^n] = 0$ for all n . Then, by taking the projective limit, we have

$$\begin{array}{ccccc} 0 \longrightarrow & \varprojlim_n M_3[p^n] & \longrightarrow & \varprojlim_n (M_1/p^n) & \longrightarrow & \varprojlim_n (M_2/p^n) \\ & & & \cong \uparrow & & \uparrow \cong \\ & & & A^\wedge \otimes_A M_1 & \longrightarrow & A^\wedge \otimes_A M_2. \end{array}$$

Since $A \rightarrow A^\wedge$ is flat ([?, Prop. 10.14]), the bottom arrow is injective and hence the vanishing $\varprojlim_n M_3[p^n] = 0$ follows. For the proof of (??), apply this to $M_3 = M$ and M_2 a free B -module of finite rank.

□

3.2. Syntomic symbol maps. We review syntomic symbol maps ([?, p.205], [?, Chapter I §3], [?, §2.2], [?, p.542]). Let (Y, M) be a fine log scheme which is flat over \mathbb{Z}_p . We write $(Y_n, M_n) = (Y, M) \otimes \mathbb{Z}/p^n\mathbb{Z}$ as before. Suppose that $\{(Y_n, M_n)\}_n \rightarrow \operatorname{Spec} \mathbb{Z}/p^n\mathbb{Z}$ satisfies Condition ?? where $\operatorname{Spec} \mathbb{Z}/p^n\mathbb{Z}$ is endowed with the trivial log structure, the identity as the Frobenius and the canonical DP-structure on $p\mathbb{Z}/p^n\mathbb{Z}$. For a sheaf M of monoid, we denote by M^{gp} the associated sheaf of abelian group.

For $0 \leq r < p$, there is the natural map

$$(3.5) \quad \Gamma(Y_n, M_{n+1}^{\text{gp}}) \longrightarrow H_{\text{syn}}^1((Y, M), \mathbb{Z}/p^n(1))$$

where we omit to write “/ $(\operatorname{Spec} \mathbb{Z}/p^n\mathbb{Z}, (\mathbb{Z}/p^n\mathbb{Z})^\times, \text{id})$ ” in the notation of the syntomic cohomology or syntomic complexes in below. This is defined in the following way. We define the complex C_n as

$$C_n := (1 + J_{D_n} \longrightarrow M_{D_n}^{\text{gp}}) \quad (1 + J_{D_n} \text{ is placed in degree } 0),$$

where $M_{D_n}^{\text{gp}}$ denotes the associated sheaf of abelian groups. We define the map of complexes

$$s : C_{n+1} \longrightarrow \mathcal{S}_n(1)_{(Y, M), (Z^\star, N^\star)}$$

as the map

$$s^0 : 1 + J_{D_{n+1}}^{[1]} \longrightarrow J_{D_n}^{[1]}, \quad a \longmapsto \log(a)$$

in degree 0, and the map

$$\begin{aligned} s^1 : M_{D_{n+1}}^{\text{gp}} &\longrightarrow (\mathcal{O}_{D_n} \otimes_{\mathcal{O}_{Z_n}} \omega_{Z_n}^1) \oplus \mathcal{O}_{D_n} \\ b &\longmapsto (d\log(b), p^{-1} \log(b^p \varphi_{n+1}(b)^{-1})) \end{aligned}$$

in degree 1. Here $\varphi_{n+1}(b)b^{-p}$ belongs to $1 + p\mathcal{O}_{D_{n+1}}$ and the logarithm $p^{-1} \log(b^p \varphi_{n+1}(b)^{-1}) \in \mathcal{O}_{D_n}$ is well-defined. One easily verifies that the maps s^0 and s^1 yield a map of complexes. Since there is a natural quasi-isomorphism $C_{n+1} \cong M_{n+1}^{\text{gp}}[-1]$, the map s induces a morphism ([?, (2.2.3)])

$$M_{n+1}^{\text{gp}}[1] \longrightarrow \mathcal{S}_n(1)_{(Y, M), (Z^\star, N^\star)}$$

in the derived category, and hence (??) is obtained.

Now suppose that M is defined by a divisor $D \subset Y$. Let $U := Y \setminus D$. Then we have $M^{\text{gp}} = j_* \mathcal{O}_U^\times$, and obtain a map

$$(3.7) \quad \mathcal{O}(U_{n+1})^\times \longrightarrow H_{\text{syn}}^1((Y, M), \mathbb{Z}/p^n(1)).$$

This map and the product structure of syntomic cohomology give rise to a map

$$(3.8) \quad [-]_{\text{syn}} : K_r^M(\mathcal{O}(U_{n+1})) \longrightarrow H_{\text{syn}}^r((Y, M), \mathbb{Z}/p^n(r))$$

for $0 \leq r \leq p-1$ (cf. [?, Proposition 3.2]), which we call *the syntomic symbol map*.

3.3. Comparison of Symbol maps for U/S with Syntomic symbol maps. Let $W = W(k)$ be the Witt ring of a perfect field k of characteristic $p > 0$, and K the fractional field. Let F_W be the p -th Frobenius on W or K . We omit to write “/ $(\mathrm{Spec} W, W^\times, F_W)$ ” in the notation of syntomic complexes or syntomic cohomology, as long as there is no fear of confusion, e.g.

$$H_{\mathrm{syn}}^i((Y, M), \mathbb{Z}/p^n(r)) = H_{\mathrm{syn}}^i((Y, M)/(\mathrm{Spec} W, W^\times, F_W), \mathbb{Z}/p^n(r)), \text{ etc.}$$

For a flat (log) W -scheme X , we write $X_K := X \otimes_W K$ and $X_n := X \otimes \mathbb{Z}/p^n\mathbb{Z}$ as before.

Let Q be a smooth affine scheme over W , and T a reduced relative simple NCD over W . Let Y be a smooth scheme over W and let

$$f : Y \longrightarrow Q$$

be a projective morphism that is smooth outside $f^{-1}(T)$. Let $D \subset Y$ be a reduced relative simple NCD over W . Put $S := Q \setminus T$ and $U := Y \setminus (D \cup f^{-1}(T))$. Let L be the log structure on Q defined by T , and M the log structure on Y defined by the reduced part of $D \cup f^{-1}(T)$. We then suppose that the following conditions hold.

- U and S are affine.
- The divisor $D \cup f^{-1}(T)$ is a relative simple NCD over W and $D \cap f^{-1}(S)$ is a relative simple NCD over S .
- $f : (Y, M) \rightarrow (Q, L)$ is smooth.
- There is a system $\sigma = \{\sigma_m : (Q_m, L_m) \rightarrow (Q_m, L_m)\}_m$ of p -th Frobenius endomorphisms compatible with F_W .
- Assumption ?? holds for U/S , namely the i -th relative rigid cohomology sheaf $R^i f_{\mathrm{rig}} j_U^\dagger \mathcal{O}_{U_K^{\mathrm{an}}}$ is a coherent $j_S^\dagger \mathcal{O}_{S_K^{\mathrm{an}}}$ -module for each $i \geq 0$ (this implies the comparison isomorphism (??) by Remark ?? (2)).

Theorem 3.7. *Under the above setting, let $\omega_{Y/Q}^\bullet$ (resp. ω_{Y_m/Q_m}^\bullet) denote the de Rham complex of $(Y, M)/(Q, L)$ (resp. $(Y_m, M_m)/(Q_m, L_m)$). Let $0 \leq n < p - 1$ be an integer which satisfies $H^n(\omega_{Y_m/Q_m}^{\geq n+1}) = H^{n+1}(\omega_{Y_m/Q_m}^{\geq n+1}) = 0$ for all $m > 0$. Then the following diagram is $(-1)^n$ -commutative,*

$$\begin{array}{ccc}
 & K_{n+1}^M(\mathcal{O}(U)) & \\
 \swarrow [-]_{U/S} & & \searrow [-]_{\mathrm{syn}} \\
 \mathrm{Ext}^1(\mathcal{O}_S, H^n(U/S)(n+1)) & & H_{\mathrm{syn}}^{n+1}((Y, M), \mathbb{Z}_p(n+1)) \\
 \downarrow R_\sigma & & \downarrow \\
 \mathcal{O}(S)_K^\dagger \otimes_{\mathcal{O}(S_K)} H_{\mathrm{dR}}^n(U_K/S_K) & & H_{\mathrm{syn}}^{n+1}((Y, M)/(Q, L, \sigma), \mathbb{Z}_p(n+1)) \\
 \downarrow \cap & & \uparrow i \\
 \mathcal{O}(S)_K^\wedge \otimes_{\mathcal{O}(S_K)} H_{\mathrm{dR}}^n(U_K/S_K) & \longleftarrow & \mathcal{O}(Q)^\wedge \otimes_{\mathcal{O}(Q)} H_{\mathrm{zar}}^n(Y, \omega_{Y/Q}^\bullet).
 \end{array}$$

Here the extension group is taken in the category $\mathrm{Fil}\text{-}F\text{-MIC}(S, \sigma)$. See (??) and (??) for the definitions of the symbol maps $[-]_{U/S}$ and $[-]_{\mathrm{syn}}$, and see (??) for R_σ . The isomorphism i follows from Proposition ?? and Lemma ?? under the assumption $H^{n+1}(\omega_{Y_m/Q_m}^{\geq n+1}) = 0$.

To prove Theorem ??, we prepare for three lemmas.

Lemma 3.8. *Let $V = \operatorname{Spec} C$ be a smooth affine W -scheme. Let $\sigma = \{\sigma_n : C_n \rightarrow C_n\}_n$ be a compatible system of p -th Frobenius. Let $V(R) = \operatorname{Hom}_W(\operatorname{Spec} R, C)$ denote the set of R -valued points for a W -algebra R . Put $W_n := W/p^n W$. Then for any $a_1 \in V(W_1)$, there is a unique W -valued point $a = (a_n) \in V(W) = \varprojlim_n V(W_n)$ which makes the diagram*

$$\begin{array}{ccc} \operatorname{Spec} W_n & \xrightarrow{a_n} & V_n \\ F_W \downarrow & & \downarrow \sigma_n \\ \operatorname{Spec} W_n & \xrightarrow{a_n} & V_n. \end{array}$$

commutative for all n .

Proof. For $a_n \in V(W_n)$, we define $\phi(a_n)$ to be the morphism which makes the following diagram commutative

$$\begin{array}{ccc} \operatorname{Spec} W_n & \xrightarrow{a_n} & V_n \\ F_W \downarrow \cong & & \downarrow \sigma_n \\ \operatorname{Spec} W_n & \xrightarrow{\phi(a_n)} & V_n. \end{array}$$

One easily verifies that $\phi(a_n)$ is a W -morphism (i.e. $\phi(a_n) \in V(W_n)$), and commutes with the reduction map $\rho_n : V(W_n) \rightarrow V(W_{n-1})$. Put

$$V(W_n)_{b_{n-1}} := \{a_n \in V(W_n) \mid \rho_n(a_n) = b_{n-1}\}$$

for $b_{n-1} \in V(W_{n-1})$. This is a non-empty set as the reduction map ρ_n is surjective (formal smoothness property). We claim that a map

$$\phi : V(W_n)_{b_{n-1}} \rightarrow V(W_n)_{\phi(b_{n-1})}$$

is a constant map. Indeed, let $C_n = W_n[t_1, \dots, t_m]/I$. One can write $\sigma_n(t_i) = t_i^p + pf_i(t)$. Let a_n be given by $t_i \mapsto \alpha_i$. Then $\phi(a_n)$ is given by the ring homomorphism

$$t_i \mapsto F_W^{-1}(\alpha_i^p + pf(\alpha_1, \dots, \alpha_m)) \mod p^n W.$$

which depends only on $\{\alpha_i \mod p^{n-1} W\}_i$, namely on $\rho_n(a_n) = b_{n-1}$. This means that $\phi(a_n)$ is constant on the set $V(W_n)_{b_{n-1}}$.

We prove Lemma ??. We have shown that if $b_{n-1} \in V(W_{n-1})$ satisfies $\phi(b_{n-1}) = b_{n-1}$, then $\phi^m(a_n) = \phi(a_n)$ for all $a_n \in V(W_n)_{b_{n-1}}$ and $m \geq 1$. Let $a_1 \in V(W_1)$ be an arbitrary element, and take an arbitrary sequence

$$(a_1, c_2, \dots, c_n, \dots) \in \varprojlim_n V(W_n) = V(W).$$

Since ϕ is the identity on $V(W_1)$, one has $\phi^2(c_2) = \phi(c_2)$ as $c_2 \in V(W_2)_{a_1}$. Then $\phi^3(c_3) = \phi^2(c_3)$ as $\phi(c_3) \in V(W_3)_{\phi(c_2)}$. Continuing this, $a_n := \phi^{n-1}(c_n)$ satisfies that $\phi(a_n) = a_n$ in $V(W_n)$ and $\rho_n(a_n) = \phi^{n-1}(c_{n-1}) = \phi^{n-2}(c_{n-1}) = a_{n-1} \in V(W_{n-1})$. Hence the sequence

$\{a_n\}_n$ defines a W -valued point $a = (a_n) \in V(W)$ which makes a diagram

$$\begin{array}{ccc} \mathrm{Spec} W_n & \xrightarrow{a_n} & V_n \\ F_W \downarrow & & \downarrow \sigma_n \\ \mathrm{Spec} W_n & \xrightarrow{a_n} & V_n. \end{array}$$

commutative for all $n \geq 1$. We show that such a is unique. Suppose that another W -valued point $a' = (a'_1, a'_2, \dots, a'_n, \dots)$ makes the above diagram commutative. This means $\phi(a'_n) = a'_n$ for all $n \geq 1$. Since ϕ is a constant map on $V(W_n)_{a'_{n-1}}$, the point $a'_n = \phi(a'_n)$ is uniquely determined by a'_{n-1} , and hence by a_1 . Therefore a' is uniquely determined by a_1 , which means $a = a'$. \square

Lemma 3.9. *Suppose that $k = W/pW$ is an algebraically closed field. Let $V = \mathrm{Spec} C$ be a smooth affine scheme over W . Let \widehat{C} be the p -adic completion, and $\widehat{C}_K := \widehat{C} \otimes_W K$. Let H_K be a locally free \widehat{C}_K -module of finite rank. Let $\{P_a\}_{a \in V(k)}$ be a set of W -valued points of V such that each $P_a \in V(W)$ is a lifting of $a \in V(k)$. For an element $x \in H_K$, let $x|_{P_a} \in \kappa(P_a) \otimes_{C_K} H_K$ denote the reduction at P_a where $\kappa(P_a) \cong K$ is the residue field. If x satisfies $x|_{P_a} = 0$ for all $a \in V(k)$, then $x = 0$.*

Proof. Since there is an inclusion $H_K \hookrightarrow E_K$ into a free \widehat{C}_K -module of finite rank which has a splitting, we may replace H_K with E_K and hence we may assume that H_K is a free \widehat{C}_K -module. Then one can reduce the proof to the case of rank one, i.e. $H_K = \widehat{C}_K$. Let $x \in \widehat{C}_K$ satisfy $x|_{P_a} = 0$ for all P_a . Suppose that $x \neq 0$. There is an integer n such that $x \in p^n \widehat{C} \setminus p^{n+1} \widehat{C}$. Write $x = p^n y$ with $y \in \widehat{C} \setminus p \widehat{C}$ that satisfies $y|_{P_a} = 0$ for all $a \in V(k)$. Then y is zero in $\widehat{C}/p \widehat{C} \cong C/pC$ by Hilbert nullstellensatz, namely $y \in p \widehat{C}$. This is a contradiction. \square

Lemma 3.10. *Let Y be a projective smooth scheme over W , D_Y a relative NCD in Y over W . Let M be the log structure on Y defined by D_Y , and put $U = Y \setminus D_Y$. Then there is a canonical isomorphism*

$$(3.10) \quad c: H_{\mathrm{syn}}^i((Y, M), \mathbb{Q}_p(j)) \longrightarrow H_{\mathrm{rig-syn}}^i(U, \mathbb{Q}_p(j)).$$

Moreover, a diagram

$$(3.11) \quad \begin{array}{ccc} K_i^M(\mathcal{O}(U)) & & \\ \mathrm{reg}_{\mathrm{rig-syn}} \downarrow & \searrow [-]_{\mathrm{syn}} & \\ H_{\mathrm{rig-syn}}^i(U, \mathbb{Q}_p(i)) & \xleftarrow[\cong]{c} & H_{\mathrm{syn}}^i((Y, M), \mathbb{Q}_p(i)) \end{array}$$

is commutative.

Proof. Write $U_K = U \times_W K$ and $(Y_K, M_K) = (Y, M) \times_W K$. There is a canonical isomorphism

$$R\Gamma_{\mathrm{log-syn}}(U, \mathbb{Q}_p(j)) \cong \mathrm{Cone}[\mathrm{Fil}^j R\Gamma_{\mathrm{log-dR}}((Y_K, M_K)) \xrightarrow{1-p^{-j}\phi_{\mathrm{crys}}} R\Gamma_{\mathrm{log-crys}}((Y_1, M_1)/K)][-1]$$

arising from (??) (see also Proposition ??) where ϕ_{crys} is the p -th Frobenius which is defined thanks to the comparison of log de Rham and log crystalline cohomology. There is also a canonical isomorphism ([?, Remark 8.7, 3])

$$(3.12) \quad R\Gamma_{\text{rig-syn}}(U, \mathbb{Q}_p(j)) \cong \text{Cone}[\text{Fil}^j R\Gamma_{\text{dR}}(U_K/K) \xrightarrow{1-p^{-j}\phi_{\text{rig}}} R\Gamma_{\text{rig}}(U_1/k)][-1]$$

where ϕ_{rig} is the p -th Frobenius which is defined thanks to the comparison of the algebraic de Rham cohomology and the rigid cohomology. Now, the isomorphism c is induced by the comparison morphism between the rigid and the log crystalline cohomology [?] and the one between the de Rham cohomology and the log de Rham cohomology.

By the construction, c is compatible with respect to the cup-product and the period map to the etale cohomology,

$$\begin{array}{ccc} H_{\text{syn}}^i(U, \mathbb{Q}_p(i)) & \xrightarrow{c} & H_{\text{rig-syn}}^i(U, \mathbb{Q}_p(i)) \\ & \searrow \rho_{\text{syn}} \quad \swarrow \rho_{\text{rig-syn}} & \\ & H_{\text{ét}}^i(U_K, \mathbb{Q}_p(i)) & \end{array}$$

where ρ_{syn} is as in [?, §3.1], and $\rho_{\text{rig-syn}}$ is as in [?, Corollary 9.10]. Moreover both of $\rho_{\text{syn}} \circ [-]_{\text{syn}}$ and $\rho_{\text{rig-syn}} \circ \text{reg}_{\text{rig-syn}}$ agree with the etale symbol map ([?, Proposition 3.2.4], [?, Corollary 9.10]). We show the commutativity of the diagram (??). Thanks to the compatibility with respect to the cup-product, one can reduce the assertion to the case $i = 1$, namely it is enough to show that for $f \in \mathcal{O}(U)^\times$ an element $u := c[f]_{\text{syn}} - \text{reg}_{\text{rig-syn}}(f)$ is zero. There is an exact sequence

$$0 \longrightarrow H_{\text{dR}}^0(U_K/K) \longrightarrow H_{\text{rig-syn}}^1(U, \mathbb{Q}_p(1)) \xrightarrow{h} \text{Fil}^1 H_{\text{dR}}^1(U_K/K)$$

from (??). By the construction of c and the definition of $[-]_{\text{syn}}$ in (??) and [?, Def.6.5 and Prop.10.3], one has $h(u) = df/f - df/f = 0$. Therefore u lies in the image of $H_{\text{dR}}^0(U_K/K)$. We claim that the composition of the maps

$$H_{\text{dR}}^0(U_K/K) \longrightarrow H_{\text{rig-syn}}^1(U, \mathbb{Q}_p(1)) \xrightarrow{\rho_{\text{rig-syn}}} H_{\text{ét}}^1(U_K, \mathbb{Q}_p(1))$$

is injective. We may assume that U is connected. Moreover we may replace W with $W(\bar{k})$, so that we may further assume that U has a W -valued point. Then $H_{\text{dR}}^0(U_K/K) \cong K$ and the injectivity can be reduced to the case $U = \text{Spec } W$, which can be easily verified. We turn to the proof of $u = 0$. It is enough to show $\rho_{\text{rig-syn}}(u) = 0$. However

$$\begin{aligned} \rho_{\text{rig-syn}}(u) &= \rho_{\text{rig-syn}}(\text{reg}_{\text{rig-syn}}(f)) - \rho_{\text{rig-syn}}(c[f]_{\text{syn}}) \\ &= \rho_{\text{rig-syn}}(\text{reg}_{\text{rig-syn}}(f)) - \rho_{\text{syn}}([f]_{\text{syn}}) = [f]_{\text{ét}} - [f]_{\text{ét}} = 0 \end{aligned}$$

where $[-]_{\text{ét}}$ is the etale symbol map, so we are done. \square

(*Proof of Theorem ??*). By replacing W with $W(\bar{k})$, we may assume that k is an algebraically closed field. We fix a p -th Frobenius φ on $\mathcal{O}(U)^\dagger$ compatible with σ . Recall the diagram (??). Let $\xi \in K_{n+1}^M(\mathcal{O}(U))$, and let

$$\langle \xi \rangle_{U/S}, \quad \langle \xi \rangle_{\text{syn}} \in \mathcal{O}(S)_K^\wedge \otimes_{\mathcal{O}(S_K)} H_{\text{dR}}^n(U_K/S_K)$$

be the elements sent along the diagram in counter-clockwise direction, clockwise direction respectively. We want to show $\langle \xi \rangle_{U/S} = (-1)^n \langle \xi \rangle_{\text{syn}}$. For a closed point $a \in S_1(k)$, we take the unique lifting $P_a \in S(W)$ as in Lemma ?? . By Lemma ?? , it is enough to show

$$\langle \xi \rangle_{U/S}|_{P_a} = \langle \xi \rangle_{\text{syn}}|_{P_a} \in \kappa(P_a) \otimes_{B_K} H_{\text{dR}}^n(U_K/S_K^*)$$

for every a where $\kappa(P_a) \cong K$ denotes the residue field of P_a . Therefore, to show the $(-1)^n$ -commutativity of the diagram (??), we may specialize the diagram at P_a , so that the proof is reduced to the case $(Q, L, \sigma) = (\text{Spec } W, W^\times, F_W)$. Summing up the above, the proof of Theorem ?? is reduced to showing $(-1)^n$ -commutativity of the following diagram,

$$(3.13) \quad \begin{array}{ccc} & K_{n+1}^M(\mathcal{O}(U)) & \\ \swarrow [-]_{U/W} & & \searrow [-]_{\text{syn}} \\ \text{Ext}^1(W, H^n(U/W)(n+1)) & & H_{\text{syn}}^{n+1}((Y, M), \mathbb{Z}_p(n+1)) \\ \cong \downarrow \phi_{F_W} & & \cong \uparrow \\ H_{\text{dR}}^n(U_K/K) & \longleftarrow & H_{\text{zar}}^n(Y, \Omega_{Y/W}^\bullet(\log D)). \end{array}$$

Here Y is a projective smooth scheme over W , D a relative NCD over W , $U = Y \setminus D$ and M is the log structure on Y defined by D . The extension group is taken in the category of Fil- F -MIC(Spec W, F_W).

To compare $[-]_{U/W}$ and $[-]_{\text{syn}}$, we use the explicit formula for $[-]_{U/W}$ (Theorem ??) and the theory of rigid syntomic regulators by Besser [?]. Consider a diagram

$$(3.14) \quad \begin{array}{ccccc} & & K_{n+1}^M(\mathcal{O}(U)) & & \\ & \swarrow \text{reg}_{U/W} & \downarrow \text{reg}_{\text{rig-syn}} & \searrow [-]_{\text{syn}} & \\ & & H_{\text{rig-syn}}^{n+1}(U, \mathbb{Q}_p(n+1)) & \xleftarrow[\cong]{c} & H_{\text{syn}}^{n+1}((Y, M), \mathbb{Z}_p(n+1)) \\ & & \cong \uparrow i & & \cong \uparrow \\ H_{\text{dR}}^n(U_K/K) & \xrightarrow{(-1)^n} & H_{\text{dR}}^n(U_K/K) & \longleftarrow & H_{\text{zar}}^n(Y, \Omega_{Y/W}^\bullet(\log D)) \end{array}$$

where $\text{reg}_{U/W} := \phi_{F_W} \circ [-]_{U/W}$. Here the isomorphism i follows from [?, (8.5)] and c is the canonical isomorphism (??) in Lemma ?? . The commutativity of the right upper triangle is proven in Lemma ?? , and that of the right lower square is immediate from the construction of c . We show the commutativity of the left. Let $\xi = \{h_0, \dots, h_n\} \in K_{n+1}^M(\mathcal{O}(U))$. Then using [?, Def.6.5 and Prop.10.3] one can show that

$$(3.15) \quad \text{reg}_{\text{rig-syn}}(\xi) = \sum_{i=0}^n (-1)^i p^{-1} \log \left(\frac{h_i^p}{h_i^\varphi} \right) \left(\frac{dh_0}{h_0} \right)^{\varphi_1} \wedge \dots \wedge \left(\frac{dh_{i-1}}{h_{i-1}} \right)^{\varphi_1} \wedge \frac{dh_{i+1}}{h_{i+1}} \wedge \dots \wedge \frac{dh_n}{h_n}.$$

under the inclusion $H_{\text{dR}}^n(U_K/K) \hookrightarrow \Omega_{A_K/K}^n / d\Omega_{A_K/K}^{n-1} \cong \Omega_{A_K^\dagger/K}^n / d\Omega_{A_K^\dagger/K}^{n-1}$ in a similar way to the proof of [?, Cor. 2.9]. This agrees with $(-1)^n \text{reg}_{U/W}(\xi)$ by Theorem ?? , and hence the commutativity of the left square follows. This completes the proof of $(-1)^n$ -commutativity of the diagram (??), and hence Theorem ?? .

4. p -ADIC REGULATORS OF K_2 OF CURVES

For a regular scheme X and a divisor D , let (X, D) denote the log scheme whose log structure is defined by the reduced part of D .

4.1. Symbol map on K_2 of a projective smooth family of curves. Let $p > 2$ be a prime and let W be the Witt ring of a perfect field k of characteristic p . Put $K = \text{Frac}(W)$ the fractional field. Let F_W be the p -th Frobenius on W . Let Q be a smooth affine curve over W , and $T \subset Q$ a closed set that is finite étale over W . Put $S := Q \setminus T$. Let Y be a smooth quasi-projective surface over W , and let

$$f : Y \longrightarrow Q$$

be a projective surjective W -morphism such that f is smooth outside $F := f^{-1}(T)$, and the fibers are connected. Let $D \subset Y$ be a reduced relative simple NCD over W . Put $X := f^{-1}(S) = Y \setminus F$, $D_X := D \cap X$ and $U := X \setminus D_X$. Suppose that the following conditions hold.

- i) There is a system $\sigma = \{\sigma_n : (Q_n, T_n) \rightarrow (Q_n, T_n)\}_n$ of p -th Frobenius endomorphisms compatible with F_W , where $(X_n, M_n) := (X, M) \otimes \mathbb{Z}/p^n\mathbb{Z}$ as before.
- ii) The divisor $D + F$ is a relative simple NCD over W , and D_X is finite étale over S .
- iii) The multiplicity of each component of F is prime to p .
- iv) Assumption ?? holds for U/S ; the i -th relative rigid cohomology sheaf $R^i u_{\text{rig}} j_U^\dagger \mathcal{O}_{U_K^{\text{an}}}$ is a coherent $j_S^\dagger \mathcal{O}_{S_K^{\text{an}}}$ -module for each $i \geq 0$, where $u : U \rightarrow S$ (this implies the comparison isomorphism (??) by Remark ?? (2)).

Here is the summary of notation,

$$(4.1) \quad \begin{array}{ccc} X & \xhookrightarrow{F} & Y \xleftarrow{D \cup F} U \\ \downarrow & & \downarrow f \\ S & \xhookrightarrow{T} & Q \end{array}$$

where the notation above \hookrightarrow shows the complement of the subscheme.

The following lemma provides a sufficient condition for that the condition iv) holds.

Lemma 4.1. *Let C be a smooth affine curve over W and $v : V \rightarrow C$ a smooth W -morphism. Suppose that there is a commutative square*

$$\begin{array}{ccc} V & \hookrightarrow & \overline{Z} \\ v \downarrow & & \downarrow g \\ C & \hookrightarrow & \overline{C} \end{array}$$

where \hookrightarrow are open immersions, that satisfies the following.

- \overline{C} (resp. \overline{Z}) is a smooth projective curve (resp. a smooth projective scheme) over W .
- Put $Z := g^{-1}(C)$. Then $Z \rightarrow C$ is projective smooth.
- Put $T' := \overline{C} \setminus C$. Then T' is finite étale over W , $\overline{Z} \setminus V$ is a relative simple NCD over W , and $Z \setminus V$ is a relative simple NCD over C .
- The multiplicity of an arbitrary component of $g^{-1}(T')$ is prime to p .

Then $R^i v_{\text{rig}} j_V^\dagger \mathcal{O}_{V_K^{\text{an}}}$ is a coherent $j_C^\dagger \mathcal{O}_{C_K^{\text{an}}}$ -module for each $i \geq 0$.

Proof. The conditions imply that the morphism

$$g : (\overline{Z}, \overline{Z} \setminus V) \longrightarrow (\overline{C}, T')$$

of log schemes is smooth, so that V/C is adapted to Setting ???. Therefore $R^i v_{\text{rig}} j_V^\dagger \mathcal{O}_{V_K^{\text{an}}}$ is a coherent $j_C^\dagger \mathcal{O}_{C_K^{\text{an}}}$ -module for each $i \geq 0$ by Proposition ???. \square

We turn to the setting (??). We have the symbol map

$$[-]_{U/S} : K_2^M(\mathcal{O}(U)) \longrightarrow \text{Ext}_{\text{Fil-}F\text{-MIC}(S)}^1(\mathcal{O}_S, H^1(U/S)(2))$$

by Theorem ??. We omit to write the subscript “Fil- F -MIC(S)” in the extension groups, as long as there is no fear of confusion.

Lemma 4.2. *The following diagram is commutative,*

$$\begin{array}{ccc} K_2^M(\mathcal{O}(U)) & \xrightarrow{\partial} & \mathcal{O}(D_X)^\times \\ [-]_{U/S} \downarrow & & \downarrow [-]_{D_X/S} \\ \text{Ext}^1(\mathcal{O}_S, H^1(U/S)(2)) & \xrightarrow{\text{Res}} & \text{Ext}^1(\mathcal{O}_S, H^0(D_X/S)(1)). \end{array}$$

Here, the right vertical arrow is the symbol map defined in Theorem ??, and ∂ is the tame symbol which is defined by

$$(4.2) \quad \partial : \{f, g\} \longmapsto (-1)^{\text{ord}_{D_X}(f)\text{ord}_{D_X}(g)} \frac{f^{\text{ord}_{D_X}(g)}}{g^{\text{ord}_{D_X}(f)}} \Big|_{D_X}.$$

Proof. We first note that $K_2^M(\mathcal{O}(U))$ is generated by symbols $\{f, g\}$ with $\text{ord}_{D_X}(f) = 0$. Suppose $\text{ord}_{D_X}(f) = 0$, namely $f \in \mathcal{O}(U \cup D_X)^\times$. The natural morphism

$$p : \text{Cone}[\mathcal{L}og f(1) \rightarrow \mathcal{L}og(g)] [-1] \rightarrow \mathcal{L}og f(1)$$

and the residue map induce the following commutative diagram

$$(4.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^1(U/S)(2) & \longrightarrow & M_{f,g}(U/S) & \longrightarrow & \mathcal{O}_{S_K} \xrightarrow{\delta_{f,g}} H^2(U/S)(2) \\ & & \parallel & & \downarrow p & & \downarrow p' \\ \mathcal{O}_S & \xrightarrow{\delta_f} & H^1(U/S)(2) & \longrightarrow & H^1(\mathcal{L}og f(1)) & \longrightarrow & H^1(U/S)(1) \longrightarrow H^2(U/S)(2) \\ & & \downarrow \text{Res} & & \downarrow \text{Res} & & \downarrow \text{Res} \\ 0 & \longrightarrow & H^0(D_X/S)(1) & \longrightarrow & H^0(\mathcal{L}og f|_{D_X}) & \longrightarrow & H^0(D_X/S) \xrightarrow{\delta_{f|D}} H^1(D_X/S)(1) \end{array}$$

with exact rows where $\delta_{f,g}(1) = \frac{df}{f} \frac{dg}{g}$, $\delta_f(1) = \frac{df}{f}$ and $\delta_{f|D_X}(1) = \frac{df}{f}|_{D_X}$. The maps “Res”s are residue maps appearing in the Gysin exact sequence; as for the middle one, we are considering the Gysin exact sequence with coefficient in $\mathcal{L}og(f)(1)$ which is constructed in the same manner as Proposition ??. We show that p' agrees with the map $\delta_g : 1 \mapsto \frac{dg}{g}$. Let $[A \rightarrow B]$ denote the complex

$$\cdots \longrightarrow 0 \longrightarrow A \longrightarrow B \longrightarrow 0 \longrightarrow \cdots$$

with A placed in degree zero. Consider a commutative diagram

$$\begin{array}{ccccc} [\mathcal{L}og(f)_{dR} \rightarrow \mathcal{L}og(g)_{dR}] & \longrightarrow & [\mathcal{L}og(f)_{dR}/\mathcal{O}_{U_K}e_{-2,f} \rightarrow \mathcal{L}og(g)_{dR}] & \xrightarrow{\cong} & [0 \rightarrow \mathcal{O}_{U_K}] \\ \downarrow p & & \downarrow p' & & \\ [\mathcal{L}og(f)_{dR} \rightarrow 0] & \longrightarrow & [\mathcal{L}og(f)_{dR}/\mathcal{O}_{U_K}e_{-2,f} \rightarrow 0] & \xrightarrow{\cong} & [\mathcal{O}_{U_K} \rightarrow 0], \end{array}$$

of complexes in $\text{Fil-}F\text{-MIC}(U)$ where the two horizontal morphisms on the left-hand side are the canonical surjections and the vertical ones are the projections. Under the identification $\mathcal{L}og(f)_{dR}/\mathcal{O}_{U_K}e_{-2,f} \cong \mathcal{O}_{U_K}$ (which sends $e_{0,f}$ to 1), the homomorphism p' is nothing but the connecting morphism arising from the extension $0 \rightarrow \mathcal{O}_{U_K} \rightarrow \mathcal{L}og(g)_{dR} \rightarrow \mathcal{O}_{U_K} \rightarrow 0$. This means $p' = \delta_g$.

Since the composition $\text{Res} \circ p'$ is multiplication by $\text{ord}_D(g)$, the diagram (??) induces

$$(4.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^1(U/S)(2) & \longrightarrow & M_{f,g}(U/S) & \longrightarrow & \mathcal{O}_{S_K} \xrightarrow{\delta_{f,g}} H^2(U/S)(2) \\ & & \downarrow \text{Res} & & \downarrow & & \downarrow \text{ord}_{D_X}(g) \quad \downarrow \text{Res} \\ 0 & \longrightarrow & H^0(D_X/S)(1) & \longrightarrow & H^0(\mathcal{L}og f|D_X)' & \longrightarrow & \mathcal{O}_{S_K} \longrightarrow H^1(D_X/S)(1). \end{array}$$

Now we show the lemma. Let $\xi \in K_2^M(\mathcal{O}(U)) \cap \text{Ker}(d\log)$ be arbitrary. Fix $t \in \mathcal{O}(U)^\times$ such that $\text{ord}_{D_X}(t) > 0$. Replacing ξ with $m\xi$ for some $m > 0$, one can express

$$\xi = \{f, t\} + \sum_j \{u_j, v_j\}$$

with $\text{ord}_{D_X}(f) = \text{ord}_{D_X}(u_j) = \text{ord}_{D_X}(v_j) = 0$. Then (??) induces a commutative diagram

$$(4.5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H^1(U/S)(2) & \longrightarrow & M_\xi(U/S) & \longrightarrow & \mathcal{O}_{S_K} \longrightarrow 0 \\ & & \downarrow \text{Res} & & \downarrow & & \downarrow \text{ord}_{D_X}(t) \\ 0 & \longrightarrow & H^0(D_X/S)(1) & \longrightarrow & H^0(\mathcal{L}og f|D_X)' & \longrightarrow & \mathcal{O}_{S_K} \longrightarrow 0. \end{array}$$

Since the tame symbol of ξ is $f|_{D_X}$, the assertion follows. \square

Proposition 4.3. *Let $K_2^M(\mathcal{O}(U))_{\partial=0}$ be the kernel of the tame symbol $\partial : K_2^M(\mathcal{O}(U)) \rightarrow \mathcal{O}(D_X)^\times$. Then the symbol map $[-]_{U/S}$ uniquely extends to*

$$[-]_{X/S} : K_2^M(\mathcal{O}(U))_{\partial=0} \longrightarrow \text{Ext}^1(\mathcal{O}_S, H^1(X/S)(2)).$$

Proof. Let N be the kernel of the connecting homomorphism $H^0(D_X/S)(-1) \rightarrow H^2(X/S)$ in the Gysin exact sequence. Since $H_{\text{rig}}^0(D_X/S)$ is an overconvergent F -isocrystal on S of weight 0, the weight of N_{rig} is -2 and therefore we have $\text{Hom}_{\text{Fil-}F\text{-MIC}(S)}(\mathcal{O}_S, N(2)) = 0$. This shows that the Gysin exact sequence induces an exact sequence

$$0 \longrightarrow \text{Ext}^1(\mathcal{O}_S, H^1(X/S)(2)) \longrightarrow \text{Ext}^1(\mathcal{O}_S, H^1(U/S)(2)) \longrightarrow \text{Ext}^1(\mathcal{O}_S, N(2))$$

where the extension groups are taken in the category of $\text{Fil-}F\text{-MIC}(S)$. Now the construction of $[-]_{X/S}$ is immediate from Lemma ?? \square

Lemma 4.4. *Let*

$$H_{\text{syn}}^{\bullet}((Y, F), \mathbb{Z}_p(r)) = \varprojlim_n H_{\text{syn}}^{\bullet}((Y, F), \mathbb{Z}/p^n(r))$$

denote the syntomic cohomology with coefficients in \mathbb{Z}_p where we omit to write “/ (W, W^{\times}, F_W) ”. Then the syntomic symbol map $[-]_{\text{syn}}$ induces a map

$$K_2^M(\mathcal{O}(U))_{\partial=0} \longrightarrow H_{\text{syn}}^2((Y, F), \mathbb{Z}_p(2)),$$

which we write by the same notation. When $Q = S = \text{Spec } W$ ($F = \emptyset$), this is compatible with the regulator map by Besser [?] (or equivalently by Nekovář-Niziol [?]), which means that a diagram

$$\begin{array}{ccc} K_2^M(\mathcal{O}(U))_{\partial=0} & \xrightarrow{[-]_{\text{syn}}} & H_{\text{syn}}^2(Y, \mathbb{Z}_p(2)) \\ \downarrow & & \downarrow \\ K_2(Y)^{(2)} & \xrightarrow{\text{reg}_{\text{rig-syn}}} & H_{\text{rig-syn}}^2(Y, \mathbb{Q}_p(2)) \end{array}$$

is commutative where $K_i(-)^{(j)} \subset K_i(-) \otimes \mathbb{Q}$ denotes the Adams weight piece.

Proof. To show the former, it is enough to show that a diagram

$$(4.6) \quad \begin{array}{ccc} K_2^M(\mathcal{O}(U)) & \xrightarrow{\partial} & \mathcal{O}(D^*)^{\times} \\ \downarrow [-]_{\text{syn}} & & \downarrow [-]_{\text{syn}} \\ H_{\text{syn}}^2((Y, D + F), \mathbb{Z}/p^n(2)) & \xrightarrow{\text{Res}} & H_{\text{syn}}^1((D, D \cap F), \mathbb{Z}/p^n(1)) \end{array}$$

is commutative. Then the required symbol map is induced from an exact sequence

$$0 \rightarrow H_{\text{syn}}^2((Y, F), \mathbb{Z}/p^n(2)) \rightarrow H_{\text{syn}}^2((Y, D + F), \mathbb{Z}/p^n(2)) \rightarrow H_{\text{syn}}^1((D, D \cap F), \mathbb{Z}/p^n(1)).$$

To show the commutativity of (4.6), it is enough to check $[\partial(\xi)]_{\text{syn}} = \text{Res}([\xi]_{\text{syn}})$ for an element $\xi = \{f, g\} \in K_2^M(\mathcal{O}(U))$ such that $\text{ord}_{D_X}(f) = 0$. One has $[\partial(\xi)]_{\text{syn}} = [(f|_{D_X})^{\text{ord}_{D_X}(g)}]_{\text{syn}} = \text{ord}_{D_X}(g)[f|_{D_X}]_{\text{syn}}$. On the other hand, one has

$$\text{Res}([f]_{\text{syn}} \cup [g]_{\text{syn}}) = [f]_{\text{syn}}|_{D_X} \cup \text{Res}'([g]_{\text{syn}}) = \text{Res}'([g]_{\text{syn}})[f|_{D_X}]_{\text{syn}}$$

where

$$\text{Res}' : H_{\text{syn}}^1((Y, D + F), \mathbb{Z}/p^n(1)) \rightarrow H_{\text{syn}}^0((D, D \cap F), \mathbb{Z}/p^n(0)) = \mathbb{Z}/p^n\mathbb{Z}.$$

By the construction of $[-]_{\text{syn}}$ in §??, one can verify $\text{Res}'([g]_{\text{syn}}) = \text{ord}_{D_X}(g)$. Hence $[\partial(\xi)]_{\text{syn}} = \text{Res}([\xi]_{\text{syn}})$ as required.

When $Q = S = \text{Spec } W$, the latter assertion can be derived from Lemma ?? as $H_{\text{syn}}^2(Y, \mathbb{Q}_p(2)) \cong H_{\text{dR}}^1(Y_K/K) \rightarrow H_{\text{syn}}^2((Y, D), \mathbb{Q}_p(2)) \cong H_{\text{dR}}^1(U_K/K)$ is injective. \square

Theorem 4.5.

$$\begin{array}{ccc}
& K_2^M(\mathcal{O}(U))_{\partial=0} & \\
[-]_{X/S} \swarrow & & \searrow [-]_{\text{syn}} \\
\text{Ext}^1(\mathcal{O}_S, H^1(X/S)(2)) & & H_{\text{syn}}^2((Y, F), \mathbb{Z}_p(2)) \\
\downarrow R_\sigma \text{ (??)} & & \downarrow \\
\mathcal{O}(S)_K^\dagger \otimes_{\mathcal{O}(S)} H_{\text{dR}}^1(X/S) & & H_{\text{syn}}^2((Y, F)/(Q, T, \sigma), \mathbb{Z}_p(2)) \\
\downarrow \cap & & \uparrow \cong \\
\mathcal{O}(S)_K^\wedge \otimes_{\mathcal{O}(S)} H_{\text{dR}}^1(X/S) & \longleftarrow & \mathcal{O}(Q)^\wedge \otimes_{\mathcal{O}(Q)} H_{\text{zar}}^1(Y, \omega_{Y/Q}^\bullet)
\end{array}$$

is (-1) -commutative where the extension group is taken in the category $\text{Fil-}F\text{-MIC}(S, \sigma)$ and $\omega_{Y/Q}^\bullet$ is the log de Rham complex of $(Y, F)/(Q, T)$.

Proof. Noticing that $H_{\text{dR}}^1(X_K/S_K) \rightarrow H_{\text{dR}}^1(U_K/S_K)$ is injective, one can derive the (-1) -commutativity from Theorem ?? \square

4.2. Syntomic regulators of K_2 of Elliptic Curves with 3-torsion points. Let

$$f_{\mathbb{Q}} : X_{\mathbb{Q}} \longrightarrow S_{\mathbb{Q}} = \text{Spec } \mathbb{Q}[t, (t - t^2)^{-1}]$$

be a family of elliptic curves given by a Weierstrass equation

$$y^2 = x^3 + (3x + 4(1 - t))^2.$$

This is the universal elliptic curve over the modular curve $X_1(3) \cong \mathbb{P}_{\mathbb{Q}}^1$ with 3-torsion points $(x, y) = (0, \pm 4(1 - t))$ and $x = \infty$. The j -invariant of the generic fiber X_t is

$$j(X_t) = \frac{27(1 + 8t)^3}{t(1 - t)^3}.$$

Let $p \geq 5$ be a prime. Let $W = W(\overline{\mathbb{F}}_p)$ be the Witt ring of the algebraic closure $\overline{\mathbb{F}}_p$, and $K = \text{Frac } W$ the fractional field. We define an elliptic fibration

$$(4.7) \quad \bar{f} : \bar{Y} \longrightarrow \mathbb{P}_W^1(t)$$

over the projective line in the following way. For $\alpha \in W \cup \{\infty\}$, let P_α denote the W -valued point of $\mathbb{P}_W^1(t)$ given by $t = \alpha$. Let $S_0 = \text{Spec } W[t] \subset \mathbb{P}_W^1(t)$, and Y_0 the subscheme in $\mathbb{P}_W^2 \times_W S_0$ defined by a homogeneous equation

$$X_2^2 X_0 = X_1^3 + X_0(3X_1 + 4(1 - t)X_0)^2$$

where (X_0, X_1, X_2) denote the homogeneous coordinates of \mathbb{P}_W^2 . We set $x = X_1/X_0$ and $y = X_2/X_0$. Let $f_0 : Y_0 \rightarrow S_0$ be the natural morphism. The singular fibers are $f_0^{-1}(P_0)$ and $f_0^{-1}(P_1)$, and their singular points are $R_0 = \{(x, y, t) = (-4, 0, 0)\}$ and $R_1 = \{(x, y, t) = (0, 0, 1)\}$ respectively. The scheme Y_0 is non-regular only at the point R_1 . The fiber $f_0^{-1}(P_0)$ is the Neron 1-gon, in particular, it is not a *simple* NCD. Let $\tilde{Y}_0 \rightarrow Y_0$ be the blowing-up at $\{R_0, R_1\}$, and then we have an elliptic fibration

$$\tilde{f}_0 : \tilde{Y}_0 \longrightarrow S_0$$

that satisfies the following.

- \tilde{Y}_0 is smooth over W .
- The singular fiber $\tilde{f}_0^{-1}(P_0)$ is a non-reduced curve with two components whose multiplicities are 1 and 2, and the reduced part $\tilde{f}_0^{-1}(P_0)_{\text{red}}$ is the Neron 2-gon over W , say $\tilde{f}_0^{-1}(P_0) = Z + 2E$.
- The singular fiber $\tilde{f}_0^{-1}(P_1)$ is the Neron 3-gon over W , say $\tilde{f}_0^{-1}(P_1) = F_1 + F_2 + F_3$.

Next, let $s = 1/t$, $z = s^2x$ and $w = s^3y$. Then the Weierstrass equation of the generic fiber of \tilde{f}_0 turns out to be $w^2 = z^3 + s^2(3z + 4s(s-1))^2$. Let $S_\infty := \text{Spec } W[s, (1-s)^{-1}]$. The homogeneous equation

$$X_2^2 X_0 = X_1^3 + s^2 X_0 (3X_1 + 4s(s-1)X_0)^2, \quad z = X_1/X_0, \quad w = X_2/X_0$$

defines a subscheme $Y_\infty \subset \mathbb{P}_W^2 \times_W S_\infty$ and a projective flat morphism $f_\infty : Y_\infty \rightarrow S_\infty$. The singular fiber is $f_\infty^{-1}(P_\infty)$ with one singular point $R_\infty = \{(z, w, s) = (0, 0, 0)\}$, and the scheme Y_∞ is non-regular only at R_∞ . It is a standard exercise to resolve it. Namely, 4-time blowing-ups $\tilde{Y}_\infty \rightarrow \cdots \rightarrow Y_\infty$ provides an elliptic fibration

$$\tilde{f}_\infty : \tilde{Y}_\infty \longrightarrow S_\infty,$$

and it satisfies the following.

- \tilde{Y}_∞ is smooth over W .
- The singular fiber $\tilde{f}_\infty^{-1}(P_\infty)$ is the (non-reduced) curve of Kodaira type IV* over W , say $\tilde{f}_\infty^{-1}(P_\infty) = C_1 + C_2 + C_3 + 2E_1 + 2E_2 + 2E_3 + 3F$ (e.g. [?, IV, Fig.4.4]). This is a relative simple NCD over W .

We define the fibration (??) to be

$$\bar{f} : \bar{Y} = \tilde{Y}_0 \cup \tilde{Y}_\infty \longrightarrow \mathbb{P}_W^1(t).$$

Let $Q := \mathbb{P}_W^1 \setminus \{P_1, P_\infty\} \supset S := \mathbb{P}_W^1 \setminus \{P_0, P_1, P_\infty\}$ and put $Y := \bar{f}^{-1}(Q)$, $X := \bar{f}^{-1}(S)$. Let $c \in 1 + pW$, and let $\sigma = \{\sigma_n : (Q_n, P_{0,n}) \rightarrow (Q_n, P_{0,n})\}_n$ be the system of p -th Frobenius endomorphisms given by $\sigma_n(t) = ct^p$. Let $\bar{D}_\pm \subset \bar{Y}$ be the closure of sections $(x, y) = (0, \pm 4(1-t))$ and \bar{D}_∞ the closure of the infinity section $x = \infty$. Put $\bar{D} = \bar{D}_+ \cup \bar{D}_- \cup \bar{D}_\infty$, $D = Y \cap \bar{D}$, $D_X = D \cap X$ and $U = X \setminus D_X$. By the construction of \bar{f} , one sees that the following holds.

- Each component of \bar{D} defines a section of \bar{f} (and hence is isomorphic to \mathbb{P}_W^1).
- In the fiber $f^{-1}(P_0) = Z + 2E$, \bar{D} meets only with Z , and each intersection is transversal.
- In the fiber $f^{-1}(P_1)$, \bar{D}_+ meets only with F_1 , \bar{D}_- meets only with F_2 , and \bar{D}_∞ meets only with F_3 . Each intersection is transversal.
- In the fiber $f^{-1}(P_\infty)$, \bar{D}_+ meets only with C_1 , \bar{D}_- meets only with C_2 , and \bar{D}_∞ meets only with C_3 . Each intersection is transversal.

Now the data $(Y/Q, S, D, \sigma)$ satisfies the conditions i), ..., iv) in the beginning of §?? (note iv) follows from Lemma ??).

Let \widehat{B} (resp. B^\dagger) denote the p -adic completion (resp. the weak completion) and write $\widehat{B}_K := \widehat{B} \otimes_W K$, $B_K^\dagger := B^\dagger \otimes_W K$ as before.

We consider a Milnor symbol

$$(4.8) \quad \xi := \left\{ \frac{y - 3x - 4(1-t)}{-8(1-t)}, \frac{y + 3x + 4(1-t)}{8(1-t)} \right\} \in K_2^M(\mathcal{O}(U)).$$

It is a simple exercise to show $\partial(\xi) = 0$ where ∂ is the tame symbol (??). Hence we have a 1-extension

$$[\xi]_{X/S} \in \text{Ext}_{\text{Fil-}F\text{-MIC}(S)}^1(\mathcal{O}_S, H^1(X/S)(2))$$

by Proposition ?? . Put $h = 3x + 4(1-t)$. One has

$$\begin{aligned} d\log(\xi) &= d\log\left(\frac{y-h}{-8(1-t)}\right) \wedge d\log\left(\frac{y+h}{8(1-t)}\right) \\ &= \frac{dy-dh}{y-h} \wedge \frac{dy+dh}{y+h} + \frac{dt}{t-1} \wedge \left(\frac{dy-dh}{y-h} - \frac{dy+dh}{y+h}\right) \\ &= \frac{6}{x^3} \left(dy \wedge dx - \frac{x}{t-1} dy \wedge dt + \frac{y}{t-1} dx \wedge dt\right) \quad (\text{by } y^2 - h^2 = x^3) \\ &= \frac{3dt}{t-1} \wedge \frac{dx}{y} \in \Gamma(X_K, \Omega_{X_K/K}^2). \end{aligned}$$

Hence

$$D_{X/S}(\xi) = -3 \frac{dt}{t-1} \otimes \frac{dx}{y} \in \Omega_{B_K}^1 \otimes H_{\text{dR}}^1(X_K/S_K).$$

by Theorem ?? (??). The purpose of this section is to describe

$$\text{reg}_{X/S}^{(\sigma)}(\xi) \in B_K^\dagger \otimes H_{\text{dR}}^1(X_K/S_K)$$

in terms of the hypergeometric functions

$${}_2F_1\left(\begin{matrix} a, b \\ 1 \end{matrix}; t\right) := \sum_{n=0}^{\infty} \frac{(a)_n}{n!} \frac{(b)_n}{n!} t^n$$

where $(a)_n := a(a+1) \cdots (a+n-1)$ denotes the Pochhammer symbol. Put

$$(4.9) \quad \omega := \frac{dx}{y}, \quad \eta := \frac{x dx}{y}$$

a B_K -basis of $H_{\text{dR}}^1(X_K/S_K)$. Let

$$F(t) = \frac{1}{2\sqrt{-3}} {}_2F_1\left(\begin{matrix} \frac{1}{3}, \frac{2}{3} \\ 1 \end{matrix}; t\right),$$

and put

$$(4.10) \quad \hat{\omega} := \frac{1}{F(t)} \frac{dx}{y}, \quad \hat{\eta} := 4(1-t)(F(t) + 3tF'(t)) \frac{dx}{y} + F(t) \frac{x dx}{y}.$$

a $K((t))$ -basis of $K((t)) \otimes H_{\text{dR}}^1(X_K/S_K)$. Let $\Delta := \text{Spec } W[[t]] \rightarrow \mathbb{P}_W^1(t)$ and $O := \text{Spec } W[[t]]/(t) \hookrightarrow \Delta$. The fibration $\mathcal{E} := f^{-1}(\Delta)$ over Δ is a Tate elliptic curve with central fiber $f^{-1}(O) = F = Z + 2E$. Let $q \in tW[[t]]$ be the Tate period of \mathcal{E}/Δ which is characterized by

$$j(X_t) = \frac{27(1+8t)^3}{t(1-t)^3} = \frac{1}{q} + 744 + 196884q + \cdots$$

$$\left(\Rightarrow q = \frac{1}{27}t + \frac{250289}{243}t^2 - \frac{5507717}{243}t^3 + \frac{25287001}{81}t^4 + \dots \right).$$

Let $\omega_{\mathcal{E}/\Delta}^\bullet$ be the de Rham complex of $(\mathcal{E}, F)/(\Delta, O)$. Thanks to the fundamental theorem on log crystalline cohomology by Kato, we have the canonical isomorphism

$$H_{\text{zar}}^i(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet) \cong H_{\text{crys}}^i((\mathcal{E}_1, F_1)/(\Delta, O))$$

([?, Theorem 6.4], see also Proposition ??) where $(\mathcal{E}_n, Z_n) = (\mathcal{E}, Z) \otimes \mathbb{Z}/p^n\mathbb{Z}$ as before.

Proposition 4.6. *Let $\nabla : K((t)) \otimes H_{\text{dR}}^1(X_K/S_K) \rightarrow K((t))dt \otimes H_{\text{dR}}^1(X_K/S_K)$ be the Gauss-Manin connection. Then*

$$(4.11) \quad \nabla(\omega) = -\frac{dt}{3t}\omega + \frac{dt}{12(t^2-t)}\eta, \quad \nabla(\eta) = \frac{4dt}{3t}\omega + \frac{dt}{3t}\eta,$$

$$(4.12) \quad \nabla(\hat{\omega}) = \frac{dq}{q} \otimes \hat{\eta}, \quad \nabla(\hat{\eta}) = 0.$$

Let $L := \text{Frac}(W[[t]])$. Then $\{\frac{1}{2}\hat{\omega}, \frac{1}{2}\hat{\eta}\}$ forms the de Rham symplectic basis of $L \otimes_{W[[t]]} H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$ in the sense of §??. Moreover $\hat{\omega}$ and $\hat{\eta}$ lie in $H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$.

Proof. One can derive (??) from the well-known formula on the Gauss-Manin connection for an elliptic fibration (e.g. [?, p.304], [?, Theorem 7.1]). We show that $\{\frac{1}{2}\hat{\omega}, \frac{1}{2}\hat{\eta}\}$ forms the de Rham symplectic basis. Then (??) follows from Proposition ??. Firstly, thanks to (??), one can directly show

$$(4.13) \quad \nabla(\hat{\omega}) = \frac{dt}{12(t^2-t)F(t)^2} \otimes \hat{\eta}, \quad \nabla(\hat{\eta}) = 0$$

where we note that the hypergeometric function $F(t)$ satisfies $t(1-t)F''(t) + (1-2t)F'(t) - \frac{2}{9}F(t) = 0$ (e.g. [?, 16.8.5]). Let $\{\hat{\omega}_{\text{dR}}, \hat{\eta}_{\text{dR}}\}$ be the de Rham symplectic basis. Then

$$(4.14) \quad \nabla(\hat{\omega}_{\text{dR}}) = \frac{dq}{q} \otimes \hat{\eta}_{\text{dR}}, \quad \nabla(\hat{\eta}_{\text{dR}}) = 0$$

by Proposition ??. Since $\ker(\nabla) \cong K$, we have $\hat{\eta} = a\hat{\eta}_{\text{dR}}$ for some $a \in K^\times$. Since $\hat{\omega}$ and $\hat{\omega}_{\text{dR}}$ are regular 1-forms, there is a $h \in L^\times$ such that $\hat{\omega} = h\hat{\omega}_{\text{dR}}$. We then have

$$\nabla(\hat{\omega}) = \nabla(h\hat{\omega}_{\text{dR}}) = dh \otimes \hat{\omega}_{\text{dR}} + h\nabla(\hat{\omega}_{\text{dR}}) = dh \otimes \hat{\omega}_{\text{dR}} + h\frac{dq}{q} \otimes \hat{\eta}_{\text{dR}}.$$

By (??), $h = b$ is a constant. Let $\mathbb{G}_m = Z \setminus (Z \cap E)$ be the reduced regular locus of F and $i : \mathbb{G}_m \hookrightarrow \mathcal{E}$ the inclusion. Put $u_0 = y/(x+4)$ and $u = (u_0 - \sqrt{-3})/(u_0 + \sqrt{-3})$. Then $\mathbb{G}_m = \text{Spec } W[u, u^{-1}]$ and

$$i^*\hat{\omega} = 2\sqrt{-3}i^*\left(\frac{dx}{y}\right) = 2\frac{du}{u} = 2i^*\hat{\omega}_{\text{dR}}.$$

This shows $\hat{\omega} = 2\hat{\omega}_{\text{dR}}$. By (??) and (??), we have

$$\frac{dt}{12(t^2-t)F(t)^2} \otimes \hat{\eta} = 2\frac{dq}{q} \otimes \hat{\eta}_{\text{dR}} = 2a^{-1}\frac{dq}{q} \otimes \hat{\eta}.$$

Take the residue at $t = 0$, and then we see $a = 2$.

There remains to show $\widehat{\omega}, \widehat{\eta} \in H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$. It is straightforward to see that $dx/y \in \Gamma(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$, so that one has $\widehat{\omega} \in \Gamma(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$. The Gauss-Manin connection induces a connection

$$\nabla : H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet) \longrightarrow \frac{dt}{t} \otimes H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet),$$

and one has from (??)

$$\widehat{\eta} = q \frac{d}{dq} \widehat{\omega} = u(t) \cdot t \frac{d}{dt} \widehat{\omega}, \quad \text{where } u(t) := t^{-1} q \frac{dt}{dq} \in W[[t]]^\times$$

(note that $W[[q]] = W[[t]]$). This shows $\widehat{\eta} \in H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$. This completes the proof. \square

Proposition 4.7. *Let $c \in 1 + pW$ and let σ be the p -th Frobenius on \widehat{B}_K defined by $\sigma(t) = ct^p$. Let Φ be the σ -linear Frobenius on $H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet)$. Put*

$$\tau^{(\sigma)}(t) = p^{-1} \log \left(\frac{q^p}{q^\sigma} \right) = -p^{-1} \log(27^{p-1}c) + p^{-1} \log(q_0^p/q_0^\sigma), \quad \left(q_0 := \frac{27q}{t} \right)$$

where $\log : 1 + pW[[t]] \rightarrow W[[t]]$ is defined by the customary Taylor expansion. Then

$$\begin{pmatrix} \Phi(\widehat{\omega}) & \Phi(\widehat{\eta}) \end{pmatrix} = \begin{pmatrix} \widehat{\omega} & \widehat{\eta} \end{pmatrix} \begin{pmatrix} p & 0 \\ -p\tau^{(\sigma)}(t) & 1 \end{pmatrix}$$

Proof. Note $W[[t]] = W[[q]]$. This is a special case of Theorem ?? in Appendix. \square

Let ξ be the symbol (??). Let $\sigma : B^\dagger \rightarrow B^\dagger$ be the p -th Frobenius defined by $\sigma(t) = ct^p$. Let

$$0 \longrightarrow H_{\text{dR}}^1(X/S)(2) \longrightarrow M_\xi(X/S) \longrightarrow \mathcal{O}_S \longrightarrow 0$$

be the 1-extension in $\text{Fil-}F\text{-MIC}(S, \sigma)$, associated to $[\xi]_{X/S}$. Let $e_\xi \in \text{Fil}^0 M_\xi(X_K/S_K)_{\text{dR}}$ be the unique lifting of $1 \in \mathcal{O}(S_K)$. Define $\varepsilon_i(t), E_i(t)$ by

$$(4.15) \quad \text{reg}_{X/S}^{(\sigma)}(\xi) = e_\xi - \Phi(e_\xi) = \varepsilon_1(t)\omega + \varepsilon_2(t)\eta$$

$$(4.16) \quad = E_1(t)\widehat{\omega} + E_2(t)\widehat{\eta}.$$

It follows that $\varepsilon_i(t) \in B_K^\dagger$. By (??), one immediately has

$$(4.17) \quad \varepsilon_1(t) = \frac{E_1(t)}{F(t)} + 4(1-t)(F(t) + 3tF'(t))E_2(t),$$

$$(4.18) \quad \varepsilon_2(t) = F(t)E_2(t).$$

The power series $E_i(t)$ are explicitly described as follows.

Theorem 4.8. *Let ξ be as in (??). Put $\nu = (-1 + \sqrt{-3})/2$. Then $E_1(t), E_2(t) \in K[[t]]$ and they are characterized by*

$$(4.19) \quad \frac{d}{dt} E_1(t) = -3 \left(F(t) \frac{dt}{t-1} - F(t)^\sigma \frac{p^{-1} dt^\sigma}{t^\sigma - 1} \right), \quad E_1(0) = 0,$$

$$(4.20) \quad \frac{d}{dt} E_2(t) = -E_1(t) \frac{q'}{q} - 3F(t)^\sigma \tau^{(\sigma)}(t) \frac{p^{-1} dt^\sigma}{t^\sigma - 1}, \quad E_2(0) = -9 \ln_2^{(p)}(-\nu).$$

Proof. Apply ∇ on (??). By Proposition ?? together with the fact that $\Phi\nabla = \nabla\Phi$, one has

$$(4.21) \quad \nabla(e_\xi) - \Phi(\nabla(e_\xi)) = dE_1(t) \otimes \widehat{\omega} + \left(E_1(t) \frac{dq}{q} + dE_2(t) \right) \otimes \widehat{\eta}.$$

Since

$$\nabla(e_\xi) = -d\log(\xi) = -3 \frac{dt}{t-1} \otimes \frac{dx}{y} = -3F(t) \frac{dt}{t-1} \otimes \widehat{\omega}$$

(cf. (??)), the left hand side of (??) is

$$\begin{aligned} & -3F(t) \frac{dt}{t-1} \otimes \widehat{\omega} + 3p^{-1}\sigma \left(F(t) \frac{dt}{t-1} \right) \otimes p^{-1}\Phi(\widehat{\omega}) \\ &= -3F(t) \frac{dt}{t-1} \otimes \widehat{\omega} + 3F(t)^\sigma \frac{p^{-1}dt^\sigma}{t^\sigma-1} \otimes (\widehat{\omega} - \tau^{(\sigma)}(t)\widehat{\eta}) \quad (\text{Proposition ??}) \\ &= -3 \left(F(t) \frac{dt}{t-1} - F(t)^\sigma \frac{p^{-1}dt^\sigma}{t^\sigma-1} \right) \otimes \widehat{\omega} - 3F(t)^\sigma \tau^{(\sigma)}(t) \frac{p^{-1}dt^\sigma}{t^\sigma-1} \otimes \widehat{\eta}. \end{aligned}$$

Therefore one has

$$(4.22) \quad \frac{d}{dt} E_1(t) = -3 \left(F(t) \frac{dt}{t-1} - F(t)^\sigma \frac{p^{-1}dt^\sigma}{t^\sigma-1} \right),$$

and

$$(4.23) \quad \frac{d}{dt} E_2(t) = -E_1(t) \frac{q'}{q} - 3F(t)^\sigma \tau^{(\sigma)}(t) \frac{p^{-1}dt^\sigma}{t^\sigma-1}.$$

The differential equation (??) implies that $E_1(t) \in K[[t]] \cap \widehat{B}_K$. It determines all coefficients of $E_1(t)$ except the constant term. Note $\tau^{(\sigma)}(t) \in W[[t]]$. Taking the residue at $t = 0$ of the both sides of (??), one concludes $E_1(0) = 0$. We thus have the full description of $E_1(t)$. Since $E_1(t) \in tK[[t]]$, the differential equation (??) implies that $E_2(t) \in K[[t]] \cap \widehat{B}_K$, and it determines all coefficients of $E_2(t)$ except the constant term.

The rest of the proof is to show $E_2(0) = -9\ln_2^{(p)}(-\nu)$. To do this, we look at the syntomic cohomology of the singular fiber F . Recall that F has two components Z, E and the multiplicity of Z (resp. E) is 1 (resp. 2). The reduced part $F_{\text{red}} = Z \cup E$ is a Neron 2-gon, and the divisor D intersects only with Z . Let F_\star be the simplicial nerve of the normalization of F_{red} . Let $i_{F_{\text{red}}} : F_{\text{red}} \hookrightarrow \mathcal{E}$ and let $i_{F_\star} : F_\star \rightarrow \mathcal{E}$. There is a commutative diagram

$$\begin{array}{ccccc} H_{\text{syn}}^2(\mathcal{E}, \mathbb{Z}_p(2)) & \xrightarrow{i_{F_\star}^*} & H_{\text{syn}}^2(F_\star, \mathbb{Z}_p(2)) & \xleftarrow{\cong} & H_{\text{dR}}^1(F_\star/W) \\ & & & & \downarrow \cong u' \\ & & & & H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}) \\ \rho_\sigma \curvearrowright & \downarrow & & & \uparrow u \\ & H_{\text{syn}}^2((\mathcal{E}, F)/(\Delta, O, \sigma), \mathbb{Z}_p(2)) & & & \\ & \uparrow \cong & & & \\ & H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet) & \xrightarrow{i_{F_{\text{red}}}^*} & H_{\text{zar}}^1(F_{\text{red}}, \omega_{\mathcal{E}/\Delta}^\bullet|_{F_{\text{red}}}) & \end{array}$$

where u and u' are the maps induced from the canonical maps $\omega_{\mathcal{E}/\Delta}^\bullet|_{F_{\text{red}}} \rightarrow \mathcal{O}_{F_\star}$ and $\Omega_{F_\star}^\bullet \rightarrow \mathcal{O}_{F_\star}$. By Theorem ??, $\text{reg}_{X/S}^{(\sigma)}(\xi)$ is related to $[\xi]_{\text{syn}} \in H_{\text{syn}}^2(\mathcal{E}, \mathbb{Z}_p(2))$ as follows,

$$(4.24) \quad \rho_\sigma([\xi]_{\text{syn}}) = -E_1(t)\hat{\omega} - E_2(t)\hat{\eta} \in H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet).$$

To compute $E_2(0)$, we shall compute the both side of

$$(4.25) \quad (u' \circ i_{F_\star}^*)([\xi]_{\text{syn}}) = -E_2(0)(u \circ i_{F_{\text{red}}}^*)(\hat{\eta}) \in H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}).$$

Let $z_0 = y/(x+4)|_Z$ be a coordinate of $Z \cong \mathbb{P}_W^1$. Let R_1, R_2 be the intersection points of Z and E given by $z_0 = \sqrt{-3}, -\sqrt{-3}$ respectively. We use another coordinate $z := \nu^2(1 - z_0)/2$ of Z , so that the rational functions

$$h_1 := \frac{y - 3x - 4(1-t)}{-8(1-t)}, \quad h_2 := \frac{y + 3x + 4(1-t)}{8(1-t)} \in \mathcal{O}(U)^\times$$

in the Milnor symbol ξ in (??) satisfy

$$(4.26) \quad h_1|_Z = z^3, \quad h_2|_Z = (1 - \nu z)^3, \quad h_1|_E = h_2|_E - 1.$$

The points R_1, R_2 are given by $z = -1, -\nu$ respectively. Let $\mathcal{S}_n(r)_V = \mathcal{S}_n(r)_{V/(W, F_W)}$ denote the syntomic complex for $V = Z, E$ or $Z \cap E$. We fix isomorphisms

$$\begin{aligned} H_{\text{syn}}^j(F_\star, \mathbb{Z}/p^n(r)) &\cong H_{\text{et}}^j(F_{\text{red}}, \mathcal{S}_n(r)_Z \oplus \mathcal{S}_n(r)_E \xrightarrow{i} \mathcal{S}_n(r)_{Z \cap E}) \\ H_{\text{dR}}^j(F_\star/W) &\cong H_{\text{zar}}^j(F_{\text{red}}, \Omega_{F_\star}^\bullet \oplus \Omega_E^\bullet \xrightarrow{i} \mathcal{O}_{Z \cap E}) \\ H_{\text{zar}}^j(F_\star, \mathcal{O}_{F_\star}) &\cong H_{\text{zar}}^j(F_{\text{red}}, \mathcal{O}_Z \oplus \mathcal{O}_E \xrightarrow{i} \mathcal{O}_{Z \cap E}), \end{aligned}$$

where $i : (f, g) \mapsto f|_{Z \cap E} - g|_{Z \cap E}$. We also fix an isomorphism

$$\alpha : H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}) \cong \text{Coker}[H^0(\mathcal{O}_Z) \oplus H^0(\mathcal{O}_E) \xrightarrow{i} H^0(\mathcal{O}_{R_1}) \oplus H^0(\mathcal{O}_{R_2})] \xrightarrow{\cong} W,$$

where the last isomorphism is given by $(c_1, c_2) \mapsto c_1 - c_2$.

We compute the right hand side of (??). The map $u \circ i_{F_{\text{red}}}^*$ agrees with the composition of the maps,

$$H_{\text{zar}}^1(\mathcal{E}, \omega_{\mathcal{E}/\Delta}^\bullet) \xrightarrow{u''} H_{\text{zar}}^1(\mathcal{E}, \mathcal{O}_{\mathcal{E}}) \xrightarrow{i_{F_\star}^*} H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}).$$

Let $y_1 = 2y$ and $x_1 = x + 3$ so that we have $y^2 = x^3 + (3x + 4 - 4t)^2 \Leftrightarrow y_1^2 = 4x_1^3 - g_2x_1 - g_3$. Let $\mathcal{E} = U_0 \cup U_\infty$ be an affine open covering given by $U_0 = \{y_1^2 = 4x_1^3 - g_2x_1 - g_3\}$ and $U_\infty = \{w^2 = 4v - g_2v^3 - g_3v^4\}$ with $v = 1/x_1$, $w = y_1/x_1^2$. Then one can compute the cohomology $H_{\text{zar}}^i(\mathcal{E}, \mathcal{F}^\bullet)$ for a bounded complex \mathcal{F}^\bullet of quasi-coherent sheaves by the total complex of Cech complex

$$\Gamma(U_0, \mathcal{F}^\bullet) \oplus \Gamma(U_\infty, \mathcal{F}^\bullet) \longrightarrow \Gamma(U_0 \cap U_\infty, \mathcal{F}^\bullet), \quad (f_0, f_\infty) \longmapsto f_0|_{U_0 \cap U_\infty} - f_\infty|_{U_0 \cap U_\infty}.$$

In particular, $H_{\text{zar}}^1(\mathcal{E}, \mathcal{F}^\bullet)$ is the kernel of the map

$$\begin{aligned} \Gamma(U_0 \cap U_\infty, \mathcal{F}^0) \times (\Gamma(U_\infty, \mathcal{F}^1) \oplus \Gamma(U_\infty, \mathcal{F}^1)) &\rightarrow \Gamma(U_0 \cap U_\infty, \mathcal{F}^1) \times (\Gamma(U_\infty, \mathcal{F}^2) \oplus \Gamma(U_\infty, \mathcal{F}^2)) \\ (f) \times (g_0, g_\infty) &\longmapsto (df - (g_0|_{U_0 \cap U_\infty} - g_\infty|_{U_0 \cap U_\infty})) \times (dg_0, dg_\infty) \end{aligned}$$

modulo the image of the map

$$\begin{aligned} \Gamma(U_\infty, \mathcal{F}^0) \oplus \Gamma(U_\infty, \mathcal{F}^0) &\longrightarrow \Gamma(U_0 \cap U_\infty, \mathcal{F}^0) \times (\Gamma(U_\infty, \mathcal{F}^1) \oplus \Gamma(U_\infty, \mathcal{F}^1)) \\ (g_0, g_\infty) &\longmapsto (g_0|_{U_0 \cap U_\infty} - g_\infty|_{U_0 \cap U_\infty}) \times (dg_0, dg_\infty) \end{aligned}$$

where $d : \mathcal{F}^i \rightarrow \mathcal{F}^{i+1}$ is the differential map. We compute

$$(4.27) \quad u \circ i_{F_{\text{red}}}^*(\widehat{\eta}) = i_{F_*}^*(u''(\widehat{\eta})) = i_{F_*}^* \circ u'' \left(\frac{1}{\sqrt{-3}} \widehat{\eta}_1 \right), \quad \widehat{\eta}_1 := \frac{x_1 dx_1}{y_1}$$

where the last equality follows from the fact that $u''(dx/y) = 0$. The cohomology class $\widehat{\eta}_1 \in H_{\text{zar}}^1(\omega_{\mathcal{E}/\Delta}^\bullet)$ is represented by a Čech cocycle

$$\left(\frac{y_1}{2x_1} \right) \times \left(\frac{x_1 dx_1}{y_1}, \frac{(g_2 v + 2g_3 v^2) dv}{4w} \right) \in \Gamma(U_0 \cap U_\infty, \mathcal{O}_{\mathcal{E}}) \times (\Gamma(U_0, \omega_{\mathcal{E}/\Delta}^1) \oplus \Gamma(U_\infty, \omega_{\mathcal{E}/\Delta}^1)).$$

Then $u''(\widehat{\eta})$ is represented by a Čech cocycle $(y_1/2x_1) = (y/(x+3)) \in \Gamma(U_0 \cap U_\infty, \mathcal{O}_{\mathcal{E}})$, and hence (??) is represented by a Čech cocycle

$$\begin{aligned} & \left(\frac{1}{\sqrt{-3}} \frac{y}{x+3} \Big|_Z, 0 \right) \times (0, 0) = \left(\frac{1}{\sqrt{-3}} \frac{z_0(z_0^2+3)}{z_0^2+2}, 0 \right) \times (0, 0) \\ & = \left(\frac{1}{\sqrt{-3}} z_0, 0 \right) \times (0, 0) - \left(\frac{1}{\sqrt{-3}} \frac{-z_0}{z_0^2+2}, 0 \right) \times (0, 0) \\ & \in \Gamma(U_0 \cap U_\infty, \mathcal{O}_Z \oplus \mathcal{O}_E) \times (\Gamma(U_0, \mathcal{O}_{Z \cap E}) \oplus \Gamma(U_\infty, \mathcal{O}_{Z \cap E})). \end{aligned}$$

Since $z_0 \in \Gamma(U_0, \mathcal{O}_Z)$ and $z_0/(z_0^2+2) \in \Gamma(U_\infty, \mathcal{O}_Z)$, this is equivalent to

$$\begin{aligned} & (0, 0) \times \left(\left(\frac{1}{\sqrt{-3}} \cdot \sqrt{-3}, \frac{1}{\sqrt{-3}} \cdot (-\sqrt{-3}) \right), \left(\frac{1}{\sqrt{-3}} \cdot \frac{-\sqrt{-3}}{-1}, \frac{1}{\sqrt{-3}} \cdot \frac{\sqrt{-3}}{-1} \right) \right) \\ & = (0, 0) \times ((1, -1), (1, -1)). \end{aligned}$$

This shows

$$(4.28) \quad \alpha \left((h \circ i_{F_{\text{red}}}^*)(\widehat{\eta}) \right) = 1 - (-1) = 2$$

giving the right hand side of (??).

We write down the left hand side of (??) explicitly. Put $D_F := D \cap F = D \cap Z$ (= three points $z = 0, \nu^2, \infty$). Let $F_U := F_{\text{red}} \cap U = (Z \setminus D_F) \cup E$ and its simplicial nerve is denoted by $F_{U,*}$. One has $h_i|_{F_U} \in \mathcal{O}(F_U)^\times$ and

$$\xi|_{F_U} = \{h_1|_{F_U}, h_2|_{F_U}\} \in K_2^M(\mathcal{O}(F_U)).$$

We think $h_i|_{F_U}$ to be an element of $K_1(F_{U,*})$ and $\xi|_{F_U}$ of being an element of $K_2(F_{U,*})$ via the canonical maps $K_i^M(\mathcal{O}(F_U)) \rightarrow K_i(F_{U,*})$. To compute the left hand side of (??), it is enough to compute

$$[\xi|_{F_U}]_{\text{syn}} = [h_1|_{F_U}]_{\text{syn}} \cup [h_2|_{F_U}]_{\text{syn}} \in H_{\text{syn}}^2((F_*, D_F), \mathbb{Z}_p(2))$$

as there is a commutative diagram

$$\begin{array}{ccccc} H_{\text{syn}}^2(F_*, \mathbb{Z}_p(2)) & \xleftarrow{\cong} & H_{\text{dR}}^1(F_*/W) & \xrightarrow[u']{\cong} & H_{\text{zar}}^1(F_*, \mathcal{O}_{F_*}) \\ \downarrow & & \downarrow & & \parallel \\ H_{\text{syn}}^2((F_*, D_F), \mathbb{Z}_p(2)) & \xleftarrow{\cong} & H_{\text{dR}}^1((F_*, D_F)/W) & \longrightarrow & H_{\text{zar}}^1(F_*, \mathcal{O}_{F_*}). \end{array}$$

We further replace the log syntomic cohomology with the rigid syntomic cohomology. Take an (arbitrary) affine open set $E' \subset E$ such that $E' \supset Z \cap E$. Put $Z' := Z \setminus D_F$ and $F'_U := Z' \cup E'$ and let $F'_{U,\star}$ be the simplicial nerve. Then there is a commutative diagram

$$\begin{array}{ccccc} H_{\text{syn}}^2((F_\star, D_F), \mathbb{Z}_p(2)) & \xleftarrow{\cong} & H_{\text{dR}}^1((F_\star, D_F)/W) & \longrightarrow & H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}) \\ \downarrow & & \downarrow & & \downarrow \\ H_{\text{rig-syn}}^2(F'_{U,\star}, \mathbb{Q}_p(2)) & \xleftarrow{\cong} & H_{\text{dR}}^1(F'_{U,\star}/K) & \longrightarrow & H_{\text{zar}}^1(F_\star, \mathcal{O}_{F_\star}) \otimes \mathbb{Q}. \end{array}$$

Thanks to the compatibility with the rigid regulator maps (Lemma ??), it is enough to compute

$$(4.29) \quad \text{reg}_{\text{rig-syn}}(h_1|_{F'_U}) \cup \text{reg}_{\text{rig-syn}}(h_2|_{F'_U}) \in H_{\text{rig-syn}}^2(F'_{U,\star}, \mathbb{Q}_p(2)).$$

We take a p -th Frobenius $\varphi_{Z'}$ on $\mathcal{O}(Z')^\dagger$ given by $\varphi_{Z'}(z) = z^p$. Note $\varphi : Z' \rightarrow Z'$ fixes R_1 and R_2 . We also take a p -th Frobenius $\varphi_{E'}$ on $\mathcal{O}(E')^\dagger$ that fixes R_1 and R_2 . Then for $(V, \bar{V}) = (Z', Z)$ or (E', E) , the rigid syntomic complex $S_{\text{rig-syn}}(r)_V := R\Gamma_{\text{rig-syn}}(V, \mathbb{Q}_p(r))$ is given as follows,

$$S_{\text{rig-syn}}(r)_V = \text{Cone}[L(\bar{V}_K, \Omega_{\bar{V}_K}^{\bullet \geq r}(\log \partial V_K)) \xrightarrow{1-p^{-r}\varphi_V} \Omega_{\mathcal{O}(V)_K^\dagger}^\bullet][-1]$$

where $\partial V = \bar{V} \setminus V$ and $\Omega_{\mathcal{O}(V)_K^\dagger}^1 = \Omega_{\mathcal{O}(V)_K^\dagger/K}^1$ denotes the module of continuous differentials. Moreover the rigid syntomic cohomology of $F'_{U,\star}$ is described as follows,

$$H_{\text{rig-syn}}^j(F'_{U,\star}, \mathbb{Q}_p(r)) \cong H^j(S_{\text{rig-syn}}(r)_{Z'} \oplus S_{\text{rig-syn}}(r)_{E'} \xrightarrow{i} S_{\text{rig-syn}}(r)_{Z \cap E}),$$

where $i : (f, g) \mapsto f|_{Z \cap E} - g|_{Z \cap E}$. Under this identification, it follows from [?, Prop.10.3] that we have cocycles

$$\begin{aligned} \text{reg}_{\text{rig-syn}}(h_1|_{F'_U}) &= \left(3 \frac{dz}{z}, 0 \right) \times (0, 0) \times (0, 0) \\ \text{reg}_{\text{rig-syn}}(h_2|_{F'_U}) &= \left(\frac{-3\nu dz}{1-\nu z}, 3p^{-1} \log \left(\frac{(1-\nu z)^p}{1-\nu^p z^p} \right) \right) \times (0, 0) \times (0, 0) \\ &\in (\Omega_{\mathcal{O}(Z')_K^\dagger}^1 \oplus \mathcal{O}(Z')_K^\dagger) \times (\Omega_{\mathcal{O}(E')_K^\dagger}^1 \oplus \mathcal{O}(E')_K^\dagger) \times (\mathcal{O}(R_1) \oplus \mathcal{O}(R_2)) \end{aligned}$$

for h_i in (??), and then

$$\begin{aligned} (??) &= \left(-9p^{-1} \log \left(\frac{(1-\nu z)^p}{1-\nu^p z^p} \right) \frac{dz}{z}, 0 \right) \times (0, 0) \times (0, 0) \\ &= \left(9d(\ln_2^{(p)}(\nu z)), 0 \right) \times (0, 0) \times (0, 0) \\ &\equiv (0, 0) \times (0, 0) \times (9 \ln_2^{(p)}(-\nu), 9 \ln_2^{(p)}(-\nu^2)) \end{aligned}$$

in $H_{\text{rig-syn}}^2(F'_{U,\star}, \mathbb{Q}_p(2))$. Here we use the fact that $\ln_2^{(p)}(z)$ is an overconvergent function on $\mathbb{P}^1 \setminus \{1, \infty\}$ (Proposition ??). This shows

$$\alpha((u' \circ i_{F_\star}^*)([\xi]_{\text{syn}})) = 9 \ln_2^{(p)}(-\nu) - 9 \ln_2^{(p)}(-\nu^2) = 18 \ln_2^{(p)}(-\nu),$$

which provides the left hand side of (??). Combining this with (??), one finally has

$$18 \ln_2^{(p)}(-\nu) = -2E_2(0).$$

This completes the proof. \square

Corollary 4.9. *Let $a \in W$ satisfy $a \not\equiv 0, 1 \pmod{p}$. Let $c = F_W(a)a^{-p}$ so that $\sigma(t) = ct^p$ satisfy $\sigma(t)|_{t=a} = F_W(a)$. Let E_a be the fiber of f at $t = a$, and put $U_a := E_a \cap U$. Let $\xi|_{E_a} \in K_2^M(\mathcal{O}(U_a))_{\partial=0}$ be the restriction of the Milnor symbol ξ in (??), and we think it to be an element of the Adams weight piece $K_2(E_a)^{(2)}$ under the natural map $K_2^M(\mathcal{O}(U_a))_{\partial=0} \rightarrow K_2(E_a)^{(2)}$. Let*

$$\text{reg}_{\text{rig-syn}} : K_2(E_a)^{(2)} \longrightarrow H_{\text{rig-syn}}^2(E_a, \mathbb{Q}_p(2)) \cong H_{\text{dR}}^1(E_a/K)$$

be the regulator map by Besser [?] or equivalently by Nekovář-Nizioł [?]. Then

$$\text{reg}_{\text{rig-syn}}(\xi|_{E_a}) = -\varepsilon_1(a) \frac{dx}{y} - \varepsilon_2(a) \frac{xdx}{y} \in H_{\text{dR}}^1(E_a/K).$$

Proof. Recall the diagram in Theorem ??, and take the pull-back of it at the point $t = a$ of S^* . Then $[-]_{\text{syn}}$ turns out to be the regulator map $\text{reg}_{\text{rig-syn}}$ (Lemma ??). Now the assertion is immediate from Theorem ??. \square

5. APPENDIX : FROBENIUS ON TOTALLY DEGENERATING ABELIAN SCHEMES

5.1. De Rham symplectic basis for totally degenerating abelian varieties. Let R be a regular noetherian domain, and I a reduced ideal of R . Let $L := \text{Frac}(R)$ be the fractional field. Let J/R be a g -dimensional commutative group scheme such that the generic fiber J_η is a principally polarized abelian variety over L . If the fiber T over $\text{Spec } R/I$ is an algebraic torus, we call J a *totally degenerating abelian scheme over (R, I)* (cf. [?] Chapter II, 4). Assume that the algebraic torus T is split. Assume further that R is complete with respect to I . Then there is the uniformization $\rho : \mathbb{G}_m^g \rightarrow J$ in the rigid analytic sense. We fix ρ and the coordinates (u_1, \dots, u_g) of \mathbb{G}_m^g . Then a matrix

$$(5.1) \quad \underline{q} = \begin{pmatrix} q_{11} & \cdots & q_{1g} \\ \vdots & & \vdots \\ q_{g1} & \cdots & q_{gg} \end{pmatrix}, \quad q_{ij} = q_{ji} \in L$$

of multiplicative periods is determined up to $\text{GL}_g(\mathbb{Z})$, and this yields an isomorphism

$$J \cong \mathbb{G}_m^g / \underline{q}^{\mathbb{Z}}$$

of abelian schemes over R where $\mathbb{G}_m^g / \underline{q}^{\mathbb{Z}}$ denotes Mumford's construction of the quotient scheme ([?] Chapter III, 4.4).

In what follows, we suppose that the characteristic of L is zero. The morphism ρ induces

$$\rho^* : \Omega_{J/R}^1 \longrightarrow \bigoplus_{i=1}^g \hat{\Omega}_{\mathbb{G}_m, i}^1, \quad \hat{\Omega}_{\mathbb{G}_m, i}^1 := \varprojlim_n \Omega_{R/I^n[u_i, u_i^{-1}]/R}^1.$$

Let

$$\text{Res}_i : \widehat{\Omega}_{\mathbb{G}_m, i}^1 \longrightarrow R, \quad \text{Res}_i \left(\sum_{m \in \mathbb{Z}} a_m u_i^m \frac{du_i}{u_i} \right) = a_0$$

be the residue map. The composition of ρ^* and the residue map induces a morphism $\Omega_{J/R}^\bullet \longrightarrow R^g[-1]$ of complexes, and hence a map

$$(5.2) \quad \tau : H_{\text{dR}}^1(J/R) := H_{\text{zar}}^1(J, \Omega_{J/R}^\bullet) \longrightarrow R^g.$$

Let U be defined by

$$0 \longrightarrow U \longrightarrow H_{\text{dR}}^1(J_\eta/L) \xrightarrow{\tau} L^g \longrightarrow 0.$$

Note that the composition $\Gamma(J_\eta, \Omega_{J_\eta/L}^1) \xhookrightarrow{\sim} H_{\text{dR}}^1(J_\eta/L) \xrightarrow{\tau} L^g$ is bijective. Let $\langle x, y \rangle$ denotes the symplectic pairing on $H_{\text{dR}}^1(J_\eta/L)$ with respect to the principal polarization on J_η . We call an L -basis

$$\widehat{\omega}_i, \widehat{\eta}_j \in H_{\text{dR}}^1(J_\eta/L), \quad 1 \leq i, j \leq g$$

a *de Rham symplectic basis* if the following conditions are satisfied.

(DS1): $\widehat{\omega}_i \in \Gamma(J_\eta, \Omega_{J_\eta/L}^1)$ and $\tau(\widehat{\omega}_i) \in (0, \dots, 1, \dots, 0)$ where “1” is placed in the i -th component. Equivalently,

$$\rho^*(\widehat{\omega}_i) = \frac{du_i}{u_i}.$$

(DS2): $\widehat{\eta}_j \in U$ and $\langle \widehat{\omega}_i, \widehat{\eta}_j \rangle = \delta_{ij}$ where δ_{ij} is the Kronecker delta.

If we fix the coordinates (u_1, \dots, u_g) of \mathbb{G}_m^g , then $\widehat{\omega}_i$ are uniquely determined by **(DS1)**. Since the symplectic pairing $\langle x, y \rangle$ is annihilated on $U \otimes U$ and $\Gamma(J_\eta, \Omega_{J_\eta/L}^1) \otimes \Gamma(J_\eta, \Omega_{J_\eta/L}^1)$, the basis $\widehat{\eta}_j$ are uniquely determined as well by **(DS2)**.

Proposition 5.1. *Let V be a subring of R . Suppose that (R, I) and V satisfy the following.*

(C): *There is a regular integral noetherian \mathbb{C} -algebra \widetilde{R} complete with respect to a reduced ideal \widetilde{I} and an injective homomorphism $i : R \rightarrow \widetilde{R}$ such that $i(V) \subset \mathbb{C}$ and $i(I) \subset \widetilde{I}$ and*

$$\widehat{\Omega}_{L/V}^1 \longrightarrow \widehat{\Omega}_{\widetilde{L}/\mathbb{C}}^1$$

is injective where we put $\widetilde{L} := \text{Frac}(\widetilde{R})$ and

$$\widehat{\Omega}_{L/V}^1 := L \otimes_R \left(\varprojlim_n \Omega_{R_n/V}^1 \right), \quad \widehat{\Omega}_{\widetilde{L}/\mathbb{C}}^1 := \widetilde{L} \otimes_{\widetilde{R}} \left(\varprojlim_n \Omega_{\widetilde{R}_n/\mathbb{C}}^1 \right), \quad R_n := R/I^n, \quad \widetilde{R}_n := \widetilde{R}/\widetilde{I}^n.$$

Let

$$\nabla : H_{\text{dR}}^1(J_\eta/L) \longrightarrow \widehat{\Omega}_{L/V}^1 \otimes_L H_{\text{dR}}^1(J_\eta/L)$$

be the Gauss-Manin connection. Then

$$(5.3) \quad \nabla(\widehat{\omega}_i) = \sum_{j=1}^g \frac{dq_{ij}}{q_{ij}} \otimes \widehat{\eta}_j, \quad \nabla(\widehat{\eta}_i) = 0.$$

Proof. By the assumption **(C)**, we may replace J_η/L with $(J_\eta \otimes_L \tilde{L})/\tilde{L}$. We may assume $R = \tilde{R}$, $V = \mathbb{C}$ and $J = J_\eta \otimes_R \tilde{R}$. There is a smooth scheme J_S/S over a connected smooth affine variety $S = \text{Spec } A$ over \mathbb{C} with a Cartesian square

$$\begin{array}{ccc} J & \longrightarrow & J_S \\ \downarrow & \square & \downarrow \\ \text{Spec } R & \longrightarrow & S \end{array}$$

such that $\text{Spec } R \rightarrow S$ is dominant. Let $D \subset S$ be a closed subset such that J_S is proper over $U := S \setminus D$. Then the image of $\text{Spec } R/I$ is contained in D since J has a totally degeneration over $\text{Spec } R/I$. Thus we may replace R with the completion \hat{A}_D of A by the ideal of D . Let $L_D = \text{Frac } \hat{A}_D$ then $\hat{\Omega}_{L_D/\mathbb{C}}^1 \cong L_D \otimes_A \Omega_{A/\mathbb{C}}^1$. Let $\mathfrak{m} \subset A$ be a maximal ideal containing I , and $\hat{A}_\mathfrak{m}$ the completion by \mathfrak{m} . Then $\hat{A}_D \subset \hat{A}_\mathfrak{m}$ and $\hat{\Omega}_{\hat{A}_D/\mathbb{C}}^1 \subset \hat{\Omega}_{\hat{A}_\mathfrak{m}/\mathbb{C}}^1$. Therefore we may further replace \hat{A}_D with $\hat{A}_\mathfrak{m}$. Summing up the above, it is enough work in the following situation.

$$R = \hat{A}_\mathfrak{m} \cong \mathbb{C}[[x_1, \dots, x_n]] \supset I = (x_1, \dots, x_n), \quad V = \mathbb{C}, \quad L = \text{Frac } R$$

$$J = \text{Spec } R \times_A J_S \longrightarrow \text{Spec } R.$$

Note $\hat{\Omega}_{L/\mathbb{C}}^1 \cong L \otimes_A \Omega_{A/\mathbb{C}}^1$. Let $h : \mathcal{J} \rightarrow S^{\text{an}}$ be the analytic fibration associated to J_S/S . Write $J_\lambda = h^{-1}(\lambda)$ a smooth fiber over $\lambda \in U^{\text{an}} := (S \setminus D)^{\text{an}}$. Let

$$\nabla : \mathcal{O}_{U^{\text{an}}} \otimes R^1 h_* \mathbb{C} \longrightarrow \Omega_{U^{\text{an}}}^1 \otimes R^1 h_* \mathbb{C}.$$

be the flat connection compatible with the Gauss-Manin connection on $H_{\text{dR}}^1(J_S/S)|_U$ under the comparison

$$\mathcal{O}_{U^{\text{an}}} \otimes R^1 h_* \mathbb{C} \cong \mathcal{O}_{U^{\text{an}}} \otimes_A H_{\text{dR}}^1(J_S/S).$$

We describe a de Rham symplectic basis in $\hat{\omega}_i^{\text{an}}, \hat{\eta}_j^{\text{an}} \in \mathcal{O}_{U^{\text{an}}} \otimes_A H_{\text{dR}}^1(J_S/S)$ and prove (??) for the above flat connection. Write $J_\lambda = (\mathbb{C}^\times)^g/q^{\mathbb{Z}}$ for $\lambda \in U^{\text{an}}$. Let (u_1, \dots, u_g) denotes the coordinates of $(\mathbb{C}^\times)^g$. Let $\delta_i \in H_1(J_\lambda, \mathbb{Z})$ be the homology cycle defined by the circle $|u_i| = \varepsilon$ with $0 < \varepsilon \ll 1$. Let $\gamma_j \in H_1(J_\lambda, \mathbb{Z})$ be the homology cycle defined by the path from $(1, \dots, 1)$ to (q_{j1}, \dots, q_{jg}) . As is well-known, the dual basis $\delta_i^*, \gamma_j^* \in H^1(J_\lambda, \mathbb{Z})$ is a symplectic basis, namely

$$\langle \delta_i^*, \delta_{i'}^* \rangle = \langle \gamma_j^*, \gamma_{j'}^* \rangle = 0, \quad \langle \delta_i^*, \gamma_j^* \rangle = \frac{1}{2\pi\sqrt{-1}} \delta_{ij}$$

where δ_{ij} denotes the Kronecker delta. We have $\hat{\omega}_i^{\text{an}} = du_i/u_i$ by **(DS1)**, and then

$$(5.4) \quad \hat{\omega}_i^{\text{an}} = 2\pi\sqrt{-1}\delta_i^* + \sum_{j=1}^g \log q_{ij} \gamma_j^*.$$

Let $\tau^B : R^1 h_* \mathbb{Z} \rightarrow \mathbb{Z}(-1)^g$ be the associated map to τ . An alternative description of τ^B is

$$\tau^B(x) = \frac{1}{2\pi\sqrt{-1}}((x, \delta_1), \dots, (x, \delta_g))$$

where (x, δ) denotes the natural pairing on $H^1(J_\lambda, \mathbb{Z}) \otimes H_1(J_\lambda, \mathbb{Z})$. Obviously $\tau^B(\gamma_j^*) = 0$. This implies that $\hat{\eta}_j$ is a linear combination of $\gamma_1^*, \dots, \gamma_g^*$ by **(DS2)**. Since $\langle \hat{\omega}_i^{\text{an}}, \hat{\eta}_j \rangle = \delta_{ij} = \langle \hat{\omega}_i^{\text{an}}, \gamma_j^* \rangle$, one concludes

$$\hat{\eta}_j^{\text{an}} = \gamma_j^*.$$

Now **(??)** is immediate from this and **(??)**.

Let $\hat{\omega}_i, \hat{\eta}_j \in H_{\text{dR}}^1(J_\eta/L)$ be the de Rham symplectic basis. Let $x \in D^{\text{an}}$ be the point associated to \mathfrak{m} . Let V^{an} be a small neighborhood of x and $j : V^{\text{an}} \setminus D^{\text{an}} \hookrightarrow S^{\text{an}}$ an open immersion. Obviously $\hat{\omega}_i^{\text{an}} = du_i/u_i \in \Gamma(V^{\text{an}}, j_* \mathcal{O}^{\text{an}}) \otimes_A H_{\text{dR}}^1(J_S/S)$. Thanks to the uniqueness property, this implies $\hat{\eta}_j^{\text{an}} = \gamma_j^* \in \Gamma(V^{\text{an}}, j_* \mathcal{O}^{\text{an}}) \otimes_A H_{\text{dR}}^1(J_S/S)$, in other words, $\gamma_j^* \in \Gamma(V^{\text{an}} \setminus D^{\text{an}}, j^{-1} R^1 h_* \mathbb{Q})$. Let $\hat{S}_{\mathfrak{m}}$ be the ring of power series over \mathbb{C} containing $\hat{A}_{\mathfrak{m}}$ and $\Gamma(V^{\text{an}}, j_* \mathcal{O}^{\text{an}})$. There is a commutative diagram

$$\begin{array}{ccc} \hat{A}_{\mathfrak{m}} \otimes_A \Omega_{A/\mathbb{C}}^1 \otimes_A H_{\text{dR}}^1(J_S/S) & \longrightarrow & \hat{S}_{\mathfrak{m}} \otimes_A \Omega_{A/\mathbb{C}}^1 \otimes_A H_{\text{dR}}^1(J_S/S) \\ \uparrow & & \uparrow \\ \Omega_{A/\mathbb{C}}^1 \otimes_A H_{\text{dR}}^1(J_S/S) & \longrightarrow & \Gamma(V^{\text{an}}, j_* \mathcal{O}^{\text{an}}) \otimes_A \Omega_{A/\mathbb{C}}^1 \otimes_A H_{\text{dR}}^1(J_S/S) \end{array}$$

with all arrows injective. Hence the desired assertion for $\hat{\omega}_i, \hat{\eta}_j$ can be reduced to that of $\hat{\omega}_i^{\text{an}}, \hat{\eta}_j^{\text{an}}$. This completes the proof. \square

5.2. Frobenius on De Rham symplectic basis. Let V be a complete discrete valuation ring such that the residue field k is perfect and of characteristic p , and the fractional field $K := \text{Frac} V$ is of characteristic zero. Let F_V be a p -th Frobenius endomorphism on V .

Let A be an integral flat noetherian V -algebra equipped with a p -th Frobenius endomorphism σ on A which is compatible with F_V . Assume that A is p -adically complete and separated and that there is a family $(t_i)_{i \in I}$ of finitely many elements of A such that it forms a p -basis of $A_n := A/p^n A$ over $V_n := V/p^n V$ for all $n \geq 1$ in the sense of [?, Definition 1.3]. The latter assumption is equivalent to that $(t_i)_{i \in I}$ forms a p -basis of A_1 over V_1 since A is flat over V (loc.cit. Lemma 1.6). Then Ω_{A_n/V_n}^1 is a free A_n -module with basis $(dt_i)_{i \in I}$ (loc.cit. Lemma 1.8). Write $A_K := A \otimes_V K$.

Definition 5.2. We define the category $F\text{-MIC}(A_K, \sigma)$ as follows. An object is a triplet (M, ∇, Φ) where

- M is a locally free A_K -module of finite rank,
- $\nabla : M \rightarrow \hat{\Omega}_{A/V}^1 \otimes_A M$ is an integrable connection where $\hat{\Omega}_{A/V}^1 := \varprojlim_n \Omega_{A_n/V_n}^1$ a free A -module with basis $\{dt_i\}_{i \in I}$,
- $\Phi : \sigma^* M \rightarrow M$ is a horizontal A_K -linear map.

A morphism in $F\text{-MIC}(A_K, \sigma)$ is an A -linear map of the underlying A -modules which is commutative with ∇ and Φ . Let $L := \text{Frac}(A)$ be the fractional field. The category $F\text{-MIC}(L, \sigma)$ is defined in the same way by replacing A with L , and $\hat{\Omega}_{A/V}^1$ with $L \otimes_A \hat{\Omega}_{A/V}^1$.

Lemma 5.3 ([?, 6.1]). *Let σ' be another F_V -linear p -th Frobenius on A . Then there is the natural equivalence*

$$F\text{-MIC}(A, \sigma) \xrightarrow{\cong} F\text{-MIC}(A, \sigma'), \quad (M, \nabla, \Phi) \mapsto (M, \nabla, \Phi')$$

of categories, where Φ' is defined in the following way. Let $(\partial_i)_{i \in I}$ be the basis of $\widehat{T}_{A/V} := \varprojlim_n (\Omega_{A_n/V_n}^1)^*$ which is the dual basis of $(dt_i)_{i \in I}$. Then

$$\Phi' = \sum_{n_i \geq 0} \left(\prod_{i \in I} (\sigma'(t_i) - \sigma(t_i))^{n_i} \right) \Phi \prod_{i \in I} \partial_i^{n_i}.$$

Let $f : X \rightarrow \operatorname{Spec} A$ be a projective smooth morphism. Write $X_n := X \times_V V/p^{n+1}V$. Then one has an object

$$H^i(X/A) := (H_{\mathrm{dR}}^i(X/A) \otimes_V K, \nabla, \Phi) \in F\text{-MIC}(A_K)$$

where Φ is induced from the Frobenius on crystalline cohomology via the comparison ([?] 7.4)

$$H_{\mathrm{crys}}^\bullet(X_0/A) \cong \varprojlim_n H_{\mathrm{dR}}^\bullet(X_n/A_n) \cong H_{\mathrm{dR}}^\bullet(X/A).$$

Assume that there is a smooth V -algebra A^a and a smooth projective morphism $f^a : X^a \rightarrow \operatorname{Spec} A^a$ with a Cartesian diagram

$$\begin{array}{ccc} X & \longrightarrow & \operatorname{Spec} A \\ \downarrow & \square & \downarrow \\ X^a & \longrightarrow & \operatorname{Spec} A^a \end{array}$$

such that $\operatorname{Spec} A \rightarrow \operatorname{Spec} A^a$ is flat. One has the overconvergent F -isocrystal $R^i f_{\mathrm{rig},*}^a \mathcal{O}_{X^a}$ on $\operatorname{Spec} A_0^a$ ([?] 3.4.8.2). Let

$$H^i(X^a/A^a) := (H_{\mathrm{rig}}^i(X_0^a/A_0^a), \nabla, \Phi) \in F\text{-MIC}^\dagger(A_K^a)$$

denote the associated object via the natural equivalence $F\text{-Isoc}^\dagger(A_0^a) \cong F\text{-MIC}^\dagger(A_K^a)$ ([?] 8.3.10). The comparison $H_{\mathrm{rig}}^i(X_0^a/A_0^a) \cong H_{\mathrm{dR}}^i(X^a/A^a) \otimes_{A^a} (A^a)_K^\dagger$ in (??) induces an isomorphism $H^i(X/A) \cong A_K \otimes_{(A^a)_K^\dagger} H^i(X^a/A^a)$ in $F\text{-MIC}(A_K)$.

Definition 5.4 (Tate objects). For an integer r , a Tate object $A_K(r)$ is defined to be the triplet (A_K, ∇, Φ) such that $\nabla = d$ is the usual differential operator, and Φ is a multiplication by p^{-r} .

We define for $f \in A \setminus \{0\}$

$$\log^{(\sigma)}(f) := p^{-1} \log \left(\frac{f^p}{f^\sigma} \right) = - \sum_{n=1}^{\infty} \frac{p^{n-1} g^n}{n}, \quad \frac{f^p}{f^\sigma} = 1 - pg$$

which belongs to the p -adic completion of the subring $A[g] \subset L$. In particular, if $f \in A^\times$, then $\log^{(\sigma)}(f) \in A$.

Definition 5.5 (Log objects). Let $\underline{q} = (q_{ij})$ be a $g \times g$ -symmetric matrix with $q_{ij} \in A^\times$. We define a log object $\mathcal{L}og(\underline{q}) = (M, \nabla, \Phi)$ in $F\text{-MIC}(A, \sigma)$ to be the following. Let

$$M = \bigoplus_{i=1}^g A_K e_i \oplus \bigoplus_{i=1}^g A_K f_i$$

be a free A_K -module with a basis e_i, f_j . The connection is defined by

$$\nabla(e_i) = \sum_{j=1}^g \frac{dq_{ij}}{q_{ij}} \otimes f_j, \quad \nabla(f_j) = 0$$

and the Frobenius Φ is defined by

$$\Phi(e_i) = e_i - \sum_{j=1}^g \log^{(\sigma)}(q_{ij}) f_j, \quad \Phi(f_j) = p^{-1} f_j.$$

It is immediate to check that the log objects are compatible under the natural equivalence in Lemma ???. In this sense, our $\mathcal{L}og(\underline{q})$ does not depend on σ . By definition there is an exact sequence

$$0 \longrightarrow \bigoplus_{j=1}^g A_K(1) f_j \longrightarrow \mathcal{L}og(\underline{q}) \longrightarrow \bigoplus_{i=1}^g A_K(0) e_i \longrightarrow 0.$$

Theorem 5.6. *Let R be a flat V -algebra which is a regular noetherian domain complete with respect to a reduced ideal I . Suppose that R has a p -th Frobenius σ . Let J be a totally degenerating abelian scheme with a principal polarization over (R, I) in the sense of §??. Let $\text{Spec } R[h^{-1}] \hookrightarrow \text{Spec } R$ be an affine open set such that J is proper over $\text{Spec } R[h^{-1}]$ and $q_{ij} \in R[h^{-1}]^\times$ where $\underline{q} = (q_{ij})$ is the multiplicative periods as in (??). Suppose that $R/pR[h^{-1}]$ has a p -basis over V/pV . Let $A = R[h^{-1}]^\wedge$ be the p -adic completion of $R[h^{-1}]$. Put $L := \text{Frac}(A)$ and $J_A := J \otimes_R A$. Let J_η be the generic fiber of J_A . Then there is an isomorphism*

$$(5.5) \quad (H_{\text{dR}}^1(J_\eta/L), \nabla, \Phi) \otimes_{A_K} A_K(1) \cong \mathcal{L}og(\underline{q}) \in F\text{-MIC}(L)$$

which sends the de Rham symplectic basis $\widehat{\omega}_i, \widehat{\eta}_j \in H_{\text{dR}}^1(J_\eta/L)$ to e_i, f_j respectively.

Proof. Let q_{ij} be indeterminates with $q_{ij} = q_{ji}$, and t_1, \dots, t_r ($r = g(g+1)/2$) are products $\prod q_{ij}^{n_{ij}}$ such that $\sum n_{ij} x_i x_j$ is positive semi-definite and they give a \mathbb{Z} -basis of the group of the symmetric pairings. Let $J_q = \mathbb{G}_m^g / q^{\mathbb{Z}}$ be Mumford's construction of the quotient group scheme over a ring $\mathbb{Z}_p[[t_1, \dots, t_r]]$ ([?] Chapter III, 4.4). Then there is a Cartesian square

$$\begin{array}{ccc} J & \xrightarrow{\quad} & J_q \\ \downarrow & \square & \downarrow \\ \text{Spec } R & \longrightarrow & \text{Spec } \mathbb{Z}_p[[t_1, \dots, t_r]] \end{array}$$

such that the bottom arrow sends t_i to an element of I by the functoriality of Mumford's construction ([?] Chapter III, 5.5). Thus we may reduce the assertion to the case of $J = J_q$, $R = \mathbb{Z}_p[[t_1, \dots, t_r]]$, $I = (t_1, \dots, t_r)$ and $h = \prod q_{ij}$. Since $\mathcal{L}og(\underline{q})$ and $H_{\text{dR}}^1(J_\eta/L)$ are compatible under the natural equivalence in Lemma ??, we may replace the Frobenius σ on R with a suitable one. Thus we may assume that it is given as $\sigma(q_{ij}) = q_{ij}^p$ and $\sigma(a) = a$ for $a \in \mathbb{Z}_p$. Under this assumption $\log^{(\sigma)}(q_{ij}) = 0$ by definition. Therefore our goal is to show

$$(5.6) \quad \nabla(\widehat{\omega}_i) = \sum_{j=1}^g \frac{dq_{ij}}{q_{ij}} \otimes \widehat{\eta}_j, \quad \nabla(\widehat{\eta}_j) = 0,$$

$$(5.7) \quad \Phi(\widehat{\omega}_i) = p\widehat{\omega}_i, \quad \Phi(\widehat{\eta}_j) = \widehat{\eta}_j.$$

Since the condition (C) in Proposition ?? is satisfied, (??) is nothing other than (??). We show (??). Let $\underline{q}^{(p)} := (q_{ij}^p)$ and $J_{q^p} := \mathbb{G}_m^g / (\underline{q}^{(p)})^{\mathbb{Z}}$. Then there is the natural morphism $\sigma_J : J_{q^p} \rightarrow J_q$ such that the following diagram is commutative.

$$\begin{array}{ccc} \mathbb{G}_m^g & \xlongequal{\quad} & \mathbb{G}_m^g \\ \downarrow & & \downarrow \\ J_{q^p} & \xrightarrow{\sigma_J} & J_q \\ \downarrow & & \downarrow \\ \text{Spec } R & \xrightarrow{\sigma} & \text{Spec } R \end{array}$$

Let $[p] : J_{q^p} \rightarrow J_{q^p}$ denotes the multiplication by p with respect to the commutative group scheme structure of J_q . It factors through the canonical surjective morphism $J_{q^p} \rightarrow J_q$ so that we have $[p]' : J_q \rightarrow J_{q^p}$. Define $\varphi := \sigma_J \circ [p]'$. Under the uniformization $\rho : \mathbb{G}_m^g \rightarrow J_q$, this is compatible with a morphism $\mathbb{G}_m^g \rightarrow \mathbb{G}_m^g$ given by $u_i \rightarrow u_i^p$ and $a \rightarrow \sigma(a)$ for $a \in R$, which we also write Φ . Therefore

$$\Phi = \varphi^* : H_{\text{dR}}^1(J_\eta/L) \longrightarrow H_{\text{dR}}^1(J_\eta/L).$$

In particular Φ preserves the Hodge filtration, so that $\Phi(\widehat{\omega}_i)$ is again a linear combination of $\widehat{\omega}_i$'s. Since

$$\rho^* \Phi(\widehat{\omega}_i) = \Phi \rho^*(\widehat{\omega}_i) = \Phi \left(\frac{du_i}{u_i} \right) = p \frac{du_i}{u_i},$$

one concludes $\Phi(\widehat{\omega}_i) = p\widehat{\omega}_i$. On the other hand, since $\Phi(\ker \nabla) \subset \ker \nabla$ and $\ker \nabla$ is generated by $\widehat{\eta}_j$'s by (??), $\Phi(\widehat{\eta}_j)$ is again a linear combination of $\widehat{\eta}_i$'s. Note

$$\langle \Phi(x), \Phi(y) \rangle = p \langle x, y \rangle.$$

Therefore

$$\langle \Phi(\widehat{\eta}_j), p\widehat{\omega}_j \rangle = \langle \Phi(\widehat{\eta}_j), \Phi(\widehat{\omega}_j) \rangle = p \langle \widehat{\eta}_j, \widehat{\omega}_j \rangle.$$

This implies $\Phi(\widehat{\eta}_j) = \widehat{\eta}_j$, so we are done. \square

REFERENCES

- [A-Bal] André, Y., Baldassarri, F., *De Rham cohomology of differential modules over algebraic varieties*. Progress in Mathematics **189**, Birkhäuser, 2000.
- [A] Asakura, M., *New p -adic hypergeometric functions and syntomic regulators*. arXiv.1811.03770.
- [A-C] Asakura, M., Chida, M., *A numerical approach toward the p -adic Beilinson conjecture for elliptic curves over \mathbb{Q}* . Res. in Math. Sci. **10** (2023) Article 11.
- [A-M] Atiyah, M., Macdonald, I., *Introduction to Commutative Algebra*. Addison-Wesley Series in Math. Westview Press, Boulder, CO, 2016.
- [Bal-Ch] Baldassarri F., Chiarellotto B., *Algebraic versus rigid cohomology with logarithmic coefficients*. In *Barsotti Symposium in Algebraic Geometry*, Perspectives in Math. **15** (1994), 11–50.
- [Ban1] Bannai, K., *Rigid syntomic cohomology and p -adic polylogarithms*. J. reine angew. Math., **529** (2000), 205–237.
- [Ban2] ———, “On the p -adic realization of elliptic polylogarithms for CM-elliptic curves”, Duke Math. Journ. **113**, No. 2 (2002), 193–236.

- [BBDG] Beilinson A., Bernstein B., Deligne P., *Faisceaux pervers*. In: Analyse et topologie sur les espaces singuliers I, Astérisque **100** (1982)
- [Ber1] Berthelot, P., *Géométrie rigide et cohomologie des variétés algébriques de caractéristique p* . Groupe de travail d'analyse ultramétrique, **9**, No. 3 (1981).
- [Ber2] ———, *Finitude et pureté cohomologique en cohomologie rigide avec un appendice par Aise Johan de Jong*. Invent. Math. **128** (1997), 329–377.
- [Ber3] ———, *Cohomologie rigide et cohomologie rigide à supports propres, Première partie.*, preprint available at <https://perso.univ-rennes1.fr/pierre.berthelot/>.
- [B-O] Berthelot, P., Ogus, A., *Notes on crystalline cohomology*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1978. vi+243
- [Be1] Besser, A., *Syntomic regulators and p -adic integration. I. Rigid syntomic regulators*. Proceedings of the Conference on p -adic Aspects of the Theory of Automorphic Representations (Jerusalem, 1998). Israel J. Math. **120** (2000), part B, 291–334.
- [Be2] ———, *Syntomic regulators and p -adic integration. II. K_2 of curves*. Proceedings of the Conference on p -adic Aspects of the Theory of Automorphic Representations (Jerusalem, 1998). Israel J. Math. **120** (2000), part B, 335–359.
- [B-K] Bloch, S., Kato, K., *L -functions and Tamagawa numbers of motives*. In: Cartier, P., Illusie, L., Katz, N. M., Laumon, G., Manin, Yu. I., Ribet, K. A. (eds.) *The Grothendieck Festschrift I*, (Progr. Math. **86**), pp. 333–400, Boston, Birkhäuser, 1990
- [Co] Colmez, P., *Fonctions L p -adiques*. Séminaire Bourbaki, Vol. 1998/99, Astérisque No. 266 (2000), Exp. No. 851, 3, 21–58.
- [Dw] Dwork, B., *p -adic cycles*. Publ. Math. IHES, tome 37 (1969), 27–115.
- [E-K] Emerton, M., Kisin, M., *An introduction to the Riemann-Hilbert correspondence for unit F -crystals*. Geometric aspects of Dwork theory. Vol. I, II, 677–700, Walter de Gruyter, Berlin, 2004.
- [Et] Étesse, J.-Y., *Images directes I: Espaces rigides analytiques et images directes*. J. Théor. Nombres Bordeaux **24** (1) (2012) 101–151.
- [F-C] Faltings, G., Chai, C-L., *Degeneration of Abelian Varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 22. Springer-Verlag, Berlin, 1990.
- [F-M] Fontaine, J.-M., Messing, W.: *p -adic periods and p -adic étale cohomology*. In: Ribet, K. A. (ed.) *Current Trends in Arithmetical Algebraic Geometry*, (Contemp. Math. 67), pp. 179–207, Providence, Amer. Math. Soc., 1987.
- [G] Gerkmann, R., *Relative rigid cohomology and point counting on families of elliptic curves*, J. Ramanujan Math. Soc. **23**, No. 1 (2008), 1–31.
- [Ka1] Kato, K., *The explicit reciprocity law and the cohomology of Fontaine-Messing*. Bull. Soc. Math. France **119** (1991), no. 4, 397–441.
- [Ka2] ———, *On p -adic vanishing cycles (application of ideas of Fontaine-Messing)*. In: Algebraic geometry, Sendai, 1985 (Adv. Stud. Pure Math. 10), pp. 207–251, Amsterdam, North-Holland, 1987.
- [Ka3] ———, *Logarithmic structures of Fontaine-Illusie*. Algebraic analysis, geometry, and number theory. Johns Hopkins University Press, Baltimore 1989, 191–224.
- [P] Perrin-Riou, B., *Fonctions L p -adiques des représentations p -adiques*. Astérisque **229** (1995).
- [Lau] Lauder, A., *Rigid cohomology and p -adic point counting*. J. Théor. Nombres Bordeaux **17** (2005), no. 1, 169–180.
- [N-N] Nekovář, J., Nizioł, W., *Syntomic cohomology and p -adic regulators for varieties over p -adic fields*. With appendices by Laurent Berger and Frédéric Déglise. Algebra Number Theory **10** (2016), no. 8, 1695–1790.
- [Sa1] Saito, M., *Modules de Hodge polarisables*. Publ. Res. Inst. Math. Sci. **24** (1988) no. 6, 849–995.
- [Sa2] ———, *Mixed Hodge modules*. Publ. Res. Inst. Math. Sci. **26** (1990), no. 2, 221–333.
- [Sas] Sasai, T., *Monodromy Representations of Homology of Certain Elliptic Surfaces*, J. Math. Soc. Japan **26** (1974), no. 2, 296–305.
- [Sh1] Shiho, A., *Crystalline Fundamental Groups II – Log Convergent Cohomology and Rigid Cohomology*. J. Math. Sci. Univ. Tokyo Vol. 9 (2002), No. 1, Page 1–163.
- [Sh2] ———, *Relative log convergent cohomology and relative rigid cohomology I*, arXiv:0707.1742.

- [Sh3] ———, *Relative log convergent cohomology and relative rigid cohomology III*, arXiv:0805.3229.
- [Si] Silverman, J., *Advanced topics in the arithmetic of elliptic curves*. Graduate Texts in Mathematics, **151**. Springer-Verlag, New York, 1994.
- [So] Solomon, N., *p -adic elliptic polylogarithms and arithmetic applications*, thesis, 2008.
- [LS] Le Stum, B., *Rigid cohomology*. Cambridge Tracts in Mathematics, 172. Cambridge University Press, Cambridge, 2007. xvi+319 pp.
- [Ts1] Tsuji, T., *p -adic étale cohomology and crystalline cohomology in the semi-stable reduction case*. Invent. Math. **137**, 233–411 (1999)
- [Ts2] ———, *On p -adic nearby cycles of log smooth families*. Bull. Soc. Math. France. **128** (2000), 529–575.
- [Tz1] Tsuzuki, N., “On the Gysin isomorphism of rigid cohomology.” Hiroshima Math J. 29(3), 479–527, 1999.
- [Tz2] ———, *On base change theorem and coherence in rigid cohomology*, Doc. Math. Extra, 2003, 891–918.
- [NIST] *NIST Handbook of Mathematical Functions*. Edited by Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert and Charles W. Clark. Cambridge Univ. Press, 2010.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810 JAPAN

Email address: asakura@math.sci.hokudai.ac.jp

SCHOOL OF SCIENCE AND TECHNOLOGY FOR FUTURE LIFE, TOKYO DENKI UNIVERSITY, TOKYO, 120-8551 JAPAN

Email address: miyatani@mail.dendai.ac.jp