

HOKKAIDO UNIVERSITY

Title	A few topics on total variation flows
Author(s)	Giga, Yoshikazu; Kuroda, Hirotoshi; Lasica, Michal
Citation	Hokkaido University Preprint Series in Mathematics, 1156, 1-43
Issue Date	2024-03-14
DOI	10.14943/109270
Doc URL	http://hdl.handle.net/2115/91283
Туре	bulletin (article)
File Information	A few topics on total variation flows_202403.pdf



A few topics on total variation flows

Yoshikazu Giga^{*}, Hirotoshi Kuroda[†], Michał Łasica[‡]

Abstract

Total variation gradient flows are important in several applied fields, including image analysis and materials science. In this paper, we review a few basic topics including definition of a solution, explicit examples and the notion of calibrability, finite time extinction, and some regularity properties of solutions. We focus on the second-order flow (possibly with weights) and the fourth-order flow. We also discuss the fractional cases.

1 Introduction

The (essential) total variation of a function u defined in a domain Ω in \mathbb{R}^n is formally of the form

$$TV(u) = \int_{\Omega} |\nabla u| \, dx.$$

Although it is a special case of p-Dirichlet energy

$$E_p(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p \, dx$$

with p = 1, there are several different features compared with 1 . This is easily observed $when one considers its variation under <math>L^2$ -metric, i.e., the L^2 -gradient. If the energy is given as

$$\mathcal{E}(u) = \int_{\Omega} f(\nabla u) \, dx,$$

its L^2 -gradient $\operatorname{grad}_{L^2} \mathcal{E}$ (the Euler–Lagrange operator) formally satisfies

$$\frac{d}{d\varepsilon} \int_{\Omega} f\left(\nabla(u+\varepsilon\varphi)\right) dx \bigg|_{\varepsilon=0} = \int_{\Omega} \left(\operatorname{grad}_{L^2} \mathcal{E}(u)\right) \varphi \, dx \quad \text{for all smooth functions } \varphi.$$

Neglecting the effect of the boundary, we see, by integration by parts,

$$\operatorname{grad}_{L^2} \mathcal{E}(u) = -\operatorname{div}\left((\nabla_q f)(\nabla u)\right)$$

^{*}Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan, labgiga@ms.u-tokyo.ac.jp

[†]Department of Mathematics, Faculty of Science, Hokkaido University, Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810, Japan, kuro@math.sci.hokudai.ac.jp

[‡]Institute of Mathematics of the Polish Academy of Sciences, ul. Śniadeckich 8, 00-656 Warszawa, Poland, mlasica@impan.pl

since

$$\frac{d}{d\varepsilon} \int_{\Omega} f\left(\nabla(u+\varepsilon\varphi)\right) dx \bigg|_{\varepsilon=0} = \sum_{i=1}^{n} \int_{\Omega} \frac{\partial f}{\partial q_{i}} (\nabla u) \frac{\partial \varphi}{\partial x_{i}}(x) dx$$
$$= -\int_{\Omega} \operatorname{div}\left((\nabla_{q} f)(\nabla u)\right) \varphi dx$$

where $\nabla_q f = (\partial f / \partial q_1, \dots, \partial f / \partial q_n)$. For the *p*-Dirichlet energy, i.e., $f(q) = |q|^p / p$

$$\operatorname{grad}_{L^2} E_p(u) = -\operatorname{div}\left(|\nabla u|^{p-2}\nabla u\right)$$

so that

$$\operatorname{grad}_{L^2} TV(u) = -\operatorname{div} \left(\nabla u / |\nabla u| \right).$$

These operators are (degenerate) elliptic but their features depend on p. We write

$$\operatorname{div}\left(|\nabla u|^{p-2}\nabla u\right) = \sum_{1 \le i,j \le n} a_{ij}(\nabla u) \frac{\partial^2}{\partial x_i \partial x_j} u.$$

Then $a_{ij}(q) \to 0$ as $q \to 0$ if p > 2 and all eigenvalues of $(a_{ij}(q))$ tend to infinity as $q \to 0$ for $1 . In other words, <math>\operatorname{grad}_{L^2} E_p$ is degenerate elliptic when p > 2 and singular for 1 . (In the case <math>p = 2, it is the Laplacian.) Nevertheless, the operator is elliptic in the region where $q \neq 0$. In the case p = 1,

$$a_{ij}(q) = \frac{1}{|q|} \left(\delta_{ij} - q_i q_j / |q|^2 \right).$$

The metric $A = (a_{ij}(q))$ is degenerate in direction q for all $q \in \mathbb{R}^n$, i.e.,

$$\sum_{j=1}^{n} a_{ij}(q)q_j = 0 \quad \text{or} \quad Aq = 0$$

because |q|A is the orthogonal projection onto the hyperplane orthogonal to q. However, in the direction orthogonal to q, A is singular at q = 0. Differently from the case p > 1, $\operatorname{grad}_{L^2} TV$ has both singular and degenerate effects. From the point of energy density, TV is significantly different from E_p for p > 1. First, the energy density $f(q) = |q|^p/p$ is non-differentiable at q = 0 for p = 1 while for p > 1 it is differentiable. This says that the singularity of A at q = 0 is very strong compared with the case p > 1. In fact, due to this singularity, the operator has nonlocal nature, as we see later. Second, the growth of f(q) as $|q| \to \infty$ is just linear while $f(q)/|q| \to \infty$ for p > 1. This means that minimizers of TV(u) under suitable supplemental condition may have jump discontinuities along some hypersurface, which does not occur for p > 1. Thus, problems involving TV should be handled separately from problems involving E_p for p > 1.

In this paper, we intend to give a theoretical introduction to total variation flows, that is, gradient flows of TV. Typically we consider the second-order problem

$$u_t + \operatorname{grad}_{L^2} TV(u) = 0 \quad \text{i.e.} \quad u_t - \operatorname{div} \left(\nabla u / |\nabla u| \right) = 0 \tag{1.1}$$

or the fourth-order problem

$$u_t + \Delta \operatorname{div} \left(\nabla u / |\nabla u| \right) = 0 \tag{1.2}$$

which is regarded as an H^{-1} -gradient flow, where H^s denotes the Sobolev space of order $s \in \mathbb{R}$. We also consider the fractional case

$$u_t - (-\Delta)^s \operatorname{div} \left(\nabla u / |\nabla u| \right) = 0. \tag{1.3}$$

As alluded before, because of the singularity at q = 0 for the energy density f(q) = |q|, div $(\nabla u/|\nabla u|)$ has nonlocal nature in the sense that its value at x cannot be determined by the value of u near x. Thus, the definition of a solution itself is nontrivial. Since TV is still convex and lower semicontinuous in many function spaces contained in the space of Schwartz distributions, we are able to apply an abstract theory based on maximal monotone operators stated by Y. Kōmura and developed by H. Brezis and others to get a solution. The equation can be interpreted as a gradient flow in a Hilbert space H

$$u_t \in -\partial_H \mathcal{E}(u),$$

where ∂_H denotes the subdifferential of \mathcal{E} (with respect to the metric in H), which is an extension of the notion of derivative.

Our first goal is to give a precise definition of a solution to total variation flows including (1.1), (1.2), (1.3). For the case (1.1), we note our formulations allow a weight, i.e., we also consider

$$bu_t = \operatorname{div}\left(a\nabla u/|\nabla u|\right). \tag{1.4}$$

The weight is sometimes important, especially when we consider the Kobayashi–Warren–Carter system [KWC] in materials science. As a singular limit, one has to consider even a discontinuous weight [GOU], [GOSU]. For (1.2) the interpretation as a gradient flow is sometimes non-trivial. In the case (1.2), as discussed in [GKL] the definition of a solution is quite involved for $\Omega = \mathbb{R}^n$ for $n \leq 2$ because interpretation as $u_t \in -\partial_H \mathcal{E}(u)$ is not clear. We then calculate the subdifferential to understand the flow in a more explicit way. This is essential for the construction of explicit solutions. When we consider a characteristic function of a set as an initial datum, we wonder whether the speed u_t is constant on the set and outside of the set. This leads to the notion of calibrability of a set.

Our second goal is to explain the notion of calibrability with special emphasis on the fourthorder problem. We also note that in the fourth-order problem (1.2), the support of the characteristic function may move, which does not occur for the second order problem (1.1). This is because the speed may contain a delta part. The existence of the delta part has already been observed in [Ka1] where the author discussed TV perturbed by E_p with some p > 1. See also [GG]. Based on calibrability, we give a few examples of explicit solutions when initial datum is a characteristic function. For the fourth-order problem the material is taken from [GKL], while for the second-order problem it is mostly taken from [ACM]. As a property of a solution, we further discuss extinction time estimates as studied in [GK], [GKM]. We discuss it for (1.3) which is new. We also present another formal argument to estimate the extinction time, which will be rigorously discussed in our forthcoming paper [GKL2].

In the last part of this paper, we review a couple of regularity results for the second-order problem. The main result is that the set of jump discontinuities of u(t) is contained in that of u_0 . Moreover, jump sizes are non-increasing in time. This has been first proved in [CCN, CJN]. Their proof is based on an analogous result for a time-discretized problem in [CCN], which relies on the connection between the total variation and surfaces of prescribed mean curvature, given by the co-area formula, and regularity theory for those surfaces. Here, we use instead a conceptually simpler technique from [CL] which does not rely on such subtle properties of TV. We also show a stronger estimate in the 1D case, appearing in [BF, BCNO], which implies non-expansion of jumps as well as pointwise estimate $|\nabla u(t)| \leq |\nabla u_0|$ (note that the latter does not hold in higher dimensions, see e.g. [ACC]). The situation is significantly different in the fourth-order case, where new jumps may be created as proved in [GG]. This behavior is also expected for (1.3) for 0 < s < 1 as numerically suggested in [GMR].

Total variation type flows find applications in many fields including image analysis [ROF], [ACM] materials science [KWC] and crystal growth problems. For the latter two topics, see recent survey [GP], where a general second-order crystalline mean curvature flow is discussed. The research on fourth-order problems is less popular. There is a review paper [GG] which includes the development before 2010. The field is growing and we do not touch many important topics such as total variation flow of maps between Riemannian manifolds, see e.g. [GMM, GLM, GSTU], flows on metric spaces [MST, BCP, GM], the Wasserstein TV flow [BCDS, CP] or TV flow with time-dependent boundary conditions [BDS, GNRS].

This paper is organized as follows. In Section 2, we recall definitions of a solution both to the second-order and the fourth-order total variation flow. We also review basic unique existence results and recall the notion of a Cahn-Hoffman vector field. In Section 3, we discuss the notion of calibrability with special emphasis on the fourth-order problem. In Section 4, we mention several explicit solutions, mainly radially symmetric piecewise constant solutions. In Section 5, we discuss several upper bounds for the extinction time. In Section 6, we discuss regularity properties of the flow.

2 Definition of a solution

2.1 Total variation flows

We begin with a standard definition of total variation [Giu] for a locally integrable function defined in Ω , where Ω is either a domain in \mathbb{R}^n or flat torus $\mathbb{T}^n = \prod_{i=1}^n (\mathbb{R}/\omega_i\mathbb{Z})$ with $\omega_i > 0$ for $i = 1, \ldots, n$. For an integrable function u, we define the total variation of u in Ω by

$$TV(u) := \sup\left\{-\int_{\Omega} u \operatorname{div} \varphi \, dx \, \middle| \, |\varphi(x)| \le 1 \text{ for all } x \in \Omega, \, \varphi \in (C_c^{\infty}(\Omega))^n\right\},$$

where $C_c^{\infty}(\Omega)$ denotes the space of all smooth functions compactly supported in Ω . If u belongs to the Sobolev space $W^{1,1}(\Omega)$ (that is, if its distributional derivative is an integrable function), then $TV(u) = \int_{\Omega} |\nabla u| dx < \infty$. More generally, TV(u) is finite if and only if the distributional derivative Du is a finite vector measure [AFP]. In this case we say that u is a function of bounded variation, $u \in BV(\Omega)$, and we can write $TV(u) = \int_{\Omega} |Du|$.

For later convenience, we also define weighted total variation. Let $a : \Omega \to [0, \infty]$ be a lower semicontinuous function. We set

$$TV_a(u) := \sup\left\{-\int_{\Omega} u \operatorname{div} \varphi \, dx \, \middle| \, |\varphi(x)| \le a(x) \text{ for all } x \in \Omega, \, \varphi \in (C_c^{\infty}(\Omega))^n\right\}.$$

This definition can be easily extended to the case when u is a Schwartz distribution, i.e., $u \in \mathcal{D}'(\Omega)$, by replacing $-\int_{\Omega} u \operatorname{div} \varphi \, dx$ by a canonical pair $-\langle u, \operatorname{div} \varphi \rangle$. Since $u_m \to u$ in $\mathcal{D}'(\Omega)$ implies $\langle u_m, \operatorname{div} \varphi \rangle \to \langle u, \operatorname{div} \varphi \rangle$, $TV_a(u)$ is a supremum of (sequentially) continuous function in $\mathcal{D}'(\Omega)$. Thus, $TV_a(u)$ is lower (sequentially) semicontinuous on $\mathcal{D}'(\Omega)$. In particular, $TV_a(u)$ is lower semicontinuous on $L^p(\Omega)$, $p \ge 1$. Since $TV_a(u)$ is a supremum of linear functionals $u \mapsto -\langle u, \operatorname{div} \varphi \rangle$ it must be convex on $\mathcal{D}'(\Omega)$. Note that our definition on $TV_a(u)$ is different from the one given by [AB] where the condition $|\varphi(x)| \le a(x)$ is only imposed for almost all $x \in \Omega$ when a is discontinuous. In their definition, $TV_a = TV$ if $a \equiv 1$ almost everywhere, while in our definition it may happen that

$$TV_a(u) < TV(u)$$

for some u even if a = 1 a.e. This is already observed in [AB]. Indeed, let us consider $\Omega = (-1, 1)$ and a(x) = 1 for $x \neq 0$, $a(0) = a_0 < 1$ ($a_0 \ge 0$). Then, we see that

$$TV_a(u) = \int_{\Omega_-} \left| \frac{du}{dx} \right| + \int_{\Omega_+} \left| \frac{du}{dx} \right| + a_0 \left| u(+0) - u(-0) \right|$$

with $\Omega_{-} = (-1,0)$, $\Omega_{+} = (0,1)$, where $u(\pm 0) = \lim_{\delta \downarrow 0} u(\pm \delta)$. Such a type of discontinuous lower semicontinuous function *a* will be important when we study a singular limit of the Kobayashi–Warren– Carter energy as discussed in [GOU] and [GOSU]; see [U] for a review. Although the total variation is defined under Finsler type metric [AB] which is important to study crystalline curvature [GP], we do not touch this problem in this paper.

We recall a classical theory for the gradient flow of a convex functional in a Hilbert space due to Y. Kōmura [Ko] and H. Brezis [Br].

Proposition 2.1. Let H be a (real) Hilbert space. Let \mathcal{E} be a lower semicontinuous functional on H with values in $(-\infty, \infty]$ and $\mathcal{E} \neq \infty$. Then for any $u_0 \in \overline{D(\mathcal{E})}$, there exists a unique $u \in C([0, \infty), H)$ with $u_t \in \bigcap_{\delta > 0} L^2((\delta, \infty), H)$ such that

$$u_t(t) \in -\partial_H \mathcal{E}\left(u(t)\right) \quad \text{for a.e. } t > 0, \quad u(0) = u_0. \tag{2.1}$$

If $u_0 \in D(\mathcal{E})$, then $\delta = 0$ is allowed.

The symbol $\partial_H E$ denotes the subdifferential of \mathcal{E} in H, i.e.,

$$\partial_H \mathcal{E}(v) = \left\{ f \in H \mid \mathcal{E}(v+h) - \mathcal{E}(v) \ge (h, f)_H \text{ for all } h \in H \right\}$$

for $v \in D(\mathcal{E}) = \{f \in H \mid \mathcal{E}(f) < \infty\}$, where $(\cdot, \cdot)_H$ denotes the inner product in H. One way to construct the solution to (2.1) is the *minimizing movements scheme*. For given $f \in H$, $\lambda > 0$, consider the functional $\mathcal{E}_f^{\lambda} \colon H \to (-\infty, \infty]$ of \mathcal{E} given by

$$\mathcal{E}_f^{\lambda}(w) = \lambda \mathcal{E}(w) + \frac{1}{2} \|w - f\|_H^2.$$
(2.2)

Like $\mathcal{E}, \mathcal{E}_f^{\lambda}$ is lower semicontinuous and satisfies $\mathcal{E}_f^{\lambda} \not\equiv \infty$. Moreover, it is coercive and strictly convex, so it has a unique minimizer. For given $N \in \mathbb{N}$, we inductively produce a sequence $(u_k^N)_{k \in \mathbb{N}}$ of elements of H by setting u_k^N to be the minimizer of $\mathcal{E}_{u_{k-1}^N}^{1/N}$, $k \in \mathbb{N}$ and $u_0^N = u_0$. Then, defining $u^N \in L^{\infty}(0, \infty; H)$ by

$$u^{N}(t) = u_{k}^{N}$$
 for $t \in [k/N, (k+1)/N),$ (2.3)

one can show that u^N converges locally uniformly on $[0, \infty)$ to a solution to (2.1). The solution is unique by monotonicity of the subdifferential of a convex functional.

For a given non-negative measurable function $b \in L^{\infty}(\Omega)$ with $1/b \in L^{\infty}(\Omega)$, we define an inner product on $L^{2}(\Omega)$ by

$$(u,v)_{L^2_b} := \int_{\Omega} b(x)u(x)v(x) \ dx.$$

This is equivalent to the standard inner product on L^2 , $(u, v)_{L^2}$, corresponding to $b \equiv 1$. Let $L_b^2(\Omega)$ be the space $L^2(\Omega)$ equipped with the inner product $(\ ,\)_{L_b^2}$. The equation $u_t \in -\partial_{L_b^2} TV_a(u)$ formally corresponds to

$$b\frac{\partial u}{\partial t} = \operatorname{div}\left(a\frac{\nabla u}{|\nabla u|}\right) \quad \text{in} \quad \Omega \times [0,\infty),$$

$$\frac{\partial u}{\partial \nu} = 0 \quad \text{on} \quad \partial\Omega \times (0,\infty) \quad \text{if there is a boundary} \quad \partial\Omega \quad \text{of} \quad \Omega.$$
(2.4)

This type of equation is discussed in [GGK] in one-dimensional periodic case, i.e., $\Omega = \mathbb{T}$. Here $\partial u / \partial \nu$ is the normal derivative of u on $\partial \Omega$.

Definition 2.2. We say that $u \in C([0,\infty), L_b^2(\Omega))$ with $u_t \in \bigcap_{\delta>0} L^2((\delta,\infty), L_b^2(\Omega))$ is a solution of (2.4) with initial datum $u_0 \in L_b^2(\Omega)$ if it satisfies (2.1) with $H = L_b^2(\Omega)$ and $\mathcal{E} = TV_a$.

We note that $D(TV_a)$ is dense in $L_b^2(\Omega)$ since $TV_a(u)$ is finite on a space $C_c^{\infty}(\Omega)$, which is dense in $L_b^2(\Omega)$. Since TV_a is lower semicontinuous in \mathcal{D}' , it is also lower semicontinuous in $L_b^2(\Omega)$. Since TV is convex in $L_b^2(\Omega)$, Proposition 2.1 yields

Theorem 2.3. Let $b \in L^{\infty}(\Omega)$ be nonnegative with $1/b \in L^{\infty}(\Omega)$. Let $a : \Omega \to [0, \infty]$ be a lower semicontinuous function. Then for any $u_0 \in L^2_b(\Omega)$, there exists a unique solution u to (2.4).

We are interested in considering a higher order total variation flow of the form

$$\frac{\partial u}{\partial t} = (-\Delta) \operatorname{div} \left(a \frac{\nabla u}{|\nabla u|} \right).$$

We first consider the case $\Omega = \mathbb{T}^n$. For simplicity, we set $\omega_i = 1$ $(1 \leq i \leq n)$. In this case, the homogeneous Sobolev space can be defined by imposing the average free condition, using Fourier series. For $s \in \mathbb{R}$, we set

$$\dot{H}^{s}_{\mathrm{av}}(\mathbb{T}^{n}) = \left\{ u = \sum_{\substack{m \in \mathbb{Z}^{n} \\ m \neq 0}} a_{m} e^{2\pi i x \cdot m} \in \mathcal{D}' \ \left| \ \|u\|^{2}_{\dot{H}^{s}_{\mathrm{av}}} := \sum_{\substack{m \in \mathbb{Z}^{n} \\ m \neq 0}} |m|^{2s} |a_{m}|^{2} < \infty \right\}.$$

This space is a (complex) Hilbert space with inner product

$$((u,v))_s := \sum_{m \neq 0} a_m \overline{b}_m |m|^{2s}, \quad u = \sum_{m \neq 0} a_m e^{2\pi i x \cdot m}, \quad v = \sum_{m \neq 0} b_m e^{2\pi i x \cdot m}$$

It is an average free space. We consider the space of all real-valued functions in \dot{H}^s_{av} which is still denoted by \dot{H}^s_{av} . The total variation TV_a is well-defined on any \dot{H}^s_{av} since \dot{H}^s_{av} can be viewed as a subspace of $\mathcal{D}'(\mathbb{T}^n)$. Since TV_a is convex and lower semicontinuous, Proposition 2.1 yields

Theorem 2.4. Let $a : \mathbb{T}^n \to [0,\infty]$ be a lower semicontinuous function. Let $s \in \mathbb{R}$. For any $u_0 \in \dot{H}^{-s}_{av}(\mathbb{T}^n)$, there is a unique $u \in C\left([0,\infty), \dot{H}^{-s}_{av}(\mathbb{T}^n)\right)$ with $u_t \in \bigcap_{\delta>0} L^2\left((\delta,\infty), \dot{H}^{-s}_{av}(\mathbb{T}^n)\right)$ satisfying

 $u_t \in -\partial_{\dot{H}_{av}^{-s}} TV_a(u), \quad a.e. \quad t > 0, \quad u(0) = u_0.$

We simply say u is a solution of

$$u_t = (-\Delta)^s \operatorname{div} \left(a \nabla u / |\nabla u| \right) \tag{2.5}$$

with initial data u_0 , where Δ denotes the Laplacian.

The operator $(-\Delta)^s$ comes from relation of $\partial_{\dot{H}_{av}^{-s}}$ and ∂_{L^2} . Let us give a formal explanation for s = 1. If $f \in \partial_{L^2} TV_a(v)$, then

$$TV_a(v+h) - TV_a(v) \ge (f,h)_{L^2}$$
 for $h \in L^2(\Omega)$.

Since $\dot{H}_{\rm av}^{-1}$ is the dual of $\dot{H}_{\rm av}^1$ and since $-\Delta$ is the canonical isometry from $\dot{H}_{\rm av}^1$ to $\dot{H}_{\rm av}^{-1}$, i.e.,

$$-\Delta: u \mapsto ((u, \cdot))_1$$

we see

$$(f,h)_{L^2} = ((-\Delta f,h))_{-1}$$

since $((u,v))_{-1} = ((-\Delta)^{-1}u,v)_{L^2}$. If $(-\Delta)f \in \dot{H}_{av}^{-1}$, by density of \dot{H}_{av}^1 in \dot{H}_{av}^{-1} , we conclude that $-\Delta f \in \partial_{\dot{H}_{av}^{-1}}TV_a(v)$.

If $\Omega = \mathbb{R}^n$, the definition of the fourth order total variation flow is more involved, especially for n = 1, 2. We consider an inner product

$$((u,v))_1 := \int_{\mathbb{R}^n} \nabla u \cdot \nabla v \ dx$$

for $u, v \in C_c^{\infty}(\mathbb{R}^n)$. Let $D_0^1(\mathbb{R}^n)$ be the completion of $C_c^{\infty}(\mathbb{R}^n)$ in the norm $||u||_1 = ((u, u))_1^{1/2}$. It is a Hilbert space equipped with $((u, v))_1$. For $n \ge 3$, this space is identified with

$$D_0^1(\mathbb{R}^n) = D^1(\mathbb{R}^n) \cap L^{2^*}(\mathbb{R}^n), \quad D^1(\mathbb{R}^n) = \left\{ u \in L^1_{loc}(\mathbb{R}^n) \mid \nabla u \in L^2(\mathbb{R}^n) \right\},$$

where $2^* = 2n/(n-2)$ is the corresponding Sobolev exponent; i.e., the exponent such that $D_0^1(\mathbb{R}^n) \subset L^{2^*}(\mathbb{R}^n)$; see e.g. [Gal]. However, for $n \leq 2$, this $D_0^1(\mathbb{R}^n)$ is not a subspace of $L_{loc}^1(\mathbb{R}^n)$. Instead, it is isometrically identified with the quotient space $D^1(\mathbb{R}^n) := D^1(\mathbb{R}^n)/\mathbb{R}$ equipped with the inner product $((u, v))_1$; see e.g. [Gal]. We need to be careful because an element of $D_0^1(\mathbb{R}^n)$ is determined up to constant. In the case $\Omega = \mathbb{T}^n$, the space $D^1(\mathbb{T}^n)$ has a direct sum decomposition

$$D^1(\mathbb{T}^n) = \dot{H}^1_{\mathrm{av}}(\mathbb{T}^n) \oplus \mathbb{R}$$

which corresponds to a decomposition of $u \in D^1(\mathbb{T}^n)$ as

$$u = (u - u_c) + u_c,$$

where u_c is the average of u over \mathbb{T}^n . Thus, the space $\dot{D}(\mathbb{T}^n) := D^1(\mathbb{T}^n)/\mathbb{R}$ is identified with a subspace $\dot{H}^1_{av}(\mathbb{T}^n)$ of $D^1(\mathbb{T}^n)$. In the case of \mathbb{R}^n , no such decomposition is available.

We are interested in the dual space $(D_0^1(\mathbb{R}^n))'$. Let $-\Delta$ denote the canonical isometry from $D_0^1(\mathbb{R}^n)$ and its dual, i.e.,

$$-\Delta: u \mapsto ((u, \cdot))_1.$$

The inner product of $(D_0^1(\mathbb{R}^n))'$ is defined as

$$((u,v))_{(D_0^1(\mathbb{R}^n))'} := (((-\Delta)^{-1}u, (-\Delta)^{-1}v))_{D_0^1(\mathbb{R}^n)}.$$

We introduce a subspace $\tilde{D}^{-1}(\mathbb{R}^n) \subset D_0^1(\mathbb{R}^n)'$ of the form

$$\tilde{D}^{-1}(\mathbb{R}^n) = \left\{ w \mapsto \int_{\mathbb{R}^n} uv \, dx \, \middle| \, u \in C_c^{\infty}(\mathbb{R}^n) \right\} \quad \text{if} \quad n \ge 3,$$

$$\tilde{D}^{-1}(\mathbb{R}^n) = \left\{ w \mapsto \int_{\mathbb{R}^n} uw \, dx \, \middle| \, u \in C_{c,\text{av}}^{\infty}(\mathbb{R}^n) \right\} \quad \text{if} \quad n = 1 \quad \text{or} \quad n = 2$$

where

$$C_{c,\mathrm{av}}^{\infty}(\mathbb{R}^n) = \left\{ u \in C_c^{\infty}(\mathbb{R}^n) \mid \int_{\mathbb{R}^n} u \, dx = 0 \right\}.$$

It is well known that $\tilde{D}^{-1}(\mathbb{R}^n)$ is dense in $(D_0^1(\mathbb{R}^n))'$; see e.g. [Gal]. In the case n = 1, 2, the restriction to the average-free space $C_{c,\mathrm{av}}^{\infty}(\mathbb{R}^n)$ is necessary for the functionals to be well-defined on $D^1(\mathbb{R}^n)/\mathbb{R}$. Since $\mathcal{D} = C_c^{\infty}(\mathbb{R}^n)$ is continuously embedded in $D_0^1(\mathbb{R}^n)$, $D^{-1}(\mathbb{R}^n) = (D_0^1(\mathbb{R}^n))'$ can be viewed as a subspace of $\mathcal{D}'(\mathbb{R}^n)$. Thus, $TV_a(u)$ for $u \in D^{-1}(\mathbb{R}^n)$ is well-defined and it is convex, lower semicontinuous on the Hilbert space $D^{-1}(\mathbb{R}^n)$ provided that a is a lower semicontinuous. Proposition 2.1 guarantees the existence of a unique solution to $u_t \in -\partial_{D^{-1}}TV_a(u)$ with initial datum $u_0 \in D^{-1}(\mathbb{R}^n)$. This is a rigorous way to interpret $u_t = -\Delta \operatorname{div}(a\nabla u/|\nabla u|)$. Unfortunately, this existence result has a drawback even if $a \equiv 1$: the characteristic function 1_K of a set K does not belong to $D^{-1}(\mathbb{R}^n)$ unless the Lebesgue measure of K equals zero for n = 1, 2, since $D^{-1}(\mathbb{R}^n)$ requires a kind of average-free condition for n = 1, 2. In fact for $u_0 \in L^2(\mathbb{R}^n)$ with compact support, $u_0 \in D^{-1}(\mathbb{R}^n)$ if and only if $\int_{\mathbb{R}^n} u_0 \, dx = 0$ as proved in [GKL, Lemma 17]. We have to extend space D^{-1} when $n \leq 2$. This is quite involved; see discussion at the end of Section 2.2. We refer to [GKL] for details when $a \equiv 1$. In the case where a depends on x, the argument in [GKL] still works provided that the approximation lemma [GKL, Lemma 6] can be extended to TV_a .

In general, we can consider the space D_0^s (with $s \in \mathbb{R}$) which is the completion of $C_c^{\infty}(\mathbb{R}^n)$ in the norm

$$||u||_{D_0^s}^2 = \int_{\mathbb{R}^n} |\xi|^{2s} |\hat{u}(\xi)|^2 d\xi,$$

where \hat{u} denotes the Fourier transform of u, i.e., $\hat{u}(\xi) = \int_{\mathbb{R}^n} e^{-ix\cdot\xi} u(x) dx$. For $0 < s \leq 1$, the space $D_0^s \subset L_{loc}^1$ for n > 2s but again this does not hold for $n \leq 2s$. In a similar way, we are able to define a total variation flow $u_t = (-\Delta)^s \operatorname{div}(\nabla u/|\nabla u|)$ for n > 2s, whose existence is proved by Proposition 2.1.

Remark 2.5. (i) We are able to consider higher-order problems in a domain with boundary. Even in the case of s = 1, there are several choices depending on what kind of boundary condition we impose for the Laplacian. If we impose the homogeneous Dirichlet boundary condition, then the resulting equation is formally of the form

$$\begin{aligned} \frac{\partial u}{\partial t} &= -\Delta \operatorname{div} \left(a \nabla u / |\nabla u| \right) \quad in \quad \Omega \times (0, \infty) \\ u &= 0, \quad \operatorname{div} \left(a \nabla u / |\nabla u| \right) = 0 \quad on \quad \partial \Omega \times (0, \infty). \end{aligned}$$

A rigorous formulation is given in [GKM]. It is enough to take $H^{-1}(\Omega) = (H_0^1(\Omega))'$ as the Hilbert space H. The Sobolev space $H_0^1(\Omega)$ can be defined similarly as $D_0^1(\mathbb{R}^n)$ by replacing \mathbb{R}^n by Ω . By the Poincaré inequality, this space $H_0^1(\Omega)$ belongs to $L^2(\Omega)$ so its dual $H^{-1}(\Omega)$ includes $L^2(\Omega)$. However, if we consider the Laplace operator with the homogeneous Neumann boundary condition, the correct choice of the space is $D_N^{-1} = (D^1(\Omega)/\mathbb{R})'$, and expected boundary condition is

$$\frac{\partial u}{\partial \nu} = 0, \quad \frac{\partial}{\partial \nu} \operatorname{div} \left(a \frac{\nabla u}{|\nabla u|} \right) = 0 \quad on \quad \partial \Omega \times (0, \infty).$$

Similar to the case $D^{-1}(\mathbb{R}^n)$, the analysis is quite involved. It will be discussed in our forthcoming paper. (ii) Even in the second-order problem, if one would like to consider the Dirichlet problem, i.e., u = 0on $\partial\Omega$, one should replace the energy functional TV_a by

$$TV_a(u) + \int_{\partial\Omega} a|u| \ d\mathcal{H}^{n-1}.$$

This type of problem is discussed in [ACM] at least for $a \equiv 1$.

2.2 Formulation by the Cahn–Hoffman vector field

It is nontrivial to characterize the subdifferential of TV_a in a given Hilbert space. For a general energy \mathcal{E} , a standard way is to propose a candidate set A(u) by calculating the Euler-Lagrange operator and prove $A(u) \subset \partial \mathcal{E}(u)$. This part is not hard. The converse inclusion is difficult. Since $\partial \mathcal{E}$ is maximal monotone, it suffices to prove that A is also maximal monotone which yields $A = \partial \mathcal{E}$. The proof of monotonicity is not difficult. To show maximality, we prove that the resolvent equation $u + \lambda Au \ni f$ is always solvable for $f \in H$ and $\lambda > 0$. This argument is often carried out by approximation of the operator A. This procedure is found for example in [GNRS].

However, if \mathcal{E} is positively homogeneous of degree one (i.e., one-homogeneous), there is an easier method due to F. Alter; see [ACM, Chapter 1]. The basic strategy is characterize the subdifferential $\partial \mathcal{E}$ by the polar \mathcal{E}^0 of $\mathcal{E} : H \to (-\infty, \infty]$

$$\mathcal{E}^{0}(v) := \sup\left\{ (u, v)_{H} \mid u \in H, \ \mathcal{E}(u) \le 1 \right\}.$$

By convex analysis, $(\mathcal{E}^0)^0 = \mathcal{E}$ if \mathcal{E} is a non-negative, lower semicontinuous, convex provided that \mathcal{E} is positively one-homogeneous, i.e.,

$$\mathcal{E}(\lambda u) = \lambda \mathcal{E}(u) \text{ for all } \lambda > 0, \quad u \in H.$$

A key observation is a simple lemma.

Lemma 2.6 ([ACM], Theorem 1.8). Let \mathcal{E} be convex and positively one-homogeneous in a Hilbert space H, then $v \in \partial_H \mathcal{E}(u)$ if and only if $\mathcal{E}^0(v) \leq 1$ and $(u, v)_H = \mathcal{E}(u)$.

It is convenient to introduce a class of vector fields

$$X_2 = \left\{ z \in L^{\infty}(\Omega, \mathbb{R}^n) \mid \operatorname{div} z \in L^2(\Omega) \right\}.$$

For $z \in X_2$, the normal trace $[z \cdot \nu]$ is well defined as an element of $L^{\infty}(\partial \Omega)$ see e.g. [ACM]. In many cases, \mathcal{E}^0 is computable. For example, if $\mathcal{E} = TV$ and $H = L^2(\Omega)$,

$$\mathcal{E}^{0}(v) = \inf \left\{ \|z\|_{L^{\infty}} \mid z \in X_{2}, \ v = -\operatorname{div} z \text{ in } \Omega, \ [z \cdot \nu] = 0 \text{ on } \partial\Omega \right\}$$

at least when $\Omega = \mathbb{R}^n$, \mathbb{T}^n or a bounded domain with Lipschitz boundary [ACM]. For higher order problem,

$$\mathcal{E}^{0}(v) = \inf \left\{ \|z\|_{L^{\infty}} \mid z \in L^{\infty}(\Omega, \mathbb{R}^{n}), \ v = \Delta \operatorname{div} z \text{ in } \mathbb{R}^{n}, \ \operatorname{div} z \in D_{0}^{1} \right\},\$$

when $\mathcal{E} = TV$ and $H = D^{-1}(\mathbb{R}^n)$; see [GKL, Theorem 12]. Although there is no explicit literature, we expect

$$\mathcal{E}^{0}(v) = \inf \left\{ \|z\|_{L^{\infty}} \mid z \in X_{2}, \ bv = -\operatorname{div}(az) \ \text{in } \Omega, \ a[z \cdot \nu] = 0 \ \text{on } \partial\Omega \right\}$$

for general TV_a and $H = L_b^2$. In the fractional case, we expect

$$\mathcal{E}^{0}(v) = \inf \left\{ \|z\|_{L^{\infty}} \mid z \in X_{2}, \ v = -(-\Delta)^{s} \operatorname{div} z, \ \operatorname{div} z \in \dot{H}^{s}(\mathbb{T}^{n}) \right\}$$

for $\mathcal{E} = TV$ and $H = \dot{H}^{-s}(\mathbb{T}^n)$. Note that the minimizer is attained, so Lemma 2.6 implies the characterization of the subdifferential. We only state precise results for $\partial_{L^2}TV$ and $\partial_{D^{-1}}TV$.

- **Theorem 2.7.** (i) [ACM, Lemma 2.4] Assume that $\Omega = \mathbb{R}^n$, \mathbb{T}^n or a bounded domain in \mathbb{R}^n with Lipschitz boundary. An element $v \in L^2(\Omega)$ belongs to $\partial_{L^2}TV(u)$ if and only if there is $Z \in X_2(\Omega)$ such that
- (ia) $|Z| \leq 1$ in Ω and $[Z \cdot \nu] = 0$ on $\partial \Omega$
- (*ib*) $v = -\operatorname{div} Z$ in Ω
- $(ic) (u, \operatorname{div} Z)_{L^2} = TV(u)$
- (*ii*) [GKL, Theorem 14] Assume that $\Omega = \mathbb{R}^n$. An element $v \in D^{-1}(\mathbb{R}^n)$ belongs to $\partial_{D^{-1}}TV(u)$ if and only if there is $Z \in L^{\infty}(\mathbb{R}^n)$ with div $Z \in D_0^1(\mathbb{R}^n)$ such that
- (iia) $|Z| \leq 1$ in Ω
- (*iib*) $v = \Delta \operatorname{div} Z$ in Ω
- $(\ddot{u}c) \langle u, \operatorname{div} Z \rangle = TV(u).$ Here \langle , \rangle denotes the duality pairing between D^{-1} and D_0^1 .

This vector field Z is often called a Cahn-Hoffman vector field. If u is Lipschitz continuous, the conditions (ia), (ic) (or (iia), (iib)) imply that $Z = \nabla u/|\nabla u|$ whenever $\nabla u(x) \neq 0$ for almost all x. Thus, for example, for the second-order problem, the subdifferential $-\operatorname{div} Z$ formally agrees with the standard Euler-Lagrange operator $-\operatorname{div}(\nabla u/|\nabla u|)$ at least where $\nabla u(x) \neq 0$. If u is a special class of functions, the subdifferential at such u is easily computable when n = 1 including the case where there are weights a, b; see [FG], [GGK]. Once the subdifferential is calculated, we are able to give an explicit formulation of a solution.

Theorem 2.8. (i) Assume that $u \in C([0,\infty), L^2(\Omega))$. Then u is a solution of $u_t \in -\partial_{L^2}TV(u)$ with $u_0 = u(0)$ if and only if there exists $Z \in L^{\infty}(\Omega \times (0,\infty))$ with

div
$$Z \in \bigcap_{\delta > 0} L^2(\delta, \infty; L^2(\Omega))$$

satisfying

$$\left\{ \begin{array}{l} u_t = \operatorname{div} Z \ in \ L^2(\Omega) \\ |Z(x,t)| \leq 1 \ for \ a.e. \ x \in \Omega, \ [Z(x,t) \cdot \nu] = 0 \ for \ a.e. \ x \in \partial \Omega \\ -(u, \operatorname{div} Z)_{L^2} = TV(u) \end{array} \right.$$

for a.e. t > 0. (If $TV(u_0) < \infty$, $\delta = 0$ is allowed.)

(ii) Assume that $u \in C([0,\infty), D^{-1}(\mathbb{R}^n))$. Then u is a solution of $u_t \in -\partial_{D^{-1}}TV(u)$ with $u_0 = u(0)$ if and only if there exists $Z \in L^{\infty}(\mathbb{R}^n \times (0,\infty))$ with

$$\operatorname{div} Z \in \bigcap_{\delta > 0} L^2\left(\delta, \infty; D^1_0(\mathbb{R}^n)\right)$$

satisfying

$$\begin{cases} u_t = -\Delta \operatorname{div} Z \text{ in } D^{-1}(\mathbb{R}^n) \\ |Z(x,t)| \leq 1 \text{ for a.e. } x \in \mathbb{R}^n, \\ \langle u, \operatorname{div} Z \rangle = -TV(u) \end{cases}$$

for a.e. t > 0. (If $TV(u_0) < \infty$, $\delta = 0$ is allowed.)

This follows from the characterization of the subdifferential (Theorem 2.7) except that the Cahn-Hoffman vector field should be chosen to be measurable both in x and t. For the second-order case [ACM, Section 2.4], this can be done by recalling that Bochner integrable functions can be well approximated by piecewise constant functions. For the fourth-order problem, although the idea is similar, the situation is slightly different as discussed in [GKL, Theorem 15, Lemma 16]. By the way this characterization of a solution using Cahn-Hoffman vector fields for the fourth-order of the problem was already proposed by [GKM]. For the case $\Omega = \mathbb{T}^n$, a similar characterization of a solution is given in [GMR, Theorem 1] for general s > 0 with $a \equiv 1$. It is of the form.

Theorem 2.9. For $s \in \mathbb{R}$, assume that $u \in C\left([0,\infty), \dot{H}_{av}^{-s}(\mathbb{T}^n)\right)$. Then u is a solution of (2.5) with $a \equiv 1$ and initial datum $u(0) = u_0 \in \dot{H}_{av}^{-s}(\mathbb{T}^n)$ if and only if there exists $Z \in L^{\infty}(\Omega \times (0,\infty))$ with

div
$$Z \in \bigcap_{\delta > 0} L^2\left(\delta, \infty; \dot{H}^{-s}_{\mathrm{av}}(\mathbb{T}^n)\right)$$

satisfying

$$\begin{cases} u_t = (-\Delta)^s \operatorname{div} Z \text{ in } \dot{H}_{\operatorname{av}}^{-s}(\mathbb{T}^n) \\ |Z(x,t)| \le 1 \text{ for a.e. } x \in \mathbb{T}^n \\ \langle u, \operatorname{div} Z \rangle = -TV(u) \end{cases}$$

for a.e. t > 0. (If $TV(u_0) < \infty$, $\delta = 0$ is allowed).

This follows from the fact that $f \in \partial_{\dot{H}_{av}^{-s}}TV(u)$ for $u \in \dot{H}_{av}^{-s}$ if and only if $TV^0((-\Delta)^{-s}f) \leq 1$ and $\int_{\mathbb{T}^n} (-\Delta)^{-s} f \cdot v \, dx = TV(u)$, when TV^0 is taken in a middle space $L^2_{av}(\mathbb{T}^n) = \dot{H}^0_{av}(\mathbb{T}^n)$. For the case s = 1, this is found in [GK, Lemma 3.1].

As mentioned before, in the case s = 1 and $\Omega = \mathbb{R}^n$, the gradient flow $u_t \in -\partial_{D^{-1}}TV(u)$ is not enough to study the evolution of a characteristic function for n = 1, 2. We have to extend the function space D^{-1} . We quickly review the way it is done in [GKL]. For this purpose, we take $\psi \in L^2(\mathbb{R}^n)$ with compact support such that $\int_{\mathbb{R}^n} \psi \, dx \neq 0$. We set

$$E_{\psi}^{-1} = \left\{ e + c\psi \mid w \in D^{-1}(\mathbb{R}^n), \ c \in \mathbb{R} \right\}.$$

This space in independent of the choice of ψ so we simply write by E^{-1} . For $n \ge 3$, $E^{-1} = D^{-1}$. We also denote $E_0^1 = D^1$ if $n \le 2$ and $E_0^1 = D_0^1$ if $n \ge 3$. The space E^{-1} can be considered as a dual space of E_0^1 . The inner product for $n \le 2$ is defined by

$$((v_1, v_2))_{E_0^1} := (([v_1], [v_2]))_{D_0^1} + \int_{\mathbb{R}^n} \psi v_1 \, dx \int_{\mathbb{R}^n} \psi v_2 \, dx$$
$$((u_1, u_2))_{E^{-1}} := ((w_1, w_2))_{D^{-1}} + c_1 c_2$$

for $u_i = w_i + c_i \psi$, $w_i \in D^{-1}$, $c_i \in \mathbb{R}$, $v_i \in E_0^1$, where $\int_{\mathbb{R}^n} \psi \, dx = 1$. It turns out that the "partial" gradient flow $u_t \in -\partial_{D^{-1}}TV(u)$ is exactly what we want. Calculating the subdifferential $\partial_{E^{-1}}TV(u)$ and projecting to D^{-1} yields $\partial_{D^{-1}}TV(u)$. It turns out the flow is independent of the choice of ψ . We have a characterization of this flow in terms of a Cahn–Hoffman vector field extending the one obtained in Theorem 2.8(ii). We prefer to use this characterization as a definition of a solution.

Definition 2.10. Assume that $u_0 \in E^{-1}$. We say that $u \in C([0,\infty), E^{-1})$ with $u(0) = u_0$ and $u_t \in L^2_{loc}((0,\infty), D^{-1})$ is a solution of

$$u_t = -\Delta \operatorname{div} \left(\nabla u / |\nabla u| \right) \quad in \quad \mathbb{R}^n \times (0, \infty) \tag{2.6}$$

with initial datum u_0 if there exists $Z \in L^{\infty} (\mathbb{R}^n \times (0, \infty))$ with

div
$$Z \in \bigcap_{\delta > 0} L^2\left(\delta, \infty; E_0^1(\mathbb{R}^n)\right)$$

satisfying

$$\begin{cases} u_t = -\Delta \operatorname{div} Z \text{ in } D^{-1}(\mathbb{R}^n) \\ |Z(x,t)| \leq 1 \text{ for a.e. } x \in \mathbb{R}^n \\ \langle u, \operatorname{div} Z \rangle = -TV(u) \end{cases}$$

for a.e. t > 0, where $\langle \ , \ \rangle$ denotes the duality pairing between E^{-1} and E_0^1 .

Although there are several technical steps, we conclude the unique existence of the solution [GKL, Theorem 2].

Theorem 2.11. For $u_0 \in E^{-1}$, there exists a unique solution u to (2.6) with initial datum u_0 .

Although sometimes it is difficult to compare with original gradient flow, it is reasonable to define a solution using a Cahn–Hoffman vector field like Definition 2.10. Here are a few examples of definitions.

Definition 2.12. Assume that $a : \Omega \to [0, \infty]$ is lower semicontinuous and that b > 0 satisfies $b \in L^{\infty}(\Omega)$ and $1/b \in L^{\infty}(\Omega)$. We say that $u \in C((0,T), \mathcal{D}'(\Omega))$ with $TV_a(u(t)) < \infty$ for all $t \in (0,T)$ is a Cahn–Hoffman solution (CH solution for short) of (2.4) if there exists a measurable function Z on $\Omega \times (0,T)$ with div $Z \in L^2_{loc}(\overline{\Omega} \times (0,T))$ such that

$$\begin{cases} bu_t = \operatorname{div} Z\\ |Z(x,t)| \le a(x) \text{ for a.e. } x \in \Omega, \ [Z,\nu] = 0 \text{ a.e. } x \in \partial\Omega \text{ for a.e. } t \in (0,T)\\ -(u,\operatorname{div} Z)_{L^2} = TV_a(u) \end{cases}$$

for a.e. $t \in (0, T)$.

We next consider

$$u_t = -\operatorname{div}\left(M\nabla\operatorname{div}\left(\nabla u/|\nabla u|\right)\right) \tag{2.7}$$

where $M \in L^{\infty}(\Omega, \mathbb{R}^{n \times n})$ is a real symmetric matrix-valued function with $M \ge c_0 I$ uniformly with some $c_0 > 0$. The boundary condition we impose is

$$\frac{\partial u}{\partial \nu} = 0, \quad M \cdot \nabla \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) = 0 \quad \text{on} \quad \partial \Omega.$$
 (2.8)

Definition 2.13. We say that $u \in C((0,T), \mathcal{D}'(\Omega))$ with $TV(u(t)) < \infty$ for all $t \in (0,T)$ is a CH solution to (2.7) with (2.8) if there exists $Z \in L^{\infty}(\Omega \times (0,\infty))$ with $\operatorname{div} Z \in L^{2}_{loc}(0,T,H^{1}_{loc}(\bar{\Omega}))$

$$\begin{cases} u_t = -\operatorname{div}(M\nabla\operatorname{div} Z) \\ |Z(x,t)| \le 1 \text{ for a.e. } x \in \Omega, \ [Z \cdot \nu] = 0 \text{ for a.e. } x \in \partial\Omega \\ -\langle u, \operatorname{div} Z \rangle = TV(u) \end{cases}$$

for a.e. t > 0; here $\langle u, \operatorname{div} Z \rangle$ should be interpreted as some "duality pair" but for a low-dimensional problem $n \leq 4$,

$$\langle u, \operatorname{div} Z \rangle = \int_{\Omega} u \operatorname{div} Z \, dx$$

when Ω is bounded. In fact, in this case, $u \in L^{n/(n-1)}$ since $TV(u) < \infty$ and $\operatorname{div} Z \in L^{\frac{n}{n-2}}$ by the Sobolev embedding so $1 - \frac{1}{n} + \frac{1}{2} - \frac{1}{n} = \frac{3}{2} - \frac{2}{n} \leq 1$ for $n \leq 4$. This implies $u \operatorname{div} Z \in L^1(\Omega)$ (for a.e. t > 0).

In this definition, we do not assume that u is in L^2 . The problem like (2.7) with weight appears in relaxation process of a surface of a crystal; see e.g. a review paper by R. V. Kohn [Kh].

We mention that in the second-order case, existence and uniqueness of a solution defined in terms of a Cahn–Hoffman field can be obtained also for initial data in the non-hilbertian space $L^1(\Omega)$ larger than $L^2(\Omega)$, owing to complete *m*-accretivity of the operator div $\frac{\nabla u}{|\nabla u|}$ [ABC1, ABC2], see also [ACM]. This does not seem to have an analogue in the higher-order cases.

3 Calibrability

The domain of TV, that is the space $BV(\Omega)$, contains functions with jump discontinuities, such as characteristic functions of sufficiently regular sets. In many cases, the evolutions of characteristic functions of sets under total variation flows are particularly simple. As a first step towards constructing such examples, we are interested in sets for which the speed $u_t(t)$ of the solution u(t) is spatially constant on the set. This leads to a geometric notion of *calibrability* of sets.

In the setting of Proposition 2.1, a general theory implies that

$$u_t(t) = -\partial_H^0 \mathcal{E}(u(t)) \quad \text{for all} \quad t > 0,$$

where $\partial^0_H \mathcal{E}$ denotes the minimal section (canonical restriction) of $\partial_H \mathcal{E}$, i.e.,

$$\partial_H^0 \mathcal{E}(u) := \arg\min\left\{ \|v\|_H \mid v \in \partial_H \mathcal{E}(u) \right\}.$$

In other words, $\partial_H^0 \mathcal{E}(u)$ is the minimizer of $||v||_H$ over $\partial_H \mathcal{E}(u)$. Since $\partial_H \mathcal{E}(u)$ is a closed convex set, there always exists a unique minimizer so $\partial_H^0 \mathcal{E}(u)$ is well defined.

To calculate the minimal section, we begin with a simple situation. Let U be a smooth open set in $\Omega = \mathbb{R}^n$. We consider a Lipschitz function u such that

$$\bar{U} = \left\{ x \in \Omega \mid u(x) = 0 \right\}$$

and u is smooth outside \overline{U} . Assume further that $\nabla u \neq 0$ outside \overline{U} .



Figure 1: Examples of U

We begin with the second-order problem. In this case, by Theorem 2.7 (i),

$$\partial_{L^2}^0 TV(u) = \arg\min\left\{ \|\operatorname{div} Z\|_{L^2} \mid |Z| \le 1 \text{ a.e. in } U, \ Z = \nabla u / |\nabla u| \text{ in } \bar{U}^c, \ Z \in X_2 \right\}.$$

Let Z_0 be a minimizer. Since the value of Z_0 outside \overline{U} is always the same, it suffices to consider its restriction to U (still denoted by Z_0). Then, Z_0 is a minimizer of

$$\left\{ \int_{U} |\operatorname{div} Z|^2 \, dx \, \middle| \, |Z| \le 1 \text{ a.e. in } U, \ [Z \cdot \nu] = \chi \text{ on } \partial U, \ Z \in X_2 \right\}.$$

Here, $\chi : \partial U \to \{-1, 1\}$ is a signature function defined by

$$\chi(x) = \begin{cases} 1 & \text{if } u > 0 \text{ outside } \bar{U} \text{ near } x \in \partial U \\ -1 & \text{otherwise} \end{cases}$$

For the fourth-order problem, by Theorem 2.7 (ii),

$$\partial_{D^{-1}}^0 TV(u) = \arg\min\left\{ \|\operatorname{div} Z\|_{D_0^1} \mid |Z| \le 1 \text{ a.e. in } U, \ Z = \nabla u / |\nabla u| \text{ in } \bar{U}^c, \ \operatorname{div} Z \in E_0^1 \right\}.$$

We argue in the same way. Let Z_0 be a minimizer of this problem, its restriction to U (still denoted by Z_0) is a minimizer of

$$\left\{ \int_{U} |\nabla \operatorname{div} Z|^2 \, dx \, \middle| \, |Z| \le 1 \text{ a.e. in } U, \, [Z \cdot \nu] = \chi \text{ on } \partial U, \, \operatorname{div} Z = \chi \kappa \text{ on } \partial U \right\},$$

where κ is the sum of all inward principal curvatures, i.e., n-1 times the mean curvature. This is because the inward and outward trace must agree for div Z since div $Z \in E_0^1$. We note that

$$\operatorname{div} Z = \operatorname{div} \left(\nabla u / |\nabla u| \right) = \chi \operatorname{div} \nu = \chi \kappa,$$

where ν is an external unit normal.

We are interested in the case where div Z_0 for the second-order problem and $\Delta \operatorname{div} Z_0$ for the fourth-order problem are constant on U. In other words, we are interested in the case that the speed of the corresponding flow on U is constant. We rather consider a more general equation with weight for the second-order problem. For later convenience, we call any continuous function $\chi : \partial U \to \{-1, 1\}$ a signature of U as in [LMM] or [GGP]. We first consider the second-order problem corresponding to $\partial_{L_t^2}^0 TV_a(u)$.

Definition 3.1. Let U be a smooth open set in \mathbb{R}^n with signature χ . Assume that $a \ge 0$ is continuous up to \overline{U} and that $b \in L^{\infty}(U)$ with $1/b \in L^{\infty}(U)$ for b > 0. We say that U is (a, b)-calibrable (with signature χ) if there exists Z_0 satisfying the constraint

$$|Z_0| \leq 1$$
 on U

with boundary condition

 $[Z_0 \cdot \nu] = \chi \quad on \quad \partial U$

with the property that $\frac{1}{b} \operatorname{div}(aZ_0)$ is constant on U. We call any such Z_0 an (a,b)-calibration for U (with signature χ).

Note that in the case $b \neq 1$, we still have the characterization of subdifferential $\partial_{L_b^2} TV_a(u)$ as in Theorem 2.7 (i) with the modification that $v = -(\operatorname{div}(aZ))/b$ instead of (ib) and $-(u, \operatorname{div}(aZ))_{L^2} = TV_a(u)$ instead of (ic). When $b \equiv 1$, the proof is given in [Mol] (for general weighted anisotropic total variation), including the case when a is discontinuous. However, as mentioned before, our definition of TV_a for discontinuous a is different from theirs. The extension to general b is straightforward. In the one-dimensional case, this type of characterization is given in [GGK] when u is piecewise linear.

Our calibration Z_0 gives a minimizer of

$$\left\{ \int_{U} b \left| \frac{\operatorname{div}(aZ)}{b} \right|^{2} dx \mid |Z| \leq 1 \text{ a.e. in } U, \ [Z \cdot \nu] = \chi \text{ on } \partial U, \ \operatorname{div}(aZ) \in L^{2}(U) \right\}.$$

Indeed, we have by integration by parts and the Schwarz inequality

$$\int_{\partial U} a\chi \ d\mathcal{H}^{n-1} = \int_{U} \operatorname{div}(aZ) \ dx = \int_{U} \frac{\operatorname{div}(aZ)}{b^{1/2}} b^{1/2} \ dx \le \left(\int_{U} \frac{\left(\operatorname{div}(aZ)\right)^{2}}{b} \ dx\right)^{1/2} \left(\int_{U} b \ dx\right)^{1/2}.$$

For $Z = Z_0$, since div $(aZ_0)/b$ is constant, we see div $(aZ)/b^{1/2}$ and $b^{1/2}$ is parallel. In this case, the Schwarz inequality becomes equality. Thus,

$$\int_{U} b \left| \frac{\operatorname{div}(aZ)}{b} \right|^{2} dx$$

attains its minimum value

$$\left(\int_{\partial U} a\chi \ d\mathcal{H}^{n-1}\right)^2 \bigg/ \int_U b \ dx$$

at $Z = Z_0$.

In the one-dimensional case, it is known that an interval (with signature χ) is (a, b)-calibrable if and only if a is concave with respect to a metric induced by b [GGK]. In particular, all intervals are (a, b)-calibrable for constant a, b > 0. In higher-dimension case, the situation is more involved even for constant a and b. Even if U is convex, it is not necessarily (1, 1)-calibrable.

There is a very related notion called a Cheeger set. We only discuss the case when $\chi \equiv 1$. For a given open set, we set

$$h(U) := \inf \left\{ \frac{P(F)}{\mathcal{L}^n(F)} \mid F \subset U \text{ Borel}, \, \mathcal{L}^n(F) \in (0,\infty) \right\},\,$$

where $P(F) = TV(\mathbf{1}_F)$ is the perimeter of F and \mathcal{L}^n denotes the Lebesgue measure. Here $\mathbf{1}_F$ denotes the characteristic function of F, i.e., $\mathbf{1}_F(x) = 1$ of $x \in F$ and $\mathbf{1}_F(x) = 0$ if $x \notin F$. This quantity h(U)is called the *Cheeger constant* and $P(F)/\mathcal{L}^n(F)$ is called the *Cheeger ratio*. A set $F \subset U$ satisfying $P(F)/\mathcal{L}^n(F) = h(U)$ is a Cheeger set. If U itself is a Cheeger set, U is called self-Cheeger. It is interesting to find the value h or characterize the Cheeger subset of U. Such problems are often called the Cheeger problem; see e.g. [Leo].

We consider the total variation flow (2.4) with $a \equiv b \equiv 1$ and $\Omega = \mathbb{R}^n$ or \mathbb{T}^n . If U with signature $\chi \equiv 1$ is (1, 1)-calibrable, then the speed of $u = 1_U$ is equal to the Cheeger ratio on U at least formally. Indeed, let Z_0 be a calibration. Then, by integration by parts, we see

$$P(U) = \int_{\partial U} \chi \ \mathcal{H}^{n-1} = \int_U \operatorname{div} Z_0 \ dx = \mathcal{L}^n(U) \cdot \operatorname{div} Z_0.$$

Thus, the speed

 $u_t = \operatorname{div} Z_0$

equals $P(U)/\mathcal{L}^n(U)$ on U.

It is not difficult to prove that if U is (1, 1)-calibrable, it is self-Cheeger; see e.g. [GP]. The converse is also true at least in n = 2. For more details, the reader is referred to [ACC] or [ACM] as well as [GP]. For weighted case, the speed of $u = \mathbf{1}_U$ on U should be the weighted Cheeger ratio

$$\int_{\partial U} a\chi d\mathcal{H}^{n-1} \bigg/ \int_{U} b \, dx$$

if U with signature χ is (a, b)-calibrable.

We next consider the fourth-order problem.

Definition 3.2 ([GKL]). Let U be a smooth open set in \mathbb{R}^n with signature χ . We say that U is (D^{-1}) -calibrable (with signature χ) if there exists Z_0 satisfying the constraint

$$|Z_0| \leq 1$$
 on U

with boundary conditions

$$[Z_0 \cdot \nu] = \chi, \quad \operatorname{div} Z_0 = \chi \kappa \quad on \quad \partial U,$$

with the property that

 $\Delta \operatorname{div} Z_0$ is constant over U.

We call any such Z_0 a (D^{-1}) -calibration for U (with signature χ).

If U is bounded, the constant $\lambda = -\Delta \operatorname{div} Z_0$ can be characterized as a solution of the Saint-Venent problem (or the torsion problem):

$$\begin{cases} -\Delta w = \lambda & \text{in } U \\ w = \chi \kappa & \text{on } U \end{cases}$$
(3.1)

by setting $w = \operatorname{div} Z_0$. The constant should be determined by the other boundary condition:

$$\int_{U} w \, dx = \int_{\partial U} \chi \, d\mathcal{H}^{n-1}. \tag{3.2}$$

In the second-order problem, the constant $\operatorname{div}(aZ_0)/b$ is determined by the Cheeger ratio. For the fourth-order problem, it is determined by using the solution of the Saint-Venant problem

$$\begin{cases} -\Delta w_{\rm sv} = 1 & \text{in } U \\ w_{\rm sv} = 0 & \text{on } \partial U \end{cases}$$
(3.3)

Proposition 3.3 ([GKL]). Let U be a smooth bounded domain in \mathbb{R}^n . Assume that Z is a calibration for U with signature χ . If

$$\lambda = -\Delta \operatorname{div} Z$$

then

$$\lambda = \left(\int_{\partial U} \chi \kappa \nu \cdot \nabla w_{\rm sv} \, d\mathcal{H}^{n-1} + \int_{\partial U} \chi \, d\mathcal{H}^{n-1} \right) \, \Big/ \, \int_{U} w_{\rm sv} \, dx$$

Proof. We first note that $w_{sv} > 0$ by the maximum principle, so the denominator is not zero. We set w = div Z and decompose

$$w = \lambda w_{\rm sv} + h.$$

Since w solves (3.1), h is a harmonic extension of $\chi\kappa$. The condition (3.2) gives

$$\chi \int_U w_{\rm sv} \, dx + \int_U h \, dx = \int_{\partial U} \chi \, d\mathcal{H}^{n-1}.$$

Thus

$$\lambda = \left(\int_{\partial U} \chi \ d\mathcal{H}^{n-1} - \int_{U} h \ dx \right) \Big/ \int_{U} w_{\rm sv} \ dx.$$

It remains to prove

$$-\int_{U} h \, dx = \int_{\partial U} \chi \kappa \nu \cdot \nabla w_{\rm sv} \, d\mathcal{H}^{u-1}$$

Indeed,

$$-\int_{U} h \, dx = \int_{U} h \Delta w_{\rm sv} \, dx = \int_{\partial U} \nu \cdot \nabla w_{\rm sv} \chi \, {\rm div} \, \nu \, d\mathcal{H}^{n-1} + \int_{U} \nabla w_{\rm sv} \cdot \nabla h \, dx$$
$$= \int_{\partial U} \nu \cdot \nabla w_{\rm sv} \chi \kappa \, d\mathcal{H}^{n-1},$$

since div $\nu = \kappa$ and

$$\int_{U} \nabla w_{\rm sv} \cdot \nabla h \, dx = \int_{\partial U} w_{\rm sv} \nu \cdot \nabla h \, d\mathcal{H}^{n-1} - \int_{U} w_{\rm sv} \Delta h \, dx = 0 - 0$$

by the definition of w_{sv} and h.

As for the second-order problem, the calibration gives a minimizer of

$$\left\{ \int_{U} |\nabla \operatorname{div} Z|^2 \, dx \, \middle| \, |Z| \le 1 \text{ a.e. in } U, \, [Z \cdot \nu] = \chi \text{ on } \partial U, \, \operatorname{div} Z = \chi \kappa \text{ on } \partial U \right\}.$$
(3.4)

(Although a minimizer Z^* is not unique, $\Delta \operatorname{div} Z^*$ is uniquely determined.)

Theorem 3.4 ([GKL]). Let U be a smooth bounded domain in \mathbb{R}^n . If Z_0 is a calibration of U with signature χ , then Z_0 is a minimizer of (3.4).

Compared with the second-order problem, the proof is more involved but it is still not difficult [GKL, Theorem 26].

We conclude this section by giving examples of radial calibrable sets.

The second-order problem. We conclude the case $a \equiv b \equiv 1$.

Theorem 3.5. (i) All balls are ((1,1))-calibrable.

- (ii) All complement of the ball are calibrable and $\lambda = 0$.
- (iii) All annuli are calibrable.

It is not difficult to find (radial) calibration Z(x) = z(r)x/r, r = |x|. In the case of an open ball B_R of radius R (centered at the origin), we take

$$z(r) = \frac{\chi r}{R}$$

so that div $Z = z'(r) + \frac{n-1}{r}z(r) = \chi n/R$ and $z(R) = \chi$, $|z| \leq 1$ on (0, R). For the complement of a ball $\overline{B_R}$, it suffices to take $z(r) = \chi R^{n-1}/r^{n-1}$ so that div Z = 0. For an annulus $A_{R_0}^{R_1} = B_{R_1} \setminus \overline{B_{R_0}}$, we have to find z such that $z' + \frac{n-1}{r}z = r^{1-n}(r^{n-1}z)' = \lambda$ satisfying $z(R_1) = \chi_1$, $z(R_0) = -\chi_0$ and $|z(r)| \leq 1$ for $r \in (R_0, R_1)$ where $\chi_i = \chi|_{|x|=R_i}$. Integrating the differential equation, we observe that $z(r) = r\lambda/n - c/r^{n-1}$ with some constant c. By an explicit calculation, we are able to take z satisfying the boundary condition as well as the constraint $|z| \leq 1$. For example, consider the case $\chi_0 = \chi_1 = 1$. We may assume that $R_0 = 1$, $R_1 > 1$ by scaling. The boundary conditions read

$$\frac{\lambda}{r} - c = -1, \quad \frac{\lambda}{n}R_1 - \frac{c}{R_1^{n-1}} = 1$$

so that $R_1(-1+c) - cR_1^{1-n} = 1$. In particular, c > 0. We know $\lambda > 0$ since $\chi \equiv 1$. Thus, z(r) is monotone increasing with $z(R_0) = -1$, $z(R_1) = 1$ so it must satisfy the constraint. The case of indefinite signature, e.g. $\chi_0 = -1$, $\chi_1 = 1$ is more involved and c must have a different sign from λ .

The fourth-order problem. We consider D^{-1} -calibrability

Theorem 3.6 ([GKL]). (i) All balls are calibrable for $n \ge 1$.

- (ii) All complements of balls are calibrable and $\lambda = 0$ except n = 2.
- (iii) If n = 2, all complement of balls are not calibrable.
- (iv) All annuli (with definite signature i.e., with $\chi \equiv 1$ or $\chi \equiv -1$) are calibrable except n = 2.
- (v) For n = 2, there is $Q_* > 1$ such that an annulus $A_{R_0}^{R_1}$ (with definite signature) is calibrable if and only if $R_1/R_0 \leq Q_*$.

Note that by taking angular averaging, we see that a D^{-1} -calibration exists if and only if a radial D^{-1} -calibration exists for radially symmetric set [GKL, Lemma 31]. So to assert non-calibrability, it suffices to prove non-existence of radial calibration. Compared with the second-order case, we see that there occurs several exceptional phenomena for two-dimensional setting.

Instead if giving a full proof, we just give a strategy of the proof by studying the case of a ball B_R . The equation $-\Delta \operatorname{div} Z = \lambda$ is a third-order differential equation of the form

$$-r^{1-n}\left(r^{n-1}\left(r^{1-n}(r^{n-1}z)'\right)'\right)' = \lambda$$
(3.5)

for a radial vector field Z(x) = z(r)x/r since div $Z = r^{1-n}(r^{n-1}z)'$. For B_R with $\chi \equiv -1$, the boundary conditions are

$$z(R) = -1, \quad x'(R) = 0 \tag{3.6}$$

since div $Z = \kappa \chi$ is equivalent to saying that z'(R) + (n-1)z(R)/R = (-1)(n-1)/R. A general solution of (3.5) is of the form

$$z(r) = c_0 r^3 + c_1 r^{3-n} + c_2 r + c_3 r^{1-n}, \quad c_0 = -\frac{\lambda}{2n(n+2)}, \quad n \neq 2,$$

$$z(r) = c_0 r^3 + c_1 r \log r + c_2 r + c_3 r^{-1}, \quad c_0 = -\lambda/16, \quad n = 2.$$

We have to find a solution satisfying (3.6) together with the constraint $|z| \leq 1$. The right choice is

$$z(r) = \frac{1}{2} \left(\frac{r}{R}\right)^3 - \frac{3}{2}\frac{r}{R}, \quad \lambda = -\frac{n(n+2)}{R^3}.$$

(The possibly singular term should be neglected so that we take $c_1 = c_3 = 0$. We determine c_2 , λ by (3.6).)

4 Some simple explicit solutions

As a simple example, we consider a solution starting from $u_0 = a_0 \mathbf{1}_{B_{R_0}}$.

For the second-order problem (2.4) with $a \equiv b \equiv 1$ or (1.1), the answer is very simple since B_{R_0} and $\mathbb{R}^n \setminus \overline{B_{R_0}}$ are calibrable according to Theorem 3.5. Let Z_0^{in} , Z_0^{out} be calibrations of B_{R_0} and $\mathbb{R}^n \setminus \overline{B_{R_0}}$ with signatures $-\operatorname{sgn} a_0$, $\operatorname{sgn} a_0$ respectively. Since the normal trace of Z_0^{in} , Z_0^{out} at ∂B_{R_0}

is continuous, there is no delta part of div Z, where $Z = Z_0^{\text{in}}$ in B_{R_0} and $Z = Z_0^{\text{out}}$ in $\mathbb{R}^n \setminus \overline{B_{R_0}}$. By definition, the speed $u_t = \text{div } Z$ is constant on B_{R_0} and its outside, so it is rather clear that the solution to (1.1) starting from u_0 is of the form

$$u(x,t) = (\operatorname{sgn} a_0) \left(|a_0| - \frac{n}{R_0} t \right)_+ \mathbf{1}_{B_{R_0}}, \quad c_+ = \max(c,0).$$

The number n/R_0 is the Cheeger ratio of B_R in \mathbb{R}^n . Note that the speed div Z outside B_{R_0} equals zero.

For the fourth-order problem, (2.6) the answer is more involved. Different from the second-order problem, for the minimal Cahn–Hoffman vector field Z_0 , $\nabla \operatorname{div} Z_0$ may have jump discontinuity on ∂B_0 so the velocity $-\Delta \operatorname{div} Z_0$ may have a non-zero singular part concentrated on ∂B_0 . In one-dimensional periodic setting, this phenomenon is already observed in [GG], [Ka1], [Ka2]. As a consequence, even if the set K of $\mathbf{1}_K$ is calibrable together with its complement, it may expand or shrink during evolution.

We shall discuss the case $n \neq 2$. We consider $B_{R(t)}$ whose radius R(t) may depend on time. Since $B_{R(t)}$ and its complement are calibrable by Theorem 3.6, we take radial calibration Z_0^{in} in $B_{R(t)}$ and Z_0^{out} in $\mathbb{R}^n \setminus B_{R(t)}$. We set

$$Z(x,t) = \begin{cases} Z_0^{\text{in}}(x), & x \in B_{R(t)} \\ Z_0^{\text{out}}(x), & \mathbb{R}^n \backslash B_{R(t)}. \end{cases}$$

As we already observed,

$$Z_0^{\text{in}}(x) = z^{\text{in}}(r)x/r, \quad z^{\text{in}}(r) = \frac{1}{2}\left(\frac{r}{R}\right)^3 - \frac{3}{2}\frac{r}{R}.$$

Similarly,

$$Z_0^{\text{out}}(x) = z^{\text{out}}(r)x/r, \quad z^{\text{out}}(r) = -\frac{n-1}{2}\left(\frac{r}{R}\right)^{3-n} + \frac{n-3}{2}\left(\frac{r}{R}\right)^{1-n}$$

This vector field Z fulfills all requirements in Definition 2.10. Although div Z is continuous across $\partial B_{R(t)}$, $\nabla \operatorname{div} Z$ may jump across $\partial B_{R(t)}$. Thus,

$$-\Delta \operatorname{div} Z = \lambda \mathbf{1}_{B_{R(t)}} + \nu \cdot \left(\nabla \operatorname{div} Z_0^{\operatorname{in}} - \nabla \operatorname{div} Z_0^{\operatorname{out}}\right) \delta_{\partial B(R(t))}$$

with $\lambda = -n(n+2)/R^3$, where ν is the exterior unit normal of $B_{R(t)}$. A direct calculation shows that

$$\nu \cdot \left(\nabla \operatorname{div} Z_0^{\operatorname{in}} - \nabla \operatorname{div} Z_0^{\operatorname{out}}\right) = -\frac{n(n-4)}{R^2}$$

If we set

$$u(x,t) = a(t)\mathbf{1}_{B_{R(t)}},$$

then $\partial_t u = \frac{da}{dt} \mathbf{1}_{B_{R(t)}} + a \frac{dR}{dt} \delta_{\partial B_{R(t)}}$. Since $u_t = -\Delta \operatorname{div} Z$, we end up with

$$\frac{da}{dt} = -\frac{n(n+2)}{R^3}, \quad \frac{dR}{dt} = -\frac{n(n-4)}{aR^2}$$

This is easy to solve. Indeed,

$$\frac{d}{dt}(aR^3) = -n(n+2) - 3n(n-4) = -n(4n-10).$$

Thus an explicit solution is given as

$$a(t) = a_0 \left(1 - \frac{n(4n-10)}{a_0 R_0^3} t \right)^{\frac{n+2}{4n-10}}, \quad R(t) = R_0 \left(1 - \frac{n(4n-10)}{a_0 R_0^3} t \right)^{\frac{n-4}{4n-10}}$$

if we start with $u_0 = a_0 \mathbf{1}_{B_{R_0}}$.

The case n = 2 is more complicated. It was nontrivial to define a solution; see Definition 2.10. Moreover, the complement of a disk is not calibrable. We expect that the solution becomes radially strictly decreasing for t > 0 outside a ball $B_{R(t)}$. If u is radially strictly decreasing outside $B_{R(t)}$, the minimal Cahn-Hoffman vector field must be $Z^{\text{out}}(x) = -\nabla u/|\nabla u| = -x/|x|$ for |x| > R(t). The solution to (2.6) must satisfy

$$u_t = -\Delta \operatorname{div} Z^{\operatorname{out}}$$

provided that $\nabla \operatorname{div} Z_{\operatorname{out}} \in L^2\left(\left(\overline{B_{R(t)}}\right)^c\right)$ with $\left(\overline{B_{R(t)}}\right)^c = \mathbb{R}^n \setminus \overline{B_{R(t)}}$ as in Definition 2.10. We observe that

div $Z^{\text{out}} = -(n-1)/|x|^2$ and $\nabla \operatorname{div} Z^{\text{out}} = (n-1)x/|x|^3$.

If $n \leq 3$, $\nabla \operatorname{div} Z^{\operatorname{out}} \in L^2\left(\left(\overline{B_{R(t)}}\right)^c\right)$ so it must agree with the (minimal) Cahn–Hoffman vector field for |x| > R(t). (For $n \geq 4$, $\nabla \operatorname{div} Z^{\operatorname{out}}$ is not in $L^2\left(\overline{B_{R(t)}}^c\right)$, so it is not a Cahn–Hoffman field. This indicates that if u_0 is radially strictly decreasing, then $u_0 \notin D(\partial_{D^{-1}}TV)$ for $n \geq 4$.) The speed is formally equal to

$$u_t(x,t) = \frac{(n-1)(n-3)}{|x|^3} \quad x \in \mathbb{R}^n \setminus \overline{B_{R(t)}}.$$

For n = 1 and 3, $u_t = 0$ so the part $\overline{B_{R(t)}}^c$ cannot move. This is consistent that the complement of the ball is calibrable for n = 1 and n = 3. For n = 2, the expected form of the solution of (2.6) is

$$u(x,t) = a(t)\mathbf{1}_{B_{R(t)}}(x) + \frac{t}{|x|^3}\mathbf{1}_{\overline{B_{R(t)}}}(x),$$
(4.1)

where

$$\frac{da}{dt} = -\frac{2 \cdot 4}{R^3}, \quad \left(a(t) - \frac{t}{R(t)^3}\right) \frac{dR}{dt} = \frac{2 \cdot 2}{R(t)^2}.$$
(4.2)

As in the case for $n \ge 3$, this system of ODEs provides several qualitative properties of the solution. Let us summarize what we observe.

Theorem 4.1 ([GKL]). Assume that the initial datum u_0 is of the form

$$u_0 = a_0 \mathbf{1}_{B_{R_0}} \quad with \quad a_0 > 0$$

If $n \geq 3$, then the solution u to (2.6) with initial datum u_0 of the form

$$u(x,t) = a(t)\mathbf{1}_{B_R(t)}$$
 for $t < t_* = \frac{a_0 R_0^3}{n(4n-10)}$

and $u(x,t) \equiv 0$ for $t \geq t_*$. Moreover, a(t) is strictly decreasing and $a(t) \downarrow 0$ as $t \uparrow t_*$. The time t_* is called the extinction time.

- (i) R(t) is strictly increasing and $R(t) \uparrow \infty$ as $t \uparrow t_*$ for n = 3.
- (ii) $R(t) \equiv R_0$ for n = 4.



Figure 2: Profile of a solution when n = 2

(iii) R(t) is strictly decreasing and $R(t) \downarrow 0$ as $t \uparrow t_*$ for $n \ge 5$.

If n = 2, the solution is of the form (4.1). The functions a and R satisfy (4.2). In particular, there is no extinction time and R(t) is strictly increasing and a(t) strictly decreasing. Moreover, $R(t) \uparrow \infty$ and $a(t) \downarrow 0$ as $t \to \infty$. The gap $a(t) - t/R(t)^3$ is always positive. See Figure 2.

If n = 1, then the solution is of the form $u(x,t) = a(t)\mathbf{1}_{B_{R(t)}}$ for t > 0. There is no extinction time. Moreover, R(t) is strictly increasing and a(t) strictly decreasing with $R(t) \uparrow \infty$, $a(t) \downarrow 0$ as $t \to \infty$.

The reason why there is no finite extinction time for $n \leq 2$ is related to the fact that 0 is not an element of the affine space $u_0 + D^{-1}$ where the flow lives if $\int_{\mathbb{R}^n} u_0 \, dx \neq 0$. We shall discuss finite extinction properties in the next section.

Similar analysis can be carried out for more general radially symmetric data. In the case of the second order problem, as we have checked in Theorem 3.5, any annulus, with any choice of signature, is calibrable. Thus, by pasting together the calibrations for the annuli $A_{R_{k-1}}^{R_k}$, ball B_{R_0} and ball exterior $\mathbb{R}^n \setminus B_{R_{m-1}}$, we can construct a Cahn-Hoffman vector field Z_0 for any piecewise constant, radially symmetric initial datum (called a *stack*), i.e.

$$u_0 = a_0^0 \mathbf{1}_{B_{R_0}} + \sum_{k=1}^{m-1} a_0^k \mathbf{1}_{A_{R_{k-1}}}^{R_k} + a_0^m \mathbf{1}_{\mathbb{R}^n \setminus B_{R_{m-1}}}$$
(4.3)

with $0 < R_0 < R_1 < \cdots < R_{m-1}$, $a_0^k \in \mathbb{R}$ $(k = 0, \ldots, m)$. Since the L^2 function div Z_0 is constant on $A_{R_{k-1}}^{R_k}$, B_{R_0} and $\mathbb{R}^n \setminus B_{R_{m-1}}$, this shows that for small t > 0

$$u(t) = a^{0}(t)\mathbf{1}_{B_{R_{0}}} + \sum_{k=1}^{m-1} a^{k}(t)\mathbf{1}_{A_{R_{k-1}}^{R_{k}}} + a^{m}(t)\mathbf{1}_{\mathbb{R}^{n}\setminus B_{R_{m-1}}},$$
(4.4)

with da^k/dt constant, depending only on R_{k-1} , R_k and the signs of $a_0^{k+1} - a_0^k$ and $a_0^k - a_0^{k-1}$ for $k = 1, \ldots, m-1$ (da^0/dt and da^m/dt are also constant, depending only on R_0 , $\operatorname{sgn}(a^1 - a^0)$, and R_m , $\operatorname{sgn}(a^m - a^{m-1})$, respectively. This determines the evolution until the first time instance t > 0 such that $a^k(t) = a^{k+1}(t)$ for some $k \in \{0, \ldots, m-1\}$ (first merging time). Then, the solution is again of form (4.4), with a smaller m, and one can repeat the procedure until the solution becomes constant.

In [GKL], the evolution of stacks under the fourth-order TV flow was studied in detail. In particular, it has been proved that in dimensions $n \neq 2$, if the initial datum is of form (4.3), i.e. if u_0 is a stack, then u(t) is also a stack for t > 0. The same does not hold in n = 2 as evidenced already by evolution of characteristic functions of balls.

In the second-order case, one can produce more complicated examples of explicit piecewise constant solutions with initial data such as characteristic functions of sums of calibrable sets which are distant enough from each other [BCN]. In the fourth-order case, we do not know about any non-radial examples of explicit solutions in $n \ge 2$.

In the case of bounded domains, the solutions can be more complicated. However, in the 1D setting (where, to be fair, the only connected bounded domains are intervals) one can produce explicit solutions for a dense set of initial data: the step functions. Let us present the construction in the case of periodic boundary condition, i.e. $\Omega = \mathbb{R}/\mathbb{Z}$, for the second-order flow. Take u_0 a step function, that is

$$u_0 = a_0^1 \mathbf{1}_{[x_0, x_1)} + a_0^2 \mathbf{1}_{[x_1, x_2)} + \ldots + a_0^m \mathbf{1}_{[x_{m-1}, x_m)},$$
(4.5)

where $x_m = x_0 + 1$. We can assume that $a_0^{k-1} \neq a_0^k$ for k = 1, ..., m and $a_0^m \neq a_0^1$. Since the intervals are calibrable, for small t > 0 (until the first merging time) we have

$$u(t) = a^{1}(t)\mathbf{1}_{[x_{0},x_{1})} + a^{2}(t)\mathbf{1}_{[x_{1},x_{2})} + \ldots + a^{m}(t)\mathbf{1}_{[x_{m-1},x_{m})}$$
(4.6)

with a^k evolving at constant speed

$$da^{k}/dt = \theta_{k}/(x_{k} - x_{k-1}) \text{ for } k = 1, \dots, m,$$

$$\theta_{k} = \begin{cases} +2 & \text{if } a_{0}^{k} < a_{0}^{k-1} \text{ and } a_{0}^{k} < a_{0}^{k+1}, \\ -2 & \text{if } a_{0}^{k} > a_{0}^{k-1} \text{ and } a_{0}^{k} > a_{0}^{k+1}, \\ 0 & \text{otherwise.} \end{cases}$$

$$(4.7)$$

Then, as in the radially symmetric case, we can show that the solution remains a step function throughout the evolution.

5 Upper bounds for the extinction time

In many examples, the solution may have a finite extinction time. We consider this problem both for the second-order and the fourth-order problem. For an initial datum u_0 , the *extinction time* of a solution u is defined as

$$T^*(u_0) = \inf \left\{ t \in (0, \infty) \mid u(x, \tau) = 0 \text{ for } \tau \ge t \right\}.$$

5.1 Second and fourth-order problems

We consider the second-order problem (1.1). We first give an upper bound for the extinction time given by [GK, Theorem 2.4, Theorem 2.5]. Let S_n be the best constant of the Sobolev (isoperimetric) inequality

$$\|v\|_{n/(n-1)} \le S_n TV(v),$$

where $||v||_p = (\int_{\mathbb{R}^n} |v|^p dx)^{1/p}$.

Theorem 5.1 (Second-order problem). (i) Assume that $n \ge 2$ and $u_0 \in L^2(\mathbb{R}^n)$. Then

$$T^*(u_0) \le S_n ||u_0||_n.$$

(iii) Assume that n = 1. Then $||u||_1(t) \le ||u_0||_1 - t$, where u is the solution to (1.1). In particular,

$$T^*(u_0) \le ||u_0||_1.$$

If $u_0 \in L^2(\mathbb{R})$ does not belong to $L^1(\mathbb{R})$, the solution u may not have finite extinction time, i.e., $T^*(u_0) = \infty$.

Idea of the proof. In the case n = 2. By the abstract definition of a solution, we know that

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^2} |u|^2 dx = (u, u_t)_{L^2} = -(u, \partial_{L^2}TV(u))_{L^2}.$$

Since TV(u) is positively one-homogeneous, we see $(u, \partial_{L^2}TV(u)) = TV(u)$ (see Lemma 2.6). In particular,

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^2} |u|^2 \, dx = -TV(u).$$

We apply the Sobolev inequality to get

$$\frac{1}{2}\frac{d}{dt}\int_{\mathbb{R}^2} |u|^2 \, dx \le -\frac{1}{S_2} \left(\int_{\mathbb{R}^2} |u|^2 \, dx\right)^{1/2},$$

which yields

$$\frac{d}{dt} \|u\|_2(t) \le -S_2^{-1} \text{ provided that } \|u\|_{L^2}(t) \ne 0.$$

This yields

$$||u||_2(t) \le ||u_0||_2 - S_2^{-1}t$$

which implies the desired estimate for $T^*(u_0)$.

In the case $n \ge 3$, we formally multiply $|u|^{n-2}u$ to (1.1) to get

$$\frac{1}{n}\frac{d}{dt}\int_{\mathbb{R}^n}|u|^n\,dx=\int_{\mathbb{R}^n}|u|^{n-2}uu_t\,dx=\int_{\mathbb{R}^n}|u|^{n-2}u\operatorname{div}\left(\nabla u/|\nabla u|\right)dx.$$

Integrating by parts, the right-hand side becomes

$$-(n-1)\int_{\mathbb{R}^n} |u|^{n-2} \nabla u \cdot \nabla u / |\nabla u| = -(n-1)\int_{\mathbb{R}^n} |u|^{n-2} |\nabla u| \, dx = -\int_{\mathbb{R}^n} \left|\nabla |u|^{n-2} u\right| \, dx.$$

Then applying the Sobolev inequality for $v = |u|^{n-2}u$ to get

$$\frac{1}{n}\frac{d}{dt}\int_{\mathbb{R}^n}|u|^n\ dx\leq -S_n^{-1}\left(\int_{\mathbb{R}^n}|u|^n\ dx\right)^{(n-1)/n}$$

i.e.,

$$\frac{1}{n}\frac{d}{dt}\|u\|_{n}^{n} \le -S_{n}^{-1}\|u\|_{n}^{n-1}.$$

Thus, as for n = 2, we have $||u||_n(t) \leq ||u_0||_n - S_n^{-1}t$ if $||u||_n(t) \neq 0$, and the desired estimate holds. The argument for $n \geq 3$ is formal because we do not know whether multiplication by $|u|^{n-2}u$ is justified since u may not be an L^n -valued absolutely continuous function of t. Fortunately, our argument is justified by approximation of the equation by smooth uniformly parabolic equations as in [GK]. In one-dimensional setting, we approximate $\int |u| dx$ by $\int f(u)$ where f is a convex function; see [GK, Section 2.5]. **Remark 5.2** ([GK]). The results still hold when $\Omega = \mathbb{T}^n$ or a smooth bounded domain with the Neumann boundary condition for average-free L^2 initial data with possibly different value of the best Sobolev constant S_n . Note that S_n may depend on the shape of Ω but it is scale invariant in the sense that S_n is invariant under dilation, i.e., $S_n(\lambda\Omega) = S_n(\Omega)$ for any $\lambda > 0$, where $\lambda\Omega = \{\lambda x \mid x \in \Omega\}$. For the Dirichlet problem, it still holds for n = 2 but there is no literature claiming the estimate $T^*(u_0) \leq S_n ||u_0||_n$ for $n \geq 3$; justification of the formal estimate will be difficult since boundary detachment phenomenon is expected unless the domain is mean-convex.

For the gradient flow of p-Dirichlet energy for p > 1, i.e., $u_t \in -\partial E_p(u)$, an extinction time estimate

$$T^*(u_0) \le C \|u_0\|_s^{2-p}, \quad s = n(2-p)/p, \quad n \ge 2, \quad 1$$

has been proved by a similar method [DiB, Proposition 2.1, Proposition 3.1 of Chapter VII] both for the Cauchy problem $\Omega = \mathbb{R}^n$ and the Dirichlet problems for a bounded domain.

Remark 5.3. Since (1.1) has a comparison principle, the estimate $T^*(u_0) < \infty$ is often proved by comparison with the evolution of the characteristic function $a(t)\mathbf{1}_{B_{R_0}}$. For example, suppose that $u_0 \geq 0$ and $u_0 \leq a_0$ and $u_0 \equiv 0$ outside B_{R_0} . Then

$$0 \le u(x,t) \le a(t)$$
 in B_{R_0}

and $a(t) = (a_0 - (n/R_0)t)_+$. Thus $T^*(u_0) \le (\sup u_0)R_0/n$.

We next study the fourth-order problem. We first calculate the growth of L^p -norm in a formal way

$$\frac{1}{p(p-1)} \frac{d}{dt} \int_{\mathbb{R}^n} |u|^p \, dx = \frac{1}{p-1} \int_{\mathbb{R}^n} |u|^{p-2} u u_t \, dx$$
$$= \frac{1}{p-1} \int_{\mathbb{R}^n} |u|^{p-2} u (-\Delta \operatorname{div} z) \, dx \quad \text{with} \quad z = \nabla u / |\nabla u|$$

since (2.6) is of the form $u_t = -\Delta \operatorname{div} z$. Integrating by parts, the right-hand side equals

$$\int_{\mathbb{R}^n} |u|^{p-2} \nabla u \cdot \nabla \operatorname{div} z \, dx = -\int_{\mathbb{R}^n} |u|^{p-2} \nabla^2 u : \nabla \otimes z \, dx - (p-2) \int_{\mathbb{R}^n} |u|^{p-4} u \nabla \otimes z : \nabla u \otimes \nabla u \, dx,$$

where we assume that effect at space infinity does not appear; here $A : B = \text{trace}(AB^T)$ for matrices A and B and $\nabla \otimes z$ for a vector field $z = (z_1, \ldots, z_n)$ denotes a matrix $(\partial_i z_j)$ for $\partial_i = \partial/\partial x_i$. Since z is a subgradient of a positively one-homogeneous function of ∇u we see that $(\nabla \otimes z)(\nabla u)^T = 0$ for $\nabla u = (\partial_1 u, \ldots, \partial_n u)$ from the Euler's identity for a positively zero-homogeneous function. Here is a more explicit argument. Since

$$\nabla \otimes z = \nabla \otimes \frac{\nabla u}{|\nabla u|} = \frac{1}{|\nabla u|} \nabla^2 u \left(I - \frac{\nabla u}{|\nabla u|} \otimes \frac{\nabla u}{|\nabla u|} \right)$$

it is rather clear that $(\nabla \otimes z)(\nabla u)^T = 0$ since $P = I - \nabla u \otimes \nabla u / |\nabla u|^2$ is a projection orthogonal to ∇u . We end up with

$$\frac{1}{p(p-1)}\frac{d}{dt}\int_{\mathbb{R}^n}|u|^p\ dx = -\int_{\mathbb{R}^n}|u|^{p-2}\nabla^2 u: \nabla\otimes z\ dx \le 0$$

since $|\nabla u| \nabla \otimes z = \nabla^2 u P$ and P is a non-negative symmetric matrix. This is a formal argument that needs to be rigorously justified, see [GKL2]. Eventually, we have

Lemma 5.4. Assume that $u_0 \in L^p(\mathbb{R}^n) \cap E^{-1}$. Let u be the solution of (2.6) with initial datum u_0 . Then $||u||_p(t)$ is non-increasing in t for $1 \leq p < \infty$. In particular, $||u||_p(t) \leq ||u_0||_p$ for all $t \geq 0$.

We now discuss an upper bound for the extinction time. Since the extinction time for $n \leq 2$ may be infinite as observed in Section 4, we assume $n \geq 3$. In this case, our fundamental identity

$$\frac{1}{2}\frac{d}{dt}\|u\|_{D^{-1}}^2(t) = -TV(u)$$

is obtained by taking inner product of u with (2.6). By the Sobolev inequality, we see that

$$\frac{1}{2}\frac{d}{dt}\|u\|_{D^{-1}}^2(t) \le -S_n^{-1}\|u\|_{n/(n-1)}$$
(5.1)

as for the second-order problem. Again by the Sobolev inequality

$$||u||_{2^*} \le C_n ||u||_{D^1_0}$$

for $2^* = 2n/(n-2)$, hence

$$\|u\|_{D^{-1}} = \sup\left\{ \langle u, v \rangle \mid \|v\|_{D_0^1} \le 1 \right\}$$

$$\leq C_n \sup\left\{ \langle u, v \rangle \mid \|v\|_{2^*} \le 1 \right\} = C_n \|u\|_{(2^*)'}$$
(5.2)

where $(2^*)' = 2n/(n+2)$. In the case n = 4, $(2^*)' = 4/3 = n/(n-1)$ so (5.1) yields

$$\frac{1}{2}\frac{d}{dt}\|u\|_{D^{-1}}^2(t) \le -S_4^{-1}C_4^{-1}\|u\|_{D^{-1}}(t)$$

and this implies

$$T^*(u_0) \le S_4 C_4 ||u||_{D_0^1}.$$

This type of estimate is already obtained in [GK] when $\Omega = \mathbb{T}^n$. For other *n*, we recall the Hölder inequality

$$\|u\|_{(2^*)'} \le \|u\|_{n/(n-1)}^{\theta} \|u\|_p^{1-\theta}$$

with $\frac{n+2}{2n} = \frac{n-1}{n}\theta + \frac{1}{p}(1-\theta)$ for $n \ge 3$. If n = 4, $(2^*)' = 4/3 = n/(n-1)$ so $\theta = 1$. If $n \ge 5$ we have to take $p > (2^*)'$, and if n = 3 we have to take $p < 6/5 = (2^*)'$ so that $0 < \theta < 1$. By (5.1) and (5.2), we now obtain

$$\frac{1}{2} \frac{d}{dt} \|u\|_{D^{-1}}^{2}(t) \leq -S_{n}^{-1} \left(C_{n}^{-1} \|u\|_{D^{-1}}\right)^{1/\theta} / \|u\|_{p}^{(1-\theta)/\theta}$$
$$\leq -S_{n}^{-1} \left(C_{n}^{-1} \|u\|_{D^{-1}}\right)^{1/\theta} / \|u_{0}\|_{p}^{(1-\theta)/\theta}.$$

In the last inequality, we invoke Lemma 5.4. In other words,

$$\frac{d}{dt} \|u(t)\|_{D^{-1}} \le -A_{\theta}^{-1} \|u(t)\|_{D^{-1}}^{1/\theta-1} / \|u_0\|_p^{1/\theta-1}, \quad A_{\theta} = S_n C_n^{1/\theta}.$$

The differential inequality

$$\frac{dy}{dt} \le -ky^{\sigma}, \quad y(0) = y_0 > 0,$$

with $\sigma \in [0, 1), k > 0$ yields the estimate

$$y(t) \le (y_0^{1-\sigma} - (1-\sigma)kt)_+^{\frac{1}{1-\sigma}}.$$

In particular, y(t) must be zero for $t \ge t_* = y_0^{1-\sigma} / (1-\sigma)k$. Applying this estimate to our differential inequality for $y(t) = ||u(t)||_{D^{-1}}$ implies that

$$\|u(t)\|_{D^{-1}} \le \left(\|u_0\|_{D_0}^{2-1/\theta} - (2-1/\theta)A_{\theta}^{-1}t \ / \ \|u_0\|_p^{(1-\theta)/\theta}\right)_+^{\theta/(2\theta-1)}$$

provided that $0 \leq \frac{1}{\theta} - 1 < 1$, which is equivalent to

$$\frac{1}{2} < \theta = \left(\frac{n+2}{2n} - \frac{1}{p}\right) \left/ \left(\frac{n-1}{n} - \frac{1}{p}\right) \le 1.$$
(5.3)

In this case, we have an upper bound for the extinction time

$$T^*(u_0) \le \frac{A_{\theta}\theta}{2\theta - 1} \|u_0\|_p^{1/\theta - 1} \|u_0\|_{D^{-1}}^{2 - 1/\theta}.$$
(5.4)

It remains to check the validity of the inequality (5.3). If n = 4 so that (n + 2)/2n = (n - 1)/n, θ must be 1 and (5.4) is reduced to what we already obtained. Since $\theta > 1/2$ in (5.3) can be written as

$$\frac{n+2}{2n} - \frac{n-1}{2n} > \frac{1}{2p}$$
, i.e., $\frac{1}{p} < \frac{3}{n}$,

the estimate (5.4) holds for all p > n/3 provided that $n \ge 5$. In the case n = 3, (5.3) yields that 1 . Summarizing what we discussed, we obtain at least formally

Theorem 5.5. Let p > n/3 for n = 5 and $p \in (1, 6/5)$ for n = 3. Assume that

$$\theta = \left(\frac{n+2}{2n} - \frac{1}{p}\right) \left/ \left(\frac{n-1}{n} - \frac{1}{p}\right)\right.$$

for $n \geq 3$ and $n \neq 4$. Then

$$T^*(u_0) \le \frac{A_{\theta}\theta}{2\theta - 1} \|u_0\|_p^{1/\theta - 1} \|u_0\|_{D^{-1}}^{2-1/\theta}$$

with $A_{\theta} = S_n C_n^{1/\theta}$ provided that $u_0 \in L^p(\mathbb{R}^n) \cap D^{-1}(\mathbb{R}^n)$. In the case n = 4,

 $T^*(u_0) \le A_1 \|u_0\|_{D^{-1}}.$

This result is consistent with our explicit solutions in Section 4. Our examples in Section 4 for $n \leq 2$ shows that Theorem 5.5 cannot be extended to $n \leq 2$. Note that our estimate is scale-invariant.

The results easily extend to the case $\Omega = \mathbb{T}^n$ by considering average free spaces. For a smooth bounded domain Ω , there are several possible boundary conditions

(DD) Dirichlet–Dirichlet: u = 0, div $\left(\frac{\nabla u}{|\nabla u|}\right) = 0$ on $\partial \Omega$;

(ND) Neumann–Dirichlet: $\frac{\partial u}{\partial \nu} = 0$, div $\left(\frac{\nabla u}{|\nabla u|}\right) = 0$ on $\partial \Omega$;

(DN) Dirichlet–Neumann: u = 0, $\frac{\partial}{\partial \nu} \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) = 0$ on $\partial \Omega$;

(NN) Neumann–Neumann: $\frac{\partial u}{\partial \nu} = 0$, $\frac{\partial}{\partial \nu} \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) = 0$ on $\partial \Omega$.

In (DD) and (DN) case, as well as in the (NN) case if Ω is convex, the boundary terms in the calculation of $\frac{d}{dt} \int_{\Omega} |u|^p dx$ vanish, so we still expect monotonicity of $||u||_{L^p(\Omega)}$. Note that the formulation of flow for (DN) and (NN) itself is non-trivial. We shall discuss the formulation in the forthcoming paper [GKL2]. In [GK], an upper bound for the extinction time in a periodic domain is obtained using a Sobolev space of negative order, and it is extended in [GKM] to a bounded domain under (DD). In these settings, even in $n \leq 2$ the solution has a finite extinction time. This is a big difference between \mathbb{R}^n case and a bounded domain (or \mathbb{T}^n).

5.2 Fractional case

We next consider the fractional case (2.5) with constant weight, i.e., $a \equiv 1$. In this case, we mimic the way to derive an upper bound for the extinction time by using negative Sobolev norm in the case s = 1 discussed in [GK] and [GKL], since it is not clear whether L^p -norms of the solution are well controlled or not.

We begin with a fundamental identity.

Proposition 5.6. Let u be a solution to (2.5) (with $a \equiv 1$) with initial datum $u_0 \in \dot{H}^{-s}_{av}(\mathbb{T}^n)$. Then

$$\frac{1}{2}\frac{d}{dt}\|u(t)\|^2_{\dot{H}^{-s}_{\rm av}(\mathbb{T}^n)} = -TV(u(t)) \quad for \ a.e. \quad t > 0.$$

This is easily obtained from $u_t \in -\partial_{\dot{H}_{av}^{-s}} TV(u)$ by taking \dot{H}^{-s} inner product with u.

The basic strategy is the same as in Section 5.1. We shall estimate the right-hand side by an interpolation inequality:

$$-TV(u) \le -||u||_{\dot{H}^{-s}_{av}}^{1+\alpha} / ||u||_X^{\alpha}$$

with a suitable norm $||u||_X$, which does not grow quickly as t increases. In Section 5.1, we take $||u||_X$ just L^p -norm since $||u(t)||_p$ is not increasing. In this section, we instead take a negative Sobolev-norm.

We begin with a simple setting when n = 2(s+1), where an interpolation inequality is unnecessary. This corresponds to the case n = 4 for s = 1. We recall a fractional Sobolev inequality

$$||u||_{L^q} \le C_{n,s} ||u||_{\dot{H}^s_{\mathrm{av}}} \quad \text{for} \quad u \in \dot{H}^s_{\mathrm{av}}(\mathbb{T}^n),$$

where $\frac{1}{q} = \frac{1}{2} - \frac{s}{n}$ for s > 0, $q < \infty$. As discussed when deriving (5.2) for D^{-1} , by duality we observe that

$$||u||_{\dot{H}_{av}^{-s}} \le C_{n,s} ||u||_{L^{q'}}, \quad 1/q + 1/q' = 1.$$

If s > 0 satisfies n = 2(s + 1), then q = n and q' = n/(n - 1). Since $||u||_{L^{n/(n-1)}} \leq S_n TV(u)$ by the Sobolev (isoperimetric) inequality, Proposition 5.6 implies that

$$\frac{1}{2}\frac{d}{dt}\|u\|_{\dot{H}^{-s}_{\mathrm{av}}(\mathbb{T}^n)}^2(t) \le -C_{n,s}^{-1}S_n^{-1}\|u\|_{\dot{H}^{-s}_{\mathrm{av}}(\mathbb{T}^n)}.$$

We thus obtain an upper bound for the extinction time.

Proposition 5.7. Let u be a solution to (2.5) (with $a \equiv 1$) with initial datum $u_0 \in \dot{H}^{-s}_{av}(\mathbb{T}^n)$. Assume that n = 2(s+1). Then,

$$T^*(u_0) \le C_{n,s} S_n \|u_0\|_{\dot{H}^{-s}_{av}(\mathbb{T}^n)}.$$

We next derive an interpolation inequality, which is an extension of the inequality obtained in [GK] for s = 1. We define a homogeneous negative Sobolev norm as

$$\|w\|_{\dot{W}_{\mathrm{av}}^{-1,p}} = \sup\left\{ \langle w, \varphi \rangle \ \Big| \ \varphi \in C^{1}_{\mathrm{av}}(\mathbb{T}^{n}), \ \|\nabla\varphi\|_{L^{p'}} \le 1 \right\}$$

when w is a distribution on \mathbb{T}^n .

Lemma 5.8. Assume that s, p, θ satisfy

$$1 \le n \le 2(s+1), \quad 0 < s \le 1, \quad 1 \le p \le \infty, \quad \frac{1}{2} \le \theta \le 1$$

and the scaling balance

$$s + \frac{n}{2} = (1 - \theta) \left(2s + 1 + \frac{n}{p} \right) + \theta(n - 1).$$

Then there is a constant C_* such that

$$\|u\|_{\dot{H}^{-s}_{\mathrm{av}}} \leq C_* \left\| (-\Delta)^{-s} u \right\|_{\dot{W}^{-1,p}_{\mathrm{av}}}^{1-\theta} TV(u)^{\theta} \quad for \ all \quad u \in \dot{H}^{-s}_{\mathrm{av}}(\mathbb{T}^n) \cap BV(\mathbb{T}^n).$$

The constant C_* is invariant under dilation in the sense that it is independent of $\lambda > 0$ if u is replaced by $u_{\lambda} (= u(\lambda x))$.

The scaling balance is a consequence of invariance of C_* . It can be rewritten as

$$\theta = \frac{1}{2} + \left\{ 2 + 2p\left(\frac{2(s+1)}{n} - 1\right) \right\}^{-1}.$$

Thus the assumption $\frac{1}{2} \le \theta \le 1$ is redundant since it follows from $n \le 2(s+1)$, $1 \le 1' \le \infty$ and the scaling balance. We note that

$$\begin{split} \theta &= 1 \Longleftrightarrow n = 2(s+1) \\ \theta &= \frac{1}{2} \Longleftrightarrow 1 \leq n < 2(s+1), \quad p = \infty, \end{split}$$

since $s \leq 1$. The idea of the proof is parallel to that of [GK]. We give the proof in the case n < 2(s+1) for the reader's convenience.

Proof. We decompose u into regular part u_{reg} and singular part u_{sing}

$$u = u_{\text{reg}} - u_{\text{sing}}, \quad u_{\text{reg}} := e^{t\Delta}u, \quad u_{\text{sing}} := \int_0^t \Delta e^{\tau\Delta}u \ d\tau$$

based on the formula

$$e^{t\Delta}u - u = \int_0^t \frac{d}{d\tau} e^{\tau\Delta}u \ d\tau = \int_0^t \Delta e^{\tau\Delta}u \ d\tau.$$

(This idea is standard to prove an interpolation inequality as in [GGS, Chapter 6].) Since

$$\|u\|_{\dot{H}^{-s}_{av}}^{2} = \int_{\mathbb{T}^{n}} (-\Delta)^{-s} u \cdot u \, dx = \int_{\mathbb{T}^{n}} (-\Delta)^{-s} u \cdot u_{\text{reg}} \, dx - \int_{\mathbb{T}^{n}} (-\Delta)^{-s} u \cdot u_{\text{sing}} \, dx,$$

we estimate each term separately.

For the regular part, by definition of $\dot{W}_{\rm av}^{-1,p}$ norm, we have

$$\left| \int_{\mathbb{T}^n} (-\Delta)^{-s} u \cdot u_{\text{reg}}(t) \, dx \right| \le \left\| (-\Delta)^{-s} u \right\|_{\dot{W}_{\text{av}}^{-1,p}} \left\| \nabla u_{\text{reg}}(t) \right\|_{L^{p'}}.$$

Invoking $L^{p'}-L^1$ estimate for the heat semigroup, we have

$$\|\nabla u_{\rm reg}(t)\|_{L^{p'}} = \|\nabla e^{t\Delta}u\|_{L^{p'}} \le C_1 t^{-\frac{n}{2}\left(1-\frac{1}{p'}\right)}TV(u) = C_1 t^{-\frac{n}{2p}}TV(u).$$

Thus

$$\left| \int_{\mathbb{T}^n} (-\Delta)^{-s} u \cdot u_{\text{reg}}(t) \, dx \right| \le C_1 t^{-n/2p} \left\| (-\Delta)^{-s} u \right\|_{\dot{W}^{-1,p}_{\text{av}}} TV(u).$$

The estimate for the singular part is more involved. We proceed

$$\begin{split} \left| \int_{\mathbb{T}^n} (-\Delta)^{-s} u \cdot u_{\operatorname{sing}}(t) \, dx \right| &= \left| \int_{\mathbb{T}^n} (-\Delta)^{-s} u \cdot \int_0^t \Delta e^{\tau \Delta} u \, d\tau dx \right| \\ &\leq \int_0^t \left| \int_{\mathbb{T}^n} (-\Delta)^{-s/2} u (-\Delta)^{1-s/2} e^{\tau \Delta} u \, dx \right| d\tau \\ &\leq \int_0^t \left\| (-\Delta)^{-s/2} u \right\|_{L^2} \left\| (-\Delta)^{1-s/2} e^{\tau \Delta} u \right\|_{L^2} \, d\tau \\ &= \left\| u \right\|_{\dot{H}^{-s}_{\operatorname{av}}} \int_0^t \left\| (-\Delta)^{1-s/2} e^{\tau \Delta} u \right\|_{L^2} \, d\tau. \end{split}$$

We note that

$$(-\Delta)^{1-s/2}e^{\tau\Delta}u = -\sum_{j=1}^{n}(-\Delta)^{-s/2}\partial_{j}e^{\tau\Delta/2}e^{\tau\Delta/2}\partial_{j}u.$$

We use L^2 - L^2 estimate for the heat semigroup to get

$$\left\| (-\Delta)^{-s/2} \partial_j e^{\tau \Delta/2} \right\|_{L^2 \to L^2} \le C \tau^{(s-1)/2}.$$

Here we invoked the assumption that $s \leq 1$. (This can be easily proved by the Parseval identity.) Thus,

$$\left\| (-\Delta)^{1-s/2} e^{\tau \Delta} u \right\|_{L^2} \le C \tau^{(s-1)/2} \sum_{j=1}^n \left\| e^{\tau \Delta/2} \partial_j u \right\|_{L^2}.$$

Using L^2 - L^1 estimate for the heat semigroup, we end up with

$$\left\| (-\Delta)^{1-s/2} e^{\tau \Delta} u \right\|_{L^2} \le C' \tau^{(s-1)/2} \tau^{-n/4} \sum_{j=1}^n \|\partial_j u\|_{L^1}$$
$$\le C'' \tau^{(2s-2-n)/4} TV(u).$$

(To be precise, we take an approximate sequence of average-free smooth functions f_k on \mathbb{T}^n so that $f_k \to u$ in L^2 and $TV(f_k) \to TV(u)$.) We thus conclude that

$$\left| \int_{\mathbb{T}^n} (-\Delta)^{-s} u \cdot u_{\text{sing}}(t) \, dx \right| \le C'' \|u\|_{\dot{H}^{-s}_{\text{av}}} \int_0^t \tau^{(2s-2-n)/4} \, d\tau \, TV(u)$$
$$= C_2 \|u\|_{\dot{H}^{-s}_{\text{av}}} TV(u) t^{(2+2s-n)/4}, \quad C_2 = \frac{4}{2s+2-n} C''$$

since (2s-2-n)/4 > -1 by our assumption n < 2(s+1).

Combining the estimate for the regular part and the singular part, we have

$$\|u\|_{\dot{H}^{-s}_{av}}^{2} \leq \left(C_{1}t^{-n/2p} \left\|(-\Delta)^{-s}u\right\|_{\dot{W}^{-1,p}_{av}} + C_{2}t^{(2+2s-n)/4} \|u\|_{\dot{H}^{-s}_{av}}\right) TV(u).$$
(5.5)

We take t so that the two terms in the right-hand side are balanced, i.e.,

$$C_1 t^{-n/2p} \| (-\Delta)^s u \|_{\dot{W}_{av}^{-1,p}} = C_2 t^{(2+2s-n)/4} \| u \|_{\dot{H}_{av}^{-s}},$$

or

$$t^{\beta} = \frac{C_1 \| (-\Delta)^s u \|_{\dot{W}_{av}^{-1,p}}}{C_2 \| u \|_{\dot{H}_{av}^{-s}}}, \quad \beta = \frac{1+s}{2} - \frac{n}{4} + \frac{n}{2p}$$

We fix this t and observe that (5.5) becomes

$$\|u\|_{\dot{H}_{av}^{-s}}^{2} \leq 2C_{1} \left(\frac{C_{1} \|(-\Delta)^{-s} u\|_{\dot{W}_{av}^{-1,p}}}{C_{2} \|u\|_{\dot{H}_{av}^{-s}}} \right)^{-n/2p\beta} \|(-\Delta)^{-s} u\|_{\dot{W}_{av}^{-1,p}} TV(u)$$

or

$$\|u\|_{\dot{H}_{av}^{-s}}^{2-(n/2p\beta)} \le C_3 \|(-\Delta)^{-s}u\|_{\dot{W}_{av}^{-1,p}}^{1-(n/2p\beta)} TV(u)$$

with $C_3 = 2C_1^{1-(n/2p\beta)}C_2^{n/2p\beta}$. If we take

$$\theta := \left(2 - \frac{n}{2p\beta}\right)^{-1},$$

we see that

$$1 - \frac{n}{2p\beta} = \frac{1}{\theta} - 1.$$

Thus

$$\|u\|_{\dot{H}^{-s}_{av}} \le C_* \|(-\Delta)^{-s}u\|_{\dot{W}^{-1,p}_{av}}^{1-\theta} TV(u)^{\theta}$$

with $C_* = C_3^{\theta}$. The definition of $\theta = \left(2 - \frac{n}{2p\beta}\right)^{-1}$ is nothing but the scaling balance. The proof is now complete.

We next prove a mild growth of $||u(t)||_{\dot{W}_{av}^{-1,p}}$ as t grows. For $\mathbb{T}^n = \prod_{i=1}^m (\mathbb{R}/\omega_i\mathbb{Z})$, we set $|\mathbb{T}^n| = \omega_i \cdots \omega_n$, which is the volume of a fundamental domain.

Lemma 5.9. Let u be a solution to (2.5) (with $a \equiv 1$) with initial datum $u_0 \in \dot{H}^{-s}_{av}(\mathbb{T}^n)$. Then,

$$\left\| (-\Delta)^{-s} u(t) \right\|_{\dot{W}_{\mathrm{av}}^{-1,p}} \le |\mathbb{T}^n|^{1/p} t + \left\| (-\Delta)^{-s} u_0 \right\|_{\dot{W}_{\mathrm{av}}^{-1,p}}$$

for $1 \leq p \leq \infty$.

Proof. We first observe that

$$\frac{d}{dt} \left\| (-\Delta)^{-s} u \right\|_{\dot{W}_{av}^{-1,p}}(t) \le \left\| (-\Delta)^{-s} u_t \right\|_{\dot{W}_{av}^{-1,p}}(t)$$

since $\frac{d}{dt} \|u\|_X \leq \|u_t\|_X$ by the triangle inequality of the norm $\|\cdot\|_X$. By Theorem 2.9, we see

$$u_t = (-\Delta)^s \operatorname{div} Z$$

with

$$||Z||_{L^{\infty}} \le 1$$
 and $(u, -(-\Delta)^s \operatorname{div} Z)_{\dot{H}^{-s}_{\operatorname{av}}} = TV(u)$

Thus

$$\begin{split} \left\| (-\Delta)^{-s} u_t \right\|_{\dot{W}_{\rm av}^{-1,p}}(t) &= \| \operatorname{div} Z \|_{\dot{W}_{\rm av}^{-1,p}} \\ &= \sup \left\{ \int_{\mathbb{T}^n} (-\nabla \varphi) \cdot Z \, dx \, \bigg| \, \| \nabla \varphi \|_{L^{p'}} \le 1 \right\} \le \| Z \|_{L^p} \le |\mathbb{T}^n|^{1/p}. \end{split}$$

We now conclude

$$\frac{d}{dt} \left\| (-\Delta)^{-s} u \right\|_{\dot{W}^{-1,p}_{\mathrm{av}}}(t) \le |\mathbb{T}^n|^{1/p}$$

which yields the desired inequality.

We are now ready to prove our upper bound for the extinction time which is an easy extension of the case s = 1.

Theorem 5.10. For $s \in (0, 1]$, assume that $1 \le n \le 2(s + 1)$, $1 \le p \le \infty$. Assume that $1/2 < \theta \le 1$ satisfies the scaling balance

$$s + \frac{n}{2} = (1 - \theta) \left(2s + 1 + \frac{n}{p} \right) + \theta(n - 1).$$

Then

$$T^*(u_0) \le \frac{A_0}{a} \left\{ \left(1 + \frac{a C_*^{1/\theta} \|u_0\|_{\dot{H}_{av}^{-s}}^{\gamma}}{A_0^{\gamma}} \right)^{1/\gamma} - 1 \right\}$$

with $a := |\mathbb{T}^n|^{1/p}$, $A_0 := ||(-\Delta)^{-s} u_0||_{\dot{W}_{av}^{-1,p}}$, $\gamma := 2 - 1/\theta$.

Proof. We set $y(t) = ||u(t)||_{\dot{H}^{-s}_{av}}$ and recall the fundamental dissipation identity

$$(y^2/2)' = -TV(u).$$

By the interpolation inequality (Lemma 5.8), we have

$$-TV(u) \le -C_*^{-1/\theta} y(t)^{1/\theta} \left\| (-\Delta)^{-s} u(t) \right\|_{\dot{W}_{av}^{-1,p}}^{1-1/\theta}.$$

We may assume $y(t) \neq 0$. We now apply our growth estimate to obtain

$$y(t)^{1-1/\theta}y'(t) \le -C_*^{-1/\theta} \left(|\mathbb{T}^n|^{1/p}t + \left\| (-\Delta)^{-s}u_0 \right\|_{\dot{W}_{av}^{-1,p}} \right)^{1-1/\theta}$$

since $1 - 1/\theta \leq 0$. In other words,

$$\frac{1}{\gamma}\frac{d}{dt}y^{\gamma} \le -C_*^{1/\theta}(at+A_0)^{\gamma-1}.$$

Note that $\gamma \in (0,1]$ since θ satisfies $1/2 < \theta \leq 1$. Integrating both sides over (0,t) we get

$$\frac{1}{\gamma} (y(t)^{\gamma} - y(0)^{\gamma}) \le -\frac{C_*^{-1/\theta}}{a\gamma} \{ (at + A_0)^{\gamma} - A_0^{\gamma} \}$$

or

$$y(t)^{\gamma} \le \|u_0\|_{\dot{H}^{-s}_{av}}^{\gamma} - \frac{C_*^{-1/\theta}}{a} \left\{ (at + A_0)^{\gamma} - A_0^{\gamma} \right\}.$$
(5.6)

Since the right-hand side is nonnegative,

$$\frac{C_*^{-1/\theta}}{a} \left\{ (at + A_0)^{\gamma} - A_0^{\gamma} \right\} \le \|u_0\|_{\dot{H}_{av}^{-s}}^{\gamma}$$

or

$$at + A \le A_0 \left(1 + \frac{aC_*^{1/\theta} \|u_0\|_{\dot{H}_{av}^{-s}}^{\gamma}}{A_0^{\gamma}} \right)^{1/\gamma}$$

Thus, the desired estimate follows from (5.6). (Note that if $\theta = 1/2$, γ must be zero so the above argument does not apply. The case $\theta = 1/2$ corresponds to the case $1 \le n < 2(s+1)$ and $p = \infty$ as we observed before.)

6 Regularity

A fundamental feature of total variation flows, that sets them apart from usually considered quasilinear parabolic equations, is that u(t) is typically only a BV function for t > 0. That is, the distributional derivative Du is in general not an integrable function, but a vector measure. In particular, as we have seen in Section 4, u can have jump discontinuities along hypersurfaces. This is a desirable feature from the point of view of applications to image processing, where jumps corresponds to sharp contours in images.

Rigorously speaking, a point $x \in \Omega$ is called an *(approximate) jump point* of $w \in L^1_{loc}(\Omega)$ if there exist real numbers $w^- = w^-(x)$, $w^+ = w^+(x)$, $w^- \neq w^+$ and a vector $\nu_w = \nu_w(x)$ such that

$$\lim_{r \to 0^+} \oint_{B_r^-(x,\nu_w)} |w(y) - w^-| dy = 0, \quad \lim_{r \to 0^+} \oint_{B_r^+(x,\nu_w)} |w(y) - w^+| dy = 0$$
(6.1)

where the symbol f denotes average integral over a set and $B_r^{\pm}(x,\nu_w)$ are the half-balls

$$B_r^-(x,\nu_w) = \left\{ y \in B_r(x) \mid (y-x) \cdot \nu_w \ge 0 \right\}, \quad B_r^+(x,\nu_w) = \left\{ y \in B_r(x) \mid (y-x) \cdot \nu_w \le 0 \right\}$$

To be precise, the triple $(w^+(x), w^-(x), \nu_w(x))$ is defined up to permutation of $w^+(x)$ and $w^-(x)$ with simultaneous multiplication of $\nu_w(x)$ by -1. We also recall the notion of approximate continuity, closely related to Lebesgue points. As in [AFP], we say that $x \in \Omega$ is a *point of approximate continuity* of w, if there exists $w^* = w^*(x) \in \mathbb{R}$ such that

$$\lim_{r \to 0^+} \oint_{B_r(x)} |w(y) - w^*| dy = 0.$$
(6.2)

The set of jump points of w is denoted by J_w , while the set of approximate discontinuity, i.e., the complement of the set of points of approximate continuity of w, is denoted by S_w . Clearly $J_w \,\subset S_w$. By the Federer–Vol'pert theorem [AFP, Theorem 3.78], if $w \in BV(\Omega)$, then S_w (and J_w) is countably \mathcal{H}^{n-1} -rectifiable, in particular, it can be covered by a countable sum of graphs of C^1 functions up to a \mathcal{H}^{n-1} -negligible set (i.e. a set of \mathcal{H}^{n-1} measure 0) [AFP, p. 80]. Moreover, J_w coincides with S_w up to a \mathcal{H}^{n-1} -negligible set, that is $\mathcal{H}^{n-1}(S_w \setminus J_w) = 0$. We also note that if (6.1) holds for a given $x \in \Omega$ with $w^- = w^+$, then also (6.2) holds with $w^* = w^{\pm}$. On the other hand, if (6.2) holds with $w^* \in \mathbb{R}$, then both equations in (6.1) are satisfied with $w^+ = w^- = w^*$. Thus, if $w \in BV(\Omega)$, then $w^{\pm}(x)$ can be defined for \mathcal{H}^{n-1} -a.e. $x \in \Omega$. Furthermore, one-sided traces of w along any C^1 hypersurface in Ω are well defined [AFP, Theorem 3.77] and coincide with w^{\pm} up to a pointwise permutation [AFP, Remark 3.79].

In this section we denote by ∇w the Radon–Nikodym derivative of Dw with respect to the Lebesgue measure \mathcal{L}^n . Thus, we have

$$Dw = \nabla w \mathcal{L}^n + D^s w,$$

where $D^s w$ is called the singular part of Dw. The singular part can be further decomposed into the jump part $D^j w = (w^+ - w^-)\nu_w \mathcal{H}^{n-1}$ supported on J_w and the remaining Cantor part $D^c w$, see [AFP] for details.

It is then natural to ask about the location of the jumps of u, or more generally, of (the support of) $D^s u$. The situation is particularly simple in the 1D setting (in order to avoid technicalities related to the boundary, we restrict ourselves to the periodic case $\Omega = \mathbb{R}/\mathbb{Z}$). In this case BV functions are well approximated by step functions. Indeed, given $w \in BV(\Omega)$ and a natural number N, we define w^N by

$$w^{N}(x) = N \int_{(k-1)/N}^{k/N} w(y) dy$$
 for $x \in [(k-1)/N, k/N), \ k = 1, \dots, N.$

If $w \in C^1(\overline{\Omega})$, then w attains its average value $N \int_{(k-1)/N}^{k/N} w(y) dy$ on [(k-1)/N, k/N) and, by the fundamental theorem of calculus, we have for $x \in [(k-1)/N, k/N)$

$$|w(x) - w^{N}(x)| = \left| w(x) - N \int_{(k-1)/N}^{k/N} w(y) dy \right| \le \int_{[(k-1)/N, k/N)} |Dw|.$$
(6.3)

It is not difficult to see that the same holds for general $w \in BV(\Omega)$, $\Omega = (a, b)$, if we identify it with its left-continuous representative which is of form

$$w(x) = c + \int_{(a,x)} Dw$$

with some $c \in \mathbb{R}$, see [AFP, Thm. 3.28]. From (6.3) we deduce

$$\begin{split} \|w - w^N\|_{L^2(\Omega)}^2 &= \sum_{k=1}^N \int_{(k-1)/N}^{k/N} \left| w(x) - N \int_{(k-1)/N}^{k/N} w(y) dy \right|^2 dx \le \frac{1}{N} \sum_{k=1}^N \left(\int_{[k-1)/N, k/N} |Dw| \right)^2 \\ &\le \frac{1}{N} \left(\sum_{k=1}^N \int_{[k-1)/N, k/N} |Dw| \right)^2 = \frac{1}{N} TV(w)^2, \end{split}$$

which clearly tends to 0 as $N \to \infty$, i.e. $w^N \to w$ in $L^2(\Omega)$.

Moreover, again identifying w with the left-continuous representative,

$$\begin{split} TV(w^N) &= \int_{\Omega} |Dw^N| = \sum_{k=1}^N \left| N \int_{k/N}^{(k+1)/N} w(y) dy - N \int_{(k-1)/N}^{k/N} w(y) dy \right| \\ &= N \sum_{k=1}^N \left| \int_{(k-1)/N}^{k/N} w(y+1/N) - w(y) dy \right| = N \sum_{k=1}^N \left| \int_{(k-1)/N}^{k/N} \left(\int_{[y,y+1/N)} Dw \right) dy \right| \\ &\leq N \sum_{k=1}^N \int_{(k-1)/N}^{k/N} \left(\int_{[y,y+1/N)} |Dw| \right) dy = \int_{\Omega} N \left(\int_{[y,y+1/N)} |Dw| \right) dy = \int_{\Omega} |Dw|, \end{split}$$

where the intuitive last equality can be justified using Fubini's theorem. Thus, by lower semicontinuity of the total variation on any open subset of Ω and [AFP, Prop. 1.80], measures $|Dw^N|$ converge weakly star to |Dw|.

We apply this approximation to the initial datum u_0 , obtaining step functions u_0^N . Let u^N be the solution to the total variation flow with initial datum u_0^N . Then $u^N(t)$ is a step function for t > 0. Precisely, between merging times, the solution is given by formulae (4.6)–(4.7). We observe that a^k is a decreasing (resp. increasing) function of time if and only if $a^k > a^{k-1}$ and $a^k > a^{k+1}$ (resp. $a^k < a^{k-1}$ and $a^k < a^{k+1}$). Thus, the functions $|a^{k+1} - a^k|$ are decreasing for $k = 1, \ldots, m$. Since

$$|Du^{N}(t)| = \sum_{k=1}^{m} |a^{k+1}(t) - a^{k}(t)|\delta_{x_{k}},$$

this implies $|Du^N(t)| \leq |Du_0^N|$ as measures, which is equivalent to saying that

$$\int_{\Omega} \varphi \, d|Du^{N}(t)| \leq \int_{\Omega} \varphi \, d|Du_{0}^{N}| \quad \text{for any } \varphi \in C_{c}(\Omega),$$
(6.4)

in other words $TV_{\varphi}(u^N(t)) \leq TV_{\varphi}(u(t))$ for $\varphi \in C_c(\Omega)$. We now want to pass to the limit $N \to \infty$ in (6.4). Since $u_0^N \to u_0$ in $L^2(\Omega)$, we have, by monotonicity of $-\partial TV$, $u^N(t) \to u(t)$ in $L^2(\Omega)$ for t > 0. By lower semicontinuity of TV_{φ} , we obtain

$$\liminf_{N \to \infty} \int_{\Omega} \varphi \, d |Du^N(t)| \ge \int_{\Omega} \varphi \, d |Du(t)|.$$

On the other hand, we have showed that measures $|Du_0^N|$ converge weakly star to $|Du_0|$, i.e.

$$\int_\Omega \varphi \, d |D u_0^N| \to \int_\Omega \varphi \, d |D u_0|$$

Thus, we have

Theorem 6.1. Let Ω be an interval and suppose that u is the solution to the total variation flow with initial datum $u_0 \in BV(\Omega)$. Then

$$|Du(t)| \le |Du_0| \quad as \ measures \ for \ t > 0. \tag{6.5}$$

Inequality (6.5) implies that $|\nabla u(t)| \leq |\nabla u_0|$ Lebesgue-almost everywhere in Ω and $|D^s u(t)| \leq |\nabla u_0|$ $|D^s u_0|$ as measures. This has quite strong consequences in terms of regularity. In particular, it implies that the total variation flow preserves Sobolev spaces $W^{1,p}(\Omega)$ for $p \in [1,\infty]$, or the space $SBV(\Omega)$ of functions $w \in BV(\Omega)$ such that $D^c w = 0$. Many of these consequences were derived by a technique similar to the one presented here in [BF]. Independently, a generalization of Theorem 6.1 was obtained in [BCNO] by a different technique. We note that one can also show that the 1D total variation flow preserves second-order BV regularity. That is, if $Du_0 \in BV(\Omega)$, then $Du(t) \in BV(\Omega)$ for t > 0 [MR]. This is the highest regularity preserved by the flow: in fact, if $u_0 \in C^{\infty}(\overline{\Omega})$ is non-monotone, then Du(t) will have jump discontinuities for any t > 0 small enough, see e.g. [KMR].

Theorem 6.1 can also be generalized to the vector-valued setting, with total variation calculated with respect to possibly non-Euclidean norms [GiL, GrL]. If the total variation is replaced by a non-homogeneous functional, the pointwise inequality $|\nabla u(t)| \leq |\nabla u_0|$ in general fails. However one can still obtain a bound on $D^{s}u$ [MS, GrL].

As for the total variation flow in higher-dimensional domains, the pointwise estimate on ∇u is known to be violated, as evidenced by examples of *facet bending* [ACC]. However one can still show an estimate on the size of jumps of u(t) in terms of u_0 . Known results of this type rely on the minimizing movements approximation briefly discussed in Section 2.1, consisting in iteratively solving a minimization problem for the functional $\mathcal{E}_{f}^{\lambda}$ given by (2.2), which in the case $\mathcal{E} = TV$ reads

$$TV_f^{\lambda}(w) = \lambda TV(w) + \frac{1}{2} \int_{\Omega} (w - f)^2 dx.$$
(6.6)

Initial works on this subject used an argument based on the fact that level sets of the minimizer of TV_f^{λ} solve a prescribed mean curvature problem [CCN, CJN]. This has been later significantly generalized in [Val, CL] by different techniques, without reference to particular structure of \mathcal{E} or level sets of the minimizer, thus allowing to handle also the vector-valued case. In fact, it is enough to assume that \mathcal{E} has a mild regularity property of differentiability along inner variations, which holds for TV [Giu, Chapter 10]. That is, setting $w_{\varphi}^{\tau}(x) = w(x + \tau \varphi(x))$ for $w \in BV(\Omega), \varphi \in C_c^{\infty}(\Omega)^n, x \in \Omega,$ $\tau \in \mathbb{R}$, the real function $R_{\varphi}: \tau \mapsto TV(w_{\varphi}^{\tau})$ is differentiable for any $\varphi \in C_c^{\infty}(\Omega)^n$. To obtain the desired assertion, we actually need to use a bit more complicated mixed variations $w_{\varphi}^{\tau,\vartheta} := (1 - \vartheta)w + \vartheta w_{\varphi}^{\tau}$. By convexity of TV we have

$$\frac{1}{\tau}(TV(w_{\varphi}^{\tau,\vartheta}) - TV(w)) \le \frac{1}{\tau}((1 - \vartheta)TV(w) + \vartheta TV(w_{\varphi}^{\tau}) - TV(w)) = \frac{\vartheta}{\tau}(R_{\varphi}(\tau) - R_{\varphi}(0)) \to \vartheta R_{\varphi}'(0),$$

 $\frac{1}{\tau}(TV(w_{\varphi}^{-\tau,\vartheta}) - TV(w)) \leq \frac{1}{\tau}((1-\vartheta)TV(w) + \vartheta TV(w_{\varphi}^{-\tau}) - TV(w)) = \frac{\vartheta}{\tau}(R_{\varphi}(-\tau) - R_{\varphi}(0)) \to -\vartheta R_{\varphi}'(0)$ as $\tau \to 0^+$. Thus,

$$\limsup_{\tau \to 0^+} \frac{1}{\tau} (TV(w_{\varphi}^{\tau,\vartheta}) - TV(w)) + \limsup_{\tau \to 0^+} \frac{1}{\tau} (TV(w_{\varphi}^{-\tau,\vartheta}) - TV(w)) \le 0.$$
(6.7)

Since the function $\tau \to TV(w_{\varphi}^{\tau,\vartheta})$ is convex, it actually follows from this inequality that it is also differentiable (at least at 0).

Now, let $v \in BV(\Omega)$ be a minimizer of TV_f^{λ} for a given $f \in BV(\Omega)$, $\lambda > 0$. We assume moreover that $f \in L^{\infty}(\Omega)$, in which case it is easy to show that $v \in L^{\infty}(\Omega)$ and $||v||_{L^{\infty}(\Omega)} \leq ||f||_{L^{\infty}(\Omega)}$. Since v is a minimizer, we have

$$\liminf_{\tau \to 0^+} \frac{1}{\tau} (TV_f^{\lambda}(v_{\varphi}^{\tau,\vartheta}) - TV_f^{\lambda}(v)) \ge 0, \quad \liminf_{\tau \to 0^+} \frac{1}{\tau} (TV_f^{\lambda}(v_{\varphi}^{-\tau,\vartheta}) - TV_f^{\lambda}(v)) \ge 0.$$

Adding the two inequalities together and taking into account (6.7), we deduce

$$0 \le \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{\tau,\vartheta} - f)^2 - (v - f)^2 dx + \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{-\tau,\vartheta} - f)^2 - (v - f)^2 dx.$$
(6.8)

Further on, we will only use (6.8). Note that it does not involve TV at all. We calculate

$$(v_{\varphi}^{\pm\tau,\vartheta} - f)^2 - (v - f)^2 = (v_{\varphi}^{\pm\tau,\vartheta} - v)(v + v_{\varphi}^{\pm\tau,\vartheta} - 2f) = \vartheta(v_{\varphi}^{\pm\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\pm\tau} - 2f).$$

Thus, dividing (6.8) by ϑ ,

$$0 \leq \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{\tau} - v)((2-\vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f)dx + \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{-\tau} - v)((2-\vartheta)v + \vartheta v_{\varphi}^{-\tau} - 2f)dx.$$
(6.9)

All the calculations so far were done for arbitrary $\varphi \in C_c^{\infty}(\Omega)$. Now let $\Gamma \subset \Omega$ be any one of the C^1 graphs that cover J_v by rectifiability, let $x_0 \in \Gamma$, and let ν_0 be a vector normal to Γ at x_0 . By isometric change of coordinates, we can assume without loss of generality that $x_0 = 0$, $\nu_0 = (0, \ldots, 0, 1)$ and $\Gamma \supset \{(x', x_n) : x' \in B_r^{n-1}, x_n = \gamma(x')\}$ where $\gamma \in C^1(B_r^{n-1})$ and $B_r^{n-1} := \{x' \in \mathbb{R}^{n-1} : |x'| < r\}$ for r > 0. Owing to differentiability of γ , possibly decreasing r we can assume that $\gamma(B_s^{n-1}) \subset (-s/2, s/2)$ for $0 < s \leq r$. Then, we take $\varphi = \nu_0 \psi = (0, \ldots, 0, \psi)$ supported in $Q_r := B_r^{n-1} \times (-r, r)$ with $\psi \in C_c^{\infty}(Q_r)$ such that $0 \leq \psi \leq 1$ and $\psi(x', x_n) = \psi_{\parallel}(x')\psi_{\perp}(x_n)$, $\psi_{\parallel} = 1$ on $B_{r-\varepsilon}^{n-1}$, $\psi_{\perp} = 1$ on $(-r + \varepsilon, r - \varepsilon)$ for a temporarily fixed $\varepsilon \in (0, r/2)$.

For $\tau \in (0, r/2)$ we rewrite

$$\frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f) dx = \frac{1}{2\tau} \int_{B_r^{n-1}} \int_{\gamma(x') - \tau \psi_{\parallel}(x')}^{\gamma(x')} (v_{\varphi}^{\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f) dx_n dx'
+ \frac{1}{2\tau} \int_{B_r^{n-1}} \int_{(-r,r) \setminus [\gamma(x') - \tau \psi_{\parallel}(x'), \gamma(x')]} (v_{\varphi}^{\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f) dx_n dx'. \quad (6.10)$$

Identifying v with its precise representative, the slicing properties of BV functions [AFP, §3.11, Theorem 3.107] ensure that for \mathcal{L}^{n-1} -a.e. $x' \in B_r^{n-1}$ the function $v_{x'}: x_n \mapsto v(x', x_n)$ is in BV((-r, r)), and we can write for \mathcal{L}^1 -a.e. $x_n \in (-r, r)$:

$$v_{\varphi}^{\tau}(x',x_n) - v(x',x_n) = v(x',x_n + \tau\psi(x',x_n)) - v(x',x_n) = \int_{(x_n,x_n + \tau\psi(x',x_n))} Dv_{x'} + \frac{1}{2} \int_{(x_n,x_n + \tau\psi(x',x_n)} Dv_{x'} + \frac{1}{2} \int_{(x_n,x_n + \tau\psi(x',x_n))} Dv_{x'} + \frac{1}$$

Since the support of ψ is contained in Q_r , for small enough $\tau > 0$ we have $x_n + \tau \psi(x', x_n) \leq 1$ $\min(x_n + \tau \psi_{\parallel}(x'), r)$ for $x_n \in (-r, r)$. Therefore

$$\begin{aligned} \left| (v_{\psi}^{\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f) \right| (x', x_n) &\leq 4 \|f\|_{L^{\infty}(\Omega)} \left| v_{\psi}^{\tau} - v \right| (x', x_n) \\ &\leq 4 \|f\|_{L^{\infty}(\Omega)} \int_{(x_n, x_n + \tau\psi(x', x_n))} |Dv_{x'}| \leq 4 \|f\|_{L^{\infty}(\Omega)} \int_{(x_n, \min(x_n + \tau\psi_{\parallel}(x'), r))} |Dv_{x'}| \end{aligned}$$

and so, by Fubini's theorem, for \mathcal{L}^{n-1} -a.e. $x' \in B_r^{n-1}$,

$$\frac{1}{\tau} \int_{(-r,r)\setminus[\gamma(x')-\tau\psi_{\parallel}(x')),\gamma(x')]} \left| (v_{\psi}^{\tau}-v)((2-\vartheta)v+\vartheta v_{\varphi}^{\tau}-2f) \right| (x',x_{n}) dx_{n} \\
\leq 4 \|f\|_{L^{\infty}(\Omega)} \frac{1}{\tau} \int \int \mathbf{1}_{(-r,r)\setminus[\gamma(x')-\tau\psi_{\parallel}(x')),\gamma(x')]} (x_{n}) \mathbf{1}_{(x_{n},\min(x_{n}+\tau\psi_{\parallel}(x'),r))}(s) d| Dv_{x'}|(s) dx_{n} \\
\leq 4 \|f\|_{L^{\infty}(\Omega)} \frac{1}{\tau} \int \int \mathbf{1}_{(s-\tau\psi_{\parallel}(x'),s))} (x_{n}) \mathbf{1}_{(-r,r)\setminus\{\gamma(x')\}} (s) dx_{n} d| Dv_{x'}|(s) \\
\leq 4 \|f\|_{L^{\infty}(\Omega)} \psi_{\parallel}(x') \int_{(-r,r)\setminus\{\gamma(x')\}} |Dv_{x'}| \leq 4 \|f\|_{L^{\infty}(\Omega)} \int_{(-r,r)\setminus\{\gamma(x')\}} |Dv_{x'}|. \quad (6.11)$$

Appealing to [AFP, Theorem 3.107],

$$\frac{1}{2\tau} \int_{B_r^{m-1}} \int_{(-r,r)\setminus[\gamma(x')-\tau\psi_{\parallel}(x'),\gamma(x')]} (v_{\varphi}^{\tau}-v)((2-\vartheta)v + \vartheta v_{\varphi}^{\tau}-2f) dx_n dx' \leq 2\|f\|_{L^{\infty}(\Omega)} \int_{Q_r\setminus\Gamma} |Dv|.$$
(6.12)

Thus we have estimated the second term on the r.h.s. of (6.10). As for the other one, using [AFP, Theorem 3.108], for \mathcal{L}^{n-1} -a.e. $x' \in B_r^{m-1}$ we have

$$\frac{1}{\tau} \int_{\gamma(x')-\tau\psi_{\parallel}(x')}^{\gamma(x')} (v_{\psi}^{\tau}-v)((2-\vartheta)v + \vartheta v_{\varphi}^{\tau}-2f) dx_n \to \psi (v^+-v^-)((2-\vartheta)v^- + \vartheta v^+ - 2f^-) \Big|_{(x',\gamma(x'))},$$

where v^- , f^- (resp. v^+ , f^+) are the approximate limits corresponding to traces of v, f along Γ "from below" (resp. "from above"). By a rough estimate in the vein of (6.11), we can show that

$$\frac{1}{\tau} \int_{\gamma(x')-\tau\psi_{\parallel}(x')}^{\gamma(x')} \left| (v_{\psi}^{\tau}-v)((2-\vartheta)v + \vartheta v_{\varphi}^{\tau}-2f) \right| (x',x_n) \, dx_n \le 4 \|f\|_{L^{\infty}(\Omega)} \int_{(-r,r)} |Dv_{x'}|.$$

Since the r.h.s. is an integrable function of x' (again by [AFP, Theorem 3.107]), we can apply dominated convergence theorem to show that the first term on the r.h.s. of (6.10) converges as $\tau \to 0^+$ to

$$\frac{1}{2} \int_{B_r^{n-1}} \psi \left(v^+ - v^- \right) \left((2 - \vartheta) v^- + \vartheta v^+ - 2f^- \right) \bigg|_{(x', \gamma(x'))} dx'.$$

Thus, recalling (6.12),

$$\begin{split} \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{\tau} - 2f)dx \\ & \leq \frac{1}{2} \int_{B_r^{n-1}} \psi \left(v^+ - v^- \right)((2 - \vartheta)v^- + \vartheta v^+ - 2f^-) \Big|_{(x', \gamma(x'))} dx' + 2\|f\|_{L^{\infty}(\Omega)} \int_{Q_r \setminus \Gamma} |Dv|. \end{split}$$

Repeating the same reasoning, we also obtain

$$\begin{split} \limsup_{\tau \to 0^+} \frac{1}{2\tau} \int_{\Omega} (v_{\varphi}^{-\tau} - v)((2 - \vartheta)v + \vartheta v_{\varphi}^{-\tau} - 2f) dx \\ & \leq \frac{1}{2} \int_{B_r^{n-1}} \psi \left(v^- - v^+ \right) ((2 - \vartheta)v^+ + \vartheta v^- - 2f^+) \Big|_{(x', \gamma(x'))} dx' + 2 \|f\|_{L^{\infty}(\Omega)} \int_{Q_r \setminus \Gamma} |Dv|. \end{split}$$

Summing these two inequalities, recalling (6.9), and passing with $\vartheta \to 0^+$, $\varepsilon \to 0^+$,

$$0 \le \int_{B_r^{n-1}} (v^+ - v^-)(v^- - v^+ + f^+ - f^-) \Big|_{(x',\gamma(x'))} dx' + 4 \|f\|_{L^{\infty}(\Omega)} \int_{Q_r \setminus \Gamma} |Dv|.$$

Finally, we divide the obtained inequality by $|B_r^{n-1}|$ and pass to the limit $r \to 0^+$. By [AFP, eq. (2.41) on p. 79], the second term vanishes in the limit for \mathcal{H}^{n-1} -a.e. $x_0 \in \Gamma$. Here \mathcal{H}^{n-1} denotes the n-1-dimensional Hausdorff measure, whose restriction to Γ corresponds to the classical surface measure on Γ . Since also \mathcal{H}^{n-1} -a.e. $x_0 \in \Gamma$ is a Lebesgue point of the function $(v^+ - v^-)(v^- - v^+ + f^+ - f^-)$ with respect to the surface measure, we obtain

$$0 \le (v^{+} - v^{-})(v^{-} - v^{+} + f^{+} - f^{-})$$

for \mathcal{H}^{n-1} -a.e. $x_0 \in \Gamma$ (see [CL] for a more detailed explanation of this part). We deduce

$$|v^+ - v^-| \le |f^+ - f^-|$$
 for \mathcal{H}^{n-1} -a.e. $x_0 \in J_v$.

In particular, J_v is contained in J_f up to a \mathcal{H}^{n-1} -negligible set.

Applying this to the minimizing movements approximation, if $u_0 \in BV(\Omega) \cap L^{\infty}(\Omega)$, we have $J_{u^N(t)} \subset J_{u_0}$ for $t > 0, N \in \mathbb{N}$ up to a \mathcal{H}^{n-1} -negligible set and

$$|u^{N}(t)^{+} - u^{N}(t)^{-}| \le |u_{0}^{+} - u_{0}^{-}| \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x_{0} \in J_{u_{0}}.$$
(6.13)

Recall that for $v \in BV(\Omega)$, the functions v^{\pm} can be defined \mathcal{H}^{n-1} -a.e. in Ω by setting $v^{+} = v^{-}$ to be the approximate limit of v at any point where it exists. Indeed, by the Federer-Vol'pert theorem, \mathcal{H}^{n-1} -a.e. point in $\Omega \setminus J_v$ is a point of approximate continuity of v [AFP].

It remains to transfer the result to the total variation flow. We will follow the approach from [CCN, CJN] relying on the theory of accretive operators on Banach spaces, see Appendix A in [ACM] and references therein. We will focus on some details related to limit passage with the bound on jump size (6.13) which seem to be omitted in [CCN, CJN]. We define A_{∞} as the restriction of $\partial_{L^2}TV$ to $L^{\infty}(\Omega)$. Using the characterization of $\partial_{L^2}TV$ in terms of Cahn–Hoffman vector fields (Theorem 2.7), it is not difficult to see that it is *accretive*, i.e.,

$$\|v_1 - v_2\|_{L^{\infty}(\Omega)} \le \|v_1 - v_2 + \lambda(w_1 - w_2)\|_{L^{\infty}(\Omega)} \quad \text{for } v_1, v_2 \in D(A_{\infty}), \ w_1 \in A_{\infty}(v_1), \ w_2 \in A_{\infty}(v_2), \ \lambda > 0,$$

where $D(A_{\infty})$ denotes the set of $v \in L^{\infty}(\Omega)$ such that $A_{\infty}(v)$ is non-empty. One can show it by approximating ∞ with finite p, using convexity of the power function and integrating by parts. Moreover, A_{∞} satisfies the range condition

$$\overline{D(A_{\infty})} \subset R(I + \lambda A_{\infty}) \quad \text{for all } \lambda > 0,$$

where $R(I+\lambda A_{\infty})$ is the range of the operator $I+\lambda A_{\infty}$. In fact $R(I+\lambda A_{\infty}) = L^{\infty}(\Omega)$ since minimizers of (2.2) with $f \in L^{\infty}(\Omega)$ belong to $L^{\infty}(\Omega)$. Thus, by the Crandall–Liggett theorem [ACM, Theorem A.28], if $u_0 \in \overline{D(A_{\infty})}$, the minimizing movements approximation $u^N(t)$ defined in (2.3) converges to u(t) in $L^{\infty}(\Omega)$ for t > 0.

Recall that u^N is piecewise constant as a function from $[0, \infty)$ to $L^2(\Omega)$, in particular its image is countable. Thus, there is a \mathcal{H}^{n-1} -full subset J of J_{u_0} whose elements are either jump points or points of approximate continuity of $u^N(t)$ for all $N \in \mathbb{N}$, t > 0. Moreover, since the jump sets of u^N are \mathcal{H}^{n-1} -almost contained in J_{u_0} , they can all be \mathcal{H}^{n-1} -almost covered by the same countable family of C^1 surfaces. Owing to this, we can assume that in case $x \in J$ is a jump point for $u^N(t)$, $N \in \mathbb{N}$, t > 0, then $\nu_{u^N(t)}(x) = \nu_{u_0}(x)$. Therefore we have

$$\lim_{r \to 0^+} \oint_{B_r^{\pm}(x,\nu_{u_0})} |u^N(t,y) - u^N(t)^{\pm}(x)| dy = 0$$
(6.14)

for all $x \in J$, $N \in \mathbb{N}$, t > 0. By the uniform bound $||u^N(t)||_{L^{\infty}(\Omega)} \leq ||u_0||_{L^{\infty}(\Omega)}$, the sequences $\{u^N(t)^{\pm}(x)\}_{N\in\mathbb{N}}$ are bounded for $x \in J$ and thus each one has a limit point a^{\pm} . Given $\varepsilon > 0$,

$$\begin{aligned} \oint_{B_{r}^{\pm}(x,\nu_{u_{0}})} |u(t,y) - a^{\pm}| dy \\ &\leq \int_{B_{r}^{\pm}(x,\nu_{u_{0}})} |u(t,y) - u^{N}(t,y)| dy + \int_{B_{r}^{\pm}(x,\nu_{u_{0}})} |u^{N}(t,y) - u^{N}(t)^{\pm}(x)| dy + |u^{N}(t)^{\pm}(x) - a^{\pm}| \\ &\leq \int_{B_{r}^{\pm}(x,\nu_{u_{0}})} |u^{N}(t,y) - u^{N}(t)^{\pm}(x)| dy + 2\varepsilon \quad (6.15) \end{aligned}$$

for N large enough, independently of r > 0. As $\varepsilon > 0$ is arbitrarily small, we deduce from (6.14)

$$\lim_{r \to 0^+} \oint_{B_r^{\pm}(x,\nu_{u_0})} |u(t,y) - a^{\pm}| dy = 0,$$

hence $x \in J_{u(t)} \cup (\Omega \setminus S_{u(t)})$ and, by (6.13),

$$|u^{+}(t) - u^{-}(t)| = |a^{+} - a^{-}| \le |u_{0}^{+} - u_{0}^{-}|$$

By an estimate similar to (6.15), using the inclusion \mathcal{H}^{n-1} -almost inclusion $S_{u^N(t)} \subset S_{u_0}$, we also show that $S_{u(t)} \subset S_{u_0}$ (equivalently, $J_{u(t)} \subset J_{u_0}$) up to a \mathcal{H}^{n-1} -negligible set.

Finally, we would like to remove the enigmatic assumption $u_0 \in \overline{D(A_{\infty})}$. Instead, let us only assume that $u_0 \in BV(\Omega) \cap L^n(\Omega)$. By the $L^n - L^\infty$ regularization property [ACM, Theorem 2.16] and the pointwise estimate $|u_t(t)| \leq 2\frac{|u(s)|}{t-s}$ for $0 \leq s < t$ following from homogeneity of TV [ACM, (2.34)], we have $u(t) \in D(A_{\infty})$ for t > 0. Thus, by the previous step, $J_{u(t)} \subset J_{u(s)}$ and the inequality $|u^+(t) - u^-(t)| \leq |u^+(s) - u^-(s)|$ holds \mathcal{H}^{n-1} -a.e. in Ω for 0 < s < t. Let us take a sequence of positive numbers s_k such that $s_k \to 0$ as $k \to \infty$. By lower semicontinuity of TV and its monotonicity along trajectories of the flow,

$$\lim_{k \to \infty} \int_{\Omega} |Du(s_k)| = \int_{\Omega} |Du_0|.$$

This improves the convergence $u(s_k) \to u_0$ in $L^2(\Omega)$ to strict convergence in $BV(\Omega)$, whence $|Du(s_k)|$ converges to $|Du_0|$ weakly* in the space of signed Radon measures $M(\Omega)$ [AFP, Proposition 1.80]. Thus, using [AFP, Proposition 1.63], given $x \in \Omega$,

$$\int_{B_r(x)} |Du(s_k)| \to \int_{B_r(x)} |Du_0|$$

for a.e. r > 0 such that $B_r(x) \subset \Omega$. For \mathcal{H}^{n-1} -a.e. $x \in J_{u(t)}$,

$$\int_{B_{r}(x)\cap J} |u^{+}(t,y) - u^{-}(t,y)| d\mathcal{H}^{n-1}(y) \leq \int_{B_{r}(x)} |u^{+}(t,y) - u^{-}(t,y)| d\mathcal{H}^{n-1}(y) \\
\leq \int_{B_{r}(x)} |u^{+}(s_{k},y) - u^{-}(s_{k},y)| d\mathcal{H}^{n-1}(y) \leq \int_{B_{r}(x)} |Du(s_{k})|, \quad (6.16)$$

where J is one of the C^1 graphs covering $J_{u(t)}$ by rectifiability such that $x \in J$. Dividing both sides of (6.16) by $|B_r^{n-1}|$, we deduce

$$|u^{+}(t,x) - u^{-}(t,x)| \le \liminf_{r \to 0^{+}} \frac{1}{|B_{r}^{n-1}|} \int_{B_{r}(x)} |Du_{0}| \quad \text{for } \mathcal{H}^{n-1}\text{-a.e. } x \in J_{u(t)}.$$
(6.17)

Thus, by [AFP, Proposition 3.92], we conclude that $J_{u(t)} \subset J_{u_0}$ up to a \mathcal{H}^{n-1} -negligible set. Then, using the Radon–Nikodym derivation theorem, we also deduce that $|u^+(t) - u^-(t)| \leq |u_0^+ - u_0^-| \mathcal{H}^{n-1}$ -a.e. on $J_{u(t)}$. Summing up, we have obtained

Theorem 6.2. Let $u_0 \in BV(\Omega) \cap L^n(\Omega)$. Then for a.e. t > 0, $J_{u(t)} \subset J_{u_0}$ up to a \mathcal{H}^{n-1} -negligible set and

$$|u^+(t) - u^-(t)| \le |u_0^+ - u_0^-|$$
 \mathcal{H}^{n-1} -a.e. on $J_{u(t)}$

The thesis of the theorem can also be rephrased as $|D^j u(t)| \leq |D^j u_0|$ in the sense of measures. As we have mentioned before, analogous inequality for the absolutely continuous part of Du fails in general if n > 1. To our knowledge it remains an open question whether $|D^c u(t)| \leq |D^c u_0|$.

As for the fourth-order case, we have seen in Section 4 that the jump inclusion $J_{u(t)} \subset J_{u_0}$ does not hold, as the jumps can move. Moreover, jump discontinuities can emerge out of Lipschitz continuous initial data, even in the 1D case [GG].

Acknowledgements

The first author was in part supported by the Japan Society for the Promotion of Science (JSPS) through grants No. 20K20342 (Kaitaku), No. 19H00689 (Kiban A), No. 18H05323 (Kaitaku) and by Arithmer Inc., Daikin Industries, Ltd. and Ebara Corporation through a collaborative grant. The second author was in part supported by JSPS through grant No. 18H05323 (Kaitaku). The third author was supported by the grant No. 2020/36/C/ST1/00492 of the National Science Centre, Poland.

References

[ACC] F. Alter, V. Caselles and A. Chambolle, A characterization of convex calibrable sets in \mathbb{R}^N . Math. Ann. **332** (2005), no. 2, 329–366. doi:10.1007/s00208-004-0628-9.

- [AB] M. Amar and G. Bellettini, A notion of total variation depending on a metric with discontinuous coefficients. Ann. Inst. H. Poincaré C Anal. Non Linéaire 11 (1994), no. 1, 91–133. doi:10.1016/s0294-1449(16)30197-4.
- [AFP] L. Ambrosio, N. Fusco and D. Pallara, Functions of bounded variation and free discontinuity problems. Oxford Math. Monogr., The Clarendon Press, Oxford University Press, New York, 2000. xviii+434 pp.
- [ABC1] F. Andreu, C. Ballester, V. Caselles and J. M. Mazón, Minimizing total variation flow. Differential Integral Equations 14 (2001), no. 3, 321–360. doi:10.57262/die/1356123331.
- [ABC2] F. Andreu, C. Ballester, V. Caselles and J. M. Mazón, The Dirichlet problem for the total variation flow. J. Funct. Anal. 180 (2001), no. 2, 347–403. doi:10.1006/jfan.2000.3698.
- [ACM] F. Andreu-Vaillo, V. Caselles and J. M. Mazón, Parabolic quasilinear equations minimizing linear growth functionals. Progr. Math., 223, Birkhäuser Verlag, Basel, 2004. xiv+340 pp. doi:10.1007/978-3-0348-7928-6.
- [BCN] G. Bellettini, V. Caselles and M. Novaga, The total variation flow in \mathbb{R}^N , J. Differential Equations 184 (2002), no. 2, 475–525. doi:10.1006/jdeq.2001.4150.
- [BCDS] M. Benning, L. Calatroni, B. Düring and C.-B. Schönlieb, A primal-dual approach for a total variation Wasserstein flow. *Geometric science of information*, 413–421, Lecture Notes in Comput. Sci., 8085, Springer, Heidelberg, 2013. doi:10.1007/978-3-642-40020-9_45.
- [BDS] V. Bögelein, F. Duzaar and C. Scheven, The total variation flow with time dependent boundary values. Calc. Var. Partial Differential Equations 55 (2016), no. 4, Art. 108, 31 pp. doi:10.1007/s00526-016-1041-4.
- [BF] M. Bonforte and A. Figalli. Total variation flow and sign fast diffusion in one dimension. J. Differential Equations 252 (2012), no. 8, 4455–4480. doi:10.1016/j.jde.2012.01.003.
- [Br] H. Brézis, Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert. North-Holland Math. Stud., No. 5, Notas Mat., No. 50, North-Holland Publishing Co., Amsterdam-LondonAmerican Elsevier Publishing Co., Inc., New York, 1973, vi+183 pp.
- [BCNO] A. Briani, A. Chambolle, M. Novaga and G. Orlandi, On the gradient flow of a one-homogeneous functional. *Confluentes Math.* **3** (2011), no. 4, 617–635. doi:10.1142/S1793744211000461.
- [BCP] V. Buffa, M. Collins and C. Pacchiano Camacho, Existence of parabolic minimizers to the total variation flow on metric measure spaces. *Manuscripta Math.* 170 (2023), no. 1–2, 109–145. doi:10.1007/s00229-021-01350-2.
- [CP] G. Carlier and C. Poon, On the total variation Wasserstein gradient flow and the TV-JKO scheme. ESAIM Control Optim. Calc. Var. 25 (2019), no. 42, 21 pp. doi:10.1051/cocv/2018042.
- [CCN] V. Caselles, A. Chambolle and M. Novaga, The discontinuity set of solutions of the TV denoising problem and some extensions. *Multiscale Model. Simul.* 6 (2007), no. 3, 879–894. doi:10.1137/070683003.

- [CJN] V. Caselles, K. Jalalzai and M. Novaga, On the jump set of solutions of the total variation flow. Rend. Semin. Mat. Univ. Padova 130 (2013), 155–168. doi:10.4171/RSMUP/130-5.
- [CL] A. Chambolle and M. Łasica, Inclusion and estimates for the jumps of minimizers in variational denoising, preprint. arXiv:2312.01900.
- [DiB] E. DiBenedetto, Degenerate parabolic equations. Universitext, Springer-Verlag, New York, 1993. xvi+387 pp. doi:10.1007/978-1-4612-0895-2.
- [FG] T. Fukui and Y. Giga, Motion of a graph by nonsmooth weighted curvature. World Congress of Nonlinear Analysts '92, Vol. I-IV (Tampa, FL, 1992), 47–56. Walter de Gruyter & Co., Berlin, 1996. doi:10.1515/9783110883237.47.
- [Gal] G. P. Galdi, An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems. Second edition, Springer Monogr. Math. Springer, New York, 2011, xiv+1018 pp. doi:10.1007/978-0-387-09620-9.
- [GiL] L. Giacomelli and M. Łasica, A local estimate for vectorial total variation minimization in one dimension. *Nonlinear Anal.* **181** (2019), 141–146. doi:10.1016/j.na.2018.11.009.
- [GLM] L. Giacomelli, M. Lasica and S. Moll, Regular 1-harmonic flow. Calc. Var. Partial Differential Equations 58 (2019), no. 2, Paper No. 82, 24 pp. doi:10.1007/s00526-019-1526-z.
- [GMM] L. Giacomelli, J. M. Mazón and S. Moll, The 1-harmonic flow with values in a hyperoctant of the N-sphere. Anal. PDE 7 (2014), no. 3, 627–671. doi:10.2140/apde.2014.7.627.
- [GG] M.-H. Giga and Y. Giga, Very singular diffusion equations: second and fourth order problems. Jpn. J. Ind. Appl. Math. 27 (2010), no. 3, 323–345. doi:10.1007/s13160-010-0020-y.
- [GGK] M.-H. Giga, Y. Giga and R. Kobayashi, Very singular diffusion equations. Taniguchi Conference on Mathematics Nara '98, 93-125. Adv. Stud. Pure Math., 31, Mathematical Society of Japan, Tokyo, 2001. doi:10.2969/aspm/03110093.
- [GGP] M.-H. Giga, Y. Giga and N. Požár, Periodic total variation flow of non-divergence type in \mathbb{R}^n . J. Math. Pures Appl. (9) **102** (2014), no. 1, 203–233. doi:10.1016/j.matpur.2013.11.007.
- [GGS] M.-H. Giga, Y. Giga and J. Saal, Nonlinear partial differential equations. Asymptotic behavior of solutions and self-similar solutions. Progr. Nonlinear Differential Equations Appl., 79 Birkhäuser Boston, Ltd., Boston, MA, 2010. xviii+294 pp. doi:10.1007/978-0-8176-4651-6.
- [GK] Y. Giga and R. V. Kohn, Scale-invariant extinction time estimates for some singular diffusion equations. *Discrete Contin. Dyn. Syst.* **30** (2011), no. 2, 509–535. doi:10.3934/dcds.2011.30.509.
- [GKL] Y. Giga, H. Kuroda and M. Lasica, The fourth-order total variation flow in \mathbb{R}^n . Math. Eng. **5** (2023), no. 6, Paper No. 091, 45 pp. doi:10.3934/mine.2023091.
- [GKL2] Y. Giga, H. Kuroda and M. Lasica, On H^{-1} total variation flows in bounded domains, in preparation.

- [GKM] Y. Giga, H. Kuroda and H. Matsuoka, Fourth-order total variation flow with Dirichlet condition: characterization of evolution and extinction time estimates. Adv. Math. Sci. Appl. 24 (2014), no. 2, 499–534.
- [GMR] Y. Giga, M. Muszkieta and P. Rybka, A duality based approach to the minimizing total variation flow in the space H^{-s} . Jpn. J. Ind. Appl. Math. **36** (2019), 261–286. doi:10.1007/s13160-018-00340-4.
- [GNRS] Y. Giga, R. Nakayashiki, P. Rybka and K. Shirakawa, On boundary detachment phenomena for the total variation flow with dynamic boundary conditions. J. Differential Equations 269 (2020), no. 12, 10587–10629. doi:10.1016/j.jde.2020.07.015.
- [GOSU] Y. Giga, J. Okamoto, K. Sakakibara and M. Uesaka, On a singular limit of the Kobayashi–Warren–Carter energy. *Indiana Univ. Math. J.*, accepted for publication.
- [GOU] Y. Giga, J. Okamoto and M. Uesaka, A finer singular limit of a single-well Modica-Mortola functional and its applications to the Kobayashi-Warren-Carter energy. Adv. Calc. Var. 16 (2023), no. 1, 163–182. doi:10.1515/acv-2020-0113.
- [GP] Y. Giga and N. Požár, Motion by crystalline-like mean curvature: a survey. Bull. Math. Sci. 12 (2022), no. 2, Paper No. 2230004, 68 pp. doi:10.1142/S1664360722300043.
- [GSTU] Y. Giga, K. Sakakibara, K. Taguchi and M. Uesaka, A new numerical scheme for discrete constrained total variation flows and its convergence. *Numer. Math.* 146 (2020), 181–217. doi:10.1007/s00211-020-01134-y.
- [Giu] E. Giusti, Minimal surfaces and functions of bounded variation. Monogr. Math., 80. Birkhäuser Verlag, Basel, 1984. xii+240 pp. doi:10.1007/978-1-4684-9486-0.
- [GrL] Z. Grochulska and M. Lasica, Local estimates for vectorial Rudin–Osher–Fatemi type problems in one dimension. *ESAIM Control Optim. Calc. Var.*, forthcoming. doi:10.1051/cocv/2023090.
- [GM] W. Górny and J. M. Mazón, The Neumann and Dirichlet problems for the total variation flow in metric measure spaces. Adv. Calc. Var. 17 (2024), no. 1, 131–164. doi:10.1515/acv-2021-0107.
- [Ka1] Y. Kashima, A subdifferential formulation of fourth order singular diffusion equations. Adv. Math. Sci. Appl. 14 (2004), no. 1, 49–74.
- [Ka2] Y. Kashima, Characterization of subdifferentials of a singular convex functional in Sobolev spaces of order minus one. J. Funct. Anal. 262 (2012), no. 6, 2833–2860. doi:10.1016/j.jfa.2012.01.005.
- [KMR] K. Kielak, P. B. Mucha, and P. Rybka, Almost classical solutions to the total variation flow. J. Evol. Equ. 13 (2013), 21–49. doi:10.1007/s00028-012-0167-x.
- [KWC] R. Kobayashi, J. A. Warren and W. C. Carter, A continuum model of grain boundaries. *Phys. D* 140 (2000), no. 1–2, 141–150. doi:10.1016/S0167-2789(00)00023-3.
- [Kh] R. V. Kohn, Surface relaxation below the roughening temperature: Some recent progress and open questions. Nonlinear Partial Differential Equations: The Abel Symposium 2010, Springer, Heidelberg, 2012, 207–221. doi:10.1007/978-3-642-25361-4_11.

- [Ko] Y. Kōmura, Nonlinear semi-groups in Hilbert space. J. Math. Soc. Japan 19 (1967), 493– 507. doi:10.2969/jmsj/01940493.
- [LMM] M. Lasica, S. Moll and P. B. Mucha, Total variation denoising in l^1 anisotropy. SIAM J. Imaging Sci. 10 (2017), no. 4, 1691–1723. doi:10.1137/16M1103610.
- [Leo] G. P. Leonardi, An overview on the Cheeger problem. New trends in shape optimization, 117–139. Internat. Ser. Numer. Math., 166, Birkhäuser/Springer, Cham, 2015. doi:10.1007/978-3-319-17563-8_6.
- [MST] J. M. Mazón, M. Solera and J. Toledo, The total variation flow in metric random walk spaces. *Calc. Var. Partial Differential Equations* 59 (2020), no. 1, Paper No. 29, 64 pp. doi:10.1007/s00526-019-1684-z.
- [Mol] J. S. Moll, The anisotropic total variation flow. Math. Ann. 332 (2005), no. 1, 177–218. doi:10.1007/s00208-004-0624-0.
- [MS] S. Moll and F. Smarrazzo, The Neumann problem for one-dimensional parabolic equations with linear growth Lagrangian: evolution of singularities. J. Evol. Equ. 22 (2022), article number 79. doi:10.1007/s00028-022-00835-1.
- [MR] P. B. Mucha and P. Rybka, Well posedness of sudden directional diffusion equations, Math. Methods Appl. Sci. 36 (2013), no. 17, 2359–2370. doi:10.1002/mma.2759.
- [ROF] L. I. Rudin, S. Osher and E. Fatemi, Nonlinear total variation based noise removal algorithms. Phys. D 60 (1992), no. 1–4, 259–268. doi:10.1016/0167-2789(92)90242-F.
- [U] M. Uesaka, On a sharp interface limit of the Kobayashi-Warren-Carter functional, in this volume, in preparation.
- [Val] T. Valkonen, The jump set under geometric regularization. Part 1: basic technique and first-order denoising. SIAM J. Math. Anal. 47 (2015), no. 4, 2587–2629. doi:10.1137/140976248.