



Title	Solid-Phase Total Synthesis of Sandramycin and Its Analogues
Author(s)	Komatani, Yuya; Momosaki, Kyoka; Katsuyama, Akira; Yamamoto, Kazuki; Kaguchi, Rintaro; Ichikawa, Satoshi
Citation	Organic letters, 25(3), 543-548 <a href="https://doi.org/10.1021/acs.orglett.2c04327">https://doi.org/10.1021/acs.orglett.2c04327</a>
Issue Date	2023-01-18
Doc URL	<a href="http://hdl.handle.net/2115/91297">http://hdl.handle.net/2115/91297</a>
Rights	This document is the Accepted Manuscript version of a Published Work that appeared in final form in Organic letters, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <a href="https://pubs.acs.org/articlesonrequest/AOR-UJZNIYD6IYBD5B6YGM">https://pubs.acs.org/articlesonrequest/AOR-UJZNIYD6IYBD5B6YGM</a> .
Type	article (author version)
File Information	Text_sasndramycin_revise_3_combine.pdf



[Instructions for use](#)

# Solid-Phase Total Synthesis of Sandramycin and Its Analogues


Yuya Komatani,<sup>1</sup> Kyoka Momosaki,<sup>1</sup> Akira Katsuyama,<sup>\*1-3</sup> Kazuki Yamamoto,<sup>1,2</sup> Rintaro Kaguchi,<sup>1</sup> and Satoshi Ichikawa<sup>\*1-3</sup>

<sup>1</sup>Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan


<sup>2</sup>Center for Research and Education on Drug Discovery, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

<sup>3</sup>Global Station for Biosurfaces and Drug Discovery, Global Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo, 060-0812, Japan

[Paste publication-size TOC graphic here]



**ABSTRACT:** Solid-phase total synthesis of sandramycin (**1**), which is a  $C_2$ -symmetric cyclic decadepsipeptide natural product, and its analogues is described. On-resin ester formation and [5+5] peptide coupling allowed to prepare a range of de-symmetrized analogues. An amino acid residue which would not hamper the biological activity of **1** was successfully identified, and probe molecules and dimeric analogue were prepared based on the result of the structure-activity relationship study.




**Figure 1.** Structure of sandramycin (**1**)

Sandramycin (Figure 1, **1**) belongs to a family of bisintercalator natural products isolated from *Nocardioides* sp. (ATCC 39419).<sup>1</sup> It contains a  $C_2$  symmetric cyclic decadepsipeptide moiety comprising glycine (Gly), sarcosine (Sar), *N*-methyl-L-valine (*N*-Me-Val), D-serine (D-Ser), and L-pipecolic acid (Pip) and possesses two 3-hydroxyquinoline-2-carboxylic acids as a chromophore. It binds to the minor groove of double-stranded DNA with the bisintercalation of its chromophores<sup>2-4</sup> and exhibits potent cytotoxicity against mouse leukemia cell lines *in vitro* and *in vivo*.<sup>1,2</sup> Because of the potent cytotoxicity of bisintercalator natural products, their derivatives have been examined as payloads for antibody-drug conjugates.<sup>5</sup> Hence, it is expected that **1** could be a lead compound as a novel anticancer agent. Two solution-phase total syntheses of **1** have been reported by Boger *et al.*<sup>2</sup> and us.<sup>6</sup> Both methods deployed the  $C_2$  symmetry of **1** to reduce the number of required synthetic steps. Therefore, accessible analogues prepared using these synthetic methods have been limited to those with  $C_2$  symmetry.<sup>6-9</sup> The same limitation has existed for the total synthesis of other bisintercalator natural products.<sup>10-14</sup> This synthetic limitation prevented us from examining the effects of altering single amino acids on the biological activity. Solid-phase peptide synthesis (SPPS) has an advantage of enabling the preparation of various analogues because the protocol simplifies the purification at each step in the linear

synthesis and the amino acids contained in the peptide sequences can be easily exchanged for different amino acids by simply changing the condensed amino acids. Consequently, SPPS is easily extended to the parallel synthesis of analogues. Herein, we report solid-phase total syntheses of **1** and its analogues, which have not been previously elucidated. The method using SPPS allows preparation of desymmetrized analogues for structure-activity relationship studies. Furthermore, chemical probes, which could be used to investigate the mode of action of **1** and the dimeric analogue, were synthesized.

## Scheme 1. Retrosynthetic analysis of **1**



Our retrosynthetic analysis of **1** is illustrated in Scheme 1. As with the solution-phase syntheses, the chromophores would be attached to macrocycle **2** in the final stage of the synthesis. We planned to synthesize the macrocycle by macrolactamization of the linear peptide, which could be obtained from resin-bound

depsipeptide **3**. Considering the synthesis of desymmetrized analogues, we envisioned that the coupling of two pentadepsipeptide units<sup>15-18</sup> (hereafter [5 + 5] peptide coupling) would be a powerful method for the synthesis of **3**. The same fragment coupling methodologies on a solid support were reported in the syntheses of peptide and depsipeptide,<sup>18</sup> which reduced the number of synthetic steps compared with the standard solid-phase synthesis in which each amino acid residue is elongated one by one. The pentadepsipeptide units could be synthesized from resin-bound Fmoc-Pip **6** using SPPS, involving the ester-bond formation step. In this synthetic strategy, the ester formation step and on-resin [5 + 5] peptide coupling would be challenging because of the chemically labile ester bond and the reduced reactivity of pentadepsipeptide due to steric hindrance, respectively. Hence, the optimization of these steps was required to develop a synthetic method, which was applicable for the preparation of various analogues.


The solid-phase total synthesis of **1** is illustrated in Scheme 2a. Fmoc-L-Pip was first loaded on a 2-chlorotrityl (2-CT) resin to afford resin-bound Fmoc-amino acid **6** (loading: 0.82 mmol/g). After the removal of the Fmoc group of **6** with piperidine in *N,N*-dimethylformamide (DMF), the free amine was condensed with **7**. In this coupling, 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMT-MM)<sup>19,20</sup> was superior to suppress the undesired *O*-acylation on the unprotected hydroxy group of **7**, and the desired alcohol **11** was cleanly obtained using this coupling method. Alcohol **11** was condensed with **8** (diisopropylcarbodiimide, 4-dimethylaminopyridine (DMAP), and DMF at 0 °C to room temperature) to afford depsipeptide **12**. Subsequently, 2-methylpiperidine, which is less likely to cause the nucleophilic substitution of the ester due to its steric hindrance,<sup>21-23</sup> was used for Fmoc deprotection instead of piperidine. Removal of the Fmoc group of **12** with 2-methylpiperidine in DMF successfully afforded the corresponding amine without the undesired transamidation. The free *N*-methylamine was further coupled with dipeptide **9** (1-[bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU), *N,N*-diisopropylethylamine (*i*Pr<sub>2</sub>NEt), and DMF) to afford resin-bound pentadepsipeptide **13a** (loading: 0.54 mmol/g, based on the UV measurement). Notably, the choice of carboxylic acid was crucial in this coupling. When Fmoc-Sar-OH was used instead of **9**, the coupling reaction proceeded smoothly to provide **13b**; however, the amine produced by the Fmoc removal of **13b** caused an unacceptable

amount of diketopiperazine formation by the nucleophilic attack of the ester bond via a 6-*exo-trig* cyclization mechanism (Scheme 2b). The side reaction could be suppressed using dipeptide **9**, which did not produce the intermediate causing 6-*exo-trig* cyclization. The


resin-bound depsipeptide **13** could be used as a common intermediate of **4** and **5**. Specifically, the Fmoc removal of **13** with 2-methylpiperidine afforded **4**, and cleavage from the solid support under a weak acidic condition yielded carboxylic acid **5** in 84% overall yield (1% trifluoroacetic acid (TFA)/CH<sub>2</sub>Cl<sub>2</sub>). The on-resin [5 + 5] peptide coupling between **4** and **5** with HATU and *i*Pr<sub>2</sub>NEt in DMF smoothly proceeded to afford **3**. The removal of the Fmoc group on the *N*-terminus of the peptide (2-methylpiperidine, DMF), followed by cleavage from the resin (1% TFA/CH<sub>2</sub>Cl<sub>2</sub>) yielded the cyclization precursor, which was successfully cyclized in the presence of diphenylphosphoryl azide (DPPA) and NaHCO<sub>3</sub> in DMF to afford macrocycle **2** in 41% yield over five steps from **13**. After the removal of the Cbz groups on the  $\alpha$ -nitrogen of the serine residues in **2**, carboxylic acid **14** was condensed to liberate the amines [1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-hydroxy-7-azabenzotriazole (HOAt), NaHCO<sub>3</sub>, and DMF] and produce **1** in 46% yield over two steps. The analytical data of **1** agreed well with those reported.

With the concise synthetic method in hand, we applied the method to the synthesis of various analogues, particularly the unsymmetrical ones. The synthesis of unsymmetrical analogues is illustrated in Scheme 3. A differently protected pentadepsipeptide, **16**, with Boc-protected D-Ser residues, was prepared using the established SPPS protocol (Scheme 3a). Specifically, Cbz-D-Ser **7** in Scheme 2 was changed to Boc-protected D-Ser to produce resin-bound pentadepsipeptide **15**, and cleavage from the resin afforded Boc- and Fmoc-protected pentadepsipeptide **16**. Similarly, resin-bound pentadepsipeptides **17–20** containing L-Val, L-Pro, L-Pip, and *N*-Me-L-Lys residues instead of *N*-Me-L-Val, respectively, were prepared (Scheme 3b). Resin-bound pentadepsipeptide **15** was also used to achieve a chromophore modification. After the Fmoc removal of **15** and **17–20** with 2-methylpiperidine in DMF, the free amines were coupled with Boc-protected depsipeptide **16** or Cbz-protected depsipeptide **5** to afford resin-bound decadepsipeptides. Sequential Fmoc deprotection, cleavage from the resins, and macrolactamization with DPPA yielded cyclic peptides **21–25**, respectively. Two Boc groups of **21–23** were removed using 4 M HCl in 1,4-dioxane to afford amines, and the coupling reaction

**Scheme 2.** Total synthesis of sandramycin (**1**)



### Scheme 3. Synthesis of sandramycin analogues



between the amines and **14** produced unsymmetrical analogues **26–28**, respectively. Similarly, the amine prepared by the Cbz removal of **24** was also condensed with **14** to afford analogue **29** in 80% over two steps. The Boc removal of **25** followed by installation of 3-hydroxyquinoline-2-carbonyl group on the amine gave **30**, and further installation of quinoxaline-2-carbonyl group yielded chromophore-modified analogue **31**. All unsymmetrical analogues **26–29** and **31** were purified by silica gel column chromatography, their chemical structures were confirmed by NMR analyses, and their purity was assured by LC-MS analysis to be sufficient for the following biological evaluation (see the Supporting Information).

The cell growth inhibitory activities of the synthesized analogues are summarized in Table 1 (for details, see the Supporting Information). All side chain-modified analogues **26–29** and

chromophore-modified analogue **31** retained the activity, indicating that the substitution of the *N*-Me-Val to other amino acids did not hamper the cell growth inhibitory activity. Notably, analogue **29** possessed a Boc-protected Lys side chain, and the bulky substituent acceptably exhibited the activity. The result motivated us to use the protected amino group as a starting point for further modifications (Scheme 4a). After the removal of the Boc group of **29**, the amine was acylated with dicarbonate **32** to produce an active ester intermediate. The active ester was further treated with biotin-PEG2-amine **33** to afford biotinylated probe **34**. The free amine corresponding to **31** was also acylated with **35** to produce **36**, which contained Yao's minimalist clickable photo-crosslinker.<sup>24</sup> The chromophore modification shown in Scheme 3b could also be applied to prepare probe molecules. The amine derived from **30** was

condensed with Hosoya's diazide carboxylic acid **37**<sup>25</sup> to afford diazide probe **38**. Analogues **34**, **36** and **38** retained the biological activity of **1** ( $IC_{50} = 29$ , 1.3 and 2.7 nM, respectively); hence, these probe molecules could be used to investigate the mode of action of **1**.

**Table 1.** Cell growth inhibitory activities of **1** and its analogues.

compounds	$IC_{50}$ (nM)
<b>1</b>	0.13
<b>26</b>	1.1
<b>27</b>	3.2
<b>28</b>	1.4
<b>29</b>	0.53
<b>31</b>	0.36
<b>34</b>	29
<b>36</b>	1.3
<b>38</b>	2.7

Dimerization of natural products is an attractive strategy for enhancing their biological activity;<sup>26-29</sup> therefore, we synthesized a homo dimer of **1**. Because the modification of the Lys side chain could be used for further derivatization without the loss of activity, we planned to conjugate two molecules of **1** using the amino groups as linking points (Scheme 4b). After removing the Boc group of **29**, the free amine was coupled with almost a half equivalent of linker compound **39** to afford dimeric analogue **40**. This example illustrates that our SPPS-based synthetic route is a powerful method for synthesizing various types of analogues of **1**. Analyses of the biological activity and the effect of dimerization on the biological activity of **40** are currently in progress.

In conclusion, we developed a method for synthesizing **1** and its analogues using SPPS. The method, involving on-resin [5 + 5]

peptide coupling, can be used for the efficient preparation of various analogues. The desymmetrization realized by the synthesis enables the pinpointing modification of the  $C_2$  symmetric macrocycle of **1**, and the synthesis illustrated here is the first example of the synthesis of an asymmetric analogue of **1**. The structure-activity relationships of the analogues of **1** demonstrated that *N*-Me-Val could be converted to another amino acid residue without the loss of activity. We also prepared chemical probes: two photoaffinity probes, a biotin analogue, and a dimeric analogue, which will be useful to elucidate the mode of action of **1**.

## ASSOCIATED CONTENT

### Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information.

### Supporting Information Statement

The Supporting Information is available free of charge on the ACS Publications website.

Experimental details; <sup>1</sup>H, <sup>13</sup>C NMR and mass spectra, and LC-MS trace (PDF)

## AUTHOR INFORMATION

### Corresponding Author


\*E-mail: [ichikawa@pharm.hokudai.ac.jp](mailto:ichikawa@pharm.hokudai.ac.jp)  
[katsuyama@pharm.hokudai.ac.jp](mailto:katsuyama@pharm.hokudai.ac.jp)

### Author Contributions


The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

## Scheme 4. Synthesis of probe molecules **34**, **36** and **38**, and dimer **40**

### a. Synthesis of probe molecules



### b. Synthesis of dimer **40**



## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENT

We thank Takamitsu Hosoya (Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University) for providing compound 37. This research was supported by JSPS KAKENHI Grant-in-Aid for Scientific Research (B) (S.I., Grant Number 22H02738 and JP19H03345), Grant-in-Aid for Scientific Research on Innovative Areas "Frontier Research on Chemical Communications" (S.I., JP20H04757), JSPS KAKENHI Grant-in-Aid for Early-Career Scientist (A.K., Grant Number JP22K15241), Japan Agency for Medical Research and Development (AMED) under Grant Number 21ak0101118h9903, JST the establishment of university fellowships towards the creation of science technology innovation (Y.K., JPMJFS2101), North Tech Foundation, Takeda Science Foundation and was partly supported by Hokkaido University, Global Facility Center (GFC), Pharma Science Open Unit (PSOU), funded by MEXT under "Support Program for Implementation of New Equipment Sharing System", Platform Project for Supporting Drug Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS)) from AMED under Grant Number JP22ama121039, 18ae0101047h0001 and 19ae0101047h0002.

## REFERENCES

- Matson, J. A.; Bush, J. A. Sandramycin, a novel antitumor antibiotic produced by a *Nocardioideis sp.* *J. Antibiot.* **1989**, *42*, 1763–1767.
- Boger, D. L.; Chen, J.-H.; Saionz, K. W. (–)-Sandramycin: total synthesis and characterization of DNA binding properties. *J. Am. Chem. Soc.* **1996**, *118*, 1629–1644.
- Dawson, S.; Malkinson, J. P.; Paumier, D.; Searcey, M. Bisintercalator natural products with potential therapeutic applications: isolation, structure determination, synthetic and biological studies. *Nat. Prod. Rep.* **2007**, *24*, 109–126.
- Zolova, O. E.; Mady, A. S. A.; Garneau-Tsodikova, S. Recent developments in bisintercalator natural products. *Biopolymers* **2010**, *93*, 777–790.
- Ratnayake, A. S.; Chang, L.-p.; Tumey, L. N.; Loganzo, F.; Chemler, J. A.; Wagenaar, M.; Musto, S.; Li, F.; Janso, J. E.; T. Ballard, Eric.; Rago, B.; Steele, G. L.; Ding, W.; Feng, X.; Hosselet, C.; Buklan, V.; Lucas, J.; Koehn, F. E.; O'Donnell, C. J.; Grazianiand, E. I. Natural product bis-intercalator depsipeptides as a new class of payloads for antibody–drug conjugates. *Bioconjugate Chem.* **2019**, *30*, 200–209.
- Katayama, K.; Nakagawa, K.; Takeda, H.; Matsuda, A.; Ichikawa, S. Total synthesis of sandramycin and its analogues via a multicomponent assemblage. *Org. Lett.* **2014**, *16*, 428–431.
- Boger, D. L.; Chen, J. H.; Saionz, K. W.; Jin, Q. Synthesis of key sandramycin analogs: systematic examination of the intercalation chromophore. *Bioorg. Med. Chem.* **1998**, *6*, 85–102.
- Boger, D. L.; Chen, J.-H. An exceptionally potent analog of sandramycin. *Bioorg. Med. Chem. Lett.* **1997**, *7*, 919–922.
- Boger, D. L.; Saionz, K. W. DNA binding properties of key sandramycin analogues: systematic examination of the intercalation chromophore. *Bioorg. Med. Chem.* **1999**, *7*, 315–322.
- Boger, D. L.; Ledebøer, M. W.; Kume, M. Total synthesis of luzopeptins A–C. *J. Am. Chem. Soc.* **1999**, *121*, 1098–1099.
- Boger, D. L.; Ledebøer, M. W.; Kume, M.; Searcey, M.; Jin, Q. Total synthesis and comparative evaluation of luzopeptin A–C and quinoxapeptin A–C. *J. Am. Chem. Soc.* **1999**, *121*, 11375–11383.
- Ciufolini, M. A.; Valognes, D.; Xi, N. Total synthesis of luzopeptin E2. *Angew. Chem. Int. Ed.* **2000**, *39*, 2493–2495.
- Valognes, D.; Belmont, P.; Xi, N.; Ciufolini, M. A. Total synthesis of luzopeptin C. *Tetrahedron Lett.* **2001**, *42*, 1907–1909.
- Boger, D. L.; Ledebøer, M. W.; Kume, M.; Jin, Q. Total synthesis of quinoxapeptin A–C: establishment of absolute stereochemistry. *Angew. Chem. Int. Ed.* **1999**, *38*, 2424–2426.
- Ichikawa, S.; Okamura, T.; Matsuda, A. Total synthesis of quinaldopeptin and its analogues. *J. Org. Chem.* **2013**, *78*, 12662–12670.
- Garcia-Martin, F.; Cruz, L. J.; Rodriguez-Mias, R. A.; Giralt, E.; Albericio, F. Design and synthesis of FAJANU: a de novo C2 symmetric cyclopeptide family. *J. Med. Chem.* **2008**, *51*, 3194–3202.
- Marcucci, E.; Tulla-Puche, J.; Albericio, F. Solid-phase synthesis of NMe-IB-01212, a highly N-methylated cyclic peptide. *Org. Lett.* **2012**, *14*, 612–615.
- Xiong, X. F.; Zhang, H.; Underwood, C. R.; Harpsoe, K.; Gardella, T. J.; Woldike, M. F.; Mannstadt, M.; Gloriam, D. E.; Brauner-Osborne, H.; Stromgaard, K. Total Synthesis and Structure-Activity Relationship Studies of a Series of Selective G Protein Inhibitors. *Nat. Chem.* **2016**, *8*, 1035–1041.
- Kunishima, M.; Kawachi, C.; Morita, J.; Terao, K.; Iwasaki, F.; Tani, S. 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride: an efficient condensing agent leading to the formation of amides and esters. *Tetrahedron* **1999**, *55*, 13159–13170.
- Kunishima, M.; Kawachi, C.; Hioki, K.; Terao, K.; Tani, S. Formation of carboxamides by direct condensation of carboxylic acids and amines in alcohols using a new alcohol- and water-soluble condensing agent: DMT-MM. *Tetrahedron* **2001**, *57*, 1551–1558.
- Hachmann, J.; Lebl, M. Alternative to piperidine in Fmoc solid-phase synthesis. *J. Comb. Chem.* **2006**, *8*, 149.
- Kralt, B.; Moreia, R.; Palmer, M.; Taylor, S. D. Total synthesis of analogs of A54145D and A54145A1 for structure–activity relationship studies. *J. Org. Chem.* **2020**, *85*, 2213–2219.
- Takashina, K.; Katsuyama, A.; Kaguchi, R.; Yamamoto, K.; Sato, T.; Takahashi, S.; Horiuchi, M.; Yokota, S.-i.; Ichikawa, S. Solid-phase total synthesis of plusbacin A<sub>3</sub>. *Org. Lett.* **2022**, *24*, 2253–2257.
- Li, Z.; Hao, P.; Li, L.; Tan, C. Y. J.; Cheng, X.; Chen, G. Y. J.; Sze, S. K.; Shen, H.-M.; Yao, S. Q. Design and synthesis of minimalist terminal alkyne-containing diazirine photo-crosslinkers and their incorporation into kinase inhibitors for cell- and tissue-based proteome profiling. *Angew. Chem. Int. Ed.* **2013**, *52*, 8551–8556.
- Hosoya, T.; Hiramatsu, T.; Ikemoto, T.; Nakanishi, M.; Aoyama, H.; Hosoya, A.; Iwata, T.; Maruyama, K.; Endo, M.; Suzuki, M. Novel bifunctional probe for radioisotope-free photoaffinity labeling: compact structure comprised of photospecific ligand ligation and detectable tag anchoring units. *Org. Biomol. Chem.* **2004**, *2*, 637–641.
- Luther, A.; Urfer, M.; Zahn, M.; Müller, M.; Wang, S.-Y.; Mondal, M.; Vitale, A.; Hartmann, J.-B.; Sharpe, T.; Monte, F.; Lo Kocherla H.; Cline, E.; Pessi, G.; Rath, P.; Modaresi, S. M.; Chiquet, P.; Stiegeler, S.; Verbree, C.; Remus, T.; Schmitt, M.; Kolopp, C.; Westwood, M.-A.; Desjonquères, N.; Brabet, E.; Hell, S.; LePoupon, K.; Vermeulen, A.; Jaisson, R.; Rithié, V.; Upert, G.; Lederer, A.; Zbinden, P.; Wach, A.; Moehle, K.; Zerbe, K.; Locher, H. H.; Bernardini, F.; Dale, G. E.; Eberl, L.; Wollscheid, B.; Hiller, S.; Robinson, J. A.; Obrecht, D. Chimeric peptidomimetic antibiotics against gram-negative bacteria. *Nature* **2019**, *576*, 452–458.
- Sundram, U. N.; Griffin, J. H.; Nicas, T. I. Novel vancomycin dimers with activity against vancomycin-resistant enterococci. *J. Am. Chem. Soc.* **1996**, *118*, 13107–13108.
- Wood, T. M.; Slingerland, C. J.; Martin, N. I. A convenient chemoenzymatic preparation of chimeric macrocyclic peptide

antibiotics with potent activity against gram-negative pathogens. *J. Med. Chem.* **2021**, *64*, 10890-10899.

(29) van Groesen, E.; Slingerland, C. J.; Innocenti, P.; Mihajlovic M.; Masereeuw, R.; Martin, N. I. Vancomyxins: Vancomycin-

polymyxin nonapeptide conjugates that retain anti-gram-positive activity with enhanced potency against gram-negative strains. *ACS Infect. Dis.* **2021**, *7*, 2746-2754.

---

