



Title	Some new Butterflies
Author(s)	Matsumura, S.
Citation	Insecta matsumurana, 2(2), 114-118
Issue Date	1927-11
Doc URL	http://hdl.handle.net/2115/9139
Type	bulletin (article)
File Information	2(2)_p114-118.pdf



[Instructions for use](#)

10. <i>Cardiophorus yanagiharae</i> MIWA (n. sp.)	ヤナギハラハナコメツキ
11. <i>Paracardiophorus nigro-apicalis</i> MIWA (n. sp.)	ツماغロハナコメツキ
12. <i>Neodiploconus angulatus</i> CANDÈZE	チヤイロホソクシコメツキ
13. <i>Neodiploconus melanopterus</i> CANDÈZE	アカアシホソクシコメツキ
14. <i>Neodiploconus cyprianus</i> CANDÈZE	アカホソクシコメツキ
15. <i>Neodiploconus ferrugineipennis</i> MIWA (n. sp.)	アカバネホソクシコメツキ
16. <i>Spheniscosomus pilosulus</i> MIWA (n. sp.)	ケブカオホクシコメツキ
17. <i>Spheniscosomus horishanus</i> MIWA (n. sp.)	ホリシヤオホクシコメツキ
18. <i>Spheniscosomus fuscus</i> FABRICIUS	ヒラタクシコメツキ
19. <i>Melanotus ventralis</i> CANDÈZE	ハラアカクシコメツキ
20. <i>Melanotus castanipes</i> PAYKULL	ハネナガクシコメツキ (改稱)
21. <i>Melanotus legatus</i> CANDÈZE	クシコメツキ (改稱)
22. <i>Cremnostethus telum</i> CANDÈZE	オホナガクシコメツキ
23. <i>Metriaulacus formosanus</i> MIWA (n. sp.)	ホソツヤクシコメツキ

SOME NEW BUTTERFLIES

By

PROF. DR. S. MATSUMURA

Papilio memnon thunbergi SIEB. (Pl. III, fig. 1, ♀)

ab. *takamukai* n. ab.

♀. This differs from *thunbergi mela* FRUHST. in the following characters:

1. Secondaries on the upperside lacks entirely the white markings.
2. Secondaries on the underside with a few whitish part in the interspace 2 at the innerside of the upper black patch, which is wanted by *thunbergi* SIEB.

This resembles also subsp. *agenor* L., but differs in lacking a broad band which runs along the inner margin of the secondaries.

Hab.—Kiushu; one single specimen was collected by T. TAMANUKI at Kagoshima.

ab. *shikokuana* n. ab. (Pl. III, fig. 6, ♀)

♀. Differs from subsp. *thunbergi* SIEB. as follows:

1. Reddish patch at the base on the upperside of the primaries smaller and narrower.
2. 4 discal white patches only on the upperside of the secondaries, while on the underside with only 3; underside of the secondaries lacks reddish lunular markings near the middle of each interspace, which are conspicuous in *thunbergi* SIEB.

Hab.—Shikoku; one single specimen was collected on the 20th of August, 1926, by K. FUJII at Uwajima.

***Papilio taiwanus watanabei* n. ab. (Pl. III, fig. 2, ♀)**

♀. Differs from the typical *taiwanus* ROTHSCH. in lacking rosy lunular spot-series which runs from the anal angle to the patch of the 5th interspace, the latter being smaller and that of the 6th truncated at the outside.

Hab.—Formosa; one single specimen was collected in 1906 at Mt. Gakoki by the late K. WATANABE, chief of policemen at Hoppo.

***Papilio maackii jezoensis* n. subsp.**

♂. Differs from the subspecies *raddei* BREM. in the following points:

Upperside—Secondaries with a much broader bluish postdiscal band, at the costal half being concolorous with the ground colour, while in the typical *raddei* the band greenish and the costal half bluish.

Underside—Primaries with the outer $\frac{1}{3}$ whitish grey, only at the extreme termen being fuscous; secondaries with a broader whitish band which is somewhat expanded at the interspace 3, being nearly twice as broad as that of *raddei*.

Hab.—Hokkaido; 2 male specimens were collected on the 28th of June, 1897, at Atsubetsu near Sapporo.

ab. ***iss hikii* n. ab.**

♂. Differs from *jezoensis* MATS. in having a greenish blue band on the upperside of secondaries and a much narrower whitish band on the underside of primaries.

Hab.—Hokkaido; one male specimen was collected on the latter part of May, 1918, by S. ISSHIKI.

***Papilio maackii mandschurica* n. subsp.**

Differs from *raddei* as follows:

♀. Upperside—Primaries dark brown, scattered with very few golden green scales; postdiscal band pale yellowish green, being somewhat broader than that of *raddei*. Secondaries with much larger reddish lunules, the greenish

postdiscal band somewhat narrower and at the costal half not bluish coloured as in *raddei*.

Underside—Primaries whitish grey, with a few fuscous scales, a narrow part of the termen and longitudinal stripes in the interspaces as well as in the discoidal cell being fuscous. Secondaries with very large reddish ocellar patches, pupilated with each a velvety black spot.

Hab.—Manchuria; one female specimen was collected on the 14th of July, 1924, by D. MIURA at Koshurei.

Hypolimnas bolina inornata n. ab. (Pl. III, fig. 8, ♂)

♂. Differs from the typical specimen in having no whitish blue patch on the upperside of secondaries.

Hab.—Formosa; one male specimen was collected on the 21st of July, 1918, by S. WATARI at Horisha.

Argynnis paphia formosicola n. subsp.

Closely allied to *paphioides* BUTL., but differs from it in the following points:

♂. Upperside—Primaries somewhat broader, deeper in colour, black patches much larger, those of the marginal series especially larger, being mostly confluent with the marginal black line; in the interspace 1^b and 2 with each 3 black patches, while in *paphioides* lacks the inner spot. Secondaries with also much larger spots in the marginal region.

Underside—Patches of primaries correspond to those of the upperside and much larger; apex dark greenish with 2 or 3 broadly yellowish patches in it. Secondaries more deeply dark green with broader silvery bands, that of the 2nd being broad; on the postmedial region with a series of 4 fuscous spots, having each a paler pupil; at the submarginal region with a dark purplish line.

Hab.—Formosa; one male specimen was collected on the 12th of July, 1925, by T. UCHIDA, Y. MIWA and H. KÔNO.

Argynnis selene chibiana n. subsp. (Pl. III, fig. 4, ♀)

Closely allied to *selene dilutior* FIX., but differs from it as follows:

♂, ♀. Upperside—Primaries with much larger black spots especially in the cell, at the base being distinctly infuscated; black spots of the marginal series larger and roundish, defined inwardly by a series of roundish yellowish spots. Secondaries at the basal $\frac{1}{3}$ infuscated, so that the basal spot of the cell being invisible.

Underside—Markings of the primaries correspond to those of the upperside, being distinctly larger; secondaries with 2 large brownish patches, respectively,

one near the apex and another near the anal angle, so that the black spots of the submarginal region not conspicuous.

Hab.—Kuriles; numerous male and female specimens were collected on the first part of August, 1927, by K. DOI at Horomoshuru.

Melitaea matura konumensis n. subsp. (Pl. III, fig. 5, ♀)

Differs from the typical specimens as follows:

♀, ♂. Upperside—Yellowish spots of the marginal series to primaries much larger, those of the female being similar to those of the submarginal series; pale yellowish spots of the marginal series to secondaries very small and not conspicuous, spots of the cell to primaries being much larger.

Underside—Postdiscal yellowish band towards the dorsum not divided into 2 as in the typical specimen.

Exp.—♂ 46 mm.; ♀ 50–56 mm.

Hab.—Saghalien; 4 (2 ♂, 2 ♀) were collected on 6th of July, 1927, by Y. MIWA and S. MINOWA at Konuma.

Nom. Jap.:—*Konuma-hyomon-modoki*.

Thecla eximia watarii n. subsp. (Pl. III, fig. 3, ♀)

Closely allied to *eximia fixeni* LEECH, but differs from it in the following points:

♀. Upperside—Primaries in the middle with a transverse, conical, orange patch which becomes suddenly narrower from the discocellulars towards the base, and which extends beyond the postmedial line and ends at the fork of the 2nd and median veins. Secondaries with 2 orange spots, respectively, one at the anal angle and another near the end of the interspace 2, that of the anal angle being obsolete.

Underside—Nearly the same with that of *fixeni* LEECH, but the submarginal series of fuscous spots to primaries prominent, which becomes smaller towards the interspace 6, that of 6 being very obsolete, and some orange spots being visible near the hind angle.

Hab.—Formosa; one male specimen was collected on the 14th of June, 1918, by S. WATARI at Horisha.

Taraka hamada formosana n. subsp. (Pl. III, fig. 7, ♂)

Closely allied to subsp. *jezoensis* MATS.*, but differs from it in the following points:

Much smaller in size: ♂ 20 mm., ♀ 25 mm.

♂, ♀. Primaries in the middle scarcely paler. Underside with much smaller

*Thousand Insects of Japan, Add. III, p. 735 (1917).

black spots, none of which are not connected with each other; termen of the primaries with only 2 or 3 spots, while that of the secondaries having none.

Antennae on the inner sides orange yellow.

Hab.—Formosa; 3 (1 ♂, 2 ♀) specimens were collected on the 25th of October, 1926, by K. KIKUCHI at Baibara in the district of Horisha.

Lycaena arionides sugitanii n. subsp. (Pl. III, fig. 10, ♂)

This differs from the typical specimens as follows:

♂. Primaries with a very broad black margin (about 3 mm. in width); discoidal lunule large and not divided into 2 as in the figure 1^b* of the typical specimen given by STAUDINGER; underside along the submarginal spot-series much infuscated.

Secondaries with a similar broad black margin as that of the primaries, so that the spot-series on the submarginal region indistinct, having a series of bluish spots on that place; black discal spots distinctly larger; underside with larger spots in a series.

Hab.—Corea; 2 male specimens were collected on 30th, July, 1927, by Prof. I. SUGITANI at Choanji.

Nom. Jap. :—*Chosen-ogoma-shijimi*.

Lycaena euphemus shiriyensis n. subsp. (Pl. III, fig. 9, ♂)

Closely allied to *euphemus ogumae* from Saghalien, but differs from it in having a broad black margin to both wings (about 3 mm. in primaries and 4 mm. in secondaries); underside much darker, but somewhat paler than that of *kazamoto* DRUCE.

Hab.—Honshu (Aomori); 6 (5 ♂, 1 ♀) specimens were collected on 8th, August, 1927, at Shiriya by H. KONO.

*Rom. Mém. III, p. 141, pl. VII, fig. 1 a, b, c (1887).

摘 要

本文には九の亞種と五の異常形を記載せり。此内コマヘウモンモドキミテフセンオホゴマシジミとは種として日本領土内で初めての發表なり。此等の標本は高椋悌吉、渡正監、杉谷岩彦、三浦道哉、内田登一、三輪勇四郎、河野廣道、土井久作、箕輪重胤、菊地寛治、藤井吉之進諸氏の採集に係るものなれば、爰に同諸氏に向つて深謝の意を公表す。

EXPLANATION OF PLATE III

1. *Papilio memnon thunbergi takamukui* MATS. (n. ab.) ♀
 2. *P. thaiwanus watanabei* MATS. (n. ab.) ♀
 3. *Thecla eximia watarii* MATS. (n. subsp.) ♀
 4. *Argynnis selene chibiana* MATS. (n. subsp.) ♀
 5. *Melitaea matura konumensis* MATS. (n. subsp.) ♀
 6. *Papilio memnon shikokuana* MATS. (n. ab.) ♀
 7. *Taraka hamada formosana* MATS. (n. subsp.) ♂
 8. *Hypolimnas bolina inornata* MATS. (n. ab.) ♂
 9. *Lycaena euphemus shiriensis* MATS. (n. subsp.) ♂
 10. *L. arionides sugitanii* MATS. (n. subsp.) ♂
-

