



Title	A new Butterfly for Japan
Author(s)	Matsumura, S.
Citation	Insecta matsumurana, 5(1-2), 100-100
Issue Date	1930-11
Doc URL	http://hdl.handle.net/2115/9211
Type	bulletin (article)
File Information	5(1-2)_p100-100.pdf



[Instructions for use](#)

weissen Ring. 3) Zwei Basalhinterleibssegmente etwas bräunlich. Körperlänge: ♀ ♂ 5–6 mm.

Fundorte: Honshu (Tokio), Shikoku (Tosa), Kiushu (Miyazaki), Korea (Suigen); gesammelt in 5 (♀ 4, ♂ 1) Exemplaren von Y. SUGIHARA, M. YAMANAKA, S. FUJII, und K. SATÔ.

Anmerkung: Das Individuum, welches sich mehr oder minder von der typischen Art unterscheidet, habe ich bis heute als eine Varietät geschrieben, von nun an aber möchte ich dieses als eine Form der Stammart behandeln.

今回本邦産姫蜂亞科の一新屬、八新種及び三新型を發見したれば、茲に公表すべし。而してこれ等の和名は次の如し。

<i>Protichneumon karenkoensis</i> UCH. (sp. nov.)	カレンコウクロヒメバチ (新稱)
<i>Coelichneumon ohtai</i> UCH. (sp. nov.)	オホタヒメバチ (新稱)
<i>Stenichneumonoides</i> UCH. (gen. nov.)	
<i>Stenichneumon nigrimarginalis</i> UCH. (sp. nov.)	クロメブチヒメバチ (新稱)
<i>S. nudicoxalis</i> UCH. (sp. nov.)	ツヤケシクロヒメバチ (新稱)
<i>Ctenichneumon maruyamensis</i> UCH. (sp. nov.)	マルヤマサキブトヒメバチ (新稱)
<i>Spilichneumon doii</i> UCH. (sp. nov.)	フキサキブトヒメバチ (新稱)
<i>S. johnsoni</i> HOLMGR. form. <i>satoi</i> UCH. (form. nov.)	フトヒゲサキブトヒメバチ (新稱)
<i>Platylabus semimarginalis</i> UCH. (sp. nov.)	ヒラタヘリヒメバチ (新稱)
<i>P. takeuchii</i> UCH. (sp. nov.)	タケウチヒメバチ (新稱)
<i>Neotypus lapidator</i> F. form. <i>orientalis</i> UCH. (form. nov.)	セアカヒメバチ (新稱)
<i>N. melanocephalus</i> GEMEL. form. <i>coreensis</i> UCH. (form. nov.)	オホセアカヒメバチ (新稱)


A NEW BUTTERFLY FOR JAPAN

By

PROF. S. MATSUMURA

Hesperia bieti OBERTH. f. *tokachiana* n.

Differs from the typical specimens in having much larger yellowish white spot-series on the submarginal region to primaries, lacking a y-shaped dark brown marking to secondaries, underside, which becomes a roundish patch provided with a paler speck in the middle.



Hesperia bieti OBERTH.
f. *tokachiana* n.
record for Japan.

Hab.—Hokkaido; two male specimens were collected by H. KINOSHITA at Taisho-mura in the Prov. Tokachi and one of them was set at my dispersal by Mr. S. KUWAYAMA.

This butterfly is very common in China, but first