



Title	New Species of Insects from Japan described during the Year 1930
Author(s)	Matsumura, S.
Citation	Insecta matsumurana, 5(4), 211-217
Issue Date	1931-07
Doc URL	http://hdl.handle.net/2115/9224
Type	bulletin (article)
File Information	5(4)_p211-217.pdf



[Instructions for use](#)

ries with a large areole, from the apex of it emitting veins $7+8+9$, the stalk of which being quite short; at the termen in the interspaces 4 and 5 to the primaries and in the interspace 4 to the secondaries somewhat excavated; veins 4 and 5 to the secondaries distinctly stalked. Legs slender, each tibia at the hind part being less and shorter pubescent, the hind tarsus with numerous longer and robuster spines.

Genotype: *Noshimea fulguralis* MATS.

NEW SPECIES OF INSECTS FROM JAPAN DESCRIBED DURING THE YEAR 1930

HYMENOPTERA

GUSSAKOVSKIJ, B. B.: Revue des espèces palaearctiques du genre *Priocnemis* Schdte. (Ann. Mus. Zool., Leningrad, Tom. XXXI, pp. 227-290).

Priocnemis japonicus, p. 252.

ONOE, T. et YUASA, H.: Ptilineurus marmoratus, an Anoblid-beetle noxious to the Rush Mat, with Descriptions of a new Bethylid Fly (Rep. Journ. Imp. Agr. Expt. Sta. Tokyo, Vol. I, No. 3, pp. 215-230).

Sclerodermus nipponicus, p. 218.

SATO, K.: A new Species of the Genus *Megaxyela* Ash. from Korea (Jour. Chosen Nat. Hist. Soc. No. XI, p. 4-8).

Megaxyela mikado, p. 4.

SONAN, J.: A few Host-known Ichneumonidae found in Japan and Formosa (Trans. Nat. Hist. Soc. Formosa, Vol. XX, pt. 30, pp. 268-273).

Charops arctornisae, p. 268; *Anilasta japonica*, p. 269; *Pezomachius lemae*, p. 271; *Cryptus flavicoxatus*, p. 272.

———: Some new Species of Hymenoptera in Japanese-Empire, with two known Species (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 111, pp. 355-360).

Odynerus (Lioutus) flaviclypeatus, p. 355; *O. (Hoplomerus) nigriclypeatus*, p. 356; *Rhynchium satsumae* p. 356; *Habrocyptus ruficoxatus*, p. 357; *Batithrix kuwanae* VIER. var. *nigrans*, p. 358; *Batithrix rufus*, p. 358; *Mecha lemae*, p. 359.

———: A few Host-known Ichneumonidae found in Formosa (2), (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 108, pp. 137-144).

Itopectis homonae, p. 137; *Iseropus heichinus*, p. 138; *Anomalon discoidellum*, p. 139; *Limnerium homonae*, p. 140; *Cremastus shirakii*, p. 142; *Gambrus homonae*, p. 143.

TAKEUCHI, K.: A revisional List of the Japanese Pamphiliidae, with Descriptions of nine new Species (Trans. Kansai Ent. Soc., No. 1, pp. 3-16).

Cephaleia issikii, p. 6; *C. variegata*, p. 7; *Neurotoma atrata*, p. 8; *Pamphilus komonensis*, p. 12; *P. minomalis* p. 12; *P. viriditibialis*, p. 13; *P. kamikochiensis*, p. 14; *P. nakagawai*, p. 15; *P. jucundus*, p. 16.

TOZAWA, N.: On *Oryssus japonicus* sp. nov. (Kontyu, Vol. IV, No. 3, pp. 160-162).

Oryssus japonicus, p. 160.

UCHIDA, T.: Beitrag zur Kenntnis der Ichneumoniden-fauna Insel Izu-Ohshima (Trans. Sapporo Nat. Hist. Soc. Nol. XI, pt. 2, pp. 81-86)).

Cryptus ohshimensis, p. 81; *Henicospilus yamanakai*, p. 83; *Campoplex ohshimensis*, p. 85; *Pimpla portheretiae* VIER. f. *neustriae*, p. 86.

_____: Vierter Beitrag zur Ichneumoniden-fauna Japans (Journ. Fac. Agr. Hokkaido Imp. Univ., Vol. XXV, pt. IV, pp. 243-298).

Sphinctus chinensis, p. 245; *Metopius coreanus*, p. 248; *M. maruyamensis*, p. 249; *M. baijarensis*, p. 250; *M. iyoensis*, p. 252; *M. takabayashii*, p. 253; *M. arakawai*, p. 253; *M. sapporensis*, p. 254; *Homocidus iwatensis*, p. 257; *H. satoi*, p. 257; *H. yokohamensis*, p. 258; *H. takaosanus*, p. 259; *Colpotrochia nipponensis*, p. 262; *Colpotrochoides* (n. g.), p. 263; *C. orientalis*, p. 264; *C. kurisuei*, p. 264; *Metacoelus formosanus*, p. 266; *Triclistus fuscofascialis*, p. 267; *Exochus oshimensis*, p. 267; *Tricamptus rufipes*, p. 269; *Eexenterus coreensis*, p. 270; *Tryphon akaashii*, p. 272; *T. semirufus*, p. 273; *T. T. jezoensis*, p. 273; *T. nagahamensis*, p. 274; *Pseudasthenara* (n. g.), p. 276; *Ctenopelma tokioensis*, p. 278; *Homaspis kuriensis*, p. 279; *Prionopoda ewaniensis*, p. 280; *P. otaruensis*, p. 281; *Periüssus sachalinensis*, p. 282; *P. miyoshii*, p. 284; *Mesoleptus shibuyai*, p. 284; *M. ogasawarai*, p. 284; *M. sapporensis*, p. 285; *Sychnoporthus nipponicus*, p. 287; *Daisetsuzania* (n. g.), p. 289; *D. albifrons*, p. 289; *Alexeter shakojiensis*, p. 292; *A. dorogawensis*, p. 292; *Mesoleius nikkoensis*, p. 294; *Scopesus sachalinensis*, p. 294.

_____: Fünfter Beitrag zur Ichneumoniden-fauna Japans (Journ. Fac. Agr., Hokkaido Imp. Univ., Vol. XXV, pt. IV, pp. 299-347).

Megaplectes monticola GRAV. var. *dentata*, p. 304; *Cryptus tenuiabdominalis*, p. 307; *C. nipponensis*, p. 308; *Goniocryptus yezoensis*, p. 311; *G. tokioensis*, p. 311; *Kaltenbachia bidentata*, p. 312; *Habrocyptus assimiliis*, p. 313; *Spilocryptus japonicus*, p. 314; *Gambrus sapporensis*, p. 315; *Listrognathus coreensis*, p. 316; *Friiona okinawana*, p. 319; *Nematopodius flavoguttatus*, p. 320; *Orientostenaraeus* (n. g.) *chinensis*, p. 312; *Piectocryptus nohirai*, p. 324; *P. jezonicus*, p. 325; *P. saitamensis*, p. 326; *P. hokkaidensis*, p. 327; *P. albibaldis*, p. 327; *Giraudia teranishi*, p. 328; *Microcryptus sapporensis*, p. 329; *Acanthocryptus coreensis*, p. 330; *Stylocryptus osakensis*, p. 332; *S. satoi*, p. 333; *Phygadeuon suigensis*, p. 336; *P. maruyamanus*, p. 336; *P. daisetsuzanus*, p. 337; *P. akaashii*, p. 338; *Ischnocryptus elongatus*, p. 338; *I. similis*, p. 339; *Hemiteles chosensis*, p. 340; *H. kumamotensis*, p. 341; *H. akannensis*, p. 341; *H. oshimensis*, p. 342; *H. narangae* = *H. (Bathythrix) kuwanae* VIER., p. 342; *H. aporiae*, p. 343; *Pezomachus asozanus*, p. 344.

COLEOPTERA

ADACHI, T.: On the Japanese Species of the Genera *Onthophilus* and *Hololepta*, with Description of a new Species (Kontyū, Vol. IV, No. 4, pp. 247-252).

Onthophilus kamiyai, p. 250.

FLEUTIAUX, M. E.: Description d'un *Campsosternus* nouveau de la collection du Muséum National d'Histoire Naturelle de Paris (Bull. Mus. Hist., No. 4, 1930).

Campsosternus mirabilis, p. 409.

KANO, T.: A List of Benthidae from the Japanese Empire, with Description of a new Species (Kontyū, Vol. IV, No. 3, pp. 151-154).

Calodromus horishanus, p. 153.

_____: Formosan Paussidae, with Description of two new Species (Kontyū, Vol. IV, No. 2, pp. 90-98).

Paussus elongatus, p. 92; *P. horikawae*, p. 93.

_____: Contribution to the Carabideous Fauna of Formosa (1) (Trans. Nat. Hist. Soc. Formosa, vol. XX, no. 101, pp. 24-31).

Nebria niitakana, p. 25; *Galerita formosana*, p. 29.

NEW SPECIES OF INSECTS FROM JAPAN DESCRIBED DURING THE YEAR 1930 213

KÔNO, H.: Die Cleriden Japans (Trans. Sapporo Nat. Hist. Soc., Vol. XI, pt. 3, pp. 134-140).
Tillus igarashii, p. 134; *Thanasimus karaftonis*, p. 135; *Tenerus castaneus*, p. 136.

— : Cerogria miwai, nova specie (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 106, pp. 18-19).

Cerogria miwai, p. 18.

— : Kurzrüssler aus dem japanischen Reich (Jour. Facul. Agr., Hokkaido Imp. Univ., Sapporo, Vol. XXIV, Pt. 5, pp. 163-237).

Scytropus iishikii, p. 169; *Calapionus viridimetallicus* MOTSCH. f. *frequens* (n. f.), 172; *C. viridimetallicus* MOTSCH. f. *naipoensis* (n. f.), *jezoensis* (n. f.), *C. oblongus*, p. 173; *Chlorophanus grandis* ROEL. f. *konumensis* (n. f.), *C. rugicollis*, p. 182; *Scepticus minowai*, p. 185; *S. uniformis*, p. 186; *S. hachijoensis*, p. 187; *Sympiezomias cribicollis*, p. 189; *Pachyrhynchus sarcitis* BEHRENS subsp. *kotoensis* (n. subsp.), p. 193; *P. sonani*, p. 194; *Phyllotius brevitarsis*, p. 205; *Myllocerus sakaguchi*, p. 212; *M. canoixoides*, p. 213; *M. rostralis*, p. 213; *M. setosus*, p. 214; *M. aceri*, p. 217; *M. tosanus*, *M. otsukai*, p. 218; *M. kuchibutonis*, p. 219; *Episomoides* (n. g.), p. 220; *Calomycterus variabilis*, p. 224.

— : Die biologischen Gruppen der Rhynchitinen, Attelabinen und Apoderinen (Journ. Facul. Agr. Hokkaido Imp. Univ., Sapporo, Vol. XXIX, Pt. 1, pp. 1-36).

Aderorhinus venustus, p. 29; *Eugnamptus* (*Eugnamptobius*) *distinctus*, p. 30; *Auletobius nitens*, p. 32.

— : Die Apoderinen aus dem japanischen Reich (Jour. Facul. Agr., Hokkaido Imp. Univ., Sapporo, Vol. XXIX, Pt. 2 pp. 35-83).

Paroplapoderus pardalis VOLLENH. f. *abdominalis* (n. f.), f. *nakamurai* (n. f.) p. 47; *Comadaranus* (n. subg.), p. 48; *Paroplapoderus* (*Erycapoderus*) *validus* VOSS *nigroguttatus* (n. subsp.), p. 51; *Apoderus jekeli* ROEL. f. *sapporensis* (n. f.), *A. okamotonis*, p. 58; *A. mushanus*, p. 59; *A. (Compsapoderus) nitens* ROEL. f. *abdominalis* (n. f.), f. *shimauchiensis* (n. f.), p. 67; *Paratrachelophorus vossi*, p. 75; *P. katonis*, p. 76.

Miwa, Y.: Elateridae of Formosa (III) (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 106, pp. 1-12).

Hypnoidus (*Zorochius*) *nitobei*, p. 2; *Quasimus shirakii*, p. 3; *Q. horishanus*, p. 3; *Cardiotarsus humeralis*, p. 6; *Paracardiophorus cryptohypnoidus*, p. 6; *Cardiophorus sonani*, p. 9; *C. baibaranus*, p. 10; *C. rufithorax*, p. 10; *Cardiohypnus sonani*, p. 11.

— : A new Formosan blister-beetle of the Genus *Meloe* (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 106, pp. 12-13).

Meloe formosensis, p. 12.

— : Elateridae of Formosa (IV) (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 107, pp. 59-69).

Neodiploconus songuinicollis, p. 59; *Sphenicosomus melanotoides*, p. 61; *S. frequens*, p. 61; *Melanotus rufiventris*, p. 64; *Athous temperatus*, p. 66; *Taiwanathous* (n. g.) *arisanus*, p. 68.

MURAYAMA, J.: Révisions des Familles des Ipides et des Platypides de Corée (Jour. Chosen, Nat. Hist. Soc., No. XI, pp. 6-29).

Scolytus seulenensis, p. 9; *Xyleborus seiryorensis*, p. 25; *Crostatarsus koryoensis*, p. 28.

YUASA, H.: Two new Species of Eumolpid-beetles noxious to the Mubberry-tree in the Liukiu Islands (Proc. Jap. Acad., VI, No. 7, pp. 293-295).

Rhyparida sakisimensis, p. 293; *Abirus yashiroi* p. 294.

DIPTERA

ALEXANDER, C. P.: New or little-known Tipulidae from eastern Asia, VI (Philipp. Jour. Science, Vol. 42, No. 1, pp. 59-82).

Tipula uenoii, p. 59; *Limonia* (*Geranomyia*) *radialis*, p. 62; *L. (Geranomyia) alpestris*, p. 63;

L. (Thrypticomyia) apicalis majuscula (n. subsp.), *Helius chikurinensis*, p. 65; *Trycyphona yakushimana*, p. 67; *Ula flavidibasis*, p. 68; *Adelphomyia ariana*, p. 69; *Pseudolimnophila chikurina*, p. 70; *Nippolimnophila* (n. g.) *yakushimensis*, *Nippolimnophila kiusiuensis*, p. 71; *Ulomorpha polytricha*, 72; *Elephantomyia serotina*, p. 74; *Ceratocheilus tinctipennis*, *Lipsothrix yakushimae*, p. 75; *Gonomyia (Ptilostena) curticula*, p. 76; *Erioptera ensifera*, p. 78; *Erioptera (Tisia) tenuisentis*, p. 79; *E. (Empeda) microtrichiatia*, p. 80; *E. (Empeda) angustistigma*, *Molophilus nigrinus*, p. 81; *M. nigratarsis*, p. 82.

_____: New or little-known Tipulidae from eastern Asia, VII (Philipp. Jour. Science, Vol. 42, No. 4, pp. 507-534).

Tipula latiligura, p. 509; *Limonia yakushimensis*, p. 511; *L. tristina*, p. 512, *Helius costosimbriatus*, 514; *Antocha suconfluenta*, p. 516; *Orimarga yakushimana*, p. 517; *Dicranota (Amalopina) gibbera karafutonis* (n. subsp.), p. 520; *Dactylolabis longicauda megastylata* (n. subsp.), *Limnophila (Dicranophragma) dorsolineata*, p. 521; *L. (Prionolabis) lipophleps*, p. 522; *L. (Prionolabis) liponeura*, p. 523; *L. (Tricholimnophila) satsumicola*, *L. (Tricholimnophila) platystyla parallela* (n. subsp.), p. 524; *Atarba (Atarbodes) issikiana*, p. 525; *A. (Atarbodes) minuticornis*, *Elephantomyia dietziana*, p. 526; *Gonomyia (Lipophleps) quadrigila*, p. 530; *Molophilus aricola*, p. 533.

_____: New or little-known Tipulidae from eastern Asia, VIII (Philipp. Jour. Science, Vol. 43, No. 4, pp. 507-534).

Dolichozeza (Nesopeza) tarsalba, p. 509; *D. (Oropeza) saitamensis*, p. 510; *Limonia (Discobola) taivanella*, p. 511; *L. (Limonia) koxinga*, p. 252; *L. (Libnotes) hassenana*, *L. (Rhipidia) triarmata*, p. 514; *L. (Dicranomyia) subpunctulata*, p. 515; *L. (Geranomyia) apicifasciata*, p. 516; *Antocha styx*, p. 517; *Helius (Eurhamphidia) perelegans*, *Dicranoptyla issikina*, p. 518; *Dicranota (Amalopina) delectata*, p. 519; *Adelphomyia issikina*, p. 520; *Limnophila (Prionolabis) serridentata*, p. 521; *Trenteophlia (Mongoma) montina*, p. 523; *Gonomyia (Gonomyia) nansei*, p. 524; *G. (Lipophleps) sauteri*, p. 525; *G. (Lipophleps) neonebulosa*, p. 526; *G. (Lipophleps) sinuosa*, p. 527; *Cryptolabis (Baeoura) trichopoda*, p. 528; *Erioptera (Empeda) litliputina*, p. 530; *Ormosia anthracopoda*, p. 531; *Styringomyia taiwanensis*, p. 532; *S. sinensis*, p. 533.

DUDA, O.: Neue und bekannte orientalische Chloropiden des Deutschen Entomologischen Instituts in Berlin-Dahlem (Stett. Ent. Zeit., 91, pp. 278-304).

Siphunculina costalis, p. 282; *Goniopsita formosana*, p. 283; *G. pallidinervis* BECK var. *angustifrons* (n. var.), *Disciphus flavitarsis*, p. 284; *Meroscinis Sauteri*, p. 286; *Conioscinella griseostriata*, p. 289; *C. nigrohalterata*, *C. opacifrons*, p. 291; *C. Sauteri*, p. 292; *Dicraeus flavipes*, p. 295; *Lioscinella aequisecta*, p. 296; *Chlorops stigmatella* BECK var. *nigricornis*, p. 297.

NINOMIYA, E.: On one new Species of Syrphidae (Oyo-Dobutsu, Vol. 2, No. 4, pp. 251-255).

Syrphus noboritoensis, p. 251.

SHIRAKI, T.: Die Syrphiden des japanischen Kaiserreichs, mit Berücksichtigung Benachbarter Gebiete (Mem. Fac. Sci.-Agr. Taihoku Imp. Univ., Vol. 1, Ent. No. 1, pp. 1-446).

Cerioides formosensis, p. 3; *Sphyximorphoides*, (n. subg.) p. 6; *Paramixogasteroides*, (n. g.) p. 8; *Myxogasteroides*, (n. g.) p. 9; *Parocyptamus*, (n. g.) p. 11; *Parocyptamus sonamii*, p. 12; *Microdon oitanus*, p. 18; *M. formosanus*, p. 22; *Chrysotoxum formosanum*, p. 27; *C. coreanum*, p. 32; *C. japonicum*, p. 35; *Pseudovolucella* (n. g.), p. 39; *P. mimica* p. 40; *Sericomyia japonica*, p. 45; *S. nigriceps*, p. 46; *Criorrhina japonica*, p. 49; *C. formosana*, p. 51; *C. apiformis*, p. 52; *C. setosa*, p. 54; *C. trifasciata* p. 56; *Cynorrhina japonica*, p. 60; *Myolepta japonica*, p. 62; *Zelima makiana*, p. 65; *Z. flavifacies*, p. 68; *Z. brachyparpoides*, p. 70; *Z. spinipes*, p. 72; *Z. sapporensis*, p. 76; *Z. simplex*, p. 81; *Z. bimaculata*, p. 85; *Z. hervei*, p. 86; *Tropidia rostrata*, p. 90; *Eumerus kongosanensis*, p. 93; *E. okinawensis*, p. 98; *Ferdinandea formosana*, p. 103; *Temnostoma taiwanum*, p. 106; *Milesia oshimaensis*, p. 113; *M. vesparia*, p. 118; *Takaomyia formosana*, p. 125; *Eristalomyia semiplendens*, p. 140; *E. formosana*, p. 144; *Pseuderistalis* (n. g.), p. 147; *P. bicolor*, 148; *Kertesziomyia* (n. g.) p. 151; *Korinchia formosana*, p. 154; *K. apicalis*, p. 157; *Eurinomyia japonica*, p. 167; *Mesembrius formosanus*, p. 172; *Paramesembrius* (n. g.), p. 176; *P. horishanus*, p. 178; *Pseudomallota* (n. subg.) p. 187; *Pseudomerodon* (n. subg.), p. 189; *Paramallota* (n. subg.), p. 191; *P. horishana*, p. 194; *P. formosana*, p. 196; *Pseudozetterstedtia* (n. subg.)

p. 199; *P. unicolor*, p. 200; *Imatisma dimorpha*, p. 202; *Azpeytia maculata*, p. 209; *Volutella taiwana*, p. 215; *V. coreana*, p. 225; *V. vespimima*, p. 228; *Graptomyza uchiyamai*, p. 232; *G. formosana*, p. 234; *G. arisana*, p. 236; *G. nitobei*, p. 238; *Paragus hokusankōensis*, p. 249; *Triglyphus formosanus*, p. 251; *Orthoneura formosana*, p. 256; *Chilosia rakurakuensis*, p. 265; *C. yoshinoi*, p. 267; *C. leptorrhyncha*, p. 269; *C. makiana*, p. 272; *C. niitakana*, p. 275; *C. eurodes*, p. 276; *C. sapporensis*, p. 279; *C. okinawae*, p. 285; *C. ochreipila*, p. 283; *C. urakawensis*, p. 287; *C. ochripes*, p. 292; *C. longipennis*, p. 293; *C. iwayakienensis*, p. 295; *C. lata*, p. 297; *C. okunii*, p. 298; *C. splendida*, p. 302; *C. nikkoensis*, p. 304; *C. josankeiana*, p. 306; *C. formosana*, p. 309; *C. diminuta*, p. 311; *C. opaca*, p. 312; *C. latifasciella*, p. 314; *C. matsumurana*, p. 316; *C. hiteipes*, p. 319; *Sonanomyia*, (n. g.) p. 320; *S. formosana*, p. 321; *Platychirus formosanus*, p. 325; *Eriozona nigroscutellata*, p. 333; *Syrphus minimus*, p. 358; *S. ambiguus*, p. 376; *S. issiki*, p. 378; *S. sonami*, p. 389; *Xanthogramma coreanum*, p. 403; *X. arisanicum*, p. 405; *X. fasciatum*, p. 410; *Sphegina varidissima*, p. 421; *S. apicalis*, p. 423; *S. nigerrima*, p. 425; *Ascia longiscutata*, p. 427; *A. breviscutata*, p. 429; *Rhingia formosana*, p. 431.

LEPIDOPTERA

ESAKI, T. and NAKAHARA, W.: Some new Aberrant Forms of Japanese and Formosan Butterflies (Zephyrus, Vol. II, No. 3, pp. 148-150).

Papilio sarpedon connectens FRUHSTORFER ab. *sasanami* (n. ab.) *Junonia lemonias* LINNÉ ab. *nirei* (n. ab.), p. 148; *Symbrenthia hippoclus formosanus* FRUHSTORFER ab. *horishana* (n. ab.), *Cureitis acuta paracuta* NICÉVILE ab. *albida* (n. ab.), 149; *Chrysophanus phlaeas daimio* SEITZ ab. *nyei* (n. ab.), p. 150.

ESAKI, T. and NOMURA, K.: Some new Aberrant Forms of the Formosan Butterflies (Zephyrus, Vol. II, No. 1, pp. 11-14).

Papilio castor formosanus ROTHSCILD ab. *apicalis* (n. ab.), p. 11; *P. doson postianus* FRUHSTORFER ab. *contaminatus* (n. ab.), p. 12.

HALL, A.: New Forms of Nymphalidae (Rhopalocera) in the Collection of the British Museum (Entomologist, Vol. LXIII, No. 806, pp. 156-160).

Euthalia thibetana insulae (n. subsp.), p. 159.

KANO, T.: On *Dodona eugenies* BATES from Formosa (Zephyrus, Vol. II, No. 3, pp. 157-159).

Dodona eugenies matsumurana (n. subsp.) p. 159.

———: Formosan "alpine" Butterflies, 2 (Zephyrus, Vol. II, No. 4, pp. 199-202).

Papilio machaon sylvia ESAKI et KANO (n. subsp.) p. 201.

KATO, M.: Two new Butterflies from Japan and Formosa (Zephyrus, Vol. II, No. 4, pp. 206-208).

Arhopaga ganesa formosana (n. subsp.), p. 206; *Daimio tethys* MÉNÉTRIÉS var. *yamashiroensis* (n. var.), p. 208.

KAWADA, A.: A List of Cochlidionid Moths in Japan, with Descriptions of two new Genera and six new Species (Jour. Imp. Agr. Exp. Stat., Tokyo, Vol. I, No. 3, pp. 231-262).

Natada basifusca, p. 238; *Miresa muramatsui*, p. 246; *Parasa shirakii*, p. 249; *Althonarosa* (n. g.) *horishaeensis*, p. 251; *Narosa edoensis*, p. 252; *Microcampa* (n. g.), p. 256; *Cherometta formosaensis*, p. 257.

MARUMO, N.: Tate-hamaki nitsuite (Öyo-Dobutsu, Vol. 2, No. 1, pp. 37-40).

Susumia (n. g.), p. 39.

MATSUMURA, S.: Two new Species of Sphingid-moths (Trans. Sapporo Nat. Hist. Soc., Vol. XI, pt. 3, pp. 119-120).

Oxyambulyx takamukui, p. 119; *Macroglossum palauensis*, p. 120.

NAKAMURA, Y.: A new Lycanid Butterfly (Lansania, Vol. 2, No. 18, pp. 113-114).

Zizeria maha kashikurai, p. 113.

NAITO, S.: A new Aberrant Form of *Eurema laeta* BOISDUVAL (Zephyrus, II No. 3, p. 151).

- Eurema laeta* BOISDUVAL ab. *eluta* (n. ab.), p. 151.
 ——— : Three new Aberrant Forms of Japanese Butterflies (Zephyrus, Vol. II, No. 4, pp. 203–205).
Papilio xuthus hondoensis FRUHSTORFER ab. *nivescens* (n. ab.), p. 203; *Colias hyale poliographus* MOTSCHULSTY ab. *taeniata* (n. ab.), *Polygonia c-aureum* LINNÉ ab. *atra* (n. ab.), p. 204.
 NOMURA, K. : A new Aberrant Form of *Hypolimnas bolina priscilla* BUTL. (Mushi, Fukuoka, Vol. III, No. 2, p. 73).
Hypolimnas bolina priscilla taihokuna, p. 73.
 SONAN, J. : Notes on some Butterflies from Formosa (Zephyrus, Vol. II, No. 3, pp. 165–176).
Appias melania minato FRUHSTORFER ab. *shima* (n. ab.), *A. melania* FABRICIUS f. *matsumurai* (n. f.), p. 168; *Catopsilia pomona* FABRICIUS f. *formosana* (n. f.), 171; *Neptis philyroides formosanus* (n. subsp.), 174; *Notocrypta arisana*, p. 175.
 ——— : Some Butterflies from Hainan Island (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 106, pp. 31–37).
Gangara thyrsis hainana, p. 37.
 TOSAWA, N. : On two new Examples of Aberrant Forms of *Papilio sarpedon* L. (Trans. Kansai Ent. Soc., No. 1, pp. 1–2).
Papilio sarpedon nipponus suzukii, p. 1; *Papilio sarpedon sarpedonides hankiensis*, p. 2,
 WILEMAN, A. E. and WEST, R. J. : Description of new Species of Japanese, Formosan and Philippine Noctuidae (Entomologist, Vol. LXIII, No. 801, pp. 35–38, No. 802, pp. 62–65, No. 804, pp. 108–110, No. 806, pp. 161–163).
Enargia flavata, p. 35; *Simplicia simplicissima*, p. 37; *Bomolocha nikkoensis*, p. 62; *Hypena innocua*, p. 63; *Ophiuches yoshinalis*, p. 64; *O. hokkaidalis*, p. 108; *Hypenagonia subsuffusata*, p. 161.

TRICHOPTERA

- IWATA, M. : Nippon-san Moshimoku Yôchu, V (Trichopterous Larvae from Japan, V) Dobutsu-gaku Zasshi, Tokyo, Vol. 42, No. 496, pp. 59–66.
Stactobia japonica, p. 59; *Erotesis japonica*, p. 60; *Ganonema uchidai*, p. 61.

RHYNCHOTA

- KATO, M. : Nippon-san Tsunozemika (Membracidae) ni tsuite (Dobutsugaku Zasshi, Vol. 42, No. 8, pp. 281–306).
Tricentrus glochidionae, p. 286; *T. albipennis*, p. 287; *T. biformis*, p. 288; *T. naifunponis*, p. 289; *T. takaoensis*, p. 290; *T. okamotoi*, p. 291; *Machaerotyphus coreanus*, p. 293; *Gargara fasciata*, p. 295; *G. suigensis*, *G. sumipennis*, *G. hyalina*, p. 299; *Arisangargara montana*, p. 300.
 ——— : Notes on the Distribution of Cicadidae in Japanese Empire (Rep. Bul. Biog. Soc. Japan, Vol. 2, No. 1, pp. 1–41).
Tanna auripennis, p. 34; *Euterpnosia suisana*, p. 36; *E. gina*, p. 37.
 SHINJI, G. : On two Species of Japanese Aphididae (Lansania, Vol. 2, No. 12, pp. 26–29).
Macrosiphum colyticola, p. 27.
 ——— : Another new Genus of the Family Aphididae (Lansania, Vol. II No. 15, pp. 78–88).
Macrocaudus (n. g.) *phaseoli*, p. 79.
 ——— : Some new Aphidids from North Eastern Japan (Lansania, Vol. II, No. 20, pp. 151–160).
Chermes laricicola, p. 151; *Trama taraxaci*, p. 152; *Cavariella azamiae*, p. 153; *Anuraphis mumecola*, p. 156; *Amphorophora katsurae*, p. 160.
 TAKAHASHI, R. : Observations on the Coccidae of Formosa, II (Dep. Agr. Gov. Research Inst.,

NEW SPECIES OF INSECTS FROM JAPAN DESCRIBED DURING THE YEAR 1930 217

Formosa, Report No. 48)

Pseudococcus bambusicola, p. 1; *Trionymus formosanus*, p. 4; *Phenacoccus bambusae*, p. 6; *As-terolecanium minutum*, p. 10; *A. pseudomiliaris* GREEN var. *bambusifoliae* (n. var.), p. 11; *Chionaspis vermiformis*, p. 12; *C. bambusifoliae*, p. 14; *Tsukushiaspis suishanus*, p. 16; *T. elongata*, p. 18; *T. neolinearis*, p. 21; *T. formosanus*, 23; *Protodiaspis nigra*, 25; *Leucaspis bambusicola*, p. 26; *Odonaspis simplex* GREEN var. *formosana* (n. var.), p. 29; *Lecaniodrosicha* (n. g.) *lithocarpi*, p. 30; *Saissetia armata*, p. 33; *Inglisia formosana*, p. 34.

_____: Some Aphididae of Loochoo (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. III, pp. 317-327).

Macrosiphum amamianum, p. 318; *Greenidea luchuana*, p. 321; *Truncaphis graminis*, p. 326.

ORTHOPTERA

SHIRAKI, T.: Some new Species of Orthoptera (Trans. Nat. Hist. Formosa, Vol. XX, No. III, pp. 327-352).

Podismopsis shareiensis, p. 327; *Arcyptera coreana*, p. 328; *Melanoplus okinawaensis*, p. 330; *Mirallia formosana*, p. 332; *Exora apicalis*, p. 333; *Hemielimaea formosana*, p. 335; *Trachyzulpha formosana*, p. 337; *Psyra japonica*, p. 339; *Pseudorhynchus japonicus*, p. 341; *Isopsera brevissima*, p. 342; *Decolya kotoshoensis*, p. 344; *Salomona ogatai*, p. 346; *Phyllophorina kotoshoensis*, p. 348; *Rhaphidophora taiwana*, p. 350.

EPHEMERIDA

IMANISHI, K.: Mayflies from Japanese Torrents (Trans. Nat. Hist. Soc. Formosa, Vol. XX, No. 110, pp. 263-267).

Acentrella japonica, p. 263; *Ameletus montanus*, p. 265.

TAKAHASHI, Y.: Kagero Ichi-Shinshu, Futaba-Kagero ni tsuite (Ins. Warld, Gifu., Vol. XXXIV, No. 4, pp. 114-118).

Siphlonurus maculosus, p. 114.

COLLEMBOLA

MATSUMOTO, S.: Mugi no hatsuga wo gaisuru Tobimushimodoki ni kansuru Kenku (Rep. Jour. Agr. Exp. Sta. Okayama, Suppl. XXXV, pp. 1-44).

Onychiurus watanabei, p. 6.