



| | |
|------------------|---|
| Title | New Species and Genera of Insects from Japan described during the Year 1932 |
| Citation | Insecta matsumurana, 7(4), 202-214 |
| Issue Date | 1933-06 |
| Doc URL | http://hdl.handle.net/2115/9260 |
| Type | bulletin (article) |
| File Information | 7(4)_p202-214.pdf |



[Instructions for use](#)

NEW SPECIES AND GENERA OF INSECTS FROM JAPAN
DESCRIBED DURING THE YEAR 1932

HYMENOPTERA

ESAKI, T. et HASHIMOTO, S.: Report on the Leaf-Hoppers injurious to the Rice-Plant and their Natural Enemies, No. 3.

Haplogonatopus atratus, p. 25; Pseudogonatopus flavifemur, p. 28; Agonatopus gracilis, p. 30.

KUWAYAMA, S.: Studies on the Morphology and Ecology of the Rice Leaf-Beetle, *Lema oryzae* KUWAYAMA, with special Reference to the Taxonomic Aspects (Journ. Fac. Agr. Hokkaido Imp. Univ., XXXIII, pp. 1-132).

Anaphes nipponicus, p. 93.

SONAN, J.: Notes on some Braconidae and Ichneumonidae from Formosa, with descriptions of 18 new species (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, Nos. 118, 119, pp. 66-87).

Opius arisanus, p. 67; Opius makii, p. 68; Chelonus formosanus, p. 70; Chelonus tabonus, p. 71; Chelonus arisanus, p. 71; Chelonus signatus, p. 72; Chelonus gohoi, p. 73; Chelonella yami, p. 74; Chelonella nitobei, p. 74; Chelonella mushana, p. 75; Chelonella rokkinia, p. 76; Chelonella nigricoxata, p. 77; Chelonella rufosignata, p. 77; Ascogaster formosensis, p. 78; Ascogaster arisanensis, p. 79; Phanerotoma taiwana, p. 81; Phancrotoma bicolor, p. 81; Chreusa taiwana, p. 85.

SUGIHARA, Y.: On a New Species of Chrysididae from Formosa (The Insect World, Gifu, XXXVI, pp. 371-374).

Chrysis (Pentachrysis) basilacuna, p. 372.

TAKEUCHI, K.: A revision of the Japanese Argidae (Trans. Kansai Entom. Soc., No. 3, pp. 27-42).

Arge captiva SMITH var. watanabei, p. 35; Arge masudai, p. 38.

UCHIDA, T.: Neue und wenig bekannte Japanische Ophioninen-Arten (Trans. Sapporo Nat. Hist. Soc., XII, pp. 73-78).

Omorgus tosenis, O. sugiharai, p. 74; Angitia argyloplevora, Cremastus (Tarytia) uchiyamai, p. 75; Epicremastus matsumuraeanus, p. 76.

—————: Beitrag zur Kenntnis der Japanischen Aulaciden (Trans. Sapporo Nat. Hist. Soc., XII, pp. 189-193).

Pristaulacus intermedius, P. rufipilosus, p. 191.

—————: H. SAUTER's Formosa-Ausbeute (Ichneumonidae) (Jour. Fac. Agr. Hokkaido Imp. Univ. Vol., XXXIII, pt. 2.)

Trogus matsumurai, p. 143; Protichneumon yayeyamensis kankauensis, p. 145; Coelichneumon hormaleoscelus, p. 146; C. sauteri, p. 147; C. taihorinus, p. 148; C. kosempensis, p. 149; C. 10-guttatus, C. ampingensis, p. 150; C. fulvibasalis, p. 151; Acanthojoppa (Akojoppa) sauteri, p. 152; Akajoppa, p. 153 & p. 217; Melanichneumon funkikonis, p. 155; M. futasujii, p. 155; M. tricolor, p. 156; M. semicircularis, M. ramellaris, p. 157; M. parvidentatus, p. 158; Barichneumon saishariensis,

B. tainanensis, p. 159; *B. hozanensis*, p. 160; *Amblyteles sauteri*, p. 161; *A. kankoensis*, p. 162; *Platylabus formosanus*, p. 162; *P. taihorinus*, p. 163; *Neotypus flavipes*, p. 164; *Takanona alboannulata*, p. 166; *Nipporicnus rufus*, p. 166; *Cryptus tuberculatus*, p. 68; *C. suisharionis*, *Apocryptus issikii*, p. 169; *Apocryptus*, p. 170; *Paracryptus formosanus*, *Caenocryptus taihorinus*, p. 171; *Goryphus robustus*, p. 172; *G. horeiensis*, p. 173; *G. kurarensis*, *G. suishanus*, p. 174; *G. taihorinus*, *G. tuberculatus*, p. 175; *G. inornatus*, p. 175; *G. taiwanellus*, p. 176; *G. sauteri*, p. 178; *Mesostenus taiwanus*, p. 179; *M. suisharionis*, p. 180; *Stenaraeoides*, p. 181; *S. ampingensis*, p. 182; *Listrognatus sauteri*, 183; *Hemiteles koshunensis*, p. 184; *Prosopostenus*, p. 184; *H. taiwanellus*, p. 184; *Koshunia*, p. 185; *Oriotohemiteles ovatus*, p. 185; *Oriotohemiteles*, p. 186; *Odontocryptus brilliantus*, p. 186; *Odontocryptus*, p. 187; *Acanthocryptus hokutensis*, *Leptocryptus pilosus*, p. 188; *L. suishariensis*, *Giraudia ranrunensis*, p. 189; *Microcryptus magnispiracularis*, p. 190; *Torbda sauteri*, 191; *T. taihokensis*, p. 192; *T. unicolor*, *Pseudotorbda geniculata*, p. 193; *Pseudotorbda*, p. 195; *Cremastus minimus*, p. 196; *Pristomerus punctatus*, *P. scutellaris*, p. 197; *Zacharops striatus*, p. 198; *Charops taiwanus*, p. 199; *Dichelobosmina tuberculatus*, p. 200; *Dichelobosmina*, p. 201; *Scenocharops longipetiolaris*, p. 201; *Scenocharops*, p. 202; *Campoplegidea melanocerus*, *C. erythra*, p. 203; *C. sauteri*, p. 204; *Diocetes nigriabdominalis*, *Casitaria insularis*, p. 205; *C. longiabdominalis*, *Limnerium sauteri*, p. 206; *Anilastus formosanus*, *A. taihoriensis*, p. 207; *Holocremus geniculatus*, p. 208; *Triclistus parallelus*, p. 209; *T. taiwanensis*, *Exochus radialis*, p. 210; *E. suishanus*, *Orthocentrus carinatus*, p. 211; *Perilissooides cubitalis*, *Perilissooides*, p. 213; *Perilissus formosensis*, *Mesoleptus sauteri*, p. 214; *M. taihorinensis*, *Thymaris taiwanensis*, p. 215; *Monoblastus nigrifacies*, p. 216; *Kobnus filicornis*, p. 217.

WATANABE, C.: A revision of Braconid-species parasitic in the injurious insects of rice plant and sugar-cane in Japan and Formosa (Trans. Sapporo Nat. Hist. Soc., XII, pp. 63-72).

Bracon onukii, p. 65.

YASUMATSU, K.: A list of the species of the genus *Homonotus* of the world, with description of a new species of the genus from Japan (Annot. Zool. Jap., 13, pp. 297-303).

Homonotus iwatai, p. 298.

YASUMATSU, K. et MASUDA, H.: On a new hunting wasp from Japan (Fukuoka Hakubutsugaku Zasshi, 1, pp. 53-65).

Alyson cameroni, p. 53.

COLEOPTERA

CHUJŌ, M.: Studies on the Chrysomelidae in Japanese Empire (I) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 121, pp. 310-317).

Colobaspis rubus, p. 313; *C. shirakii*, p. 315.

—————: l. c., (II) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 122, pp. 337-345).

Pedrillia nitida, p. 340; *P. ruficollis*, p. 342.

DOI, H.: A new species of *Luciola* from Korea (Journ. Chosen Nat. Hist. Soc., 14, p. 63).

Luciola papariensis, p. 63.

KANO, T.: Description of a new *Carabus* from Formosa (Kontyu, Vol. VI, p. 10).

Carabus nankototaizanus, p. 10.

———: New and unrecorded Longicorn-beetles from Japan and its Adjacent Territories (Kontyu, Vol. VI, pp. 259-291).

Megopsis mushensis, p. 259; *Psephactus remiger* HAROLD subsp. *taiwanus*, p. 260; *Trachylophus piyananensis*, p. 26; *Rhagium inquisitor* L. subsp. *morrisonense*, p. 262; *Toxotinus auripilosus*, p. 263; *Leptura niitakana*, p. 264; *Strangalia Rahoarei*, p. 264; *S.* (*Typocerus*) *Kappanzanensis*, p. 265; *S.* (*Typocerus*) *Kinoshitai*, p. 265; *S.* (*Strangalia*) *Tattakana*, p. 266; *S.* (*Strangalia*) *Tattakana* KANO subsp. *piyanan*, p. 267; *S.* (*Strangalia*) *Taranan*, p. 267; *S.* (*Strangalia*) *formoso-montana*, p. 268; *S.* (*Strangalia*) *formoso-montana* KANO subsp. *Masegakii*, p. 269; *S.* (*Strangalia*) *marginale*, p. 269; *Ephies dilaticornis* PASCOE subsp. *apicalis*, p. 270; *Necydalis formosanus*, p. 270; *N. giganteus*, p. 271; *Corennyss sanguineus*, p. 271; *Aphrodisium horishanense*, p. 272; *A. yugaii*, p. 272; *Semanotus chinensis* PIC subsp. *Watanabei*, p. 273; *Callidium nigromaculatum*, p. 274; *Clytus Yokoyamai*, p. 275; *Aglaophis Arakawae*, p. 276; *Anaglyptus Kamiyai*, p. 276; *Cleomenes auricollis*, p. 277; *Sternoplistes takasagoensis*, p. 278; *Psacotha hilaris* PASCOE subsp. *albomaculata*, p. 278; *Melanauster elegans*, p. 279; *Ipocregyes watantakkuni*, p. 280; *I. Arisanus*, p. 281; *Apalimna formosana*, p. 282; *A. liturata* BATES form *grabriventre*, p. 282; *Asaperda takushaensis*, p. 283; *Acanthocinus gundaiensis*, p. 285; *Rondibilis horiensis*, p. 285; *Clytosemia Kuwayamana*, p. 286; *Agapanthia (Agapanthia) melanolopha* FAIRMAIRE subsp. *japonica*, p. 286; *Cagosima sanguinolenta* THOMSON subsp. *oshimaensis*, p. 287; *Serixia testaceicollis*, p. 287; *S. botelensis*, p. 288; *Stenostola chichibuensis*, p. 289; *S. unicolor*, p. 290; *Chreonoma Fortunei* THOMSON subsp. *flavicornis*, p. 291.

KAMIYA, K.: Five new species of Dytiscidae from Japan and the Bonin Islands (*Mushi, Fukuoka*, V, pp. 4-7).

Hydrovatus adachii, *Bidessus yoshimurai*, p. 4; *B. tokunagai*, p. 5; *Copelatus ogasawarensis*, *Agabus sawadai*, p. 6.

KOBAYASHI, K.: A new longicorn-beetle from the Bonin Islands (*Mushi, Fukuoka*, V, pp. 1-3).

Merionoeda tosawai, p. 1.

KONO, H.: Die Mordelliden Japans (Dritter Nachtrag) (*Trans. Sapporo Nat. Hist. Soc.*, XII, pp. 152-160).

Glipa ishigakiana, p. 152; *Mordella kanoi*, p. 153; *Mordellistena matsumurai*, *M. takaosana*, p. 154; *M. luteola*, p. 155; *M. palliata*, *M. chibi*, p. 156; *M. takizawai*, *M. awana*, p. 157; *M. menoko*, *M. galloisi*, p. 158.

MATSUSHITA, M.: Über die auf Insel Sado gesammelten Bockkäfer (*Dobutsugaku Zasshi*, Tokyo, XXXIV, pp. 188-191).

Strangalia xanthoma Bates *sadoensis*, n. var., *S.* (s. str.) *kinhokuana*, p. 191.

MINOWA, S.: New and hitherto-unrecorded Carabidae from Formosa (I) (*Trans. Nat. Hist. Soc. Formosa*, Vol. XXII, No. 121, pp. 281-292).

Leistus (Leistidius) nōkōensis, p. 281; *Leistus (Leistidius) niitakaensis*, p. 282; *Scarites urbanus*,

p. 283; *S. lubricipennis*, p. 284; *S. unicus*, p. 285; *S. mandibularis*, p. 286; *Asaphidion dōmonense*, p. 287; *Tachys dubius*, p. 289; *T. fumigatoides*, p. 290; *T. badius*, p. 291.

—————: Two new Tiger-Beetles from Formosa and Loo-choo (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 121, pp. 293-195)

Cicindela formosana, p. 293; *C. rara*, p. 294.

MIWA, Y., CHUJŌ, M. et MITONO, T.: An Enumeration of Coleoptera from Kōtōsho (Botel-Tobago) with the Description of New Species (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 121, pp. 296-309).

Aprophata pachyrrhynchoides, pp. 306.

MORI, T.: A New Species of Dytiscidae from Korea (Journ. Chosen Nat. Hist. Soc., 14, p. 49).

Copelatus koreanus, p. 49.

MURAYAMA, J.: A new species of Platypodidae from Formosa (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 123, pp. 485-486).

Crossotarsus taiheizanensis, p. 485.

VOSS, E.: Weitere Curculioniden aus Yunnan und Szetschwan der Sammlung Hauser (Col. Curc.) (38 Beitrag zur Kenntnis der Curculioniden) (Wien. Entom. Zeit., 49, pp. 57-76).

Baris tainanica, p. 68.

—————: Weitere Curculioniden aus Yunnan und Szetschwan der Sammlung Hauser (Col. Curc.) (43 Beitrag zur Kenntnis der Curculioniden) (Wien. Entom. Zeit., 49, pp. 285-302.)

Alcides Kōnoi, p. 296.

YOSHIMURA, S.: On a new species of Haliplus from Japan (Mushi, Fukuoka, V, pp. 102-103).

Haliplus tsukushiensis, p. 102.

ZUMPT, F.: Neue Rüsselkäfer aus dem Japanischen Reich und Bemerkungen zu einigen Hylobius- (*Pagiophiloeus*-) Arten (Koleopt. Rundsch., 18, pp. 126-135).

Tenguzo freyi, p. 126; *Hylobius freyi*, p. 128; *H. desbrochersi*, n. n. (*H. roelofsi* Desbr.).

H. orientalis formosanus, p. 131; *Cryptorrhynchus japonicus*, p. 134.

DIPTERA

ALEXANDER, C. P.: New or little-known Tipulidae from Eastern Asia (Diptera), X, (Philip. Journ. Sci., Vol. 49, No. 1, pp. 105-133).

Tipula seticellula, p. 106; *T. (Tipula) okinawensis*, p. 108; *Liogma brevipecten*, p. 110; *Liogma fuscipennis*, p. 111; *Limonia (Dicranomyia) trifilamentosa*, p. 112; *Limonia (Idioglochina) tokunagai*, p. 113; *Proantocha quadrivittata*, p. 115; *Antocha sagana*, p. 116; *Dicranoptycha machidana*, p. 117; *Polymera parvicornis*, p. 119; *Limnophila imanishii*, p. 122, *Eriocera nigrotrochanterata*, p. 123; *E. kamiyai*, p. 125; *Chionea nipponica*, p. 127; *Ormosa togunagai*, p. 129.

—————: The Dipterous family Tanyderidae in Japan (Annot. Zool. Jap., 13, pp. 273-281, 1932).

Protanyderus esakii, p. 276.

CZERNY, L.: Tyliden und Neriiden des zoologischen Museums in Hamburg (Stett. Ent. Zeit., 93, pp. 267-302).

Cothornobata (n. g.) striatifrons, p. 267; Cyclospen formosanus, p. 269; Rainieria leucochira, p. 274; Systellapha niveitarsis, p. 279; Oocephala (n. g.) grata, p. 292.

JANTI, J. V. de: Descriptions de Myodaires supérieurs (Larvaevoridae) nouveaux de Formose (Bull. Soc. Ent. Fr., XXXVII, pp. 268-274).

Platyhira cyanicolor, Echinomyia (Larvaevora) rubrapex p. 268; Linnaemyia (Micropalpus) formosensis, Henilinnaemyia (n. g.) decorata, p. 269; Weberia rubiginans, p. 270.

TOKUNAGA, M.: A remarkable dipterous insect from Japan, Nymphomyia alba, gen. et sp. nov. (Annot. Zool. Jap., 13, pp. 559-569).

Nymphomyia (n. g.) alba, p. 561.

TOKUNAGA, M.: A new biting midge from Japan (Diptera, Ceratopogonidae), with anatomical notes on the larval head-capsule and mouth-parts (Trans. Kansai Entom. Soc., No. 3, pp. 1-12).

Dasyhelea crinume, p. 1.

SHIRAKI, T.: Some Diptera in the Japanese Empire, with Descriptions of New Species (1) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 121, pp. 259-280).

Silvius nigripennis, p. 259; Chrysozona yamadai, p. 262; C. antennata, p. 265; C. koryoensis, p. 268; Tabanus coreanus, p. 270; T. karenkoensis, p. 277.

—————: l. c. (2) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 122, pp. 331-336)

Oncodes nigratarsis, p. 333; O. trifasciatus, p. 335.

—————: l. c. (3) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 123, pp. 487-492).

Anacanthaspis bifasciata var. japonica, p. 489; Pseudoerinna (n. g.) fuscata, p. 491.

LEPIDOPTERA

ISSIKI, S.: On new oriental genera and species of the Hepialoidea (Stylops, 1, pp. 71-80).

Genustes (n. g.) lutata, p. 71; Ogygioses (n. g.) caliginosa, O. eurata, p. 72.

—————: Two new genera and one new species of Japanese and Formosan Tortricidae (Stylops, 1, pp. 134-136).

Gnorismoneura (n. g.) exulis, Ergasia (n. g.), p. 135.

NAKAHARA, W. et ESAKI, T.: A new form of Hebomoia glaucippe Linné from Formosa (Zephyrus, IV, pp. 95-96).

Hebomoia glaucippe formosana Fruhstorfer f. subgrisea, n. f., p. 95.

NAKAHARA, W.: Eine neue Form von Papilio paris Linné aus Formosa (Zephyrus, IV, pp. 243-243).

Papilio paris hermosanus REBEL f. luli, n. f., p. 243.

NOMURA, K.: A new aberrant form of Hestia leuconoë clara Butler and a gynandromorph of Huphina nandina eunama Fruhstorfer (Zephyrus, IV, pp. 10-13).

Hestia leuconoë clara BUTLER ab. kanamarui, n. ab., p. 10.

MARUMO, N.: Nippon san Shachihoko ga no 1 shin-shu oyobi 1 mikiroku-shu (Kontyu, Vol. VI, pp. 292-293).

Shaka amagisana, p. 292.

MEYRICK, E.: Exotic Microlepidoptera, IV, pp. 193-320.

Telphusa necromantis, T. *nephomicta*, p. 194; *Gelechia acanthopis*, G. *mesostrepta*, p. 195; *Thiotricia pontifera*, p. 196; *Compsolechia homoplasta*, p. 197; *Tricyanaula hoplocratus*, p. 198; *Pedioxestis isomorpha*, p. 199; *Chelaria bipinnata*, p. 200; *Dactyletha tegulifera*, p. 201; *Dichomeris leptosaris*, D. *horoglypta*, *Gaesa atomogypsa*, p. 202; *Cymotricha loxospila*, *Lysipatha diaxantha*, *Homaloxestis myeloxesta*, p. 203; *Lecithocera leucoceros*, L. *parthenopis*, p. 204; L. *apostatis*, *Deltoplastis commatopa*, p. 205; *Brachmia macroscopa*, p. 206; *Tinea contributa*, p. 209; *Nothogenes citrocraua*, *Machaeropteris petalacma*, p. 211; *Labdia sarcogypsa*, p. 213; *Passalotis irianthes*, p. 215; *Rhadinastis serpula*, p. 216; *Argyresthia subrimosa*, *Acrolepia argolitha*, p. 227; *Saridoscelis synodias*, p. 228; *Eulia dryonephela*, p. 256; *Periacma lagophthalma*, p. 281; *Odites lividula*, p. 286.

PROUT, L. B.: Seitz Macrolepidoptera, Fauna indoaustratica, 189.

Dindica wilemani, p. 58.

SONAN, J.: A new variety of *saletara panda* Godart from the Ishigaki Island (*Zephyrus*, IV, pp. 238-239).

Saletara panda GODART var. *iwasakii*, p. 238.

SUGITANI, I.: Some new aberrant forms of Formosan and Japanese butterflies (*Zephyrus*, IV, pp. 5-9).

Papilio cloanthus kuge FRUHSTORFER ab. *sumizome*, n. ab., p. 5; *Euploea mulciber barsine* FRUHSTORFER *awayuki*, n. ab., *Cyrestis thyodamus formosana* FRUHSTORFER f. *temporis siccitalis* ab. *nigri-radiata*, n. ab., p. 6; *Symbrenthia brabira scatinia* FRUHSTORFER ab. *mediochracea*, n. ab., *Cocconympha oedippus annulifer* BUTLER ab. *oiwakensis*, n. ab., p. 8.

—————: Some new aberrant forms of Japanese and Formosan butterflies, II (*Zephyrus*, IV, pp. 88-94).

Papilio sarpedon connectens FRUHSTORFER ab. *usuzumi*, n. ab., p. 89; *Ypthima argus* BUTLER ab. *narensis*, n. ab., *Neope goschkevitschii* MÉNÉTRIÈS ab. *tristis*, n. ab., p. 90; *Araschnia burejana* BREMER ab. *azumiana*, n. ab., p. 91.

UCHIDA, S.: On two remarkable aberrant forms of Japanese butterflies (*Zephyrus*, IV, pp. 231-232).

Papilio bianor japonica BUTLER ab. *minoensis*, n. ab., p. 231; *Zephyrus fujisanus* MATSUMURA ab. *yamamotoi*, n. ab., p. 232.

WATARI, M.: On two new forms of *Zephyrus butleri* Fenton (*Zephyrus*, IV, pp. 235-237).

Zephyrus butleri FENTON f. *melanochloë*, n. f., p. 236; *Z. butleri* FENTON f. *connexa*, n. f., p. 237.

RHYNCHOTA

ESAKI, T.: A new *Notonecta* from Formosa (Hemiptera; Notonectidae) (Trans. Nat. Hist. Soc. Formosa, Vol. XXII, No. 123, pp. 493-495).

Notonecta saramao, p. 493.

KANDA, S.: A new species of the genus *Pseudococcus* (Annot. Zool. Jap., 13, pp. 337-390).

Pseudococcus kaiensis, p. 387.

—————: A new species of the genus *Kermes* (Annot. Zool. Jap., 13, pp. 551-557).

Kermes kuwanae, p. 551.

KUWANA, I.: Nippon san Kaigaramushi no 2 shinshu ni tsukite (Kontyu, Vol. VI, pp. 143-148).

Eriococcus tokaedae, p. 143; *Lepidosaphes ezokihadae*, p. 144.

—————: Two new conifer-infesting Scale-Insects from Japan (Philipp. Journ. Sci. Vol. 48, No. 1, pp. 51-54).

Aspidiotus makii, p. 51; *Aspidiotus pseudomeyeri*, p. 52.

TAKAHASHI, R.: Records and Descriptions of the Coccidae from Formosa (Journ. Soc. Trop. Agr. Taihoku Imp. Univ., IV, pp. 41-48).

Trionymus taiwanus, p. 41; *Phenacoccus asteri*, p. 43; *Mixaspis* (n. g.), *Formosaspis* (n. g.), *Lepidosaphes glaucae*, p. 47.

—————: Aleyrodidae of Formosa, I (Departm. Agr. Govern. Research Inst., Formosa, Report 59).

Aleurodicus formosanus, p. 7; *Dialeurodes tuberculatus*, p. 9; *D. daphniphylli*, p. 10; *D. (Gigaleurodes) multipori*, p. 11; *D. (Gigaleurodes) cinnamomi*, p. 13; *D. (Dialeuropora) setigerus*, p. 14; *D. (Dialeuropora) brideliae*, p. 15; *D. (Rachisphora) machili*, p. 16; *D. (Rachisphora) maesae*, p. 18; *D. (Rachisphora) fici*, p. 19; *Aleurotuberculatus* (n. g.) *gordoniae*, p. 21; *A. guyavae*, p. 22; *A. elatostemae*, p. 23; *A. ficicola*, p. 24; *A. malloti*, p. 25; *A. jasmini*, p. 26; *A. uraianus*, p. 27; *Taiwanaleyrodes* (n. g.) *meliosmae*, p. 28; *Tuberaleyrodes* (n. g.) *machili*, p. 29; *Aleuroplatus* (*Aleuroplatus*) *fici*, p. 31; *Aleurolobus citri*, p. 34; *Aleyrodes cinnamomi*, p. 36; *A. liquidambari*, p. 37; *Trialeurodes mori*, p. 38; *T. elatostemae*, p. 39; *Pealius celti*, p. 40; *P. longispinus*, p. 41; *Aleurotrachelus elatostemae*, p. 42; *A. micheliae*, p. 43; *A. turpiniae*, *A. taiwanus*, p. 45; *Siphonaleyrodes* (n. g.) *formosanus*, p. 48.

—————: Additions to the Aphid-Fauna of Formosa (Philipp. Journ. Science, 48, pp. 69-73).

Lachnus nigripes, p. 69; *Ceratovacuna arundinariae*, p. 70; *Thoracaphis setigerus*, p. 72.

TAKEYA, C.: Some Corean lace-bugs (Mushi, Fukuoka, V, pp. 8-13).

Tingis (*Tingis*) *populi*, p. 10; *T. (Birgitta) crispifolii*, p. 12.

EPHEMERIDA

IMANISHI, K.: Mayflies from Japanese torrents, II, Further notes on the genus *Ameletus* (Annot. Zool. Jap., 13, pp. 525-531).

Ameletus kyotensis, p. 525; *A. croceus*, p. 527; *A. subalpinus*, p. 528.

THYSANURA

KUROSAWA, M.: Descriptions of three new Thrips from Japan (Kontyu, Vol. V, pp. 230-242).

Astrothrips aucubae, p. 230; *Pygothrips nogutii*, p. 234; *Ecacanthothrips inarmatus*, p. 238.

INDEX

| | PAGE | | PAGE | PAGE | |
|-----------------------------------|------|----------------------------------|------|-----------------------------------|-----|
| HYMENOPTERA | | | | | |
| affinis (Apanteles) | 87 | heinrichi (Heresiarches) | 31 | taprobanae (Apanteles)... .. | 83 |
| alboguttatus (Automalus) | 27 | heterusiae (Apanteles) | 98 | tatehae (Apanteles) | 82 |
| annulatus (Eupalamus | | igae (Apanteles) | 97 | tamarii (Callajoppa pepsoides) | 27 |
| lamentator) 29 | | iwatensis (Plectochorus) | 163 | tricolor (Charitojoppa)... .. | 25 |
| Apanteles | 75 | japonicus (Phaeogenes) | 155 | turionellae (Pimpla) | 161 |
| apoderi (Bracon) | 180 | kurarensis (Charitojoppa) | 25 | vulnerator (Pristomerus) | 162 |
| arrogans (Callajoppa | | kuwayamai (Apanteles)... .. | 93 | yagoi (Habrocryptus)... .. | 156 |
| pepsoides) 27 | | lacticolor (Apanteles)... .. | 96 | yezoensis (Callajoppa | |
| asotae (Apanteles)... .. | 98 | lamborini (Apanteles) | 91 | exaltatorius) 27 | |
| bicolor (Apanteles) | 88 | laspeyresiae (Ephialtes) | 160 | zygaenarum (Apanteles) | 80 |
| bicolorinus (Hemiteles | | liparidis (Apanteles) | 89 | COLEOPTERA | |
| (Rhadiurgus)) 158 | | mikado (Callajoppa lutorius) . | 27 | Aclees | 182 |
| bifascialis (Ichneumon) | 29 | minor (Apanteles)... .. | 99 | acuminatus (Hydrovatus) | 166 |
| brunicornis (Herpestomus) | 31 | miokensis (Hoplismenus) | 28 | adachii (Hydrovatus) | 166 |
| byctisci (Calyptus) | 181 | miyoshii (Apanteles) | 85 | affinis (Dictyopterus | |
| Charitojoppa | 25 | molestae (Dioctes) | 162 | (Pyropterus)) 56 | |
| chilonis (Apanteles flavipes) | 84 | molestae (Hemiteles | | anchoralis (Deronectes | |
| congestus (Apanteles) | 79 | (Isadelphus)) 158 | | (Potamodytes)) 176 | |
| coreanus (Cremastus (Tarytia) | | nawai (Apanteles) | 100 | antennatus (Bidessus | |
| flavo-orbitalis)) 163 | | nigrus (Tricholabus) | 31 | (Bidessus)) 170 | |
| coreanus (Tricholabus nigrus) | 31 | Nipponaëtes | 160 | atorufus (Mesolytus) | 55 |
| cyanea (Charitojoppa) | 25 | odaiensis (Stenichneumon | | aurora (Dictyopterus | |
| cymolomiae (Glypta) | 162 | (Stenichneumon)) 29 | | (Dictyopterus)) 56 | |
| epinotiae (Itopectis) | 161 | okamotoi (Apanteles) | 86 | bellus (Anaglyptus) | 105 |
| erythrothoracis (Pristomerus | | ordinarius (Apanteles) | 79 | Bidessus | 169 |
| vulnerator) 161 | | pallidipes (Automalus) | 28 | Bidessus (Bidessus) | 169 |
| falcatus (Apanteles) | 92 | pepsoides (Callajoppa) | 26 | bioculatus (Exocentrus) | 108 |
| femoratus (Apanteles) | 88 | planus (Apanteles) | 84 | bipustulatus (Dysceroides) | 187 |
| flavibasalis (Gathetus) | 25 | Plectochorus | 163 | boninense (Trinophylum) | 72 |
| flavipes (Apanteles) | 83 | politus (Apanteles) | 99 | caenosa (Pterophia) | 107 |
| flavo-orbitalis (Cremastus | | ruficrus (Apanteles) | 78 | Canthydrus | 19 |
| (Tarytia)) 162 | | sasakii (Apanteles) | 91 | Ceresium | 66 |
| fulvipes (Apanteles) | 91 | sonani (Amblyteles) | 31 | chinensis (Laccophilus) | 21 |
| gastropachae (Apanteles) | 87 | sonani (Apanteles) | 96 | Chuzenjianus | 61 |
| glomeratus (Apanteles)... .. | 80 | sugae (Apanteles) | 94 | clavigera (Stenodryas) | 72 |
| grapholithae (Spilocryptus)... .. | 157 | suzumei (Apanteles) | 81 | Clypeodytes | 169 |
| hamakii (Apanteles) | 95 | | | Coelambus | 171 |
| haeusseri (Hemiteles | | | | conjunctum (Stenygrinum | |
| (Nipponaëtes)) 159 | | | | quadrinotatum) 66 | |
| | | | | coracinus (Platerus) | 60 |

| | PAGE | | PAGE | | PAGE |
|--|------|--|------|--|------|
| coreana (Strangalia) | 104 | japonicus (Noterus) | 18 | sachalinensis (Leptura)... .. | 104 |
| daisenensis (Glenea) | 109 | kanoi (Hylobius) | 184 | saghalienis (Hydroporus) | 173 |
| Deronectes | 175 | kikuchii (Kobuzo) | 188 | sagulatus (Pristolycus) | 60 |
| difficilis (Laccophilus) | 22 | kobensis (Laccophilus)... .. | 23 | saipanicum (Paraceresium) | 71 |
| doii (Strangalia) | 105 | Kobuzo | 188 | sanguineus (Lygisterus) | 60 |
| dominator (Macrolycus) | 54 | Laccophilus | 20 | sanmarki (Deronectes (Oreodytes)) | 178 |
| Dysceroides | 187 | leucosticticum (Ceresium) | 69 | sapporensis (Anaglyptus) | 106 |
| Euthycus... .. | 189 | lewisius (Laccophilus) | 23 | sapporensis (Dictyopterus) | 57 |
| excisus (Euthycus)... .. | 189 | lineatus (Aplatopterus)... .. | 58 | septentrionalis (Aplatopterus) | 58 |
| flabellatus (Macrolycus) | 55 | longicorne (Ceresium) | 69 | sharpi (Laccophilus) | 23 |
| flavipes (Ceresium) | 67 | loochooana (Abryna coenosa) | 108 | signaticolle (Ceresium)... .. | 69 |
| flavus (Canthydrus) | 19 | lyratus (Hyphydrus) | 167 | simile (Ceresium)... .. | 70 |
| formosanus (Hylobius orientalis) | 186 | megecephalus (Bidessus (Bidessus)) | 170 | simplicipes (Deronectes (Potamodytes)) | 178 |
| formosanus (Laccophilus) | 22 | minowai (Hylobius) | 185 | sinicum (Ceresium) | 70 |
| frontalis (Bidessus (Clypeodytes)) | 169 | modestus (Lycostomus) | 55 | spinicoxis (Dictyopterus (Benibotarus)) | 56 |
| fujiyamai (Exocentrus)... .. | 108 | nasuta (Platycis) | 58 | Stenodryas | 72 |
| fuscum (Ceresium) | 67 | uatrix (Graptodytes) | 175 | Stenygrinum | 65 |
| geometricus (Cladophorus) | 59 | niitakensis (Hylobius) | 184 | subuniforme (Ceresium) | 70 |
| gorhami (Dictyopterus (Dictyopterus)) | 57 | nitidulus (Contydrus) | 20 | taiwana (Platycis nasuta) | 59 |
| Graptodytes | 175 | nipponicus (Deronectes (Potamodytes)) | 177 | taiwanensis (Aclees) | 182 |
| guttula (Canthydrus) | 20 | Noterus | 18 | tenuis (Chuzenjianus) | 61 |
| hachijoense (Ceresium) | 68 | oculatus (Dictyopterus)... .. | 57 | tokunagai (Bidessus (Bidessus)) | 170 |
| hachijonis (Glenea) | 110 | ogumae (Xylobanus) | 60 | Trinophylum | 72 |
| hirayamai (Aclees) | 183 | Oreodytes | 178 | umbrosus (Hydroporus) | 174 |
| horophaeum (Ceresium) | 68 | orientalis (Hyphydrus)... .. | 168 | vittatus (Coelambus) | 172 |
| hostilis (Deronectes (Potamodytes)) | 176 | orientalis (Pidonia) | 103 | watanabei (Hydroporus) | 174 |
| Hydroporus | 173 | otome (Platycis) | 59 | yoshimurai (Bidessus (Bidessus)) | 170 |
| Hydrovatus | 165 | palauense (Ceresium) | 70 | zeyldnicum (Ceresium)... .. | 70 |
| Hylobius | 184 | Paraceresium | 71 | | |
| Hyphoporus | 172 | perforatus (Hylobius) | 186 | | |
| Hyphydrus | 167 | pictus (Conderis) | 61 | | |
| impressopunctatus (Coelambus) | 171 | politus (Conthydrus) | 20 | | |
| japonicum (Ceresium) | 68 | Potamodytes | 176 | | |
| japonicus (Bidessus (Bidessus)) | 170 | pustulatus (Hylobius) | 186 | | |
| japonicus (Hydrovatus)... .. | 166 | quadrinotatum (Stenygrinum) | 65 | | |
| | | rotundatus (Hyphoporus) | 172 | | |
| | | ruficollis (Mimistena setigera) | 107 | | |

LEPIDOPTERA

| | |
|---------------------------------|-----|
| akanensis (Palimpsestis) | 196 |
| alba (Arctornis) | 111 |
| alba (Dura)... .. | 122 |
| albibasalis (Dasychira)... .. | 114 |
| albipes (Pantana) | 143 |

| PAGE | | PAGE | | PAGE | |
|--------------------------------|-----|---------------------------------|-----|--------------------------------|-----|
| albisparsa (Pachyparoides) ... | 50 | conspersana (Euproctis) ... | 124 | flexuosana (Euproctis) ... | 126 |
| albolineatus (Dendrolimus)... | 36 | coreana (Aroa) ... | 111 | formosae (Pseudarguda) ... | 51 |
| albomaculata (Cosmotriche) . | 34 | coreana (Dasychira) ... | 116 | formosana (Cosmotriche) ... | 35 |
| alikangiae (Euproctis) ... | 123 | coreana (Euproctis) ... | 125 | Formotogaria... .. | 192 |
| ampla (Kunugia) ... | 46 | coreana (Gastropacha) ... | 43 | fumida (Lymantria) ... | 136 |
| Amurilla... .. | 33 | Cosmotriche ... | 34 | furva (Maimaia) ... | 140 |
| angulata (Euproctis) ... | 123 | croceola (Euproctis) ... | 125 | Gastropacha | 43 |
| angulosa (Dasychira) ... | 114 | cruda (Dasychira)... | 116 | gigantea (Laelia) ... | 133 |
| antiqua (Orgyia) ... | 142 | curvata (Euproctis) ... | 125 | gonostigma (Orgyia) ... | 142 |
| arakawae (Dasychira) ... | 115 | cygna (Stilpnotia)... | 150 | grotei (Dasychira) ... | 117 |
| Arctornis | 111 | Dactylorhynchides ... | 114 | hirayamae (Dasychira)... | 117 |
| argentata (Dasychira) ... | 120 | daisensis (Eriogaster) ... | 42 | hirayamai (Palimpsestis) ... | 196 |
| argtenata (Euproctis) ... | 123 | daisetsuzana (Konokareha) ... | 132 | hopponis (Porthesia) ... | 146 |
| Arguda | 33 | daisetsuzana (Palimpsestis) ... | 196 | Horipsestis | 193 |
| arizana (Dasychira) ... | 115 | Dasychira | 114 | horishana (Gastropacha) ... | 44 |
| arizana (Kunugia) ... | 47 | dasychiroides (Pseudodura)... | 150 | horishana (Syrastenoides) ... | 52 |
| Aroa | 111 | decolorata (Pida) ... | 145 | horishanella (Dasychira) ... | 117 |
| aurifera (Dasychira) ... | 115 | Dendrolimus | 36 | horishanella (Lymantria) ... | 136 |
| aurata (Euproctis)... | 123 | devestita (Laelia) ... | 132 | Horithyatira | 193 |
| auripes (Ivela) | 131 | diplagum (Medama) ... | 145 | hyrtaca (Metanastria) ... | 50 |
| baibarana (Baipsestis) ... | 190 | discincta (Wilemaniella) ... | 54 | idiota (Bhima) | 34 |
| baibarana (Dasychira) ... | 115 | dispar (Lymantria) ... | 135 | ilicifolia (Epicnaptera)... | 41 |
| baibarana (Porthesia) ... | 145 | disparilis (Numenes) ... | 141 | infuscata (Pantana) ... | 144 |
| Baipsestis | 190 | dissimilis (Euproctis) ... | 125 | inornata (Euproctis) ... | 126 |
| bantaizana (Lymantria) ... | 134 | dissoluta (Lymantria) ... | 136 | insulata (Euproctis) ... | 126 |
| bhana (Dasychira) ... | 115 | divisa (Cosmotriche) ... | 34 | invaria (Dasychira) ... | 117 |
| Bhima | 34 | doii (Stilpnotia) ... | 150 | Ivela | 131 |
| bimaculata (Euproctis)... | 123 | Dura | 122 | iwasakii (Kunugia) ... | 47 |
| bipunctapex (Euproctis) | 124 | endoplagia (Porthesia)... | 145 | jankowskii (Aroa)... | 112 |
| brevivenis (Eriogaster)... | 42 | Epeuproctis | 122 | japonibia (Laelia)... | 133 |
| brunnea (Kunugia) ... | 47 | Epicnaptera | 41 | jezoensis (Dendrolimus) ... | 37 |
| c-album (Saronaga) ... | 198 | Episaronaga | 191 | kagina (Dasychira) ... | 118 |
| Caragora... .. | 112 | Eriogaster | 42 | kan (Euproctis) | 126 |
| catacooides (Eriogaster)... | 42 | Estigena... .. | 43 | kanshireia (Euproctis)... | 127 |
| centrofascia (Euproctis) | 124 | Euproctis | 123 | kanshireiensis (Dasychira) ... | 118 |
| centropallida (Euproctis) | 124 | eurydice (Cifuna) ... | 113 | karapina (Euproctis) ... | 127 |
| chibiana (Euproctis) ... | 124 | excisa (Takanea) ... | 53 | Karenkonio | 45 |
| Chibidokuga | 113 | exclamationis (Laelia)... | 133 | kibarae (Dasychira) ... | 118 |
| chichibensis (Stilpnotia) | 150 | Exothyatira | 192 | kibunensis (Eriogaster) ... | 42 |
| Cifuna | 113 | fascelina (Dasychira) ... | 116 | kikuchii (Dasychira) ... | 118 |
| citrinella (Dasychira) ... | 116 | flava (Dasychira) ... | 117 | kikuchii (Kunugia) ... | 48 |
| clara (Caragola) | 112 | flava (Euproctis) ... | 125 | Konokareha | 132 |
| coenosia (Laelia) | 122 | flavinata (Euproctis) ... | 126 | | |
| concolor (Lymantria) ... | 135 | flavovittata (Arguda) ... | 33 | | |
| conjuncta (Dasychira) ... | 116 | | | | |

| | PAGE | | PAGE | | PAGE |
|----------------------------------|------|------------------------------------|------|----------------------------------|------|
| Kononia | 46 | nobunaga (Lymantria) | 138 | saitonis (Dasychira) | 121 |
| Kunugia | 46 | nuda (Perina) | 144 | saitonis (Neotogaria) | 195 |
| kurilensis (Palimpsestis) | 197 | Numenes | 141 | sakaguchii (Lymantria) | 139 |
| labecula (Euproctis) | 127 | ochreipuncta (Cosmotriche) . | 35 | sangaica (Laelia)... .. | 133 |
| Laelia | 132 | ochropicta (Aroa)... .. | 112 | scintellans (Porthesia) | 147 |
| latifascia (Euproctis) | 127 | Odonestis | 50 | Selenephera... .. | 51 |
| Lebeda | 48 | okinawana (Porthesia)... .. | 146 | sericea (Caragola)... .. | 113 |
| limacodiformis (Dactylor- | | olivacea (Dasychira) | 120 | sericea (Porthesia) | 148 |
| hynchides) | 114 | Orgyia | 142 | serva (Lymantria) | 139 |
| I-nigtum (Arctornis) | 111 | Pachyparoides | 50 | shirakii (Formotogaria) | 192 |
| locuples (Cifuna) | 114 | Pantana | 143 | shirakii (Euproctis) | 129 |
| lucescens (Lymantria) | 136 | Paralebeda | 51 | shironis (Euproctis) | 130 |
| lunigera (Selenephera)... .. | 51 | pardalis (Estigena) | 43 | similis (Porthesia) | 148 |
| lunulata (Dasychira) | 118 | patrana (Numenes) | 141 | simplex (Euproctis) | 130 |
| Lymantria | 134 | Perina | 144 | simplex (Pantana)... .. | 144 |
| Maimaia | 140 | Pida | 145 | sinensis (Laelia) | 133 |
| Malacosoma | 49 | pinivora (Kononia) | 46 | sparsa (Euproctis)... .. | 130 |
| mathura (Lymantria) | 137 | piperita (Porthesia) | 147 | spectabilis (Dendrolimus) | 39 |
| Medama | 140 | plagifera (Poralebeda) | 51 | sinica (Lymantria) | 139 |
| mendosa (Dasychira) | 119 | populifolia (Gastropacha) | 44 | staudingeri (Euproctis)... .. | 130 |
| Metanastria | 50 | Porthesia | 145 | Stilpnolia | 150 |
| Meteuproctis | 141 | postalba (Pida) | 145 | striata (Euproctis).. .. | 130 |
| minosa (Porthesia) | 146 | postalbata (Euproctis) | 128 | striata (Laelia) | 134 |
| minomonis (Lymantria) | 137 | postfusca (Dasychira) | 120 | strigipennis (Pida) | 145 |
| minor (Syrastrina) | 52 | postfusca (Euproctis) | 129 | subpurpurea (Amurilla)... .. | 33 |
| Miothyatira | 194 | postica (Orgyia) | 143 | Sugiploca | 199 |
| misaona (Neoploca) | 194 | potatoria (Cosmotriche) | 35 | sigitanii (Dendrolimus) | 198 |
| miyakei (Takanea) | 52 | pruni (Odonestis)... .. | 50 | superans (Sugiploca) | 41 |
| monacha (Lymantria) | 137 | Pseudarguda | 51 | suzukii (Baipsestis) | 191 |
| mushana (Palimpsestis) | 197 | pseudo-abietis (Dasychira) | 120 | suzukii (Laelia) | 134 |
| nachiensis (Dasychira)... .. | 119 | pseudoconsersa (Euproctis) . | 129 | Suzupsestis | 199 |
| nagoyana (Dasychira) | 119 | Pseudodura | 150 | Syrastenoides... .. | 52 |
| nantonis (Orgyia)... .. | 145 | pterofera (Euproctis) | 129 | Syrastrena | 52 |
| Neopsestis | 195 | pubibanda (Dasychira)... .. | 120 | taiwana (Dasychira) | 121 |
| Neotogaria | 195 | pulverea (Porthesia) | 147 | taiwana (Gastropacha)... .. | 44 |
| Neodaruma | 194 | punctatus (Dendrolimus) | 38 | taiwana (Karenkonio) | 45 |
| neustrium (Malacosoma) | 49 | purpureofasciata (Porthesia)... .. | 147 | taiwana (Porthesia) | 149 |
| nigra (Chibidokuga) | 113 | pyriformis (Cosmotriche) | 35 | taiwana (Saronaga) | 198 |
| nigricauda (Euproctis)... .. | 123 | quercifolia (Gastropacha) | 44 | takamukuana (Dasychira) | 121 |
| nigricosta (Lymantria)... .. | 138 | riukiwana (Porthesia) | 147 | takamukuana (Amurilla) | 33 |
| nigropuncta (Cosmotriche)... .. | 35 | roseola (Lymantria) | 138 | takamukui (Eriogaster) | 43 |
| nigropuncta (Euproctis) | 120 | sachalinensis (Dasychira) | 121 | takamukui (Lymantria) | 139 |
| niphonis (Euproctis) | 128 | | | takamukui (Numenes) | 142 |
| nobilis (Lebeda) | 48 | | | Takanea... .. | 52 |
| | | | | takaazana (Togaria) | 200 |
| | | | | Takapsestis | 200 |

| | PAGE | | PAGE | | PAGE |
|-------------------------------|------|--------------------------------|------|----------------------------------|------|
| tamahonis (Cosmotriche) ... | 36 | undans (Kunugia) ... | 48 | yamadai (Kunugia) ... | 48 |
| tamahonis (Epeuproctis) ... | 123 | unifascia (Euproctis) ... | 131 | yatsugadakensis (Eriogaster) ... | 43 |
| tamanukii (Neodaruma) ... | 194 | urocoma (Porthesia) ... | 149 | | |
| tamanukii (Trichiura) ... | 53 | usukia (Euproctis) ... | 131 | | |
| teikichiana (Horipsestis) ... | 193 | | | THYSANOPTERA | |
| tenebrosa (Dasychira) ... | 122 | varians (Euproctis) ... | 131 | femoralis (Machatothrips) ... | 6 |
| thyllina (Orgyia) ... | 143 | virginea (Dasychira) ... | 122 | ipomoeae (Machatothrips) ... | 12 |
| tomponis (Baipsestis) ... | 191 | vishnou (Trabala) ... | 53 | issikii (Machatothrips) ... | 7 |
| tomponis (Meteproctis) ... | 141 | visum (Pantana) ... | 144 | | |
| tomponis (Porthesia) ... | 141 | watanabei (Palimpsestis) ... | 197 | kawamurai (Idolothrips) ... | 2 |
| Trabala ... | 53 | Wilemaniella ... | 54 | ohtai (Machatothrips) ... | 9 |
| Trichiura ... | 53 | wilemaniella (Takapsestis) ... | 199 | yashiroi (Idolothrips) ... | 4 |
| tsingtauca (Gastropacha) ... | 45 | xanthocampa (Porthesia) ... | 149 | | |
| | | xylina (Lymantria) ... | 140 | | |
| uchidai (Porthesia) ... | 149 | | | | |
| umbrifera (Lymantria) ... | 139 | | | | |

BOOK-REVIEW

"Medical Entomology" edited by **Dr. Robert Matheson**, Professor of New York State College of Agriculture, Cornell University (1932). pp. 485. Text-figures 211.

This book is divided into 20 Chapters and all the known insects which may play a part in the transmission, causation and spread of human diseases are described and illustrated. The introductory text is offered to the physician, entomologist, the public worker, the student, and layman in order that may inform and arouse a keener interest in the problems involved with insect-borne diseases. Although this book is published in United States of America, but it mentions all the world known medical insects, even the Oriental species which were studied by our entomologists and physicians. The satisfactory information of controlling the insects involved in disease transmission or causation is at hand, but the discussion of treatment is omitted. At the end of the first Chapter is enumerated a rather comprehensive list of text books, journals, and other publications which enable the student to gain access to this rapidly growing field of research. Furthermore, each Chapter is provided with a selected bibliography and exact references bearing on the statements made in the text.

This book is highly recommended as an important up-to-date reference book to our entomologists and physicians.

(Prof. Dr. S. MATSUMURA)