



Title	Identification of cheese rancidity-related lipases in <i>Aspergillus oryzae</i> AHU 7139
Author(s)	Chintagavongse, Napaporn; Kumura, Haruto; Hayakawa, Toru; Wakamatsu, Jun-ichi; Tamano, Koichi
Citation	Journal of bioscience and bioengineering, 137(5), 381-387 <a href="https://doi.org/10.1016/j.jbiosc.2024.01.016">https://doi.org/10.1016/j.jbiosc.2024.01.016</a>
Issue Date	2024-03-01
Doc URL	<a href="http://hdl.handle.net/2115/92618">http://hdl.handle.net/2115/92618</a>
Rights	© <2024>. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Rights(URL)	<a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Type	article (author version)
File Information	JBB R2-marked up.pdf



[Instructions for use](#)

1 **Identification of cheese rancidity-related lipases in *Aspergillus oryzae***  
2 **AHU 7139**

3 **A short title: Identification of cheese rancidity-related lipases**

4 Napaporn Chintagavongse<sup>1</sup>, Haruto Kumura<sup>1,\*</sup>, Toru Hayakawa<sup>1</sup>, Jun-  
5 ichi Wakamatsu<sup>1</sup>, Koichi Tamano<sup>2</sup>

6 <sup>1</sup>*Laboratory of Applied Food Science, Graduate School and Research Faculty of*  
7 *Agriculture, Hokkaido University, 060-8589, N9, W9, Sapporo, Japan,*<sup>2</sup>*Bioproduction*  
8 *Research Institute, National Institute of Advanced Industrial Science and Technology*  
9 *(AIST), 2-17-2-1 Tsukisamu-Higashi, Toyohira-ku, Sapporo, Hokkaido 062-8517,*  
10 *Japan*

11

12

13

14

15

16

17

18 \* Corresponding author: Haruto Kumura

19 Mailing address: Laboratory of Applied Food Science, Graduate School and Research  
20 Faculty of Agriculture, Hokkaido University, 060-8589, N9, W9, Sapporo, Japan

21 Tel: +81-11-706-3642, E-mail: kumura@agr.hokudai.ac.jp

22

23 **ABSTRACT**

24           The adjunct product with enzymatic activity from *Aspergillus oryzae* is  
25 beneficial for flavor enrichment in the ripened cheese. However, an excessive lipolytic  
26 reaction leads to the release of volatile free fatty acids. Accordingly, a strong off-flavor  
27 (i.e., rancidity) has been detected when *A. oryzae* AHU 7139 is used. To identify the  
28 rancidity-related lipase from this strain, we evaluated the substrate specificity and  
29 lipase distribution using five mutants cultured on a whey-based solid medium under  
30 different initial pH conditions. The results showed a higher diacylglycerol lipase  
31 activity than triacylglycerol lipase activity. Moreover, an initial pH of 6.5 for the  
32 culture resulted in higher lipolytic activity than a pH of 4.0, and most of the activity  
33 was found in the extracellular fraction. Based on the gene expression analysis by RT-  
34 PCR and location and substrate specificity, five genes (No. 1, No. 19, *mdlB*, *tgla*, and  
35 *cutL*) were selected among 25 annotated lipase genes to identify the respective  
36 knockout strains. Because  $\Delta tgla$  and  $\Delta mdlB$  showed an outstanding involvement in  
37 the release of free fatty acids, these strains were applied to *in vitro* cheese curd  
38 experiments. In conclusion, we posit that triacylglycerol lipase (Tgla) plays a key role  
39 as the trigger of rancidity and the resulting diglycerides have to be exposed to  
40 diacylglycerol lipase (MdlB) to stimulate rancidity in cheese made with *A.*  
41 *oryzae* AHU 7139. This finding could help screen suitable *A.oryzae* strains as cheese  
42 adjuncts to prevent the generation of the rancid-off flavor.

43 **KEYWORDS:** *Aspergillus oryzae*, lipase, substrate, free fatty acids, rancidity, *in vitro*  
44 curds

45

## 46 INTRODUCTION

47 The filamentous fungus *Aspergillus oryzae* is used in traditional Japanese  
48 cuisine (washoku). *A. oryzae* is recognized as safe because it is missing the aflatoxin  
49 synthesis gene (1) and is assumed to encode as much as 135 protease genes based on  
50 its genomic analysis (2).

51 In contrast to washoku, *A. oryzae* is rarely used in Western foods. Kumura et  
52 al. (3) focused on the high proteolytic potential of *A. oryzae* and attempted to apply its  
53 culture products (CP) as adjuncts for cheese flavor enrichment. As solid-state  
54 fermentation is known to be superior to liquid fermentation for abundant enzyme  
55 production (4), Kumura et al. (3) prepared solid CP of *A. oryzae* AHU 7139 as adjuncts  
56 for making Gouda-type cheese and confirmed the enhancement of proteolysis during  
57 cheese ripening. Moreover, a remarkable content of free fatty acid (FFA) due to  
58 lipolysis was detected. They reported that the application of CPs whose fermentation  
59 had started with a pH of 6.5 resulted in a pronounced increase in FFA compared to the  
60 fermentation that started with a pH of 4.0, implying the initial pH of the culture may  
61 affect the profile of the lipases produced.

62 Although proper lipolysis provides the characteristic flavor and aroma to  
63 cheese, excess FFA accumulation during cheese ripening causes an undesirable off-  
64 flavor known as rancidity (5) by the release of volatile short-chain fatty acids over an  
65 organoleptic threshold. Furthermore, once milk triacylglycerols are degraded by  
66 lipases (the first step of lipolysis), the resulting diglycerides are exposed and become  
67 the subsequent substrates for diacylglycerol lipase (the second step of lipolysis). Thus,  
68 the adjunct materials of *A. oryzae* should be prepared with high proteolytic but limited  
69 lipase activity.

70           According to the genomic analysis of a wild-type strain, about 30 genes are  
71 annotated to encode lipase genes (FungiDB database,  
72 <https://fungidb.org/fungidb/app/>). Expression of some lipases is likely to be dependent  
73 on the culture conditions, and the identification of the rancid-inducible lipase  
74 molecules could provide valuable information to screen suitable *A. oryzae* strains for  
75 preventing flavor defects in dairy application.

76           Hence, we compared lipase gene expression under different pH conditions and  
77 selected some candidate lipase genes. Then, the influence of mutations on the lipase  
78 activity was evaluated. Finally, we identified the most influential lipase molecules on  
79 the rancidity, using *in vitro* cheese curd experiments.

## 80 MATERIALS AND METHODS

### 81 Strain and culture condition

82 The strain used in this study was *A. oryzae* AHU 7139 from the culture collection of  
83 Hokkaido University and was grown on potato dextrose agar (PDA) (Merck KGaA,  
84 Darmstadt, Germany) at 30°C for 7 days. Spore suspensions were prepared by adding  
85 9.0 g L<sup>-1</sup> sodium chloride (NaCl) solution into the grown culture on PDA and diluted  
86 to the concentration of 2.5×10<sup>5</sup> spores mL<sup>-1</sup>, which was counted by using  
87 haemocytometer (NanoEntek, Seoul, Korea). The spore suspension (150 µL) was  
88 inoculated on whey solid medium whose pH was adjusted to 4.0 or 6.5, and cultivated  
89 at 20°C for 7 days (6).

### 90 RNA isolation and RT-PCR

91 The solid culture product (CP) was recovered and immediately freeze-dried and  
92 ground with a mortar and pestle to obtain freeze-dried powder (FDP). On average,  
93 0.27 g FDP was obtained from 1 g CP. The FDP (0.5 mg) was transferred into 5 mL  
94 of neutralized phenol-saturated water containing 2 mol L<sup>-1</sup> NaCl, mixed well, and  
95 incubated in a waterbath 55°C for 10 min. The sample was centrifuged at 10,000 ×g,  
96 4°C for 15 min and the upper aqueous phase was transferred to a new tube followed  
97 by the addition of chloroform:isoamylalcohol (49:1; CIA). After centrifugation at  
98 10,000 ×g, 4°C for 10 min, the upper phase was transferred to a new tube followed by  
99 addition of water-saturated acidic phenol/CIA (1:1) and centrifuged again (this step  
100 was repeated two times). Then, the upper phase was poured into a new tube and CIA  
101 was added into the tube, followed by centrifugation for 5 min. After transferring the  
102 upper phase to a new tube, 0.5 volume of 7.5 mol L<sup>-1</sup> LiCl (Kanto Chemical, Tokyo,

103 Japan) was added to the tube. The mixture was incubated at 4°C overnight, followed  
104 by centrifugation at 12,000 ×g, 4°C for 30 min. The supernatant was removed and the  
105 pellet was rinsed with 2.5 mol L<sup>-1</sup> LiCl and centrifuged twice at 12,000 ×g , 4°C for 5  
106 min. The air-dried pellet was finally dissolved in formamide to obtain total RNA and  
107 stored at -80°C. The RNA concentration was estimated by measuring the absorbance  
108 at 260 nm with a NanoDrop spectrophotometer (Thermo scientific, US). The recovery  
109 of total RNA was evaluated by electrophoresis using a commercial kit (Dynamarker  
110 RNA High for Easy Electrophoresis, Biodynamics Laboratory, Tokyo, Japan)  
111 according to the manufacturer's instructions (Fig. S1).

112 The mRNA was isolated from the total RNA using a commercial kit (Pharmacia  
113 Biotech, Stockholm, Sweden) according to the manufacturer's instructions. The  
114 mixture containing 50 ng of mRNA sample, random primer (Promega, Wisconsin, US),  
115 and dNTP was heated at 65°C for 5 min and immediately put on ice. Then, the mixture  
116 containing 5x RT buffer, RNase inhibitor (Toyobo, Osaka, Japan), and reverse  
117 transcriptase (M-MLV, Nippon Gene, Tokyo, Japan) was added and incubated at 37°C  
118 for 60 min according to the manufacturer's instructions. The diluted cDNA sample  
119 (0.5 μL) was used as a template and mixed with the designed primers (1.5 μL),  
120 nuclease-free water (3 μL), and 5 μL of the PCR mixture solution (GoTaq® DNA  
121 polymerase and 5x green reaction buffer, Promega, Wisconsin, US). The PCR  
122 amplification involved denaturing at 95°C for 2 min, followed by 40 cycles at 95°C for  
123 45 s, annealing at 55°C for 30 s for all candidate lipase genes, and an extension at 73  
124 °C for 1 min. A final extension at 73°C for 10 min was followed by cooling to 4°C. All  
125 the primers used in this study are listed in Table S1. The reliability of all primer sets  
126 was checked by PCR using genomic DNA from *A. oryzae* AHU 7139 and RIB40 as  
127 the template (data not shown).

128 **Disruption of individual genes in *A. oryzae* AHU 7139**

129 Five candidate genes (AO090012000690, AO090005001319, AO090701000644,  
130 AO090003001507, and AO090005000029, named No.1, No. 19, *mdlB*, *tglA*, and *cutL*,  
131 respectively) were selected to detect the lipase activity after gene disruption. All the  
132 primers used throughout the gene disruption are shown in Table S2, based on the  
133 genomic sequence of *A. oryzae* wild-type strain RIB40 (2), used as a reference for  
134 designing primers.

135 Prior to the candidate gene disruption, the *pyrG* gene (AO090011000868) was  
136 knocked out in AHU 7139 by homologous recombination under selective pressure by  
137 5-fluoroorotic acid (5-FOA). The *pyrG* knockout was aimed at conferring the uracil  
138 auxotrophy to AHU 7139 so as to make the *pyrG* available as a selectable marker. The  
139 DNA cassette for *pyrG* knockout was prepared by fusion PCR. Briefly, the promoter  
140 and terminator regions of *pyrG* were amplified using LU*pyrG*/LL*pyrG* and  
141 RU*pyrG*/RL*pyrG* primer pairs, respectively. The AHU 7139 chromosomal DNA was  
142 used as the template, and the KOD-PLUS DNA polymerase was used for DNA  
143 amplification (Toyobo Co. Ltd., Osaka, Japan). The amplified DNA fragments were  
144 purified by gel extraction using the Wizard SV Gel and PCR Clean-Up System  
145 (Promega Co., Madison, WI, USA) according to the manufacturer's instructions. Two  
146 types of purified DNA fragments were then mixed and joined by fusion PCR using a  
147 LU*pyrG*/RL*pyrG* primer pair and KOD-PLUS (Fig. S2). The resulting 2035 bp-long  
148 DNA fragment was purified by gel extraction and then applied to the transformation  
149 of AHU 7139. Protoplasts of AHU 7139 were prepared as reported previously (7) and  
150 used for the transformation. To generate transformants, Czapek–Dox (CD) minimal  
151 agar medium supplemented with 1.2 mol L<sup>-1</sup> sorbitol, 1 mg mL<sup>-1</sup> 5-FOA, 5 mmol L<sup>-1</sup>  
152 uridine, and 10 mmol L<sup>-1</sup> uracil was used. Because the non-homologous end-joining



153 (NHEJ) activity is originally high in filamentous fungi including *A. oryzae*, about 20  
154 single colonies of transformants were screened. After single-spore isolation of  
155 transformants was performed three times using CD agar supplemented with 1 mg mL<sup>-1</sup>  
156 5-FOA, 5 mmol L<sup>-1</sup> uridine, and 10 mmol L<sup>-1</sup> uracil, the *pyrG* knockout was checked  
157 by PCR using a LUpyrG/cLpyrG primer pair. Only one positive homokaryon clone of  
158 the *pyrG* knockout mutant was acquired and named AHU 7139 [pyrG<sup>-</sup>] (Fig. S3).

159 Then, five candidate genes were individually disrupted using AHU 7139 [pyrG<sup>-</sup>] as a  
160 host. To increase the probability of the locus-specific homologous recombination, the  
161 *pyrG* marker-split method was applied to the candidate gene disruption (8). The former  
162 and latter parts of *pyrG* were amplified by PCR using PU/PLsplit and PUsplit/PL  
163 primer pairs, respectively. The 548 bp-long DNA region overlapped between them.  
164 Approximately 1 kb of promoter and terminator of the candidate genes were also  
165 amplified by PCR, using LU/LL and RU/RL primer pairs, respectively. After  
166 purification of these four types of DNA fragments by gel extraction, both pairs of  
167 promoter/*pyrG* (former part) and *pyrG* (latter part)/terminator were concatenated by  
168 fusion PCR, using LU/PLsplit and PUsplit/RL primer pairs, respectively. As a result,  
169 two types of fusion PCR DNA fragments were prepared as the gene disruption  
170 cassettes per candidate gene. The triple crossover between the chromosomal DNA and  
171 the two cassettes that occurred in the marker-split method is illustrated in Fig. S4A–E.

172 Protoplasts of AHU 7139 [pyrG<sup>-</sup>] were prepared as mentioned above. The protoplasts  
173 were applied to co-transformation with the two types of DNA fragments as gene  
174 disruption cassettes. In other words, two types of DNA fragments were simultaneously  
175 introduced into the protoplasts for each candidate gene disruption. Consequently, the  
176 transformants were generated on CD agar supplemented with 1.2 mol L<sup>-1</sup> sorbitol, and  
177 five single colonies were selected per candidate gene. The selected clones were

178 subjected to single-spore isolation on CD agar at least three times, followed by  
179 evaluating the candidate gene disruption by PCR, using cU/cL primer pairs (data not  
180 shown).

181 The candidate gene disruption was further confirmed by Southern hybridization (Fig.  
182 S5A–E). Aliquots (8 µg) of genomic DNA were digested with each restriction enzyme  
183 (*HincII*, *PstI*, *SphI*, and *PvuII*), fractionated on 0.6% agarose gel, and transferred onto  
184 a Nytran SuPerCharge membrane (GE Healthcare Co., Piscataway, NJ, USA).  
185 Hybridization and signal detection were performed using a digoxigenin (DIG) system  
186 according to the manufacturer's instructions (GE Healthcare Co.). Briefly, DIG-  
187 labeled probes (300–800 bases long) were prepared using a PCR DIG Probe Synthesis  
188 Kit (Roche Applied Science AG, Mannheim, Germany) with SBU/SBL primer pairs,  
189 followed by agarose gel extraction. The probes were hybridized to the DNA fragments  
190 bound to the membrane, and the signals of the probes were detected via  
191 chemiluminescence arising from the anti-DIG Fab fragment alkaline phosphatase  
192 conjugate (Roche Applied Science AG), using the CDP-Star Detection Reagent (GE  
193 Healthcare Co.). A WSE-6100H LuminoGraph I gel imager (ATTO Co. Ltd., Tokyo,  
194 Japan) was used for the signal detection of Southern hybridization. The acquired  
195 disruptants of the candidate genes ( $\Delta 1$ ,  $\Delta 19$ ,  $\Delta mdlB$ ,  $\Delta tglA$ , and  $\Delta cutL$ ) and the  
196 parent strain were used to measure lipase activities and FFA levels in the *in vitro*  
197 cheese model.

#### 198 **Biomass measurement by glucosamine assay**

199 The glucosamine content was determined to estimate the fungal biomass in the solid  
200 medium, using the method of Fuji et al. (9) and Kasuga (10) with some modifications.  
201 The CP (0.5 g) was washed with 50 mmol L<sup>-1</sup> sodium phosphate buffer pH 7.0 and

202 shake vigorously by hand to obtain a homogeneous dispersion. Then, the samples were  
203 centrifuged at 1,070 ×g for 10 min at room temperature, and the supernatant was  
204 discarded. After washing three times, 15 mL of the same buffer was added to disperse  
205 the solid culture, followed by the addition of 20 mg of Yatalase (Takara Bio, Tokyo,  
206 Japan). Then, the tube was shaken at 60 rpm, 37°C for 1.5 h, followed by centrifugation.  
207 The supernatant (10 mL) was filtered through a 0.45 µm filter and collected in a 15  
208 mL conical tube. The samples were kept at -20°C until use.

209 One hundred microliters of the sample or N-acetylglucosamine (GlcNAc) standard  
210 solution were transferred to a 1.5 mL microtube and then mixed with 40 µL of 0.4 mol  
211 L<sup>-1</sup> potassium tetraborate tetrahydrate solution. The sample tubes were heated at 100°C  
212 for 3 min and were then cooled down at room temperature. Then, 600 µL of p-  
213 dimethylaminobenzaldehyde (DMAB) solution (0.1 g of DMAB powder dissolved in  
214 a mixture of 17.3 mol L<sup>-1</sup> acetic acid and 0.37 mol L<sup>-1</sup> HCl) was added to the tube and  
215 incubated at 37°C for 20 min. Finally, 150 µL of each sample solution was transferred  
216 in duplicate into a 96-well microplate to measure absorbance at 595 nm, using a  
217 microplate reader (Infinite F200 PRO, Tecan, Switzerland). One milligram of dried  
218 koji mycelium contains 0.628 µmol of GlcNAc according to Fujii et al. (9).

### 219 **Preparation of the extracellular and intracellular enzymes**

220 The CPs were mixed with an equal weight of deionized water and treated by a  
221 stomacher for 5 min. The materials were then transferred to a centrifugal tube and  
222 centrifuged at 21,130 ×g, 4°C, 10 min. The supernatant was recovered and the  
223 precipitate was re-extracted with deionized water another two times. The pooled  
224 supernatants were used as the extracellular enzyme. The precipitate was washed once  
225 more with deionized water and freeze-dried. The freeze-dried sample was ground into

226 a fine powder with a mortar and pestle and then dispersed in a simulated milk  
227 ultrafiltrate (SMUF) at pH 5.5 (11), which was used as the intracellular enzyme.

### 228 **Measurement of fractionated lipase activity**

229 The extracellular enzyme (70  $\mu\text{L}$ ) was mixed with 230  $\mu\text{L}$  of SMUF buffer pH 5.5,  
230 while the suspension comprising 0.05 g of intracellular FDP dispersed in 300  $\mu\text{L}$  of  
231 SMUF pH 5.5 was used as the intracellular enzyme. The lipase activity was determined  
232 as previously described, with some modifications (12). In brief, one gram of purified  
233 triolein or diolein emulsified with 100 mL of 2% polyvinyl alcohol (degree of  
234 polymerization 2000, Kishida Chemical, Osaka, Japan) was used as the substrate.  
235 Purification of triolein was carried out by loading commercial triolein (Kanto  
236 Chemical, Tokto, Japan) on the column packed with silica gel 60 N (Kanto Chemical,  
237 Tokyo, Japan) pre-equilibrated with an organic solvent mixture of hexane and  
238 chloroform (5:1). Triolein was recovered by the isocratic elution of the same solvent.  
239 Purification of diolein was carried out by loading commercial diolein (Kanto Chemical,  
240 Tokyo, Japan) on the same column pre-equilibrated with the organic solvent mixture  
241 of hexane and chloroform (5:1). Then, diolein was recovered by the isocratic elution  
242 of another solvent mixture of hexane and chloroform (1:1). The purity of these lipids  
243 was confirmed by conventional thin layer chromatography (TLC) (Fig. S6).

244 The substrate was divided into 2 mL for each test tube and subjected to pre-incubation  
245 at 30°C for 5 min, followed by the addition of the enzyme to be tested (300  $\mu\text{L}$ ). The  
246 reaction was carried out at 30°C for 30 min. The reaction was terminated by adding  
247 7.5 mL of extract solution (heptane : isopropanol : 0.5 mol L<sup>-1</sup> sulfuric acid = 48:48:4).  
248 FFA measurement was performed using a spectrophotometer at 570 nm, and the values  
249 were corrected by subtracting the value obtained from the blank, which was subjected

250 to the same treatment except for the incorporation of the enzyme after the addition of  
251 the extract solution. Enzyme activity was expressed as  $\mu\text{mol}$  of released oleic acid  
252 from the substrate per 1 h at the defined temperature from one gram of the CPs ( $1 \mu\text{mol}$   
253 oleic acid/h/g-CP = 1 lipase unit: LU). The specific lipase activity was calculated based  
254 on the biomass and expressed as the total lipase unit per mg of dried koji biomass (LU/  
255 mg koji).

### 256 **Study of the FFA levels of mutants in a cheese curd model**

257 The *in vitro* cheese curds used in this study were prepared as previously described (6)  
258 with the addition of 2% NaCl. After producing the curds, 0.2 g of FDP prepared using  
259 the culture with an initial pH of 4.0 or 6.5 was mixed with 200 g of curds, and 20 g  
260 was divided into nine bags, and then vacuum-sealed. The culture with the parent strain  
261 (P4, P6.5) or mutants, including  $\Delta\text{mdlB}$  (M4, M6.5) and  $\Delta\text{tglA}$  (T4, T6.5), were used.  
262 The control cheese (C) had no addition of any adjunct material. The curd packs were  
263 ripened at 12°C and sampled at 4 weeks.

264 For FFA determination, the triplicate curd sample (0.2 g) was transferred to 2 mL of  
265  $7.7 \text{ mol L}^{-1}$  HCl in a test tube and placed into boiling water to dissolve the sample.  
266 FFAs, extracted by the same extraction reagent as was used in the lipase activity  
267 measurement were determined by the phenol-red method (13) and its content was  
268 expressed as millimoles per 100 g of curds using oleic acid as the standard.

### 269 **Statistical analysis**

270 FFA content in the curds was analyzed using Tukey-Kramer's multiple comparison  
271 test. The data were analyzed by JMP software (Pro 17; SAS Institute, Inc., Tokyo,  
272 Japan). Differences were considered to be statistically significant at  $p < 0.05$ .

## 273 RESULTS AND DISCUSSION

### 274 Qualitative gene expression analysis by RT-PCR to pre-screen of candidate 275 lipase genes for mutation

276 The results of RT-PCR performed with twenty-five candidate primers revealed  
277 that initial cultures with a pH of either 4.0 or 6.5 resulted in the expression of eight  
278 and six genes respectively (Fig. 1). gDNA contamination in cDNA was unlikely  
279 because no amplification was found when the mRNA preparation was used as a  
280 template (Fig. S7). Annotated lipase genes Nos. 1, 20, 22, *mdlB*, *tglA*, and *cutL* were  
281 expressed in *A. oryzae* AHU 7139 with either an initial pH 4.0 or 6.5, whereas Nos.  
282 19 and 21 showed limited expression in the culture with an initial pH of 4.0.

283 To narrow down the candidates for mutation, we evaluated the substrate  
284 specificity and distribution of the lipase in the culture products (CPs)  
285 of *A. oryzae* AHU 7139 to compare with the gene expression profile. Introducing  
286 purified substrates revealed that the parent strain of *A. oryzae* AHU 7139 produced  
287 DG lipolytic activity approximately twenty times higher than TG lipase activity at both  
288 initial pHs (Fig. 2). The activity was predominantly found in the extracellular fraction  
289 although the CPs prepared with an initial pH of 4.0 showed a higher ratio of  
290 intracellular DG lipase than those prepared at pH 6.5. It should be noted that the  
291 number of genes expressed at pH 4.0 was greater than at pH 6.5 (Fig. 1) despite the  
292 lower total lipase activity of the initial culture at pH 4.0 compared to pH 6.5 (Fig. 2).  
293 The gene No.1 (AO090012000690) annotated as a triglyceride lipase and the gene No.  
294 19 (AO090005001319), annotated as a lipase and carboxyl ester hydrolase, were the  
295 focus of further study because of the limited information available regarding substrate  
296 specificity. On the other hand, Nos. 20 (AO090005001602), 21 (AO090023000717)

297 and 22 (AO090003000839) were screened out because they were reported to be  
298 intracellular TG lipases (14) while we found limited intracellular lipase activity toward  
299 triolein under either initial pH (Fig. 2). The expression of the *cutL* (15), *mdlB* (16),  
300 and *tglA* (17) genes has been demonstrated in liquid culture and this study  
301 demonstrated their expression in a solid-state medium as well. Taking the lipase  
302 activity and the gene expression into account, two annotated lipase genes (Nos. 1 and  
303 19), *mdlB*, *tglA*, and *cutL* were selected to obtain mutants.

#### 304 **Lipase production and distribution in the mutant strains of *A. oryzae* AHU** 305 **7139**

306 We introduced a single-gene knockout of *A. oryzae* AHU 7139 to confirm the  
307 lipase productivity. Considering the growth rate difference of the mutants, the specific  
308 lipase activity that represent the activity based on the biomass was compared (Table  
309 1). Reduced specific TG lipase activity was observed in  $\Delta 1$  (694.1 LU/mg koji),  $\Delta 19$   
310 (430.3 LU/mg koji) and  $\Delta cutL$  (1079.8 LU/mg koji) when cultivation was initiated at  
311 pH 4.0, but the activity was elevated when cultivation was initiated at pH 6.5 (9165.5,  
312 20209.0, and 12047.9 LU/mg koji, respectively) compared to the parent strain (1934.7  
313 and 5570.3 LU/mg koji at pH 4.0 and 6.5, respectively). CutL from *A. oryzae* was  
314 assumed to be one of the factors responsible for rancidity in dairy products since it  
315 showed higher activity on esters of short-chain fatty acids (15). However, CutL as well  
316 as Nos. 1 and 19, were unlikely to be related to rancidity because the rancidity was  
317 more pronounced in cheese prepared using the adjuncts produced at an initial culture  
318 pH of 6.5 rather than pH 4.0. In contrast,  $\Delta mdlB$  showed the opposite response because  
319 specific TG lipase activity at pH 6.5 was reduced to 61.9% (3450 LU/mg koji)  
320 compared with that of the parent strain with slight increase of 2153.7 LU/mg koji at

321 pH 4.0. Although no substrate specificity of MdlB toward triacylglyceride has been  
322 reported (16). Lan et al. (18, 19) reported that the single point mutation of mono- and  
323 diacylglyceride lipase increased the lipolytic activity toward TG and DG making it  
324 function as a TG lipase. Thus, we cannot rule out that certain modifications of *mdlB*  
325 in *A. oryzae* AHU 7139 conferred hydrolytic capability toward triacylglycerides. On  
326 the other hand, an 83% reduction of TG lipase activity was detected in  $\Delta tglA$  (968.1  
327 LU/mg koji) when the cultivation was initiated at pH 6.5 whereas only a 25% reduction  
328 of TG lipase activity was found in  $\Delta tglA$  when the cultivation was initiated at pH 4.0  
329 (1451 LU/mg koji). Regarding the DG lipase, Nos. 1, 19, and CutL were proven to be  
330 active toward DG as well as TG since the specific lipase activity of the culture initiated  
331 at pH 4.0 was reduced by 59%, 77% and 34.6% in the  $\Delta 1$  (19889.9 LU/mg koji),  $\Delta 19$   
332 (11003.0 LU/mg koji) and  $\Delta cutL$  (31720.8 LU/mg koji) mutants, respectively. We  
333 initially speculated that the biological significance of these lipases under acidic  
334 circumstance was related to the transition into the sporulation phase of *A. oryzae* (14)  
335 because the spore-forming ability was recognized in the CP initiated at pH 4.0 but not  
336 in that initiated at pH 6.5 (Fig. S8). In fact, it has been suggested that the metabolism  
337 of storage lipids (20) and two intracellular lipases in *A. oryzae* BCC7051 (14) are  
338 involved in fungal spore production. However,  $\Delta 1$ ,  $\Delta 19$  and  $\Delta cutL$  exhibited similar  
339 spore formation at the end of the cultivation (data not shown), which indicated that  
340 these three lipases are independent on spore formation. It is unclear that an increase in  
341 the specific DG and TG lipase activity in  $\Delta 19$  in the culture initiated at pH 6.5, despite  
342 No. 19 was unexpressed under the initial culture condition at pH 6.5 (Fig.1).  $\Delta mdlB$   
343 showed the greatest lipase reduction (84%) compared to the parent strain under either  
344 pH condition (from 48519.8 LU/mg koji to 7844.1 LU/ mg koji at pH 4.0 and from  
345 104114.6 LU/ mg koji to 16393.0 LU/ mg koji at pH 6.5). Thus, major DG lipase



346 activity could be ascribed to MdlB. Furthermore, major intracellular DG lipase found  
347 in the culture at 4.0 was regarded as MdlB because  $\Delta mdlB$  showed limited intracellular  
348 lipase accumulation (Fig. 2). The MdlB found in the intracellular fraction was  
349 probably due to the limited secretion period, followed by the rapid life cycle transition  
350 into conidiation. As Toida et al. (17) reported the hydrolytic ability of TglA toward  
351 DG, partial reduction of specific DG lipase activities of  $\Delta tglA$  was confirmed in both  
352 initial pHs. Overall, in the culture initiated at pH 4.0, the predominant TG lipase  
353 activity was derived from Nos. 1, 19 and CutL, whereas that was due to TglA in the  
354 culture initiated at pH 6.5 (Table 1). These results demonstrate that the production of  
355 triglyceride lipase molecules depends on the initial culture pH. On the other hand,  
356 MdlB was the major DG lipase irrespective of the initial culture pH. Considering the  
357 higher level of total lipase activity in the culture initiated at pH 6.5, TglA and MdlB  
358 were more likely to be related to rancidity. Therefore,  $\Delta tglA$  and  $\Delta mdlB$  were selected  
359 for further study.

### 360 ***In vitro* studies of FFA profiles in the cheese curd model**

361 Finally, we conducted an *in vitro* cheese curd experiment to evaluate the effects  
362 of triacylglycerol lipase (TglA) and mono-, diacylglycerol lipase (MdlB) on the free  
363 fatty acid (FFA) levels in the curds. Table 2 shows the whole lipase activity of freeze-  
364 dried powder (FDP), used for mixing with curds. The lipase activity was higher in all  
365 CPs with an initial pH of 6.5 compared to those with an initial pH of 4.0, except for  
366 the activity of mutant  $\Delta tglA$  toward TG. The triacylglycerol lipase activity in all  
367 mutants was lower than that in the parent strain for both initial culture pHs. Moreover,  
368 lipase activity toward DG was not detected in  $\Delta mdlB$  initially cultured at pH 4.0, and  
369 low activity was observed with an initial pH of 6.5 (302.3 LU). After 4 weeks of

370 ripening, the addition of the CPs from the mutant  $\Delta tglA$  resulted in a significantly  
371 lower amount of FFA than that from the parent strain ( $p < 0.05$ ) and retained a  
372 comparable FFA level to that of the control sample (Fig. 3). Moreover, the addition of  
373 CPs from the mutant  $\Delta mdlB$  initially cultured at pH 6.5 resulted in a significantly lower  
374 FFA level than that of the parent, although higher than that of the control cheese (Fig.  
375 3).

376 FFA accumulation during cheese ripening is mainly attributed to the action of  
377 exogenous lipases because the endogenous milk lipase is heat-labile and can be  
378 inactivated by pasteurization (21). We found that the released FFA in the *in vitro* curds  
379 seemed to correlate with the TG lipase activity levels in the adjuncts (Fig.3, Table 2).  
380 Since the FFA accumulation in  $\Delta tglA$  was significantly reduced in the ripening curds  
381 regardless of the initial pH, we suggested that TglA triggers hydrolysis that results in  
382 rancidity. As described above, lipases No. 1, No. 19, and CutL should still be active in  
383 either CPs from  $\Delta tglA$  and  $\Delta mdlB$ . However, the CPs from  $\Delta tglA$  showed comparable  
384 FFA levels to the control, which demonstrated that those lipases were not as important  
385 as TglA under the cheese ripening circumstance. This is probably due to the reduced  
386 catalytic activity of other lipases at the temperature of cheese ripening (12°C) and/ or  
387 due to the degradation by concomitant rennet, which is supplied as the milk  
388 coagulation enzyme. The reduction of the FFA levels of  $\Delta mdlB$  indicates that MdlB  
389 lipase should not be overlooked when considering rancidity. Okumura et al. (22)  
390 reported that mono- and diacylglycerol lipases, from *Penicillium cyclopium* M1, more  
391 easily hydrolyze acylglycerols than triacylglycerol lipase, and the synergistic action of  
392 mono- and diacylglycerol lipase with triacylglycerol lipase could accelerate lipolysis.  
393 Wong et al. (23) found elevated FFA productivity in *A. oryzae* RIB 40 when they

394 applied the overexpressed strain AO090701000644 (*mdlB*). Based on these results, we  
395 postulated a rancidity process for ripened cheese caused by *Aspergillus* lipolysis (Fig.  
396 4). Extracellular triacylglycerol lipase (TglA) is crucial for the degradation of  
397 triglycerides in milk and the release of FFA and DG as the first step of rancidity.  
398 Subsequently, the resulting DGs are provided as the substrate for MdlB lipase. The  
399 dual effect of TglA, followed by MdlB, is likely to cause an excess amount of volatile  
400 fatty acids and rancidity in cheese products made with *A. oryzae*.

401           It might be possible to distinguish MdlB from TglA only if highly purified  
402 triolein is used as the substrate. Suzuki et al. (24) found a poor correlation between  
403 lipase activity and butyric acid levels in cheese products made with *A. oryzae* when  
404 they used a synthetic substrate of *p*-nitrophenol butyrate, which might represent whole  
405 esterase activity. Even though butter oil was used as the substrate for lipase activity  
406 measurement, FFA levels in the resulting cheese products were not correlated with the  
407 lipase activities in the adjuncts prepared using *Aspergillus* CPs (6). As butter oil  
408 included TG as the major component, with DG and MG as minor components, it could  
409 be as the substrate not only for TG lipase but also for the abundant DG lipase. Thus,  
410 when suitable strains of *A. oryzae* are screened with respect to lipase activity for the  
411 preparation of adjuncts for cheesemaking, a strict evaluation of TG activity should be  
412 performed.

413           In conclusion, *A. oryzae* is expected to generate adjuncts to enrich the flavor  
414 of ripened cheese if the excessive accumulation of volatile fatty acids due to lipase  
415 activity is reduced. In this study, we demonstrated that *tglA* is the most pivotal lipase  
416 gene encoded in *A. oryzae*. Therefore, a TglA-inactive strain might be a candidate for  
417 use as an adjunct for cheese ripening to prevent the production of rancid-off flavor.

418 We are currently investigating the reduction of TglA catalytic activity by substituting  
419 amino acid(s) and inserting or deleting nucleotides, leading to a frameshift  
420 of *tglA* transcripts. Furthermore, genome editing might be an alternative to obtain  
421 desirable strains.

## 422 **ACKNOWLEDGEMENTS**

423 This research was partially supported by grants from The Tojuro Iijima  
424 Foundation for Food Science and Technology and Sapporo Bioscience Foundation.  
425 The authors are very grateful to Yu Toba and Hiroki Matsumo, Field Science Center  
426 for Northern Biosphere, Hokkaido University, for their assistance with cheese curds  
427 preparation. We also thank Professor Satoshi Koike for the experimental support.

## 428 **REFERENCES**

- 429 1. Barbesgaard, P., Heldt-Hansen, H. P., and Diderichsen, B.: On the safety  
430 of *Aspergillus oryzae*: a review. *Appl. Microbiol. Biotechnol.*, **36**, 569–572 (1992).
- 431 2. Machida, M., Asai, K., Sato, M., Tanaka, T., Kumagai, T., Terai, G., Kusumoto, K.,  
432 Arima, T., Akita, O., Kashiwagi, Y., and other 53 others: Genome sequencing and  
433 analysis of *Aspergillus oryzae*, *Nature*, **438**, 1157–1161 (2005).
- 434 3. Kumura, H., Saito, C., Taniguchi, Y., Machiya, T., and Takahashi, Y.: Adjunctive  
435 application of solid-state culture products from *Aspergillus oryzae* for semi-hard  
436 cheese, *J Adv Dairy Res*, **5**(3), 1000188 (2017).
- 437 4. Viniegra-González, G., Favela-Torres, E., Aguilar, C. N., Romero-Gomez, S. J.,  
438 Díaz-Godínez, G., and Augur, C.: Advantages of fungal enzyme production in solid  
439 state over liquid fermentation systems, *Biochem. Eng. J.*, **13**, 157–167 (2003).

- 440 5. Ray, P. R., Chatterjee, K., Chakraborty, C., and Ghatak, P. K.: Lipolysis of milk: A  
441 Review, *Int. J. Agric. Vet. Sci.*, **1**, 58-74 (2013).
- 442 6. Chintagavongse, N., Takiguchi, H., Ming-Hsuan, C., Tamano, K., Hayakawa, T.,  
443 Wakamatsu, J., Mitani, T., and Kumura, H.: A study of lipolysis induced by adjuncts  
444 from edible *Aspergillus* sp. solid culture products on ripened semi-hard cheese, *J. Sci.*  
445 *Food Agric.*, **102**, 4355-4362 (2022).
- 446 7. Tamano, K., Satoh, Y., Ishii, T., Terabayashi, Y., Ohtaki, S., Sano, M., Takahashi,  
447 T., Koyama, Y., Mizutani, O., Abe, K., and Machida, M.: The beta-1,3-exoglucanase  
448 gene *exgA* (*exg1*) of *Aspergillus oryzae* is required to catabolize extracellular glucan,  
449 and is induced in growth on a solid surface, *Biosci. Biotechnol. Biochem.*, **71**(4), 926-  
450 934 (2007).
- 451 8. Niu, J., Arentshorst, M., Seelinger, F., Ram, A.F., and Ouedraogo, J.P.: A set of  
452 isogenic auxotrophic strains for constructing multiple gene deletion mutants and  
453 parasexual crossings in *Aspergillus niger*, *Arch. Microbiol.*, **198**(9), 861-868 (2016).
- 454 9. Fujii, F., Ozeki, K., Kanda, A., Hamachi, M., and Nunokawa, Y.: A simple method  
455 for the determination of grown mycelial content in ricekoji using commercial cell wall  
456 lytic enzyme, Yatalase. *J. Brew. Soc. Jpn.*, **87**(10), 757-759 (1992) (in Japanese).
- 457 10. Kasuga, M.: A simple method for measuring of mycelial content in soy sauce koji  
458 culture, *J. Brew. Soc. Jpn.*, **94**(2), 122-125 (1999) (in Japanese).
- 459 11. Jenness, R. and Koops, J.: Preparation and properties of a salt solution which  
460 simulates milk ultrafiltrate, *Neth. Milk Dairy J.*, **16**, 153-164 (1962).

- 461 12. Kumura, H., Satoh, M., Machiya, T., Hosono, M., Hayakawa, T., and Wakamatsu,  
462 J.: Lipase and protease production of dairy *Penicillium* sp. on milk-protein-based solid  
463 substrates, *Int. J. Dairy Technol.*, **72**, 403-408 (2019).
- 464 13. Saito, Z.: Application of the phenol-red method for investigations on the lipolysis  
465 of raw milk, *Jap J Zootech Sci*, **50**, 710–715 (1979) (in Japanese).
- 466 14. Anantayanon, J., Jeennor, S., Panchanawaporn, S., Chutrakul, C., and Laoteng, K.:  
467 Significance of two intracellular triacylglycerol lipases of *Aspergillus oryzae* in lipid  
468 mobilization: A perspective in industrial implication for microbial lipid production,  
469 *Gene*, **793**, 145745 (2021).
- 470 15. Ohnishi, K., Yoshida, Y., and Sekiguchi, J.: Lipase production of *Aspergillus*  
471 *oryzae*, *J. Biosci. Bioeng.*, **77**(5), 490-495 (1994).
- 472 16. Toida, J., Kondoh, K., Fukuzawa, M., Ohnishi, K., and Sekiguchi, J.: Purification  
473 and characterization of a lipase from *Aspergillus oryzae*, *Biosci. Biotechnol. Biochem.*,  
474 **59**(7), 1199-1203 (1995).
- 475 17. Toida, J., Arikawa, Y., Kondou, K., Fukuzawa, M., and Sekiguchi, J.: Purification  
476 and characterization of triacylglycerol lipase from *Aspergillus oryzae*, *Biosci.*  
477 *Biotechnol. Biochem.*, **62**(4), 759-763 (1998).
- 478 18. Lan, D., Popowicz, G.M., Pavlidis, I.V., Zhou, P., Bornscheuer, U.T., and Wang,  
479 Y.: Conversion of a mono- and diacylglycerol lipase into a triacylglycerol lipase by  
480 protein engineering, *ChemBioChem*, **16**, 1431-1434 (2015).
- 481 19. Lan, D., Zhao, G., Holzmann, N., Yuan, S., Wang, J., and Wang, Y.: Structure-  
482 guided rational design of a mono- and diacylglycerol lipase from *Aspergillus oryzae*:

- 483 a single residue mutant increases the hydrolysis ability, *J. Agric. Food Chem.*, **69**(18),  
484 5344-5352 (2021).
- 485 20. Engebrecht, J.A.: Cell signaling in yeast sporulation, *Biochem. Biophys. Res.*  
486 *Commun.*, **306**(2), 325-328 (2003).
- 487 21. Shipe, W. F. and Senyk, G. F.: Effects of processing conditions on lipolysis in milk,  
488 *J. Dairy Sci.*, **64**(11), 2146-2149 (1981).
- 489 22. Okumura, S., Iwai, M., and Tsujisaka, Y.: Purification and properties of partial  
490 glyceride hydrolase of *Penicillium cyclopium* M1, *J. Biochem.*, **87**(1), 205–211 (1980).
- 491 23. Wong, P.S., Tamano, K., and Aburatani, S.: Improvement of free fatty acid  
492 secretory productivity in *Aspergillus oryzae* by comprehensive analysis on time-series  
493 gene expression, *Front. Microbiol.*, **12**, 605095 (2021).
- 494 24. Suzuki, S., Ohmori, H., Hayashida, S., Nomura, M., Kobayashi, M., Hagi, T.,  
495 Narita, T., Tomita, S., Yamashita, H., Arakawa, Y., and other 3 authors: Lipase and  
496 protease activities in Koji cheeses surface-ripened with *Aspergillus* strains, *Food Sci.*  
497 *Technol. Res.*, **27**(3), 543-549 (2021).

498 **Figure Legends**

499 **Fig. 1.** Agarose gel (1%) electrophoresis of RT-PCR products with cDNA from CP  
500 initial pH 4.0 (A) and pH 6.5 (B). M = DNA ladder marker, lane no.1-22 = annotated  
501 lipase no. 1-22, m = *mdlB*, t = *tglA*, and c = *cutL* gene. Red box shows the expected  
502 amplified band while others are non-specific bands.

503 **Fig. 2.** Lipase activities of *A. oryzae* AHU 7139 (parent) and five selected mutants  
504 ( $\Delta mdlB$ ,  $\Delta tglA$ ,  $\Delta cutL$ ,  $\Delta 1$ ,  $\Delta 19$ ) toward purified triolein (TG) and diolein (DG) from  
505 the solid culture products (CP) initiated with pH 4.0 and pH 6.5. **Values are expressed**  
506 **in average value (n = 2).**

507 **Fig. 3.** Free fatty acid (FFA) profiles in cheese model at 4 weeks.

508 C = curds only (control), P4 = parent AHU 7139 with initial culture pH 4, M4 = mutant  
509 AHU 7139  $\Delta mdlB$  with initial pH 4, T4 = mutant AHU 7139  $\Delta tglA$  with initial pH 4,  
510 P6.5 = parent AHU 7139 with initial pH 6.5, M6.5 = mutant AHU 7139  $\Delta mdlB$  with  
511 initial pH 6.5, T6.5 = mutant AHU 7139  $\Delta tglA$  with initial culture pH 6.5. Results of  
512 FFA level are expressed as the mean  $\pm$  SE (n = 3). The different letters indicate a  
513 significant difference between the samples using Tukey-Kramer test (p < 0.05)

514 **Fig. 4.** The proposed rancidity process in the ripened cheese caused by *Aspergillus*  
515 lipolysis. As triacylglycerol (TG) is the major lipid in milk, triacylglycerol lipase  
516 (TglA) hydrolyzes TG at any fatty acid position ((A) sn-1, 3 or (B) sn-2) and generates  
517 diacylglycerol (DG) and free fatty acids (FAs) during ripening. Subsequently,  
518 diacylglycerol lipase (MdlB and, to some extent, TglA) hydrolyzes DG and produces  
519 FFA. The predominant lipases (TglA or MdlB) at each hydrolytic step are shown in  
520 bold, although other genes might be involved. The excess amounts of free volatile fatty  
521 acids released by fungal lipolysis could cause rancidity.



522 **Supplementary Figure Legends**

523 **Fig. S1.** RNA integrity check by non-denaturing agarose gel electrophoresis. Total  
524 RNA was isolated from *A. oryzae* AHU 7139 inoculated on whey solid culture  
525 adjusted to pH 4.0 and pH 6.5. +con = positive control of 28S and 18S RNA bands  
526 from bovine endometrial stromal cells.

527 **Fig. S2.** Construction of the DNA fragment for *pyrG* (AO090011000868) knockout  
528 in *A. oryzae* AHU 7139. The 2035 bp-long DNA fragment was constructed for the  
529 knockout. Primers used for the construction and clone check are shown as colored  
530 and black arrows, respectively.

531 **Fig. S3.** Clone check of the *pyrG* knockout mutant isolated after transformation.  
532 Transformant clone Nos. 1-3 are shown. Lane M: 1 kb DNA ladder marker [1-10 kb],  
533 lane R: RIB40 strain as a negative control, lane A: AHU 7139 strain as a negative  
534 control. Amplified DNA size: Positive clone, 1.0 kb; Negative clone, 2.1 kb. The clone  
535 No. 3 in red was considered a positive homokaryon, which was named AHU 7139  
536 [*pyrG*<sup>-</sup>].

537 **Fig. S4A.** Construction of the DNA fragment for disrupting AO090003001507 in *A.*  
538 *oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2048 bp-long and 2375 bp-long DNA fragments were  
539 constructed for the disruption. Primers used for the construction and clone check are  
540 shown as colored and black arrows, respectively.

541 **Fig. S4B.** Construction of the DNA fragment for disrupting AO090701000644 in *A.*  
542 *oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2021 bp-long and 2377 bp-long DNA fragments were  
543 constructed for the disruption. Primers used for the construction and clone check are  
544 shown as colored and black arrows, respectively.

545 **Fig. S4C.** Construction of the DNA fragment for disrupting AO090005000029 in *A.*  
546 *oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2035 bp-long and 2392 bp-long DNA fragments were

547 constructed for the disruption. Primers used for the construction and clone check are  
548 shown as colored and black arrows, respectively.

549 **Fig. S4D.** Construction of the DNA fragment for disrupting AO090012000690 in *A.*  
550 *oryzae* AHU 7139 [pyrG<sup>-</sup>]. The 2031 bp-long and 2403 bp-long DNA fragments were  
551 constructed for the disruption. Primers used for the construction and clone check are  
552 shown as colored and black arrows, respectively.

553 **Fig. S4E.** Construction of the DNA fragment for disrupting AO090005001319 in *A.*  
554 *oryzae* AHU 7139 [pyrG<sup>-</sup>]. The 2045 bp-long and 2408 bp-long DNA fragments were  
555 constructed for the disruption. Primers used for the construction and clone check are  
556 shown as colored and black arrows, respectively.

557 **Fig. S5A.** Southern hybridization analysis of AO090003001507 disruptant constructed  
558 from AHU 7139. 1: AHU 7139; 2: AO090003001507 disruptant.

559 **Fig. S5B.** Southern hybridization analysis of AO090701000644 disruptant constructed  
560 from AHU 7139. 1: AHU 7139; 2: AO090701000644 disruptant.

561 **Fig. S5C.** Southern hybridization analysis of AO090005000029 disruptant constructed  
562 from AHU 7139. 1: AHU 7139; 2: AO090005000029 disruptant.

563 **Fig. S5D.** Southern hybridization analysis of AO090012000690 disruptant constructed  
564 from AHU 7139. 1: AHU 7139; 2: AO090012000690 disruptant.

565 **Fig. S5E.** Southern hybridization analysis of AO090005001319 disruptant constructed  
566 from AHU 7139. 1: AHU 7139; 2: AO090005001319 disruptant.

567 **Fig. S6.** Thin layer chromatography of the purified lipids. M = control monoglyceride,  
568 D = control diglyceride, T = control triglyceride (tributylin), PD = purified commercial  
569 diolein using silica gel chromatography, PT = purified commercial triolein using silica  
570 gel chromatography, CD = commercial diolein, CT = commercial triolein. Migration  
571 of the sample on the silica gel plate was carried out using chloroform : methanol :

572 formic acid : DI water (45:20:2.5:1), stained with 0.05% primulin dissolved in  
573 acetone : water (4:1 v/v), and observed under UV light.

574 **Fig. S7.** Agarose gel (1%) electrophoresis of PCR using mRNA from CP with initial  
575 pH 4.0 (A) and pH 6.5 (B) as a template with all candidate lipase gene primers. M =  
576 DNA ladder marker, lane no.1-22 = annotated lipase no. 1-22, m = *mdlB*, t = *tglA*, and  
577 c = *cutL* gene. Confirming no gDNA in the mRNA sample.

578 **Fig. S8.** Growth and appearance of *A. oryzae* AHU 7139 on the WPC media incubated  
579 at 20°C for 7 days. The initial culture pH were of (A) pH 4.0 and (B) pH 6.

**Table 1.** Specific lipase activity of *A. oryzae* AHU 7139 and its mutants toward triacylglycerol (TG) and diacylglycerol (DG).

Sample	dried biomass (mg)	TG LU <sup>(1)</sup> / mg koji	DG LU <sup>(1)</sup> / mg koji
CP initial pH 4.0			
parent	0.008	1934.7	48519.8
Δ1	0.014	694.1	19889.9
Δ19	0.027	430.3	11003.0
Δ <i>mdlB</i>	0.004	2153.7	7844.1
Δ <i>tgla</i>	0.013	1451.0	32850.0
Δ <i>cutL</i>	0.015	1079.8	31720.8
CP initial pH 6.5			
parent	0.016	5570.3	104114.6
Δ1	0.018	9165.5	99601.9
Δ19	0.005	20209.0	330880.8
Δ <i>mdlB</i>	0.013	3450.7	16393.0
Δ <i>tgla</i>	0.017	968.1	88648.6
Δ <i>cutL</i>	0.017	12047.9	108810.5


Specific lipase activity (LU/ mg koji) is calculated by total LU based on fungal biomass (**n = 2**).

<sup>(1)</sup> LU represents the sum of total extracellular and intracellular lipase activity.


**Table 2.** Whole lipase activity of freeze-dried powder (FDP) for *in vitro* cheese model

Sample	Whole lipase activity	
	TG	DG
P4	25.6	1632.4
M4	15.4	0.0
T4	8.9	808.9
P6.5	86.6	2433.3
M6.5	66.8	302.3
T6.5	4.8	2226.6


P = parent strain of *A. oryzae* AHU 7139, M =  $\Delta mdlB$ , T =  $\Delta tglA$ . FDP recovered from cultures initial pH 4.0 (4) and pH 6.5 (6.5). Lipase activity is shown as total LU per 100 g of curds (n = 2).



**Fig. 1.** Agarose gel (1%) electrophoresis of RT-PCR products with cDNA from CP initial pH 4.0 (A) and pH 6.5 (B). M = DNA ladder marker, lane no.1-22 = annotated lipase no. 1-22, m = *mdlB*, t = *tglA*, and c = *cutL* gene. Red box shows the expected amplified size band while others are non-specific bands.




**Fig. 2.** Lipase activities of *A. oryzae* AHU 7139 (parent) and five selected mutants ( $\Delta mdlB$ ,  $\Delta tglA$ ,  $\Delta cutL$ ,  $\Delta 1$ ,  $\Delta 19$ ) toward purified triolein (TG) and diolein (DG) from the solid culture products (CP) initiated with pH 4.0 and pH 6.5. Values are expressed in average value (n = 2).



**Fig. 3.** Free fatty acid (FFA) profiles in cheese model at 4 weeks.

C = curds only (control), P4 = parent AHU 7139 with initial culture pH 4, M4 = mutant AHU 7139  $\Delta mdlB$  with initial pH 4, T4 = mutant AHU 7139  $\Delta tglA$  with initial pH 4, P6.5 = parent AHU 7139 with initial pH 6.5, M6.5 = mutant AHU 7139  $\Delta mdlB$  with initial pH 6.5, T6.5 = mutant AHU 7139  $\Delta tglA$  with initial culture pH 6.5. Results of FFA level are expressed as the mean  $\pm$  SE (n = 3). The different letters indicate a significant difference between the samples using Tukey-Kramer test ( $p < 0.05$ )





**Fig. 4.** The proposed rancidity process in the ripened cheese caused by *Aspergillus* lipolysis. As triacylglycerol (TG) is the major lipid in milk, triacylglycerol lipase (TglA) hydrolyzes TG at any fatty acid position ((A) sn-1, 3 or (B) sn-2) and generates diacylglycerol (DG) and free fatty acids (FAs) during ripening. Subsequently, diacylglycerol lipase (MdlB and, to some extent, TglA) hydrolyzes DG and produces FFA. The predominant lipases (TglA or MdlB) at each hydrolytic step are shown in bold, although other genes might be involved. The excess amounts of free volatile fatty acids released by fungal lipolysis could cause rancidity.

<b>Table S1. Primers used for gene expression analysis.</b>				
Gene	Primer	Sequence (5' - 3')	Predicted amplified size (bp)	
			genomic DNA	cDNA
Triacylglycerol lipase ( <i>tglA</i> ) (AO090003001507)	Forward	CTTTCACGGAGCCCTTCCAC	957	765
	Reverse	CGTAAATTAGTTCGCAGCCGC		
Mono-, di-acylglycerol lipase ( <i>mdlB</i> ) (AO090701000644)R	Forward	TGAGTAGACCCTGCGAAGCAC	1024	921
	Reverse	CGAATTAGCGCAATGGCAATCCAG		
Cutinase 1 ( <i>cutL</i> ) (AO090005000029)	Forward	GCTTTGCCCCAGGAAGAAT	987	834
	Reverse	ACTGACCTCCATCAAATGGGAG		
AO090012000690 (1)	Forward	GAGATTTGATCGCACTGATGGC	1003	864
	Reverse	GCCAAGCTTTTGGCTTTCAC		
AO090701000692 (2)	Forward	CTACCTTAGCCCGCAAGTGC	1245	1038
	Reverse	CCACTAACAAACCATCCCTTCAAC		
AO090003001432 (3)	Forward	GCCGTATGTCCTCTGCCTCTG	1362	1362
	Reverse	CCACAATCAAGGAATTAGCCGACG		
AO090003001315 (4)	Forward	TGAGCTTTTCACCGTCTCCC	1269	1269
	Reverse	ACACATTCGTCCAAACATGGC		
AO090001000143 (5)	Forward	GCTGACTTCGCAAGGGATATTC	1056	888
	Reverse	TCAAGGCTGGCAGAAGTCTC		
AO090010000619 (6)	Forward	CGGCACTGAAGTGTCTGCATAATC	1505	1368
	Reverse	TAGATGGTGGGGTCTCTGGGAG		
AO090012000103 (7)	Forward	AGTGCGATCAATTCCCTACCC	1420	1353
	Reverse	CCCACATAACTCCCCTTTGAG		
AO090124000015 (8)	Forward	CCTTCTGTCCCTTGGACCTG	1102	894
	Reverse	CAGTACTGTAGGTCAGCTTTCC		

**Table S1.** (continue) Primers used for gene expression analysis.


Gene	Primer	Sequence (5' - 3')	Predicted amplified size (bp)	
			genomic DNA	cDNA
AO090103000172 (9)	Forward	CGGGCAGTTCCCATGTCATC	1344	1344
	Reverse	GGCACCACAAAGAAGCGAAAG		
AO090138000167 (10)	Forward	TCCCGACCTCACCAGGATAG	1009	894
	Reverse	CGATCTTTCTCAGCTATAATTCGGC		
AO090038000214 (11)	Forward	CTGTTGCTTTGTATAGTGTCACG	1153	1065
	Reverse	TCTGTCACCCAGAGCAGATG		
AO090020000171 (12)	Forward	AAGAGATACCCAACCCAC	1131	957
	Reverse	CCCAGTGACCCGATGTATC		
AO090020000609 (13)	Forward	GCACGCCAAAGACGCTTAATG	1588	1356
	Reverse	CACAATCTGACCACATCATTCCGG		
AO090003000022 (14)	Forward	ACTGCGCTTTCCCAACAAAC	1327	1263
	Reverse	TCTATCACCCCGTCATTGCG		
AO090023000125 (15)	Forward	AACCACGCCACACCCATATC	1781	1644
	Reverse	GGGAAAGTGTTGGTTTAGACTGG		
AO090138000014 (16)	Forward	CCCATATAACGCTCGAAACATC	1054	915
	Reverse	ACGGATAAGGGTTGTAACCTTAGG		
AO090003000851 (17)	Forward	TCGACCAGCATTACTCCGC	1033	933
	Reverse	TAAACCCATGTGGTAGCCCC		
AO090005000930 (18)	Forward	TGACCCAAGTTCACATCCGC	1190	1125
	Reverse	CATCCAAGACATTGATCCCGC		
AO090005001319 (19)	Forward	CAGCCCACCCCTTCAATAG	918	918
	Reverse	ACGGAACGATTCTTGACGAAAAC		
AO090005001602 (20)	Forward	GGTTGCTCGTGGTAGGAATG	425	284
	Reverse	CAGTCGGCCCAGAATGAGTC		
AO090023000717 (21)	Forward	GGTGCTGTATCTGAAAATGCG	449	361
	Reverse	ATCACAACGACCTTTCGCTG		
AO090003000839 (22)	Forward	GCGCACCGATATCCCTATTAG	837	780
	Reverse	CACTCGACCTTCTTGTCATCC		

No. 1-22 in the bracket represents the annotated lipase genes from the bioinformatics database of *A. oryzae* RIB 40.


<b>Table S2. Primers used for candidate gene disruption.</b>			
Use application		Name	Sequence (5' to 3')
Knockout of AO090011000868 ( <i>pyrG</i> )	Front arm	LUpyrG	CGACTAAGCCACGATCTCGATCAT
		LLpyrG	aaaggagtagctatccaccactacCAATTGCCGCGAAAAATTAA
	Rear arm	RUpyrG	GTAGTGGTGGATACGTA
		RLpyrG	CTCCGTTGCGGATCTTGCTGCTTG
	Clone check	cLpyrG	GAGTACGTATCCACCACTAC
<i>pyrG</i> _former part		PU	GTCCATATATCGAGGCAGGT
		PLsplit	ATTGACCTACAGCGCACGC
<i>pyrG</i> _latter part		PUsplit	CCGGTAGCCAAAGATCCCTT
		PL	TCCTCATTTACTCCCGAGAT
Disruption of AO090003001507	Front arm	LU1	CGACACGGAGAATTTCCCGATGTA
		LL1	agacacctgctcgatatatggacAAGTGGAAGGGCTCCGTGAA
	Rear arm	RU1	gcagatctcgggagtaaatgaggaTTTACGATAAGGGCTCCATG
		RL1	ACGCTTTTGTAAGAGCCAGCCAC
	Clone check	cU1	TCCGAAGTCAGTAGATCGAC
cL1		AGCCACTTATTTCCGTGACC	
Disruption of AO090701000644	Front arm	LU2	GAGGAAATTCTGGACGATTCTCG
		LL2	agacacctgctcgatatatggacCTTTGCCAGTGTGCTTCGCA
	Rear arm	RU2	gcagatctcgggagtaaatgaggaCTTACATGATTTGGACGGAC
		RL2	GCATCACTCGGCAATCCTACCTAA
	Clone check	cU2	CTCGACTACGTCAGAGGAGC
		cL2	CAACCACAGAATCTATCACC
Disruption of AO090005000029	Front arm	LU3	CTGTCTTCTTTTATTCCGCTCACCAC
		LL3	agacacctgctcgatatatggacGAGGAAAGTGTTTTAGAAAGCG
	Rear arm	RU3	gcagatctcgggagtaaatgaggaTCTGCTGCCTTGCGTCCGAT
		RL3	TTGCTGAACCATGCCCTGCTATCG
	Clone check	cU3	CAAGCTACTATGGTGTGGAT
		cL3	GCAACCTCCACATCATCTCC

<b>Table S2. (continued) Primers used for candidate gene disruption.</b>			
Use application		Name	Sequence (5' to 3')
Disruption of AO090012000690	Front arm	LU4	CTCTCTCGTGGAAGTATGTAAGCG
		LL4	agacacctgcctcgatatatggacCAGTGCGATCAAATCTCAAAC
	Rear arm	RU4	gcagatctcgggagtaaagggaATTCTCATCGCCATCACTA
		RL4	CCAGGGCTCACTATTGAGGTATTG
	Clone check	cU4	GGGTTTAGCGAGATCTTATC
		cL4	TATCATCAGGGTCTGGTAGA
Disruption of AO090005001319	Front arm	LU5	TACCAAAGTGCCCGTCACCTCATT
		LL5	agacacctgcctcgatatatggacTGTGATATCGATCACGGTTTC
	Rear arm	RU5	gcagatctcgggagtaaagggaCAAATTCAAGTCAAGGCTATCG
		RL5	GTGCTGAACAGTAGCCTCAATCCA
	Clone check	cU5	GCATTCAACAATGGCGATGC
		cL5	TTAGCAGCACCATCTAGTGC
Southern hybridization	Probe 1	003-1507 SBU	CCAACATACTGTTGTCACAC
		003-1507 SBL	TCAACAGTACGGTTTCACTC
	Probe 2	701-0644 SBU	CTTACATGATTTGGACGGAC
		701-0644 SBL	GTGTAGTGTGCTTGGCCGAC
	Probe 3	005-0029 SBU	TCTCCTCTCTCTGTGTCT
		005-0029 SBL	CTCGCTTCGGATTGTATGAT
	Probe 4	012-0690 SBU	CGTGAGCGACATGACTAAGT
		012-0690 SBL	CCGTTGTTGAGCTGGAATGT
	Probe 5	005-1319 SBU	GTGGATCAAATCGGGATGAA
		005-1319 SBL	GAATCTTGGACAGGAATCGT

Tails of primers used for overlapping in fusion PCR are shown in lower case.




**Fig. S1.** RNA integrity check by non-denaturing agarose gel electrophoresis. Total RNA was isolated from *A. oryzae* AHU 7139 inoculated on whey solid culture adjusted to pH 4.0 and pH 6.5. +con = positive control of 28S and 18S RNA bands from bovine endometrial stromal cells.




**Fig. S2.** Construction of the DNA fragment for *pyrG* (AO090011000868) knockout in *A. oryzae* AHU 7139. The 2035 bp-long DNA fragment was constructed for the knockout. Primers used for the construction and clone check are shown as colored and black arrows, respectively.


M R A 1 2 3 M




**Fig. S3.** Clone check of the *pyrG* knockout mutant isolated after transformation. Transformant clone Nos. 1-3 are shown. Lane M: 1 kb DNA ladder marker [1-10 kb], lane R: RIB40 strain as a negative control, lane A: AHU 7139 strain as a negative control. Amplified DNA size: Positive clone, 1.0 kb; Negative clone, 2.1 kb. The clone No. 3 in red was considered a positive homokaryon, which was named AHU 7139 [*pyrG*<sup>-</sup>].



**Fig. S4A.** Construction of the DNA fragment for disrupting AO090003001507 in *A. oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2048 bp-long and 2375 bp-long DNA fragments were constructed for the disruption. Primers used for the construction and clone check are shown as colored and black arrows, respectively.




**Fig. S4B.** Construction of the DNA fragment for disrupting AO090701000644 in *A. oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2021 bp-long and 2377 bp-long DNA fragments were constructed for the disruption. Primers used for the construction and clone check are shown as colored and black arrows, respectively.




**Fig. S4C.** Construction of the DNA fragment for disrupting AO090005000029 in *A. oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2035 bp-long and 2392 bp-long DNA fragments were constructed for the disruption. Primers used for the construction and clone check are shown as colored and black arrows, respectively.







**Fig. S4D.** Construction of the DNA fragment for disrupting AO090012000690 in *A. oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2031 bp-long and 2403 bp-long DNA fragments were constructed for the disruption. Primers used for the construction and clone check are shown as colored and black arrows, respectively.



**Fig. S4E.** Construction of the DNA fragment for disrupting AO090005001319 in *A. oryzae* AHU 7139 [*pyrG*<sup>-</sup>]. The 2045 bp-long and 2408 bp-long DNA fragments were constructed for the disruption. Primers used for the construction and clone check are shown as colored and black arrows, respectively.


**A**

**Fig. S5A.** Southern hybridization analysis of AO090003001507 disruptant constructed from AHU 7139. 1: AHU 7139; 2: AO090003001507 disruptant.

**B**


**Fig. S5B.** Southern hybridization analysis of AO090701000644 disruptant constructed from AHU 7139. 1: AHU 7139; 2: AO090701000644 disruptant.

C




**Fig. S5C.** Southern hybridization analysis of AO090005000029 disruptant constructed from AHU 7139. 1: AHU 7139; 2: AO090005000029 disruptant.

D




**Fig. S5D.** Southern hybridization analysis of AO090012000690 disruptant constructed from AHU 7139. 1: AHU 7139; 2: AO090012000690 disruptant.


E




**Fig. S5E.** Southern hybridization analysis of AO90005001319 disruptant constructed from AHU 7139. 1: AHU 7139; 2: AO90005001319 disruptant.



**Fig. S6.** Thin layer chromatography of the purified lipids. M = control monoglyceride, D = control diglyceride, T = control triglyceride (tributylin), PD = purified commercial diolein using silica gel chromatography, PT = purified commercial triolein using silica gel chromatography, CD = commercial diolein, CT = commercial triolein. Migration of the sample on the silica gel plate was carried out using chloroform : methanol : formic acid : DI water (45:20:2.5:1), stained with 0.05% primulin dissolved in acetone : water (4:1 v/v), and observed under UV light.



**Fig. S7.** Agarose gel (1%) electrophoresis of PCR using mRNA from CP with initial pH 4.0 (A) and pH 6.5 (B) as a template with **all candidate lipase gene primers**. M = DNA ladder marker, **lane no.1-22 = annotated lipase no. 1-22, m = *mdlB*, t = *tglA*, and c = *cutL* gene. Confirming no gDNA in the mRNA sample.**



**Fig. S8.** Growth and appearance of *A. oryzae* AHU 7139 on the WPC media incubated at 20°C for 7 days. The initial culture pH were of (A) pH 4.0 and (B) pH 6.5.