Title	A Supplementary Note on the Genus Streblocera Westwood (Hymenopter : Braconidae)
Author(s)	Watanabe, Chihisa
Citation	Insecta matsumurana, 16(3-4), 158-158
Issue Date	1942-12
Doc URL	http://hdl.handle.net/2115/9494
Туре	bulletin (article)
File Information	16(3-4)_p158-158.pdf



	of mesonotum not produced as in chamorro 2
2.	Body bluish in general, with some metallic green reflections. Antennae
	brown, scape paler. Smooth and polished area on mesepisterna broad
	Schizaspidia ponapensis Ishii
-	Body with strong metallic green reflections. Antennae dark brown, scape
	brownish. Smooth and polished area on mesepisterna narrow

A SUPPLEMENTARY NOTE ON THE GENUS STREBLOCERA WESTWOOD

(HYMENOPTERA: BRACONIDAE)

Ву

Chihisa Watanabe (渡邊千旬)

As a supplement to "A preliminary revision of the genus Streblocera Westwood",* there will be given a new locality and a host record of Streblocera okadai Watanabe.

Streblocera okadai WATANABE

Streblocera okadai WATANABE, Ins. Mats., XVI, p. 10, Q &, Fig. 2, B & C (1942).

This species is new to Nippon: I have received a female collected at Hikosan, Kyūsyū, Nippon. It agrees well enough with the type series, except the 1st tergite is fuscous.

Host: Monolepta nigrobilineata Motschulsky (Coleoptera: Chrysomeridae). So far as I am aware, nothing is known concerning the host records of the congeneric species. Monolepta nigrobilineata is the first specific host record: when I visited the Agricultural Experiment Station of Kösyurei, Manchoukuo, in the last summer, I had an opportunity to examine two females of this species bred from adults of Monolepta nigrobilineata, which is a serious soybean pest in Manchoukuo.

Habitat: Manchoukuo [Kōsyurei (公主衛), 2♀♀, 13. VIII, 1939, I. Окада & I. Оіке leg.]; Nippon [Hikosan (彥山), Kyūsyū, т♀, 28. VII, 1939, K. Yasumatsu leg.].

General Distribution: Manchoukuo; Nippon.

*Ins. Mats., XVI, pp. 1-12 (1942).

[Ins. Mats., Vol. XVI, No. 3 & 4, December, 1942]