



Title	Neoparacryptus, a new name for Paracryptus Uchida (Hymenoptera, Ichmeumonidae)
Author(s)	Uchida, Toichi
Citation	Insecta matsumurana, 22(3-4), 87-87
Issue Date	1959-03
Doc URL	http://hdl.handle.net/2115/9639
Type	bulletin (article)
File Information	22(3-4)_p87-87.pdf



[Instructions for use](#)

of fore and middle ones, hind trochanter, apical third of hind tibia, and hind tarsus throughout black or fuscous.

♂. Unknown.

Holotype: ♀, Kozagawa, Wakayama, 20/ix. 1957, S. MOMOI leg.

Paratype: 1♀, Hirooka, Kochi, 25/xi. 1934, H. OKAMOTO leg.

Distribution: Japan (Honshu, Shikoku).

NEOPARACRYPTUS, A NEW NAME FOR
PARACRYPTUS UCHIDA

(Hymenoptera, Ichmeumonidae)

By TOICHI UCHIDA

Entomological Institute, Hokkaido University

Through the kind notice of Dr. H. TOWNES, as I have been able to know the fact that *Paracryptus* UCHIDA (1932) is preoccupied by *Paracryptus* SZÉPLIGETI (1916, Ann. Mus. Nat. Hung., 14: 251), the following new name is given for this genus:-

Genus *Neoparacryptus* nom. nov.

Paracryptus UCHIDA, 1932, Ins. Mats., 6: 149 (nec SZÉPLIGETI, 1916).

Type: *Paracryptus orientalis* UCHIDA, Original designation.