



Title	On the Neuroptera of the Ryukyus
Author(s)	Kuwayama, Satoru
Citation	<i>Insecta matsumurana</i> , 27(1), 38-48
Issue Date	1964-08
Doc URL	<a href="http://hdl.handle.net/2115/9720">http://hdl.handle.net/2115/9720</a>
Type	bulletin (article)
File Information	27(1)_p38-48.pdf



[Instructions for use](#)

# ON THE NEUROPTERA OF THE RYUKYUS\*

By SATORU KUWAYAMA

Hokkaido National Agricultural Experiment Station  
Sapporo, Japan

Our knowledge on the fauna of the Neuroptera in the Ryukyus which geographically covers Okinawa and Sakishima (Miyako and Yaeyama) Archipelagoes and the Daito Islands is largely due to the works of Banks, Matsumura, Nakahara, Okamoto, Walker and others. However, there is room for further research. This paper is based chiefly on the material in the Hokkaido National Agricultural Experiment Station. The material has been forwarded especially from Mr. S. Higashihirachi of the Ryukyu Plant Protection Station. In addition, Dr. J. L. Gressitt and Prof. K. Yasumatsu have generously lent for this study the valuable specimens of the Ryukyus from the collections of the Bernice P. Bishop Museum and the Faculty of Agriculture, Kyushu University, respectively. Also Prof. C. Watanabe kindly permitted me to examine the specimens identified by the late Prof. S. Matsumura and the late Dr. H. Okamoto which are preserved at the Entomological Institute, Hokkaido University. The material thus available to me numbers more than 220 specimens. I should like to express my sincere appreciation to the above mentioned gentlemen for their kindness during my course of this study.

In my previous paper "A Revisional Synopsis of the Neuroptera in Japan" [Pacific Insects, IV (2): 325-412], I objectively followed the classification system accomplished by Brues, Melander and Carpenter (1954). In this paper, however, I treated the material by adopting some modern taxonomic opinions. I have enumerated in this paper 35 species of the Neuroptera belonging to seven families, in which one newly described and eight unrecorded species are included.

The following symbols indicate the museums or institutions, in which the specimens are preserved:

BM (Bernice P. Bishop Museum, Honolulu)

HAES (Hokkaido National Agricultural Experiment Station, Sapporo)

HU (Entomological Institute, Hokkaido University, Sapporo)

KU (Entomological Laboratory, Faculty of Agriculture, Kyushu University, Fukuoka)

## Family *Corydalidae*

### 1. *Neochauliodes sinensis meridionalis* van der Weele

*Neochauliodes sinensis meridionalis* van der Weele, Notes Leyd. Mus. XXX: 260 (1909); ditto, Coll. Zool. Selys Longch. V(1): 64, Pl. IV, fig. 33 (1910); Esben-Petersen, Ent. Mitt. II: 262 (1913).

\* This work was supported by a grant in aid from the Ministry of Education of Japan. (Principal Coordinator: Prof. Keizo Yasumatsu).

*Neochauliodes sinensis* Walker f. *meridionalis*: Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1171, Fig. (1931).

*Chauliodes formosanus* Okamoto, Wien. Ent. Zeit. XXIX: 263 (1910).

*Neochauliodes formosana*: Banks, Phil. Jour. Sci. LXII: 275 (1937).

Specimen examined:—Ishigaki I. (1 ♀, July, 1925, S. Hirayama leg., HAES).

Distribution:—Ryukyus, Taiwan, Southern China.

This species is new to the fauna of the Ryukyus. Kimmins (1954) describes in detail the genital features of Chauliodini. According to my examination, the feature of female genitalia of this subspecies is near to that of *N. sinensis sinensis* (Walker) rather than that of *N. sinensis occidentalis* Weele shown by Kimmins.

#### Family **Sialidae**

##### 2. ***Sialis kumejimae*** Okamoto

*Sialis kumejimae* Okamoto, Wien. Ent. Zeit. XXIX: 257 (1910); Banks, Phil. Jour. Sci. LXII: 275 (1937).

Specimen examined:—Okinawa I.: Nakijin (1 ♀, June 2, 1962, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Taiwan.

Specimen at my disposal is to be identified as *S. kumejimae*. Banks (1937) recorded this species from Taiwan.

#### Family **Hemerobiidae**

##### 3. ***Psectra iniqua* (Hagen)**

*Hemerobius iniquus* Hagen, Verh. Zool.-bot. Gesel. Wien IX: 208 (1859).

*Annandalia curta* Needham, Rec. Ind. Mus. III: 208, Pl. XXI, figs. 2-4 (1909); Banks, Phil. Jour. Sci. LXII: 278 (1937).

*Annandalia iniqua*: Banks, Psyche XXXIX: 104 (1932); Nakahara, Mushi XXXIV: 13, Figs. 17-21, Pl. IV, fig. 8 (1960); Nakahara and Kuwayama, Nat. & Life S. Asia I: 262 (1961).

Specimen examined:—Ishigaki I.: (1 ?\*, November 10-20, 1952, G. K. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan, Thailand, Ceylon, India.

This species is new to the Ryukyus. On the systematic position of this species, Tjeder (1961) recently discussed in detail and came to the conclusion that *Annandalia* was the same as *Psectra* and *A. curta* should be considered as a synonym of *H. iniquus*. I follow the authority of Tjeder.

##### 4. ***Sympherobius okinawensis*, sp. nov.**

Head shining testaceous, sparingly with yellowish hairs; genae and postocular space dark brown. Mouth-parts dark brown, with concolorous palpi. Antennae with the two basal segments testaceous; flagellum fulvous.

Pronotum testaceous, with yellowish hairs at the sides. Meso- and metathorax dark brown, with an interrupted fulvous dorsal vitta. Legs uniformly fulvous; tarsi slightly darker.

\* Abdomen is damaged. The same applies to those that follow.

Foreswing pale fuscous, mottled with paler whitish-grey spaces which are centered on the pale interruptions of the longitudinal veins; somewhat darker towards the inner and outer margins. Longitudinal veins fuscous, variegated with short whitish streaks. Only two branches to Rs. Margins and veins clothed with yellowish hairs. Cross-veins intensively dark brown; those of the two gradate series narrowly bordered with fuscous shading; outer gradate series of four cross-veins usually arranged in two pairs between the branches of




Fig. 1. *Sympherobius okinawensis*, sp. nov.  
Holotype ( $\delta$ ),  $\times 11$ .

R and Rs; inner series of five cross-veins, of which the lower four are usually placed in a straight line.

Hindwing pale translucent grey, slightly tinged with brownish grey in the region of the pterostigma. Venation mainly fuscous, paler in the discal and basal areas; gradate cross-veins absent.




Fig. 2. *Sympherobius okinawensis*, sp. nov. Apex of abdomen of  $\delta$  from lateral side.

Abdomen almost wholly fuscous, with long yellowish hairs sparingly. Male genitalia: Ninth tergite triangular, pointed downwards, produced at the ventro-lateral part into a dark brownish process. Anal plate, or ectoproct, broad and angulated, each more or less transversely divided into a basal and distal portions, with three prominent dark brownish distal processes; the longest one arising from the external dorsal part of the plate and the next longest from the external ventral part; the shortest arises from the internal distal portion; all the processes with simple sharply pointed apices, slightly curved except the longest one; cercal callus with 11 trichobothria. Ninth sternite broad, but not very elongate; the apical part tapers into a short apex. Aedeagus large, blackish, in lateral view smoothly curved, bristled at the apex.

Size: Length of body 4 mm, of forewing 5 mm, of hindwing 4.5 mm.

Specimen examined:—Okinawa I.: Shuri (1 $\delta$ , Holotype, April 11, 1960, S. Higashi-

hirachi leg., HAES).

Distribution:—Ryukyus.

This species somewhat resembles *S. pygmaeus* (Rambur) of Europe, but easily distinguished from it by the coloration and mottling phase of the forewing. The male genitalia are also distinctive.

#### 5. *Notiobiella subolivacea* Nakahara

*Notiobiella subolivacea* Nakahara, Ann. Zool. Jap. IX: 20 (1915); Banks, Phil. Jour. Sci. LXII: 278 (1937); Nakahara, Mushi XXXIV: 8, Figs. 6–9 (1960); Kuwayama, Pac. Ins. IV: 359 (1962).

Distribution:—Ryukyus, Taiwan, Kyushu, Shikoku, Honshu.

Banks (1937) recorded the distribution of this species to the Iriomote Island, but I have not yet examined any specimens from the Ryukyus.

#### 6. *Micromusigorotus* Banks

*Micromusigorotus* Banks, Bull. Mus. Comp. Zool. LXIV: 335 (1920).

*Micromus (Archaeomicromus)igorotus*: Banks, Phil. Jour. Sci. LXII: 138 (1937).

*Eumicromusigorotus*: Nakahara and Kuwayama, Nat. & Life S. Asia I: 263 (1961).

*Pseudomicromusigorotus*: Nakahara, Mushi XXXIV: 33, Figs. 65–67 (1960).

*Eumicromusokinawanus* Nakahara, Kontyû XXIV: 189, Pl. XXI, fig. 9 (1956).

Specimen examined:—Okinawa I.: (1♀, no date, S. Sakaguchi leg., HAES).

Distribution:—Ryukyus, Taiwan, Philippines, Thailand, Sumatra.

This seems to be widely distributed in the South-east Asia.

#### 7. *Micromustimidus* Hagen

*Micromustimidus* Hagen, Ber. Verh. K. Preuss. Akad. Wiss. Berlin 1853: 481 (1853); Tjeder, S. Afr. Anim. Life VIII: 313, Figs. 458, 466, 484–513 (1961).

*Micromusnavigatorium* Brauer, Verh. Zool.-bot. Gesel. Wien XVII: 508 (1867).

*Micromuspusillus* Gerstaeker, Mitt. naturw. Ver. Neuvorp. u. Rügen XVI: 171 (1894).

*Micromusaauteri* Esben-Petersen, Ent. Mitt. I: 198 (1912); ditto, Ent. Mitt. II: 228 (1913); Banks, Phil. Jour. Sci. LXII: 280 (1937).

*Eumicromusdiminutus* Nakahara, Kontyû XXIV: 188, Fig. 6, Pl. XX, fig. 8 (1956).

*Archaeomicromustimidus*: Krüger, Stett. Ent. Zeit. LXXXIII: 171 (1922).

*Archaeomicromusnavigatorium*: Esben-Petersen, Ins. Samoa VII: 93, Pl. II, fig. 3 (1928).

*Eumicromusnavigatorium*: Kimmins, Ann. Mag. Nat. Hist. Ser. 12, VI: 245 (1953); Nakahara, Bull. Osaka Mus. Nat. Hist. 12: 39 (1960); ditto, Mushi XXXIV: 25, Figs. 38–40 (1960); Nakahara and Kuwayama, Nat. & Life S. Asia I: 262 (1961).

*Nesomicromusnavigatorium*: Zimmerman, Ins. Hawaii VI: 63 (1957).

*Eumicromusaauteri*: Nakahara, Kontyû XXIV: 188, Fig. 5, Pl. XX, fig. 7 (1956); ditto, Mushi XXXIV: 24 (1960).

Specimens examined:—Okinawa I.: Shuri (1♀, June, 1958, N. L. H. Krauss leg., BM; 2♀♀, June 4–8, 1959, S. Higashihirachi leg., HAES; 1♂, May 14, 1960, S. Higashihirachi leg., HAES); Koza (1♂ 1♀, June 8, 1961, S. Higashihirachi leg., HAES); Mt. Goga (1♀, May 9, 1961, S. Higashihirachi leg., HAES); Chizuka (1♀, September 2, 1945, J. L. Gressitt leg., BM); Yona (1♂, November 13, 1960, K. Yasumatsu leg., KU); Osato-Yonahara (1♂, November 8, 1960, K. Yasumatsu leg., KU). Ishigaki I.: Okawa (1?, March 23, 1958, S. Higashihirachi leg., HAES); (2♂♂ 1♀, December 1–15, 1952, G. E. Bohart leg., BM). Minami-

Daito I.: (1♂, March 13, 1960, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Kyushu, Honshu, Hachijo I., Taiwan, Philippines, Thailand, Malaya, Sumatra, Java, Ceylon, South India, Buru I., New Guinea, New Caledonia, New Hebrides, Fiji Is., Samoa Is., Australia, Seychelles, Madagascar, Africa, Hawaii (introduced).

This is common in the Ryukyus. Banks (1937) recorded from the Iriomote Island. According to the precise studies of Tjeder (1961), *Micromus navigatorum*, *M. pusillus*, *M. sauteri*, and *Eumicromus diminutus* are the same species as *M. timidus*. He states that the very extended distribution of this species is the widest of any intertropical species of this order.

#### Family Chrysopidae

##### 8. *Ankylopteryx octopunctata* (Fabricius)

*Hemerobius octopunctata* Fabricius, Ent. Syst. II: 85 (1793).

*Ankylopteryx octopunctata*: van der Weele, Notes Leyd. Mus. XXXI: 57, Pl. IV, fig. 21 (1909); Banks, Phil. Jour. Sci. LXII: 280 (1937).

*Ancylopteryx octopunctata*: Okamoto, Rep. Hokkaido Agr. Exp. Sta. 9: 22, Pl. V, fig. 3 (1919); Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1163, Fig. (1931).

Specimens examined:—Okinawa I.: (2♂♂ 1♀ 1?; no date, S. Sakaguchi leg., HAES, HU). Ishigaki I.: (1♂, July, 1925, S. Hirayama leg., HAES; 1♂, November 10–15, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan, Philippines, Viet-nam, Ceylon, India, Java, Sumatra, Borneo, Celebes.

Banks (1937) reported this species from the Iriomote Island.

##### 9. *Ankylopteryx delicatula* Banks

*Ankylopteryx delicatula* Banks, Phil. Jour. Sci. LXII: 280 (1937).

Distribution:—Ryukyus.

Banks (1937) described this species from the Okinawa Island, but no specimen to be identified as this species is at my disposal.

##### 10. *Ankylopteryx doleschali* Brauer

*Ankylopteryx doleschali* Brauer, Verh. Zool.-bot. Gesel. Wien XIV: 901 (1864); van der Weele, Notes Leyd. Mus. XXXI: 60 (1909).

*Ankylopteryx doleschali*: Banks, Phil. Jour. Sci. LXII: 280 (1937).

Distribution:—Ryukyus, Amboina, Celebes.

Banks (1937) recorded the distribution of this species from the Iriomote Island, but I have not examined this species.

##### 11. *Ankylopteryx gracilis* Nakahara

*Ankylopteryx gracilis* Nakahara, Kontyû XXIII: 143, Pl. XXI, fig. 1 (1955).

Specimen examined:—Ishigaki I.: (1♀, December 10–15, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan.

This is a new record for the Ryukyus.

12. *Sencera exquisita* Nakahara

*Sencera exquisita* Nakahara, Kontyû XXIII: 143, Pl. XXI, fig. 2 (1955); Nakahara and Kuwayama, Nat. & Life S. Asia I: 259 (1961).

Specimens examined:—Ishigaki I.: (1♂ 1♀, December 10–15, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan, Thailand.

This species is also first known from the Ryukyus.

13. *Chrysopa cognatella* Okamoto

*Chrysopa cognatella* Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 70 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 46, Pl. I, fig. 10, Pl. VII, fig. 4 (1919); Banks, Phil. Jour. Sci. LXII: 282 (1937); Kuwayama, Pac. Ins. IV: 366 (1962).

Specimens examined:—Okinawa I.: Koza (1♂, May 18, 1961, S. Higashihirachi leg., HAES). Ishigaki I.: (1?, January 1–5, 1953, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan, China, Kyushu, Shikoku, Honshu, Hokkaido, Sachalin.

14. *Chrysopa decorata* Esben-Petersen

*Chrysopa decorata* Esben-Petersen, Ent. Mitt. II: 260, Fig. 10 (1913); Okamoto, Rep. Hokkaido Agr. Exp. Sta. 9: 47, Pl. I, fig. 11 (1919); Banks, Phil. Jour. Sci. LXII: 282 (1937); Kuwayama, Pac. Ins. IV: 368 (1962).

Distribution:—Ryukyus, Taiwan, Honshu.

Banks (1937) recorded this species from the Okinawa Island. I have seen no specimen of this species from there.

15. *Chrysopa nipponensis* Okamoto

*Chrysopa nipponensis* Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 65 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 50, Pl. 7, fig. 7 (1919); Kuwayama, Pac. Ins. IV: 366 (1962).

Specimens examined:—Miyako I.: (1♂ 1♀, October 27, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Kyushu, Shikoku, Honshu, Hokkaido.

This species is new to the fauna of the Ryukyus.

16. *Chrysopa formosana* Matsumura

*Chrysopa formosana* Matsumura, Inj. & Benef. Ins. Sugarcane Formosa: 78, Pl. XXVIII, fig. 4 (1910); Okamoto, Rep. Hokkaido Agr. Exp. Sta. 9: 56, Pl. VI, fig. 11 (1919); Banks, Phil. Jour. Sci. LXII: 282 (1937); Kuwayama, Pac. Ins. IV: 364 (1962).

Distribution:—Ryukyus, Taiwan, Kyushu, Shikoku, Honshu, Hokkaido.

Banks (1937) reported the occurrence of this species in the Iriomote Island. I could not find any specimens to be identified as this species from the Ryukyus.

17. *Chrysopa astur* Banks

*Chrysopa astur* Banks, Phil. Jour. Sci. LXII: 283 (1937); Adams, Ins. Micronesia VIII (2): 31, Fig. 11 a-e (1959).

Specimen examined:—Okinawa I.: Osato-Yonahara (1♂, November 8, 1960, K. Yasumatsu leg., KU).

Distribution:—Ryukyus, Micronesia.

Type locality of this species is the Iriomote Island.

18. *Chrysopa boninensis* Okamoto

*Chrysopa boninensis* Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 62 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 61, Pl. I, fig. 16, Pl. IV, figs. 19-20, Pl. V, fig. 7 (1919); Adams, Ins. Micronesia VIII (2): 28, Fig. 7 d-g (1959); Kuwayama, Pac. Ins. IV: 365 (1962).

*Chrysopa boninensis* (!): Sakaguchi, Prov. List Ins. Okinawa Is.: 33 (1927).

Specimens examined:—Okinawa I.: Shuri (1♂ 1♀, June 4-9, 1959, S. Higashihirachi leg., HAES); Koza (1♂, May 18, 1961, S. Higashihirachi leg., HAES); Yona (1♀, August 16-18, 1958, T. Hidaka leg., KU); (1♀, no date, S. Sakaguchi leg., HAES); (2♀♀, no date, S. Sakaguchi leg., HU). Ishigaki I.: (1♀, April, 1937, A. Tamanaha leg., KU).

Distribution:—Ryukyus, Taiwan, Bonin Is., Shikoku, Honshu.

19. *Chrysopa suzukii* Okamoto

*Chrysopa suzukii* Okamoto, Rep. Hokkaido Agr. Exp. Sta. 9: 60, Pl. VI, fig. 10 (1919); Kuwayama, Pac. Ins. IV: 365 (1962).

Specimen examined:—Miyako I.: (1♀, October 27, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Shikoku, Honshu.

This is a new record to the Ryukyus.

20. *Chrysopa furcifera* Okamoto

*Chrysopa furcifera* Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 61 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 56, Pl. I, fig. 15, Pl. VII, fig. 5 (1919); Banks, Phil. Jour. Sci. LXII: 285 (1937); Adams, Ins. Micronesia VIII (2): 30, Fig. 10 (1959); Kuwayama, Pac. Ins. IV: 369 (1962).

Specimens examined:—Okinawa I.: Shuri (2♂♂ 1♀, May 14-July 15, 1960, S. Higashihirachi leg., HAES); Koza (3♂♂ 5♀♀, May 18-June 8, 1961, S. Higashihirachi leg., HAES); Ōgimi (1♂ 1♀, January 22, 1961, S. Higashihirachi leg., HAES); (2♂♂ 1♀, May, 1906, H. Kuroiwa leg.; 1?, no date, S. Sakaguchi leg., HU). Irabu I.: (1♀, May 30, 1960, S. Higashihirachi leg., HAES). Ishigaki I.: (1♂, May 15, 1911, T. Iwasaki leg., HU). Iriomote I.: Ohara (2♀♀, August 24-25, 1958, T. Hidaka leg., KU). Kita-Daito I.: (2♂♂, April 8, 1936, M. Goto leg., KU; 1♀, February 20, 1960, S. Higashihirachi leg., HAES). Minami-Daito I.: (12♂♂ 17♀♀, February 23-April 16, 1960, S. Higashihirachi leg., HAES; 3♂♂ 21♀♀, February 27, 1961, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Bonin Is., Kyushu, Shikoku, Honshu, Taiwan, Philippines.

This species is very common in the Ryukyus, especially abundant in the Daito Islands.

21. *Chrysopa alcestes* Banks

*Chrysopa alcestes* Banks, Proc. Ent. Soc. Wash. XIII: 102 (1911); Adams, Ins. Micronesia VIII (2): 32, Fig. 11 f-n (1959).

Specimens examined:—Okinawa I.: Naha (1♂, June 12, 1961, S. Higashihirachi leg., HAES; 1♂ 1♀, June, 1958, N. L. H. Krauss leg., BM); Shuri (5♂♂ 4♀♀, June 4-24, 1959, S. Higashihirachi leg., HAES; 12♂♂ 10♀♀, June 9-12 & August 25, 1960, S. Higashihirachi leg., HAES; 1♀, June, 1958, N. L. H. Krauss leg., BM); Koza (1♂, December 2, 1960, S. Higashihirachi leg., HAES; 3♂♂ 6♀♀, May 18-June 8, 1961, S. Higashihirachi leg., HAES); Yona (1♀, August 16-18, 1958, T. Hidaka leg., KU); Nago (2♂♂ 3♀♀, June 28-

July 8, 1961, S. Higashihirachi leg., HAES); Ôgimi (1♀, January 22, 1961, S. Higashihirachi leg., HAES); (1♂, no date, S. Sakaguchi leg., HAES). Miyako I.: (1♀, May 27, 1960, S. Higashihirachi leg., HAES; 1♀, October 27, 1952, G. E. Bohart leg., BM). Ishigaki I.: (2♀♀, November 25–December 10, 1952, G. E. Bohart leg., BM). Minami-Daito I.: (1♀, February 27, 1961, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Bonin Is., India.

Though hitherto unrecorded from the Ryukyus, this species is very common throughout the Islands.

## 22. *Chrysopa scelestes* Banks

*Chrysopa scelestes* Banks, Proc. Ent. Soc. Wash. XIII: 103 (1911); Adams, Ins. Micronesia VIII (2): 28, Fig. 8 (1959).

Specimens examined:—Miyako I.: Former Navy Barrack (1♀, December 14, no record of collector, BM; 2♂♂, October 10, 1952, G. E. Bohart leg., BM).

Distribution:—Ryukyus, Micronesia, India.

This species is new to the Ryukyus. This species is allied to *C. basalis* Walker, but the male genitalia are quite distinctive.

## 23. *Chrysopa basalis* Walker

*Chrysopa basalis* Walker, List Neur. Ins. Coll. Brit. Mus. II: 239 (1853); Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 65 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 58 (1919); Esben-Petersen, Bishop Mus. Bul. 142: 13, Figs. 1–2; l. c.: 137, Fig. 1 b (1935); Banks, Phil. Jour. Sci. LXII: 282 (1937); Adams, Ins. Micronesia VIII (2): 24, Figs. 5–6 (1959); Nakahara and Kuwayama, Nat. & Life S. Asia I: 260 (1961).

*Chrysopa formosana* Esben-Petersen (nec Matsumura), Ent. Mitt. II: 257, Fig. 7 (1913).

*Chrysopa peterseni* Okamoto (nec Navás), Rep. Hokkaido Agr. Exp. Sta. 9: 63 (1919).

*Anisochrysa paradoxa* Nakahara, Kontyû XXIII: 146, Pl. XXIII, fig. 5 (1955).

Specimen examined:—Okinawa I.: Shuri (1♀, June 6, 1959, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Taiwan, Philippines, Micronesia, Polynesia, Australia, Hawaii.

This widely distributed species seems to be uncommon in the Ryukyus.

## 24. *Chrysopa remota* Walker

*Chrysopa remota* Walker, List Neur. Ins. Coll. Brit. Mus. II: 238 (1853); Okamoto, Jour. Coll. Agr., Tohoku Imp. Univ. VI: 65 (1914); ditto, Rep. Hokkaido Agr. Exp. Sta. 9: 57, Pl. VI, fig. 7 (1919); Esben-Petersen, Ins. Samoa VII (3): 101, Pl. III, fig. 10 (1928); Banks, Phil. Jour. Sci. LXII: 285 (1937).

Distribution:—Ryukyus, Taiwan, Samoa, New Hebrides.

Originally described from the Ryukyus and Navigator's Island (Samoa). But I have not seen any specimen of this species from the Ryukyus. Banks (1937) had a doubt on occurrence in the Ryukyus.

## 25. *Nothochrysa japonica* MacLachlan

*Nothochrysa japonica* MacLachlan, Trans. Ent. Soc. Lond. 1875: 182 (1875); Okamoto, Rep. Hokkaido Agr. Exp. Sta. 9: 26, Fig. 3, Pl. II, fig. 8, Pl. V, fig. 9 (1919); Banks, Phil. Jour. Sci. LXII: 280 (1937); Kuwayama, Pac. Ins. IV: 374 (1962).

*Nothochrysa modesta* Nakahara, Kontyū XXIII: 146, Pl. XXIII, fig. 6 (1955).

Specimens examined:—Okinawa I.: Koza (1♂, May 18, 1961, S. Higashihirachi leg., HAES). Minami-Daito I.: (1♂ 2♀♀, April 14–15, 1960, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Kyushu, Shikoku, Honshu, Taiwan, China.

#### Family **Mantispidae\***

##### 26. ***Mantispa transversa* (Stitz)**

*Mantispilla transversa* Stitz, Mitt. Zool. Mus. Berlin VII: 5, Fig. 5 (1913).

*Mantispa transversa*: Kuwayama, Jour. Coll. Agr., Hokkaido Imp. Univ. XV: 256 (1925); Banks, Phil. Jour. Sci. LXII: 289 (1937).

Specimen examined:—Ishigaki I.: (1♂, October, 1951, R. H. Bohart leg., BM).

Distribution:—Ryukyus, Taiwan.

Banks (1937) recorded the distribution of this species to the Iriomote Island.

#### Family **Myrmeleontidae**

##### 27. ***Glenuroides okinawensis* Okamoto**

*Glenuroides okinawensis* Okamoto, Wien. Ent. Zeit. XXIX: 296 (1910); Nakahara, Ent. News XXIV: 301 (1913); Okamoto, Zool. Mag. XXVI: 249 (1914); Sakaguchi, Prov. List Ins. Okinawa Is.: 33 (1927); Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1157, Fig. (1931); Banks, Phil. Jour. Sci. LXII: 287 (1937).

Specimens examined:—Okinawa I.: (1♂, no date, S. Sakaguchi leg., HAES; 1♂, no date, H. Kuroiwa leg., HU; 1♂, 2♀♀, no date, S. Sakaguchi leg., HU). Ishigaki I.: Yonehara-Hoshino (1♂, August 27, 1958, T. Hidaka leg., KU).

Distribution:—Ryukyus.

This pretty antlion is indigenous to the Ryukyus.

##### 28. ***Pseudoformicaleo jacobsoni* van der Weele**

*Pseudoformicaleo jacobsoni* van der Weele, Notes Leyd. Mus. XXXI: 25, Fig. 11, Pl. II, fig. 8 (1909); Adams, Ins. Micronesia VIII (2): 15 (1959); Kuwayama, Pac. Ins. IV: 384 (1962).

*Creagris matsuokae* Okamoto, Wien. Ent. Zeit. XXIX: 282, Fig. 3 (1910); ditto, Zool. Mag. XXVI: 249 (1914); ditto, Icon. Ins. Jap. Ed. Prim.: 1529, Fig. 3019 (1932).

*Gama matsuokae*: Banks, Phil. Jour. Sci. LXII: 287 (1937).

Specimen examined:—Ishigaki I.: (1♀, May 21, 1934, K. Baba leg., HAES).

Distribution:—Ryukyus, Kyushu, Shikoku, Honshu, Taiwan, China, Micronesia, Malaya, Java, Ceylon.

##### 29. ***Distoleon parvulus* (Okamoto)**

*Myrmecaelurus parvulus* Okamoto, Wien. Ent. Zeit. XXIX: 293, Fig. 7 (1910); Nakahara, Ent. News XXIV: 301 (1913); Okamoto, Zool. Mag. XXVI: 250 (1914).

*Distoleon parvulus*: Banks, Phil. Jour. Sci. LXII: 287 (1937); Kuwayama, Pac. Ins. IV: 386 (1962).

\* Besides *M. transversa*, *Climaciella habutsuella* Okamoto, which is synonymous with *C. quadrituberculata* (Westwood), was reported from the Ryukyus by some research workers. But the type locality of *C. habutsuella* is the Yakushima Island, adjacent to the southern Kyushu. I have not yet examined this species from the Ryukyus.

Specimens examined:—Okinawa I.: (1♂, June, 1945, G. E. Bohart leg., BM; 1♀, July, 1905, H. Kuroiwa leg., HU; 2??, S. Sakaguchi leg., HU).

Distribution:—Ryukyus, Kyushu, Bonin Is.

### 30. *Distoleon bistrigatus* (Rambur)

*Myrmeleon bistrigatus* Rambur, Hist. Nat. Ins. Nevropt.: 391 (1842).

*Distoleon bistrigatus*: Banks, Ann. Ent. Soc. Am. III: 43 (1910); Adams, Ins. Micronesia VIII (2): 15 (1959).

*Eidoleon bistrigatus*: Esben-Petersen, Arkiv Zool. XI (26): 15 (1918).

*Formicaleo acuminatus* Okamoto, Wien. Ent. Zeit. XXIX: 290, Fig. 6 (1910); Nakahara, Ent. News XXIV: 301 (1913).

*Formicaleon acuminatus*: Okamoto, Zool. Mag. XXVI: 250 (1914).

*Formicaleo yayeyamensis* Sakaguchi (nom. nud.), Prov. List Ins. Okinawa Is.: 33 (1927); Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1156, Fig. (1931). **New Synonymy.**

Specimens examined:—Okinawa I.: (1♀, no date, S. Sakaguchi leg., HAES; 1♂, no date, S. Sakaguchi leg., HU). Yaeyama Is.: (1♀, August, 1903, H. Kuroiwa leg., HU; 1♀, 1910, S. Matsumura leg., HU). Irabu I.: (1♀, May 30, 1960, S. Higashihirachi leg., HAES). Minami-Daito I.: (1♂ 1♀, March 29–30, 1960, S. Higashihirachi leg., HAES).

Distribution:—Ryukyus, Bonin Is., Micronesia, India, Australia.

### 31. *Distoleon contubernalis* (MacLachlan)

*Formicaleo contubernalis* MacLachlan, Trans. Ent. Soc. Lond. 1875: 175 (1875); Okamoto, Wien. Ent. Zeit. XXIX: 289 (1910).

*Formicaleon contubernalis*: Okamoto, Zool. Mag. XXVI: 250 (1914).

*Distoleon contubernalis*: Okamoto, Ins. Mats. I: 19 (1926); Kuwayama, Pac. Ins. IV: 386 (1962).

Specimens examined:—Okinawa I.: (1♀ 1?, no date, S. Sakaguchi leg., HU).

Distribution:—Ryukyus, Kyushu, Honshu, Korea.

### 32. *Hagenomyia micans* (MacLachlan)

*Myrmeleon micans* MacLachlan, Trans. Ent. Soc. Lond. 1875: 176 (1875); Okamoto, Wien. Ent. Zeit. XXIX: 299 (1910); Nakahara, Ent. News XXIV: 301 (1913).

*Hagenomyia micans*: Okamoto, Zool. Mag. XXVI: 250 (1914); Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1158, Fig. (1931); Kuwayama, Pac. Ins. IV: 387 (1962).

Specimen examined:—Okinawa I.: (1?, August, 1904, T. Miyagi leg., HU).

Distribution:—Ryukyus, Kyushu, Shikoku, Honshu, Hokkaido, Korea, Taiwan, China.

### 33. *Myrmeleon formicarius* Linnaeus

*Myrmeleon formicarius* Linnaeus, Syst. Nat. ed. XII: 914 (1767); Esben-Petersen, Ent. Meddel. XII: 124 (1918); Kuwayama, Kontyû XXVII: 67, Pl. V, figs. A 1–4 (1959); ditto, Pac. Ins. IV: 388 (1962).

*Myrmeleon formicarius*: Okamoto (in part), Wien. Ent. Zeit. XXIX: 298 (1910); Nakahara, Ent. News XXIV: 301 (1913); Okamoto, Zool. Mag. XXVI: 250 (1914).

Distribution:—Ryukyus ?, Korea ?, Kyushu, Shikoku, Honshu, Hokkaido, Sachalin, Siberia, Caucasus, Europe.

Although this species has been recorded from the Ryukyus by Okamoto, Nakahara and others, I could not verify this as I could not encounter any specimen to be identified with *M. formicarius* from there.

34. *Grocus acer* (Walker), comb. nov.

*Myrmeleon acer* Walker, List Neur. Ins. Coll. Brit. Mus. II: 348 (1853); van der Weele, Notes Leyd. Mus. XXXI: 41, Pl. III, Fig. 14 (1909); Esben-Petersen, Ent. Mitt. XV: 22 (1926).

*Myrmeleon acer*: Adams (in part), Ins. Micronesia VIII (2): 18 (1959).

Specimen examined:—Okinawa I.: (1♀, no date, S. Sakaguchi leg., HAES).

Distribution:—Ryukyu, Micronesia, Indonesia, Australia.

So far as I am aware, this is the first record of this species from the Ryukyu.

## Family Ascalaphidae

35. *Suphalomitus okinavensis* (Okamoto), comb. nov.

*Ogcogaster okinawensis* Okamoto, Zool. Mag. XXI: 505, Pl. XI, fig. 1 (1909).

*Ogcogaster okinawensis* (!): Okamoto, Wien. Ent. Zeit. XXIX: 62 (1910); Sakaguchi, Prov. List Ins. Okinawa Is.: 34 (1927); Matsumura, 6000 Illus. Ins. Jap.-Emp.: 1160, Fig. (1931).

Specimens examined:—Okinawa I.: Shuri (1♀, September 14, 1959, S. Higashihirachi leg., HAES); Nago (1♀, June 21, 1961, S. Higashihirachi leg., HAES); Nakasone (1♂ 1♀, May 27, 1957, K. Inami leg., HAES); (1♀, no date, S. Sakaguchi leg., HAES; 4♂♂ 4♀♀, no date, S. Sakaguchi leg., HU; 1♀, July, 1905, T. Miyagi leg., HU). Ishigaki I.: Takada (1♂, June 20, 1955, S. Higashihirachi leg., HAES); (1♀, July, 1925, S. Hirayama leg., HAES; 2♂♂ 1♀, June, 1922, S. Hirayama leg., HU); (1♀, June 20, 1958, K. Takara leg., KU). Iriomote I.: (1♂, September 26, 1958, S. Higashihirachi leg., KU).

Distribution:—Ryukyu.

Although this species is indigenous to the Ryukyu, it is rather common throughout the Islands.

## Literature

- Adams, P. A. 1959. Neuroptera: Myrmeleontidae and Chrysopidae. Ins. Micronesia VIII (2): 13-33.
- Banks, N. 1937. Neuropteroid Insects from Formosa. Phil. Jour. Sci. LXII: 255-291.
- Kuwayama, S. 1962. A Revisional Synopsis of the Neuroptera in Japan. Pac. Ins. IV: 325-412.
- Kuwayama, S. 1964. On the Neuroptera from Amami-Oshima and Yakushima. Mushi XXXVIII. (In the press).
- Matsumura, S. 1931. 6000 Illustrated Insects of Japan-Empire. 1497 pp.
- Nakahara, W. 1955. New Chrysopidae from Formosa. Kontyū XXIII: 143-147.
- Nakahara, W. 1956. New or Little Known Hemerobiidae from Japan and Adjacent Territories (Neuroptera). Kontyū XXIV: 182-191.
- Okamoto, H. 1909. On the Japanese Ascalaphidae. Zool. Mag. XXI: 499-508.
- Okamoto, H. 1910. Die Ascalaphiden Japans. Wien. Ent. Zeit. XXIX: 57-65.
- Okamoto, H. 1910. Die Sialiden Japans. Wien. Ent. Zeit. XXIX: 255-263.
- Okamoto, H. 1910. Die Myrmeleoniden Japans. Wien. Ent. Zeit. XXIX: 275-300.
- Okamoto, H. 1914. Verzeichnis der japanischen Myrmeleoniden. Zool. Mag. XXVI: 249-250.
- Sakaguchi, S. 1927. The Provisional List of the Insects collected in Okinawa Islands. 34 pp.
- Walker, F. 1853. Sialdes-Nemopterides. Catalogue of the Specimens of Neuropterous Insects in the Collection of the British Museum. II: 193-476.