



HOKKAIDO UNIVERSITY

Title	Histological, Histochemical and Biochemical Changes in the Anthers of <i>Solanum nigrum</i> L. Plants Infested by <i>Aphis spiricola</i> Patch
Author(s)	Chauhan, S.V.S.; Rathore, Rajesh K.S.; Shrivastava, J.N. et al.
Citation	Journal of the Faculty of Agriculture, Hokkaido University, 61(2), 200-207
Issue Date	1983-03
Doc URL	https://hdl.handle.net/2115/12981
Type	departmental bulletin paper
File Information	61(2)_p200-207.pdf



HISTOLOGICAL, HISTOCHEMICAL AND BIOCHEMICAL
CHANGES IN THE ANTHERS OF *SOLANUM*
NIGRUM L. PLANTS INFESTED BY
APHIS SPIRALCOLA PATCH.

S. V. S. CHAUHAN, Rajesh K. S. RATHORE
and J. N. SHRIVASTAVA

(Department of Botany, R. B. S. College, Agra-282002, India)

Toshiro KINOSHITA

(Plant Breeding Institute, Faculty of Agriculture,
Hokkaido University, Sapporo, Japan)

(Received August 21, 1982)

Introduction

Substantial literature has appeared on histopathological and biochemical changes induced in the vegetative plant parts infested by aphids^{1,6,17}. However, the manifestation of aphids on floral parts has received less attention^{18,19}. The present investigation was undertaken with the aim of finding histopathological and biochemical changes in the anthers of *Solanum nigrum* plants infested by a sap sucking insect *Aphis spiraeicola*.

Material and Methods

The seeds of *S. nigrum* collected from the fields of Agra district were sown in earthen pots. The plants thus raised were inoculated with the nymphs of the aphid *Aphis spiraeicola* patch (Order: Hemiptera; family: Aphididae) a fortnight before floral bud initiation. Some plants were left to serve as controls. The pollen sterility of these plants was checked by the procedure after ALEXANDER³. For histopathological and histochemical studies, the floral buds of infested and control plants were fixed in formalin-acetic-alcohol. These were processed by customary schedules and sections were cut at 6-12 μ m. For histopathological studies, these sections were stained with Heidenhain's iron-alum haematoxylin. For histochemical localization of total carbohydrates of insoluble polysaccharides in the microtomed sections, periodic Schiff's (PAS) test as described by JENSEN¹⁴ was followed.

Total proteins in the anthers of infested and control plants at the different stages of anther development were estimated quantitatively by LOWRY *et al.*¹⁰ in weighed dry and ground samples using a standard curve prepared from Bovine serum albumin.

Free proline in the anthers of infested and control plants at different stages of anther development was quantitatively estimated by the colorimetric method of BATES *et al.*⁴. Standard curve was made by using pure proline (B. D. H.).

Experimental Results

I. HISTOPATHOLOGICAL:

A. *Pollen Sterility*: The infested plants exhibited variable degrees of pollen sterility. The extent of sterility was directly proportional to the intensity of infestation, markedly influenced by humidity and shade (Figs. 1, 2). Based on the extent of pollen sterility caused by the aphid infestation, the aphid infested and control plants were grouped into four classes, namely; (i) Control (C) plants with dehiscent anthers containing 0-20% non-viable pollen grains, (ii) plants infested by 5-15 insects per plant were more or less normal (N). The anthers of such plants were dehiscent with 0-20% sterile pollen, (iii) semi-sterile (S. S.) type of plants infested by 16-100 aphids per plant. Such plants exhibited most the variable degrees of pollen sterility ranging between 21-95%. The anthers of these plants were dehiscent, partially dehiscent or non-dehiscent and (iv) Complete sterile (C. S.) plants infested severely by more than 100 aphids per plants. The anthers of these plants were usually indehiscent and possessed 96-100% sterile pollen grains.

B. *Anther Development*: Control and N type infested plants exhibited normal development. However, anther wall layers and connective parenchyma in S. S. and C. S. type of infested plants exhibited abnormalities. In the following paragraphs, the description is limited to these groups where they exhibited deviations from normal course.

a. *S. S. Type*: The behaviour of various anther wall layers, especially that of endothecium and tapetum and anther connective was quite variable. The abnormalities exhibited by various parts of an anther were directly proportional to the extent of pollen sterility. The anthers with pollen sterility ranging between 21-50% exhibited more or less normal development (Fig. 3). However, the degeneration of tapetum was delayed. The cells in the endothecium became very tangentially stretched and formation of characteristic fibrous bands on their radial walls was completely inhibited to make the anthers indehiscent (Fig. 4). The anthers exhibiting 51-65% pollen sterility

were partially dehiscent (Fig. 5). In such anthers, the microsporangia situated towards the center were dehiscent, while the microsporangia on the corolla side were indehiscent. The behaviour of tapetum in the dehiscent microsporangia resembled that of completely dehiscent anthers of this group of plants, while the degeneration of tapetum in the indehiscent sporangia was further delayed and only when the viable pollen grains were engorged with reserves, was the tapetal breakdown complete (Fig. 6).

The completely non-dehiscent anthers exhibited high degree of pollen sterility (66–95%) among this group of plants. The degeneration of tapetum was further delayed. The formation of fibrous thickenings on the radial walls of endothelial cells in such anthers was completely inhibited (Figs. 7, 8).

b. *C. S. Type*: Anthers of this group of plants exhibited various abnormalities in both pre- and post-meiotic stages. The anther lobes in severely infested floral buds at early sporogenous tissue along with the tapetum in such anthers degenerated much prior to the onset of meiosis (Fig. 8). On the other hand, in most of the anthers, the tapetal cells exhibited radial enlargement at meiosis I and II stage (Fig. 10). The abnormal enlargement of tapetal cells continued up to microspore tetrad stage (Figs. 11, 12). Beyond this stage, the tapetal cells showed tangential elongation and their protoplasts remained intact up to anthesis (Figs. 13, 14). The endothelial cells in such anthers elongated radially but formation of fibrous bands on their radial walls was fully inhibited (Fig. 13). Resorption tissue also failed to function and made the anthers indehiscent (Fig. 14). The vascular strand in the anthers of this group of plants remained procambial through out and parenchyma cells in the connective region showed signs of degeneration and possessed small hyaline granules (Figs. 15, 16).

II. HISTOCHEMICAL:

A. *Localization of total carbohydrates of insoluble polysaccharides (PAS Test) in the anthers of control and infested plants:*

The evaluation of PAS reaction in the anthers of various groups of infested as well as control plants at various stages of anther development is given in Table 1. The intensity of the reaction is arbitrarily divided into four parts, viz, low or slight (+), moderate (++) , intense or high (###) and most intense or highest (####).

As is evident from Table 1, the PAS reaction in various parts of an anther in control (C) and normal (N) plants at all stages was more or less similar. In the early stages, the reaction was slight in all the parts except the outer tangential walls of epidermal cells showing moderate reaction. The

TABLE 1. Evaluation of PAS reaction in the anthers of control and infested plants at different stages of development. The reaction is divided arbitrarily into four parts viz. slight or low (+), moderate (++) , intense or high (###) and highest or most intense (####)

Stage of Development	Type of plant	Cuticle	Epidermis	Endothecium	Middle layers	Tape-tum	Sporogenous tissue	Pollen mother cells	Microspore tetrads	Microspores	Pollen grains	Anther connective
Premeiotic	<i>C & N</i>		+	+	+	+	+					+
	<i>S. S.</i>		+	+	+	+	+					+
	<i>C. S.</i>		+	+	+	+	+					+
Meiosis I & II	<i>C & N</i>		##	+	+	+		+				+
	<i>S. S.</i>		##	+	+	+		+				+
	<i>C. S.</i>		##	+	+	+		+				+
Microspore Tetrad	<i>C & N</i>	##	##	+	+	##		##				##
	<i>S. S.</i>		##	+	+	+			+			+
	<i>C. S.</i>		##	+	+	+			+			+
Microspore	<i>C & N</i>	##	##	##		##				##		##
	<i>S. S.</i>	+	##	+		+				+		+
	<i>C. S.</i>		##	+		+				+		+
Pollen	<i>C & N</i>	##	##	##		##					##	##
	<i>S. S.</i>		##	##	+	+					##	##
	<i>C. S.</i>		##	+	+						+	+

intensity of the reaction increased with age and reached its maximum at microspore tetrad stage. The microspores, soon after their release from the callose wall, showed a steady increase of the reaction and pollen grains, ready to shed, possessed a large amount of total carbohydrates of insoluble polysaccharides (TCIP). The anther wall layers and connective parenchyma cells showed the presence of TICP grain (Figs. 17, 18).

On the other hand, the intensity of PAS reaction in various parts of an anther of *S. S.* and *C. S.* type of infested plants failed to accelerate and was far below as compared to that of control plants (Figs. 19, 20). The connective parenchyma cells and anther wall layers either possess a limited number of TICP grains (Fig. 19) or these grains are completely absent (Fig. 20).

III. BIOCHEMICAL:

A. *Quantitative estimation of total proteins and free proline in the anthers of control and infested plants:*

The quantitative estimation of total proteins and free proline in the anthers of control and infested plants at different stages of development is given in Table 2.

TABLE 2. Quantitative estimation of total proteins and free proline in the anthers of control and *C. S.* type of infested plants at different stages of development

Stages of development	Total proteins (% dry wt.)		Free proline (μ g/mg fresh wt.)	
	Control	<i>C. S.</i>	Control	<i>C. S.</i>
a. Pre-meiotic	2.5	9.5	Trace	Trace
b. Meiosis I & II, Tetrad and Microspore	3.4	1.2	0.097	Trace
c. Pollen grain prior to anther dehiscence	5.2	1.6	0.34	Trace

As is evident from Table 2, the quantity of proteins in the anthers of control plants increased considerably with age and anthers about to dehisce exhibited highest protein concentration. On the other hand, the anthers of *C. S.* type of infested plants were markedly deficient in proteins at all stages of development. Similarly, the amount of free proline in the anthers of control plants increase gradually with age and maximum was recorded at mature pollen grain stage, while only traces of free proline were found in the mature anthers of *C. S.* type of infested plants.

Discussion

From the foregoing description it is evident that *Solanum nigrum* plants infested by *Aphis spiraeicola* exhibited pollen sterility of variable degrees. The extent of pollen sterility was directly proportional to the degree of infestation. Anther ontogeny exhibited that pollen abortion in infested plants was associated with abnormalities in tapetal behaviour. Similar observations in *Raphanus sativus* plants infested by an aphid *Lipaphis erysimi* KALT. have also been recorded earlier²⁰. Malfunctioning of tapetum in infested plants was similar to that observed in a large number of cytoplasmic, genic and chemically induced male sterile plants. In the opinion of the present authors, the tapetal abnormalities, in all probabilities, resulted due to starvation caused by vascular inhibition in the anther connective. This is also corroborated by the present histochemical and biochemical findings showing marked deficiency of total carbohydrates of insoluble polysaccharides, and total proteins in the anthers of infested plants exhibiting a high degree of pollen sterility. Deficiency of total carbohydrates of insoluble polysaccharides, proteins, histones and nucleic acids in the anthers of several cytoplasmic, genic as well as chemically induced male sterile plants have also been reported^{8,10}. Thus the effect of the aphid infestation on anther development and microsporogenesis is similar to that of cytoplasmic, genic factors and gametocidal compounds.

The anthers of heavily infested plants also exhibited marked deficiency of free proline. Proline is one of the major amino acids present in the pollen grains of most plants and is connected with the pollen fertility and sexual process in plants.

Another notable finding of the present study is the confirmation of DE FOSSARD's, concept of programmed control of tapetum on endothelial development. According to DE FOSSARD,¹² the development of endothecium is controlled by tapetum during major course of anther development and only after tapetal breakdown, the characteristic fibrous thickenings appear on the radial walls of endothelial cells. In the presently studied material, endothelial thickenings developed only after the tapetal breakdown in the anthers of infested plants irrespective of the degree of pollen sterility induced.

Summary

Solanum nigrum plants infested by *Aphis spiraeicola* exhibited various degrees of pollen sterility associated with malfunctioning of tapetum caused by vascular inhibition. Impairment in the tapetal development is also re-

flected by inhibition of endothecium development. Histochemical and biochemical findings showed a marked deficiency of total carbohydrates of insoluble polysaccharides, total proteins and free proline in the anthers of heavily infested plants.

Acknowledgements

Sincere thanks are due to Dr. Bahadur SINGH, Professor Emeritus for his valuable help; to Dr. S. N. CHATURVEDI, Reader and Head for encouragement and to Dr. Roshan SINGH, Principal, R. B. S. College for the facilities. One of us (RKS) is grateful to the C. S. I. R., New Delhi for the award of Post Doctoral Fellowship.

Literature Cited

1. AGRIOS, G. N.: Plant Pathology. Academic Press, New York. 1978
2. ALAM, S. and SANDAL, P. C.: Electrophoretic analyses of anther proteins from male-sterile and male-fertile sudangrass, *Sorghum vulgare* L. var. *sudanese* (Piper). *Crop Sci.* **9**: 157-159. 1969
3. ALEXANDER, M. P. L.: Differential staining of aborted and non-aborted pollen. *Stain Tech.* **44**: 117-22. 1969
4. BATES, L. S., WALDREN, R. P. and TEARE, I. D.: Rapid determination of free proline for water-stress studies. *Plant and Soil* **39**: 205-207. 1973
5. BRITIKOV, E. A., MUSATAVA, N. A., VLADIMIRTSEVA, S. V. and PROTSENKO, M. A.: Proline in the reproductive system of plants. In: Pollen Physiology and Fertilization, LINSKENS, H. F. (Ed.) North-Holland Publishing Co. Amsterdam. 1964. 77-85. 1964
6. CARTER, W.: Insects in relation of plant diseases. John Wiley, New York. 1973
7. CHAUHAN, S. V. S.: Development of endothecium in relation to tapetal behaviour in some male-sterile plants. *Phytomorph.* **29**: 245-251. 1979
8. CHAUHAN, S. V. S. and KINOSHITA, T.: Histochemical localization of histones, DNA, proteins in the anthers of male-fertile and male-sterile plants. *Japan J. Breed.* **29**: 287-293. 1979
9. CHAUHAN, S. V. S. and KINOSHITA, T.: Anther ontogeny in genic, cytoplasmic and chemically induced male sterile plants. *Japan J. Breed.* **30**: 117-124
10. CHUHAN, S. V. S. and KINOSHITA, T.: Cyto-histological and biochemical studies on pollen abortion in *Datura alba* L. plants treated with gametocidal compounds. *Proc. Japan Acad. Sci. Ser. B.* **56**: 344-349. 1980 b
11. CHAUHAN, S. V. S. and RATHORE, RAJESH K. S.: Histochemical evaluation of PAS reaction in the anthers of male-fertile and male-sterile plants. *Curr. Sci.* **49**: 433-436. 1980
12. DE FOSSARD, R. A.: Development and histochemistry of the endothecium *in vitro* grown *Chenopodium rubrum* L. *Bot. Gaz.* **180**: 10-20. 1969

13. GOTTSCHALK, W. and KAUL, M. I. H. L.: The genetic control of microsporogenesis in higher plants. *Nucleus* (Calcutta). **17**: 133-166. 1974
14. JENSEN, W. A.: Botanical Histochemistry, Principles and Practice. H. W. Freeman & Co., San Francisco. 1962
15. LASER, K. D. and LERSTEN, N. R.: Anatomy and cytology of microsporogenesis in cytoplasmic male sterile angiosperms. *Bot. Rev.* **38**: 425-454. 1972
16. LOWRY, O. H., ROSENBOUGH, N. J., FARR, A. L. and RANDALL, R. J.: Protein measurement with the Folin-phenol reagent. *J. Biol. Chem.* **193**: 265, 1951
17. MEHROTRA, R. S.: Plant pathology. Tata McGraw Hill Publishing Co., Ltd. New Delhi. 1980
18. MEYER, V. G.: Floral abnormalities. *Bot. Rev.* **52**: 165-218. 1966
19. SRIVASTAVA, J. N. and CHAUHAN, S. V. S.: Floral abnormalities in infected plants. *B. V. J. Agric. Sci. Res.* **20**: 69-78. 1978
20. SRIVASTAVA, J. N. and CHAUHAN, S. V. S.: Studies in *Raphanus sativus* L. plants infested by *Lipaphis erysimi* KALT. *Curr. Sci.* **48**: 639-641. 1979
21. TUPY, J.: Metabolism of proline in styles and pollen tubes of *Nicotiana glauca*. In: Pollen Physiology and Fertilization, LINSKENS, H. F. (Ed.). North-Holland Publishing Co. Amsterdam. p. 86-94. 1964

Legend for Plate I

Figs. 1-2. *Solanum nigrum* plants infested by *Aphis spiraeicola*.

Fig. 1. Infested plant. Note the presence of aphids on stem, leaves, flowers and fruits.

Fig. 2. Magnified view of the infested plant shown in Fig. 1.

Figs. 3-10. Transverse anthers of *S. S.* and *C. S.* type of infested plants.

Fig. 3. *S. S.* anthers with lower range of pollen sterility. 120 \times .






Fig. 4. Magnified view of the anther shown in Fig. 3. 310 \times .

Fig. 5. *S. S.* partially dehiscent anthers. 120 \times .

Fig. 6. Magnified view of the anther shown in Fig. 5. 410 \times .








Fig. 7. *S. S.* non-dehiscent anthers. 120 \times .

Fig. 8. Magnified view of the anther shown in Fig. 7. 410 \times .



Legend for Plate II

- Fig. 9. *C. S.* Anthers showing degeneration of sporogeneous tissue and tapetal cells. 120 \times .
- Fig. 10. *C. S.* Anthers showing tapetal enlargement in radial direction at pollen mother cell stage. 380 \times .
- Fig. 11-16. Transverse anthers of *C. S.* type of infested plants.
- Fig. 11 & 12. Microspore tetrad stage showing tapetal enlargement. 380 \times .
- Fig. 13 & 14. Pollen grain stage showing intact tapetal protoplast. Note the absence of fibrous bands in endothelial cells and non-functional resorption tissue (RT). 380 \times .
- Fig. 15 & 16. Pollen grain stage showing poorly developed vascular tissue and presence of hyaline granules in the connective parenchyma. 410 \times .



Legend for Plate III

- Figs. 17-20. Transverse anthers of *N*, *S. S.* and *C. S.* type of plants showing PAS reaction. 410×.
- Fig. 17. *N* pollen grains with TCIP (total carbohydrates of insoluble polysaccharides) grains.
- Fig. 18. *N* anther connective and wall layers showing the presence of TCIP grains.
- Fig. 19. *S. S.* anther showing poor PAS reaction and reduction in the TCIP grains.
- Fig. 20. *C. S.* anther showing poor PAS reaction and complete absence of TCIP grains.

