



Title	Gauge theory on a non-simply connected domain and representations of canonical commutation relations
Author(s)	Arai, Asao
Citation	Journal of Mathematical Physics, 36(6), 2569-2580 https://doi.org/10.1063/1.531051
Issue Date	1995-06
Doc URL	https://hdl.handle.net/2115/13684
Rights	Copyright © 1995 American Institute of Physics
Type	journal article
File Information	jmp36-6.pdf



Gauge theory on a non-simply connected domain and representations of canonical commutation relations

Asao Arai

Department of Mathematics, Hokkaido University, Sapporo 060, Japan

(Received 17 January 1995; accepted for publication 10 February 1995)

A quantum system of a particle interacting with a (non-Abelian) gauge field on the non-simply connected domain $M = \mathbf{R}^2 \setminus \{\mathbf{a}_n\}_{n=1}^N$ is considered, where \mathbf{a}_n , $n = 1, \dots, N$, are fixed isolated points in \mathbf{R}^2 . The gauge potential A of the gauge field is a $p \times p$ anti-Hermitian matrix-valued 1-form on M , and may be strongly singular at the points \mathbf{a}_n , $n = 1, \dots, N$. If A is flat, then the physical momentum and the position operators $\{P_j, q_j\}_{j=1}^2$ of the particle satisfy the canonical commutation relations (CCR) with two degrees of freedom on a suitable dense domain of the Hilbert space $L^2(\mathbf{R}^2; \mathbf{C}^p)$. A necessary and sufficient condition for this representation to be the Schrödinger 2-system is given in terms of the Wilson loops of the rectangles not intersecting \mathbf{a}_n , $n = 1, \dots, N$. This also gives a characterization for the representation to be non-Schrödinger. It is proven that, for a class of gauge potentials, which is not necessarily flat, P_j is essentially self-adjoint. Moreover, an example, which gives a class of non-Schrödinger representations of the CCR with two degrees of freedom, is discussed in some detail. © 1995 American Institute of Physics.

I. INTRODUCTION

In Ref. 1, the author considered a quantum system of a charged particle with charge $q \in \mathbf{R} \setminus \{0\}$ moving in the plane \mathbf{R}^2 under the influence of a perpendicular magnetic field, which may be singular at given fixed isolated points $\mathbf{a}_n = (a_{n1}, a_{n2}) \in \mathbf{R}^2$, $n = 1, \dots, N$. In this quantum system, the position operators q_j and the physical momentum operators $P_j = p_j - qA_j$ ($j = 1, 2$), where p_j is the canonical momentum operator with respect to q_j and (A_1, A_2) is a vector potential of the magnetic field, satisfy, on a dense domain of $L^2(\mathbf{R}^2)$, the canonical commutation relations (CCR) with two degrees of freedom if and only if the magnetic field is concentrated on the set $\{\mathbf{a}_n\}_{n=1}^N$.

A set $\{\hat{P}_1, \dots, \hat{P}_n; \hat{Q}_1, \dots, \hat{Q}_n\}$ of self-adjoint operators on a Hilbert space \mathcal{H} is called a *Schrödinger n -system* if there exist mutually orthogonal closed subspaces \mathcal{H}_α of \mathcal{H} , such that $\mathcal{H} = \bigoplus_\alpha \mathcal{H}_\alpha$ with the following properties: (i) each \mathcal{H}_α reduces all \hat{P}_j , \hat{Q}_j ; (ii) the set $\{\hat{P}_1, \dots, \hat{P}_n; \hat{Q}_1, \dots, \hat{Q}_n\}$ is, in each \mathcal{H}_α , irreducible and unitarily equivalent to the Schrödinger system $\{P_1^s, \dots, P_n^s; Q_1^s, \dots, Q_n^s\}$, where P_j^s and Q_j^s are self-adjoint operators on $L^2(\mathbf{R}^n)$ defined by $P_j^s = -i\partial_j$ [∂_j denotes the generalized partial differential operator in the j th variable x_j of $\mathbf{x} = (x_1, \dots, x_n) \in \mathbf{R}^n$] and $Q_j^s = x_j$ (the multiplication operator by x_j) (Ref. 2, p. 81). We call a representation of the CCR with n degrees of freedom *non-Schrödinger* if it is not a Schrödinger n -system.

It was proven in Ref. 1 that the above mentioned representation $\{\bar{P}_j, q_j\}_{j=1}^2$ of CCR, where \bar{P}_j denotes the closure of P_j , is a Schrödinger 2-system if and only if the magnetic flux is locally quantized [i.e., the magnetic flux of every rectangle not intersecting \mathbf{a}_n ($n = 1, \dots, N$) is an integer multiple of $2\pi/q$]. This result, which generalizes the main result of Ref. 3 concerning a special example in the case $N = 1$, shows that, if the magnetic field is concentrated on $\{\mathbf{a}_n\}_{n=1}^N$ and the magnetic flux is not locally quantized, then $\{\bar{P}_j, q_j\}_{j=1}^2$ is a non-Schrödinger representation of CCR. Thus, a class of non-Schrödinger representations of CCR is constructed. Of interest concerning these non-Schrödinger representations is their relation to the Aharonov-Bohm effect⁴ in an idealized sense. From a mathematical point of view, the non-simply connectedness of the base manifold,

$$M = \mathbf{R}^2 \setminus \{\mathbf{a}_n\}_{n=1}^N, \quad (1.1)$$

is essential for the analysis just mentioned to be non-trivial. Further properties of the quantum system were investigated in terms of the Dirac–Weyl operator.⁵

An operator-theoretical analysis related to Refs. 1 and 3 has been made by Kurose and Nakazato,⁶ who have constructed a $*$ representation of the Weyl algebra with two degrees of freedom induced by a one-dimensional representation of the fundamental group of M , and proved that the $*$ representation is unitarily equivalent to the $*$ algebra generated by $\{P_j, q_j\}_{j=1}^2$.

The quantum system discussed in Ref. 1 is an example of Abelian gauge theory with gauge group $U(1)$, the one-dimensional unitary group. It is natural and interesting to extend the analysis made in Ref. 1 to the case of non-Abelian gauge theory. With this motivation, we consider in this paper a quantum mechanical particle moving in M under the influence of a (non-Abelian) gauge field. Indeed, in this case too, the position and the physical momentum operators of the particle give a representation of the CCR with two degrees of freedom if the gauge field strength is concentrated on $\{\mathbf{a}_n\}_{n=1}^N$. We formulate a necessary and sufficient condition for the representation to be a Schrödinger 2-system. As in the case of Ref. 1, this result gives a class of non-Schrödinger representations of CCR, which may give a form of *non-commutative Aharonov–Bohm effect*. Our analysis is general, in that it applies to any gauge theory on M with a finite-dimensional unitary representation of a Lie group. The Dirac–Weyl operator in the present case will be discussed in a separate paper.

In Sec. II, under the assumption that the physical momentum operators P_1, P_2 are essentially self-adjoint, we compute the commutation relations (in the strong sense) of the position operators q_1, q_2 and \bar{P}_1, \bar{P}_2 . It is shown that $\{\bar{P}_j, q_j\}_{j=1}^2$ is a Schrödinger 2-system if and only if the Wilson loop of every rectangle not intersecting \mathbf{a}_n ($n = 1, \dots, N$) is equal to the identity. In Sec. III, we derive, in terms of the gauge potential, a condition equivalent to the condition for the Wilson loops just mentioned. This can be done by employing the theory of product integrals. Section IV is devoted to the proof of essential self-adjointness of P_j for a wide class of gauge potentials. In the last section we discuss an example in some detail.

II. CCR IN (NON-ABELIAN) GAUGE THEORY

Let M be given by (1.1). We denote by $M_p^{\text{ah}}(\mathbf{C})$ the set of $p \times p$ anti-Hermitian matrices ($p \in \mathbf{N}$). Let $A_j, j = 1, 2$, be $M_p^{\text{ah}}(\mathbf{C})$ -valued continuously differentiable functions on M , and set

$$A(\mathbf{r}) = A_1(\mathbf{r})dx + A_2(\mathbf{r})dy, \quad \mathbf{r} = (x, y) \in M,$$

an $M_p^{\text{ah}}(\mathbf{C})$ -valued 1-form on M . This 1-form may be regarded as a gauge potential in a gauge theory on M with a p -dimensional unitary representation of a Lie group.

We shall use the system of physical units where \hbar , the Planck constant divided by 2π , is equal to 1. Let ∂_1 and ∂_2 be the generalized partial differential operators in x and y , respectively, and set

$$p_j = -i\partial_j, \quad j = 1, 2.$$

Then the physical momentum operator of a quantum mechanical particle interacting with the gauge potential A may be given by $\mathbf{P} = (P_1, P_2)$, with

$$P_j = p_j - iA_j, \quad j = 1, 2,$$

acting in the Hilbert space $L^2(\mathbf{R}^2; \mathbf{C}^p) [\simeq L^2(M; \mathbf{C}^p)]$ of \mathbf{C}^p -valued square integrable functions on \mathbf{R}^2 . We denote by $C_0^m(M; \mathbf{C}^p)$ the set of \mathbf{C}^p -valued m times continuously differentiable functions on M with compact support. Each P_j is a symmetric operator with $C_0^2(M; \mathbf{C}^p) \subset D(P_1 P_2) \cap D(P_2 P_1)$, and

$$[P_1, P_2]\psi = -F_{12}\psi, \quad \psi \in C_0^2(M; \mathbf{C}^p),$$

where

$$F_{12} := \partial_1 A_2 - \partial_2 A_1 + [A_1, A_2]$$

is the component of the gauge field strength 2-form,

$$F(A) := dA + A \wedge A = F_{12} dx \wedge dy.$$

We say that A is flat on M if $F(A) = 0$ on M .

We denote by q_1 and q_2 the multiplication operators by x and y , respectively. The following proposition is easily proven.

Proposition 2.1: Suppose that A is flat on M . Then $\{P_j, q_j\}_{j=1}^2$ satisfies the CCR with two degrees of freedom on $C_0^2(M; \mathbf{C}^p)$: for all $\psi \in C_0^2(M; \mathbf{C}^p)$,

$$[P_j, P_k]\psi = 0, \quad [q_j, q_k]\psi = 0, \quad [q_j, P_k]\psi = i\delta_{jk}\psi, \quad j, k = 1, 2.$$

For a continuous, piecewise continuously differentiable path C in M with parametrization $\gamma(\tau) = (\gamma_1(\tau), \gamma_2(\tau))$, $\tau \in [a, b]$ ($a < b, a, b \in \mathbf{R}$), we define the Wilson loop $W_A(C)$ by

$$W_A(C) = \prod_a^b e^{-\{A_1(\gamma(\tau))\dot{\gamma}_1(\tau) + A_2(\gamma(\tau))\dot{\gamma}_2(\tau)\}d\tau},$$

where $\dot{\gamma}_j(\tau) = d\gamma_j(\tau)/d\tau$, $j = 1, 2$, and the right-hand side (RHS) is the product integral of the matrix-valued function, $-\{A_1(\gamma(\tau))\dot{\gamma}_1(\tau) + A_2(\gamma(\tau))\dot{\gamma}_2(\tau)\}$ on the interval $[a, b]$ (see Ref. 7, Sec. 1.1).⁸ It can be shown that $W_A(C)$ is independent of the choice of parametrizations of C (cf. Ref. 7, Sec. 2.1). It follows from the anti-Hermiticity of A_j that $W_A(C)$ is in $U(p)$, the set of $p \times p$ unitary matrices.

For $x, y, s, t \in \mathbf{R}$, we define two hook-shaped paths $C_{x,y,s,t}^\pm$ from (x, y) to $(x + s, y + t)$, with parametrizations $\gamma_{x,y,s,t}^\pm : [0, 1] \rightarrow \mathbf{R}^2$ given by

$$\gamma_{x,y,s,t}^-(\tau) = \begin{cases} (x + 2\tau s, y), & 0 \leq \tau \leq \frac{1}{2}, \\ (x + s, y + (2\tau - 1)t), & \frac{1}{2} \leq \tau \leq 1; \end{cases} \tag{2.1}$$

$$\gamma_{x,y,s,t}^+(\tau) = \begin{cases} (x, y + 2\tau t), & 0 \leq \tau \leq \frac{1}{2}, \\ (x + (2\tau - 1)s, y + t), & \frac{1}{2} \leq \tau \leq 1; \end{cases} \tag{2.2}$$

and set

$$C_{x,y,s,t} = \{C_{x,y,s,t}^+\}^{-1} \circ C_{x,y,s,t}^-$$

which is the rectangle: $(x, y) \rightarrow (x + s, y) \rightarrow (x + s, y + t) \rightarrow (x, y + t) \rightarrow (x, y)$.

For each $s, t \in \mathbf{R}$, let

$$\mathbf{R}_s^{(1)} = \mathbf{R} \setminus \{a_{n1}, a_{n1} - s\}_{n=1}^N, \quad \mathbf{R}_t^{(2)} = \mathbf{R} \setminus \{a_{n2}, a_{n2} - t\}_{n=1}^N,$$

and $M_{s,t} = \mathbf{R}_s^{(1)} \times \mathbf{R}_t^{(2)}$. If $(x, y) \in M_{s,t}$, then $C_{x,y,s,t}^\pm$ do not intersect \mathbf{a}_n , $n = 1, \dots, N$. Hence, for each $s, t \in \mathbf{R}$, we can define $U(p)$ -valued functions $W_{s,t}^{A,\pm}$, $W_{s,t}^A$ on $M_{s,t}$ by

$$W_{s,t}^{A,\pm}(x, y) = W_A(C_{x,y,s,t}^\pm), \quad W_{s,t}^A(x, y) = W_A(C_{x,y,s,t}), \quad (x, y) \in M_{s,t}.$$

The two-dimensional Lebesgue measure of the set $\mathbf{R}^2 \setminus M_{s,t}$ is zero. Hence, $W_{s,t}^{A,\pm}$ and $W_{s,t}^A$ can be regarded as almost everywhere (a.e.) finite $U(p)$ -valued functions on \mathbf{R}^2 , so that they define unitary operators on $L^2(\mathbf{R}^2; \mathbf{C}^p)$ as multiplication operators. We denote these unitary multiplication operators by the same symbols $W_{s,t}^{A,\pm}$, $W_{s,t}^A$, respectively.

Throughout the rest of this section, we assume the following.

Assumption: The operators P_1 and P_2 are essentially self-adjoint.

Theorem 2.2: For all $s, t \in \mathbf{R}$,

$$e^{is\bar{P}_1} e^{it\bar{P}_2} = (W_{s,t}^A)^* e^{it\bar{P}_2} e^{is\bar{P}_1}. \tag{2.3}$$

Proof: The method of proof is similar to the proof of Theorem 2.1 in Ref. 1. By the present assumption, we can apply the Trotter product formula to obtain

$$\langle \phi, e^{it\bar{P}_j} \psi \rangle = \lim_{m \rightarrow \infty} \langle \phi, (e^{itp_j/m} e^{tA_j/m})^m \psi \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product of $L^2(\mathbf{R}^2; \mathbf{C}^p)$. We denote by $L_{(x,y);(x',y')}$ the straight line from (x, y) to (x', y') . The straight line $L_{(x+s,y);(x,y)}$ is parametrized by $\gamma(\tau) = (x + (1 - \tau)s, y)$, $\tau \in [0, 1]$. Then $\dot{\gamma}(\tau) = -s(1, 0)$. In terms of $\gamma(\tau)$ and $\dot{\gamma}(\tau)$, we can write

$$\begin{aligned} & ((e^{isp_1/m} e^{sA_1/m})^m \psi)(x, y) \\ &= \exp\left(-A_1\left(\gamma\left(\frac{m-1}{m}\right)\right)\dot{\gamma}_1\left(\frac{m-1}{m}\right)\frac{1}{m}\right) \cdots \exp\left(-A_1\left(\gamma\left(\frac{1}{m}\right)\right)\dot{\gamma}_1\left(\frac{1}{m}\right)\frac{1}{m}\right) \\ & \quad \times \exp\left(-A_1(\gamma(0))\dot{\gamma}_1(0)\frac{1}{m}\right) \psi(x+s, y), \end{aligned}$$

which converges to $W_A(L_{(x+s,y);(x,y)})\psi(x+s, y)$ a.e. $(x, y) \in M_{s,t}$ as $m \rightarrow \infty$. Moreover, by the fact that $e^{-A_1(\gamma(\tau))\dot{\gamma}_1(\tau)/m} \in U(p)$, we have

$$\|\phi(x, y)(e^{isp_1/m} e^{sA_1/m})^m \psi(x, y)\|_{\mathbf{C}^p} \leq \|\phi(x, y)\|_{\mathbf{C}^p} \|\psi(x+s, y)\|_{\mathbf{C}^p}$$

and

$$\int_{\mathbf{R}^2} \|\phi(x, y)\|_{\mathbf{C}^p} \|\psi(x+s, y)\|_{\mathbf{C}^p} d\mathbf{r} \leq \left(\int_{\mathbf{R}^2} \|\phi(\mathbf{r})\|_{\mathbf{C}^p}^2 d\mathbf{r}\right)^{1/2} \left(\int_{\mathbf{R}^2} \|\psi(\mathbf{r})\|_{\mathbf{C}^p}^2 d\mathbf{r}\right)^{1/2} < \infty.$$

Hence, by the dominated convergence theorem, we obtain for all $\phi, \psi \in L^2(\mathbf{R}^2; \mathbf{C}^p)$,

$$\langle \phi, e^{is\bar{P}_1} \psi \rangle = \int_{\mathbf{R}^2} \langle \phi(x, y), W_A(L_{(x+s,y);(x,y)}) \psi(x+s, y) \rangle_{\mathbf{C}^p} d\mathbf{r},$$

which implies that

$$(e^{is\bar{P}_1} \psi)(x, y) = W_A(L_{(x+s,y);(x,y)}) \psi(x+s, y), \quad \text{a.e. } (x, y) \in \mathbf{R}^2.$$

Similarly, we can show that

$$(e^{it\bar{P}_2} \psi)(x, y) = W_A(L_{(x,y+t);(x,y)}) \psi(x, y+t), \quad \text{a.e. } (x, y) \in \mathbf{R}^2.$$

Combining these formulas, we obtain

$$e^{is\bar{P}_1}e^{it\bar{P}_2}=(W_{s,t}^{A,-})^{-1}e^{isp_1}e^{itp_2}, \quad e^{it\bar{P}_2}e^{is\bar{P}_1}=(W_{s,t}^{A,+})^{-1}e^{isp_1}e^{itp_2},$$

which, together with the fact $(W_{s,t}^{A,-})^{-1}W_{s,t}^{A,+}=(W_{s,t}^A)^{-1}=(W_{s,t}^A)^*$, imply (2.3). ■

As for the commutation relations of $\exp(isq_j)$ and $\exp(it\bar{P}_j)$ ($s, t \in \mathbf{R}$), we have the following.

Lemma 2.3: For all $s, t \in \mathbf{R}$,

$$e^{isq_j}e^{it\bar{P}_k}=e^{-ist\delta_{jk}}e^{it\bar{P}_k}e^{isq_j}, \quad j, k=1, 2.$$

Proof: Similar to the proof of Theorem 2.1. Note that $\{e^{itq_j}, e^{itp_j}|t \in \mathbf{R}, j=1, 2\}$ satisfies the Weyl relations with two degrees of freedom (e.g., Ref. 2, p. 81). ■

Theorem 2.2 and Lemma 2.3 imply the following.

Theorem 2.4: The set $\{e^{itq_j}, e^{it\bar{P}_j}|t \in \mathbf{R}, j=1, 2\}$ satisfies the Weyl relations with two degrees of freedom if and only if $W_{s,t}^A=I$ for all $s, t \in \mathbf{R}$, where I is the identity operator on $L^2(\mathbf{R}^2; \mathbf{C}^p)$.

As a corollary, we obtain the following.

Corollary 2.5: Suppose that A is flat on M . Then $\{\bar{P}_j, q_j\}_{j=1}^2$ is a Schrödinger 2-system if and only if $W_{s,t}^A=I$ for all $s, t \in \mathbf{R}$.

Proof: We need only apply the well-known fact that a representation of the CCR with n degrees of freedom, satisfying the Weyl relations with the same degrees of freedom, is a Schrödinger n -system (von Neumann's theorem,⁹ Ref. 2, p. 81, Theorem 4.11.1). ■

III. THE WILSON LOOPS $W_{s,t}^A$

In view of Corollary 2.5 we derive a condition equivalent to the condition that $W_{s,t}^A=I$. Let

$$\delta_0 = \min_{n \neq m; n, m=1, \dots, N} |\mathbf{a}_n - \mathbf{a}_m|. \tag{3.1}$$

For a positive constant $\epsilon < \delta_0$ and $\mathbf{r} \in M$ with $|\mathbf{r} - \mathbf{a}_n| = \epsilon$, we denote by $C_\epsilon^r(\mathbf{a}_n)$ the circle of radius ϵ with center \mathbf{a}_n and initial point \mathbf{r} , where the direction of the circle is taken to be anticlockwise. In this section we prove the following theorem.

Theorem 3.1: The equality $W_{s,t}^A=I$ holds for all $s, t \in \mathbf{R}$ if and only if A is flat on M , and there exists a constant $\delta \in (0, \delta_0)$, such that for all $\epsilon < \delta$ and some $\mathbf{r}_n \in M$ with $|\mathbf{r}_n - \mathbf{a}_n| = \epsilon$,

$$W_A(C_\epsilon^{\mathbf{r}_n}(\mathbf{a}_n))=I, \quad n=1, \dots, N. \tag{3.2}$$

Remark: One can easily show that, if $W_A(C_\epsilon^{\mathbf{r}_n}(\mathbf{a}_n)) = I$ for some \mathbf{r}_n with $|\mathbf{r}_n - \mathbf{a}_n| = \epsilon$, then $W_A(C_\epsilon^{\mathbf{r}}(\mathbf{a}_n)) = I$ for all \mathbf{r} , with $|\mathbf{r} - \mathbf{a}_n| = \epsilon$.

We denote by $D_{x,y,s,t}$ the interior domain of the closed path $C_{x,y,s,t}$.

Lemma 3.2: For all $(x, y) \in M$,

$$W_{t,t}^A(x, y) = I - \int_{D_{x,y,t,t}} F_{12}(\mathbf{r}') d\mathbf{r}' + O(|t|^3), \tag{3.3}$$

as $t \rightarrow 0$.

Proof: Informal proofs of this lemma can be found in the physics literature (e.g., Ref. 10, pp. 52–53). For the sake of completeness, we give a rigorous proof of it. Let $\gamma_{x,y,s,t}: [0, 1] \rightarrow M$ be the parametrization of the path $C_{x,y,s,t}$, such that $\gamma_{x,y,s,t}(\tau) = \gamma_{x,y,s,t}^-(2\tau)$ for $0 \leq \tau \leq \frac{1}{2}$ and $\gamma_{x,y,s,t}(\tau) = \gamma_{x,y,s,t}^+(1 - 2(\tau - \frac{1}{2}))$ for $\frac{1}{2} \leq \tau \leq 1$, where $\gamma_{x,y,s,t}^\pm(\tau)$ are given by (2.1) and (2.2). Let

$$B_{x,y,s,t}(\tau) = A_1(\gamma_{x,y,s,t}(\tau))(\dot{\gamma}_{x,y,s,t})_1(\tau) + A_2(\gamma_{x,y,s,t}(\tau))(\dot{\gamma}_{x,y,s,t})_2(\tau), \quad 0 \leq \tau \leq 1,$$

and, for $k \geq 1$, define

$$J_k(x, y; s, t) = \int_0^1 \int_0^{\tau_1} \cdots \int_0^{\tau_{k-1}} B_{x,y;s,t}(\tau_1) B_{x,y;s,t}(\tau_2) \cdots B_{x,y;s,t}(\tau_k) d\tau_k \cdots d\tau_2 d\tau_1.$$

Then, applying Theorem 4.3 in Sec. 1.4 of Ref. 7 (p. 31), we have for all $s, t \in \mathbf{R}$,

$$W_{s,t}^A(x, y) = I + \sum_{k=1}^{\infty} (-1)^k J_k(x, y; s, t), \quad (x, y) \in M_{s,t}. \tag{3.4}$$

There is a positive constant $t_0 < 1$, such that for all $|t| \in (0, t_0)$, $\mathbf{a}_n \notin D_{x,y;t,t}$, $n = 1, \dots, N$. Let $0 < |t| < t_0$ and $a = \sup\{\|A_j(\mathbf{r}')\|; \mathbf{r}' \in D_{x,y;t_0,t_0} \cup D_{x,y;-t_0,-t_0}, j = 1, 2\}$. Then we have $\|B_{x,y;t,t}(\tau)\| \leq 8a|t|$, which implies that $\|J_k(x, y; t, t)\| \leq (8a|t|)^k/k!$, $k \geq 1$. Hence, we have

$$\sum_{k=3}^{\infty} \|J_k(x, y; t, t)\| \leq Ct^3, \tag{3.5}$$

where $C = \sum_{k=3}^{\infty} (8a)^k/k! < \infty$. By Stokes' theorem, we have

$$J_1(x, y; t, t) = \int_{D_{x,y;t,t}} (\partial_1 A_2(\mathbf{r}') - \partial_2 A_1(\mathbf{r}')) d\mathbf{r}'. \tag{3.6}$$

We can write

$$J_2(x, y; t, t) = \frac{1}{2} (J_+ + J_-),$$

where

$$J_{\pm} = \int_0^1 \int_0^{\tau_1} (B_{x,y;t,t}(\tau_1) B_{x,y;t,t}(\tau_2) \pm B_{x,y;t,t}(\tau_2) B_{x,y;t,t}(\tau_1)) d\tau_2 d\tau_1.$$

By symmetry, we have $J_+ = J_1(x, y; t, t)^2/2$. By (3.6), we have $\|J_1(x, y; t, t)\| \leq bt^2$, $0 < |t| < t_0$, where $b = \sup\{\|\partial_1 A_2(\mathbf{r}') - \partial_2 A_1(\mathbf{r}')\|; \mathbf{r}' \in D_{x,y;t_0,t_0} \cup D_{x,y;-t_0,-t_0}\}$. Hence,

$$\|J_+\| \leq \frac{b^2}{2} t^4. \tag{3.7}$$

To estimate J_- , we note that

$$B_{x,y;t,t}(\tau) = A_1(x, y)(\dot{\gamma}_{x,y;t,t})_1(\tau) + A_2(x, y)(\dot{\gamma}_{x,y;t,t})_2(\tau) + O(t^2)$$

as $t \rightarrow 0$ uniformly in $\tau \in [0, 1]$. Hence,

$$\begin{aligned} & B_{x,y;t,t}(\tau_1) B_{x,y;t,t}(\tau_2) - B_{x,y;t,t}(\tau_2) B_{x,y;t,t}(\tau_1) \\ &= \sum_{\mu, \nu=1}^2 [A_{\mu}(x, y), A_{\nu}(x, y)] (\dot{\gamma}_{x,y;t,t})_{\mu}(\tau_1) (\dot{\gamma}_{x,y;t,t})_{\nu}(\tau_2) + O(|t|^3). \end{aligned}$$

It follows that

$$\begin{aligned}
 J_- &= -[A_1(x,y), A_2(x,y)] \int_{C_{x,y;s,t}} (x' dy' - y' dx') + O(|t|^3) \\
 &= -2 \int_{D_{x,y;s,t}} [A_1(\mathbf{r}'), A_2(\mathbf{r}')] d\mathbf{r}' + O(|t|^3).
 \end{aligned}
 \tag{3.8}$$

Substituting (3.5)–(3.8) into (3.4), we obtain (3.3). ■

The following lemma is well known.¹¹

Lemma 3.3: Suppose that A is flat on M and each component A_j is m times continuously differentiable on M . Then, for every simply connected domain D of M , there exists a $U(p)$ -valued, $m + 1$ times continuously differentiable function g on D , such that $A_j = g^{-1} \partial_j g$ on D .

Lemma 3.4: Suppose that A is flat on M . Then the following (i)–(iii) hold.

(i) Let C be any continuous, piecewise continuously differentiable closed path in M , which is contractible to a point within M . Then

$$W_A(C) = I. \tag{3.9}$$

(ii) Let $C_\epsilon^{\mathbf{r}_1}(\mathbf{a}_n) \subset D_{x,y;s,t}$, with $|\mathbf{r}_1 - \mathbf{a}_n| = \epsilon < \delta_0$ and $D_{x,y;s,t} \cap \{\mathbf{a}_1, \dots, \mathbf{a}_N\} = \{\mathbf{a}_n\}$. Then there exists a unitary matrix U , such that

$$W_{s,t}^A(x,y) = U W_A(C_\epsilon^{\mathbf{r}_1}(\mathbf{a}_n)) U^{-1}. \tag{3.10}$$

(iii) Let $0 < \epsilon_1 < \epsilon_2 < \delta_0$ and $\mathbf{r}_j \in \mathbf{R}^2$, $j = 1, 2$, be, such that $|\mathbf{r}_j - \mathbf{a}_n| = \epsilon_j$. Then

$$W_A(C_{\epsilon_2}^{\mathbf{r}_2}(\mathbf{a}_n)) = W_A(L_{\mathbf{r}_2;\mathbf{r}_1})^{-1} W_A(C_{\epsilon_1}^{\mathbf{r}_1}(\mathbf{a}_n)) W_A(L_{\mathbf{r}_2;\mathbf{r}_1}), \tag{3.11}$$

where $L_{\mathbf{r}_2;\mathbf{r}_1}$ denotes the straight line from \mathbf{r}_2 to \mathbf{r}_1 .

Proof: (i) The path C is included in a simply connected domain D of M . By Lemma 3.3, there exists a $U(p)$ -valued twice differentiable function g on D , such that $A_j = g^{-1} \partial_j g$ on D ($j = 1, 2$). In terms of $h := g^{-1}$, we can write $A_j = -(\partial_j h) h^{-1}$. Let $\gamma: [0, 1] \rightarrow M$ be a parametrization of C ($\gamma(0) = \gamma(1)$). Then we have

$$-\{A_1(\gamma(\tau)) \dot{\gamma}_1(\tau) + A_2(\gamma(\tau)) \dot{\gamma}_2(\tau)\} = \frac{dh(\gamma(\tau))}{d\tau} h(\gamma(\tau))^{-1}.$$

Hence, applying Theorem 3.1 on p. 20 in Ref. 7, we have $W_A(C) = h(\gamma(1))h(\gamma(0))^{-1} = I$. Thus we obtain (3.9).

(ii) We decompose $D_{x,y;s,t}$, as is shown in Fig. 1.

Let $V_j = W_A(C_j)$, $W_j = W_A(L_j)$, $j = 1, 2$. Then, by part (i), we have

$$W_2 V_1 W_1 W_{s,t}^{A,-}(x,y) = I, \quad W_1^{-1} V_2^{-1} W_2^{-1} (W_{s,t}^{A,+})^{-1} = I,$$

which imply $W_2 V_1 V_2^{-1} W_2^{-1} W_{s,t}^{A,+}(x,y) = I$. We have $V_2 V_1^{-1} = W_A(C_\epsilon^{\mathbf{r}_1}(\mathbf{a}_n))$. Thus, taking $U = W_2$, we obtain (3.10).

(iii) We need only repeat the proof of part (ii) with ϵ and $D_{x,y;s,t}$ replaced by ϵ_1 and $C_{\epsilon_2}^{\mathbf{r}_2}(\mathbf{a}_n)$, respectively. ■

Proof of Theorem 3.1: Suppose that $W_{s,t}^A = I$ holds for all $s, t \in \mathbf{R}$. Then Lemma 3.2 implies that $F_{12}(\mathbf{r}) = 0$ for all $\mathbf{r} \in M$. Hence, A is flat on M . Then, using Lemma 3.4 (i) and (ii), we obtain (3.2).

Conversely, suppose that A is flat on M and (3.2) holds. Then, by Lemma 3.4(i), we have $W_{s,t}^A(x,y) = I$ for all $C_{x,y;s,t}$ contractible to a point within M . Let $\mathbf{a}_n \in D_{x,y;s,t}$, but $\mathbf{a}_m \notin D_{x,y;s,t}$ for $m \neq n$. Then, by Lemma 3.4 (ii), we have




FIG. 1. The decomposition of $D_{x,y;s,t}$ for proof of Lemma 3.4 (ii).

$$W_{s,t}^A(x,y) = UW_A(C_\epsilon^{r_1}(\mathbf{a}_n))U^{-1} = UIU^{-1} = I.$$

Hence, in this case, $W_{s,t}^A(x,y) = I$. Finally, we consider the case where $D_{x,y;s,t}$ includes $\{\mathbf{a}_{i_1}, \dots, \mathbf{a}_{i_k}\}$, where $2 \leq k \leq N$ and $\{i_1, \dots, i_k\}$ is a subset of $\{1, \dots, N\}$. In this case we decompose $D_{x,y;s,t}$, as is shown in Fig. 2, where we set $\mathbf{b}_j = \mathbf{a}_{i_j}$, $j = 1, \dots, k$.

Then, by Lemma 3.4(ii), there exist $U_j \in U(p)$, $j = 1, \dots, k$, such that for $j = 1, \dots, k$,

$$U_j W_A(C_\epsilon^{r_j}(\mathbf{b}_j))U_j^{-1} = W_A(D_{j-1}^{-1} \circ F_j \circ D_j \circ C_j).$$




FIG. 2. The decomposition of $D_{x,y;s,t}$ for proof of Theorem 3.1.

By (3.2), each of the RHSs of these equalities turns out to be the identity. Then the resulting equalities give $W_{s,i}^A(x,y)=I$. ■

IV. ESSENTIAL SELF-ADJOINTNESS OF P_j AND NON-SCHRÖDINGER REPRESENTATIONS

For $j=1,2$, we set $S_j=\mathbf{R}\{a_{nj}\}_{n=1}^N$. Let

$$\mathcal{L}_1^m = C_0^m(\mathbf{R} \times S_2; \mathbf{C}^p), \quad \mathcal{L}_2^m = C_0^m(S_1 \times \mathbf{R}; \mathbf{C}^p).$$

For a subset V of $M_p(\mathbf{C})$ (the set of $p \times p$ complex matrices) and an open subset D of M , we denote by $C^m(D;V)$ the set of V -valued, m times continuously differentiable functions on D . We introduce a class of gauge potentials.

Definition 4.1: We say that a 1-form A on M is in the set \mathcal{A}_m if there exist $g_1 \in C^{m+1}(\mathbf{R} \times S_2; U(p))$ and $g_2 \in C^{m+1}(S_1 \times \mathbf{R}; U(p))$, such that

$$A_1 = g_1^{-1} \partial_1 g_1, \quad \text{on } \mathbf{R} \times S_2; \quad A_2 = g_2^{-1} \partial_2 g_2, \quad \text{on } S_1 \times \mathbf{R}.$$

Theorem 4.2: Assume that $A \in \mathcal{A}_{m-1}$ ($m \geq 1$). Then, each P_j is essentially self-adjoint on \mathcal{L}_j^m .

Proof: Let g_j be as in Definition 4.1. Then g_j is a bijective mapping from \mathcal{L}_j^m to itself; we have $P_j \psi = g_j^{-1} p_j g_j \psi$, $\psi \in \mathcal{L}_j^m$. Hence, we need only to show that p_j is essentially self-adjoint on \mathcal{L}_j^m . But this is a well-known fact (see the proof of Theorem 3.2 in Ref. 1). ■

Theorem 4.3: Suppose that $A_j \in C^m(M; M_p^{\text{ah}}(\mathbf{C}))(j=1,2)$ ($m \geq 1$) and $A = A_1 dx + A_2 dy$ is flat on M . Then $A \in \mathcal{A}_m$. In particular, P_j is essentially self-adjoint on \mathcal{L}_j^{m+1} .

Proof: Let b_1, \dots, b_k ($k \leq N$) be the numbers different from each other in the set $\{a_{n2}\}_{n=1}^N$, with $b_1 < b_2 < \dots < b_k$. Let $M_l = \{(x,y) \in \mathbf{R}^2 | b_{l-1} < y < b_l\}$, $l=1, \dots, k+1$, with $b_0 = -\infty$, $b_{k+1} = +\infty$. Then, each M_l is simply connected and $\mathbf{R} \times S_2 = \cup_{l=1}^{k+1} M_l$. By Lemma 3.3, there exists a function $h_l \in C^{m+1}(M_l; U(p))$, such that $A_j = h_l^{-1} \partial_j h_l$ ($j=1,2$) on M_l . Defining $g_1 \in C^{m+1}(\mathbf{R} \times S_2; U(p))$ by $g_1(\mathbf{r}) = h_l(\mathbf{r})$ if $\mathbf{r} \in M_l$, we have $A_j = g_1^{-1} \partial_j g_1$ on $\mathbf{R} \times S_2$. In particular, $A_1 = g_1^{-1} \partial_1 g_1$ on $\mathbf{R} \times S_2$. Similarly, we can show that there exists a function $g_2 \in C^{m+1}(S_1 \times \mathbf{R}; U(p))$, such that $A_2 = g_2^{-1} \partial_2 g_2$ on $S_1 \times \mathbf{R}$. ■

Corollary 2.5 and Theorems 3.1 and 4.3 yield the following result.

Theorem 4.4: Suppose that $A_j \in C^m(M; M_p^{\text{ah}}(\mathbf{C}))(j=1,2)$ ($m \geq 1$) and A is flat on M . Then, the representation $\{\bar{P}_j, q_j\}_{j=1}^2$ of CCR is a Schrödinger 2-system if and only if there exists a constant $\delta \in (0, \delta_0)$, such that for all $\epsilon < \delta$ and some $\mathbf{r}_n \in M$ with $|\mathbf{r}_n - \mathbf{a}_n| = \epsilon$, $W_A(C_\epsilon^{\mathbf{r}_n}(\mathbf{a}_n)) = I$, $n=1, \dots, N$.

This theorem also gives a characterization for $\{\bar{P}_j, q_j\}_{j=1}^2$ to be non-Schrödinger.

V. EXAMPLE

In this section we discuss a class of $M_p^{\text{ah}}(\mathbf{C})$ -valued, flat 1-forms on M . Let S_n and T_n be $p \times p$ Hermitian constant matrices, such that, for all $n \neq m$ ($n, m = 1, \dots, N$),

$$[S_n, S_m] = 0, \quad [T_n, T_m] = 0, \quad [S_n, T_m] = 0,$$

but S_n does not necessarily commute with T_n . Let ϕ_n be a real-valued, continuously differentiable function on \mathbf{R}^2 ($n=1, \dots, N$) (so that ϕ_n and $\partial_j \phi_n$ have no singularity at $\mathbf{r} = \mathbf{a}_n$, $n=1, \dots, N$). Then, for each $\mathbf{r} \in M$, $e^{\pm i S_n \phi_n(\mathbf{r})}$ are in $U(p)$ and the matrix

$$K_n(\mathbf{r}) := e^{-i S_n \phi_n(\mathbf{r})} T_n e^{i S_n \phi_n(\mathbf{r})}$$

is Hermitian. Hence

$$A_1(\mathbf{r}) := -i \sum_{n=1}^N \left\{ \frac{(y-a_{n2})}{|\mathbf{r}-\mathbf{a}_n|^2} K_n(\mathbf{r}) - S_n \partial_1 \phi_n(\mathbf{r}) \right\}$$

and

$$A_2(\mathbf{r}) := i \sum_{n=1}^N \left\{ \frac{(x-a_{n1})}{|\mathbf{r}-\mathbf{a}_n|^2} K_n(\mathbf{r}) + S_n \partial_2 \phi_n(\mathbf{r}) \right\}$$

are in $M_p^{\text{ah}}(\mathbf{C})$. It is easy to see that A_1 and A_2 satisfy the nonlinear partial differential equation,

$$\partial_1 A_2(\mathbf{r}) - \partial_2 A_1(\mathbf{r}) + [A_1(\mathbf{r}), A_2(\mathbf{r})] = 2\pi i \sum_{n=1}^N K_n(\mathbf{a}_n) \delta(\mathbf{r}-\mathbf{a}_n),$$

where $\delta(\mathbf{r})$ is the two-dimensional Dirac delta distribution. Hence, the 1-form $A = A_1 dx + A_2 dy$ is flat on M . For $p \geq 2$, this example is a non-Abelian generalization of examples in Ref. 1 and Ref. 3.

We want to compute the Wilson loop $W_A(C_\epsilon^{\mathbf{r}}(\mathbf{a}_n))$ in the present case. We first note that we can write A_j in the form

$$A_1(\mathbf{r}) = -\frac{i(y-a_{n2})}{|\mathbf{r}-\mathbf{a}_n|^2} K_n(\mathbf{a}_n) + F_n^{(1)}(\mathbf{r}), \quad A_2(\mathbf{r}) = \frac{i(x-a_{n1})}{|\mathbf{r}-\mathbf{a}_n|^2} K_n(\mathbf{a}_n) + F_n^{(2)}(\mathbf{r}),$$

where $F_n^{(j)}$ ($j=1,2$) is a function continuous in the domain

$$D_n = \{\mathbf{r} \in \mathbf{R}^2 \mid 0 < |\mathbf{r}-\mathbf{a}_n| < \delta\},$$

($\delta < \delta_0$) with $\sup_{\mathbf{r} \in D_n} \|F_n^{(j)}(\mathbf{r})\| < \infty$, $j=1,2$.

Lemma 5.1: Let $B = B_1 dx + B_2 dy$ be an $M_p(\mathbf{C})$ -valued, continuously differentiable 1-form on M . Suppose that there exist a constant matrix $S \in M_p(\mathbf{C})$ and $M_p(\mathbf{C})$ -valued continuous functions G_1, G_2 on D_n , such that $C_j := \sup_{\mathbf{r} \in D_n} \|G_j(\mathbf{r})\| < \infty$, $j=1,2$, and, for all $\mathbf{r} \in D_n$,

$$B_1(\mathbf{r}) = -\frac{i(y-a_{n2})}{|\mathbf{r}-\mathbf{a}_n|^2} S + G_1(\mathbf{r}), \quad B_2(\mathbf{r}) = \frac{i(x-a_{n1})}{|\mathbf{r}-\mathbf{a}_n|^2} S + G_2(\mathbf{r}).$$

Then

$$\lim_{\epsilon \downarrow 0} W_B(C_\epsilon^{\mathbf{r}}(\mathbf{a}_n)) = e^{-2\pi i S},$$

independently of the choice of the initial point \mathbf{r} , with $|\mathbf{r}-\mathbf{a}_n| = \epsilon$.

Proof: We parametrize the circle $C_\epsilon^{\mathbf{r}}(\mathbf{a}_n)$ by $\gamma(\theta) = \mathbf{a}_n + (\epsilon \cos(\theta + \theta_0), \epsilon \sin(\theta + \theta_0))$, $0 \leq \theta \leq 2\pi$, where $\mathbf{r} = \mathbf{a}_n + (\epsilon \cos \theta_0, \epsilon \sin \theta_0)$ ($\epsilon < \delta$). Then we have

$$B_1(\gamma(\theta)) \dot{\gamma}_1(\theta) + B_2(\gamma(\theta)) \dot{\gamma}_2(\theta) = iS + \epsilon F(\epsilon, \theta),$$

where $F(\epsilon, \theta) = G_2(\gamma(\theta)) \cos(\theta + \theta_0) - G_1(\gamma(\theta)) \sin(\theta + \theta_0)$. Hence, we have

$$W_B(C_\epsilon^{\mathbf{r}}(\mathbf{a}_n)) = \lim_{m \rightarrow \infty} e^{-2\pi[iS + \epsilon F(\epsilon, \theta_m)]/m} e^{-2\pi[iS + \epsilon F(\epsilon, \theta_{m-1})]/m} \dots e^{-2\pi[iS + \epsilon F(\epsilon, \theta_1)]/m},$$

where $\theta_j = 2\pi j/m$, $j = 1, \dots, m$. By the condition for G_j , $j = 1, 2$, we have $\|F(\epsilon, \theta)\| \leq C$, where $C = C_1 + C_2$. In general, we can show that, for all $M_j, N_j \in M_\rho(\mathbf{C})$, $j = 1, \dots, k$ ($k = 1, 2, \dots$),

$$\|e^{M_1+N_1} \dots e^{M_k+N_k} - e^{M_1} \dots e^{M_k}\| \leq \left(\sum_{j=1}^k \|N_j\| \right) \exp \left(2 \sum_{j=1}^k \|M_j\| e^{\sum_{j=1}^k \|N_j\|} \right).$$

Applying this estimate to $M_j = 2\pi i S/m$, $N_j = 2\pi \epsilon F(\epsilon, \theta_j)/m$, we obtain

$$\|e^{-2\pi[iS + \epsilon F(\epsilon, \theta_m)]/m} e^{-2\pi[iS + \epsilon F(\epsilon, \theta_{m-1})]/m} \dots e^{-2\pi[iS + \epsilon F(\epsilon, \theta_1)]/m} - e^{-2\pi i S}\| \leq C' \epsilon,$$

with $C' = 2\pi C \exp(4\pi\|S\| + 2\pi C\epsilon)$. Hence, $\|W_B(C_\epsilon^r(\mathbf{a}_n)) - e^{-2\pi i S}\| \leq C' \epsilon$, which implies the desired result. ■

Applying Lemma 5.1 to the present example, we obtain the following.

Lemma 5.2: For all $n = 1, \dots, N$,

$$\lim_{\epsilon \downarrow 0} W_A(C_\epsilon^r(\mathbf{a}_n)) = e^{-2\pi i K_n(\mathbf{a}_n)},$$

independently of the choice of the initial point \mathbf{r} with $|\mathbf{r} - \mathbf{a}_n| = \epsilon$.

Lemma 5.3: Let $0 < \epsilon_1 < \epsilon_2 < \delta_0$ and $\mathbf{r}_j \in \mathbf{R}^2$, $j = 1, 2$, be such that $|\mathbf{r}_j - \mathbf{a}_n| = \epsilon_j$ and $\mathbf{r}_1 - \mathbf{a}_n = \alpha(\mathbf{r}_2 - \mathbf{r}_1)$ with a constant $\alpha > 0$. Then

$$W_A(L_{\mathbf{r}_2; \mathbf{a}_n}) = \lim_{\epsilon_1 \downarrow 0} W_A(L_{\mathbf{r}_2; \mathbf{r}_1})$$

exists.

Proof: The straight line $L_{\mathbf{r}_2; \mathbf{a}_n}$ is parametrized by $l(\tau) = (1 - \tau)\mathbf{r}_2 + \tau\mathbf{a}_n$, $0 \leq \tau \leq 1$. There exists a number $\tau_1 \in (0, 1)$, such that $\mathbf{r}_1 = l(\tau_1)$. Then we have, for $\tau \in [0, 1]$,

$$A_1(l(\tau))\dot{l}_1(\tau) + A_2(l(\tau))\dot{l}_2(\tau) = f_n(\tau),$$

where $f_n(\tau) = (F_n^{(1)}(l(\tau)), F_n^{(2)}(l(\tau))) \cdot (\mathbf{a}_n - \mathbf{r}_2)$. It is easy to see that $C_n := \lim_{\tau \downarrow 1} f_n(\tau)$ exists. Hence f_n can be extended to a continuous function on $[0, 1]$ with $f_n(1) = C_n$. We have $W_A(L_{\mathbf{r}_2; \mathbf{r}_1}) = \prod_0^{\tau_1} e^{-f_n(\tau)d\tau}$. On the other hand, $\prod_0^t e^{-f_n(\tau)d\tau}$ is continuous in $t \in [0, 1]$. Thus, the desired result follows. ■

Lemma 5.4: Let $0 < \delta < \delta_0$ be fixed. Then, for all $\epsilon \in (0, \delta)$, $n = 1, \dots, N$, and all $\mathbf{r} \in M$ with $|\mathbf{r} - \mathbf{a}_n| = \epsilon$,

$$W_A(C_\epsilon^r(\mathbf{a}_n)) = W_A(L_{\mathbf{r}; \mathbf{a}_n})^{-1} e^{-2\pi i K_n(\mathbf{a}_n)} W_A(L_{\mathbf{r}; \mathbf{a}_n}).$$

Proof: By Lemmas 5.2 and 5.3, we can take the limit $\epsilon_1 \downarrow 0$ of the RHS of (3.11) to obtain

$$W_A(C_{\epsilon_2}^{r_2}(\mathbf{a}_n)) = W_A(L_{\mathbf{r}_2; \mathbf{a}_n})^{-1} e^{-2\pi i K_n(\mathbf{a}_n)} W_A(L_{\mathbf{r}_2; \mathbf{a}_n}).$$

Thus, the desired result follows. ■

By Lemma 5.4 and Theorem 4.4, we obtain the following theorem.

Theorem 5.5: In the present example, the representation $\{\bar{P}_j, q_j\}_{j=1}^2$ of CCR is a Schrödinger 2-system if and only if, for all $n = 1, \dots, N$, all the eigenvalues of T_n are integers.

Proof: We need only consider the condition that $e^{-2\pi i K_n(\mathbf{a}_n)} = I$ for all $n=1, \dots, N$, which is equivalent to the condition that $e^{2\pi i T_n} = I$ for all $n=1, \dots, N$ [note that $K_n(\mathbf{a}_n)$ is unitarily equivalent to T_n]. Since T_n is Hermitian, $e^{2\pi i T_n} = I$ if and only if all the eigenvalues of T_n are integers. ■

Theorem 5.5 implies the following: Let

$$\mathfrak{A} = \{A = A_1 dx + A_2 dy \mid \text{at least one } T_n \text{ has a non-integer eigenvalue}\}.$$

Then, for each $A \in \mathfrak{A}$, $\{\bar{P}_j, q_j\}_{j=1}^2$ is a non-Schrödinger representation of the CCR, with two degrees of freedom. Thus, we obtain a class of non-Schrödinger representations of CCR associated with $M_p^{\text{ab}}(\mathbb{C})$ -valued, flat 1-forms on M .

ACKNOWLEDGMENT

This work is supported by the Grant-In-Aid No. 06640188 for science research from the Ministry of Education, Japan.

- ¹A. Arai, "Momentum operators with gauge potentials, local quantization of magnetic flux, and representation of canonical commutation relations," *J. Math. Phys.* **33**, 3374–3378 (1992).
- ²C. R. Putnam, *Commutation Properties of Hilbert Space Operators* (Springer, Berlin, 1967).
- ³H. Reeh, "A remark concerning canonical commutation relations," *J. Math. Phys.* **29**, 1535–1536 (1988).
- ⁴Y. Aharonov and D. Bohm, "Significance of electromagnetic potentials in the quantum theory," *Phys. Rev.* **115**, 485–491 (1959).
- ⁵A. Arai, "Properties of the Dirac–Weyl operator with a strongly singular gauge potential," *J. Math. Phys.* **34**, 915–935 (1993).
- ⁶H. Kurose and H. Nakazato, "Geometric construction of *-representations of the Weyl algebra with degree 2," preprint, 1994.
- ⁷J. D. Dollard and C. N. Friedman, "Product integration," *Encyclopedia of Mathematics and Its Applications Vol. 10: Analysis* (Addison-Wesley, Reading, Massachusetts, 1979).
- ⁸In the notation used in physics, $W_A(\mathbb{C}) = P \exp(-\int_{\mathcal{C}} A)$.
- ⁹J. von Neumann, "Die Eindeutigkeit der Schrödingerschen Operatoren," *Math. Ann.* **104**, 570–578 (1931).
- ¹⁰M. Göckeler and T. Schücker, *Differential Geometry, Gauge Theories, and Gravity* (Cambridge U.P., Cambridge, 1987).
- ¹¹L. Schlesinger, "Weitere Beiträge zum Infinitesimalkalkül der Matrizen," *Math. Zeit.* **35**, 485–501 (1932).