



Title	Blow-up of viscous heat-conducting compressible flows
Author(s)	Cho, Yonggeun; Jin, Bum Ja
Citation	Journal of Mathematical Analysis and Applications, 320(2), 819-826 https://doi.org/10.1016/j.jmaa.2005.08.005
Issue Date	2006-08-15
Doc URL	https://hdl.handle.net/2115/14420
Type	journal article
File Information	JMathAnalAppl_v320p819.pdf



Blow-up of viscous heat-conducting compressible flows

Yonggeun Cho^{*}

Department of Mathematics, Hokkaido University, Sapporo, 060-0810, Japan

Bum Ja Jin

Department of Mathematics, Seoul National University, Seoul 151-747, Korea

Abstract

We show the blow-up of strong solution of viscous heat-conducting flow when the initial density is compactly supported. This is an extension of Z. Xin's result[5] to the case of positive heat conduction coefficient but we do not need any information for the time decay of total pressure nor the lower bound of the entropy. We control the lower bound of second moment by total energy and obtain the exact relationship between the size of support of initial density and the existence time. We also provide a sufficient condition for the blow-up in case that the initial density is positive but has a decay at infinity.

Key words: blow-up, smooth solution, viscous heat-conducting compressible flow
2000 MSC: 35Q30, 76N10

^{*} corresponding author

Email addresses: ygcho@math.sci.hokudai.ac.jp (Yonggeun Cho),
bumjajin@hanmail.net (Bum Ja Jin).

¹ The first author was supported by Japan Society for the Promotion of Science under JSPS Postdoctoral Fellowship For Foreign Researchers.

1 Introduction

In this paper, we consider the following equations for a compressible fluid in $\mathbb{R}^n \times \mathbb{R}_+$ ($n \geq 1$):

$$\rho_t + \operatorname{div}(\rho u) = 0, \quad (1)$$

$$(\rho u)_t + \operatorname{div}(\rho u \otimes u) + Lu + \nabla p = 0, \quad (2)$$

$$(\rho e)_t + \operatorname{div}(\rho e u) - \kappa \Delta \theta + p \operatorname{div} u = Q(\nabla u), \quad (3)$$

where

$$Lu = -\mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u, \quad \text{and} \quad Q(\nabla u) = \frac{\mu}{2} |\nabla u + \nabla^t u|^2 + \lambda (\operatorname{div} u)^2.$$

Here $\rho = \rho(x, t)$, $u = (u_1, \dots, u_n)$, θ , p and e denote the density, velocity, absolute temperature, pressure and specific internal energy per unit mass, respectively. $\nabla^t u$ is the transpose of ∇u . If we denote the total energy per unit mass E by $E = \frac{1}{2}|u|^2 + e$, then the energy equation (3) can be rewritten by

$$(\rho E)_t + \operatorname{div}(\rho E u) = -Lu \cdot u + Q(\nabla u). \quad (4)$$

The viscosity coefficient μ and λ are assumed to be constant satisfying $\mu \geq 0$, $\lambda + \frac{2}{n}\mu \geq 0$ from the physical point of view. We also denote by $\kappa \geq 0$ the coefficient of heat conduction.

If $\mu = \lambda = \kappa = 0$, then we call the equations as compressible Euler equations for gas. On the other hand, if $\mu > 0$ and $\lambda + \frac{2}{n}\mu \geq 0$, then we call the equations as compressible Navier-Stokes equations. In particular, we call the equations as heat-conducting compressible Navier-Stokes equations if $\mu > 0$, $\lambda + \frac{2}{n}\mu \geq 0$ and $\kappa > 0$. A polytropic gas is a gas satisfying the following state of equations:

$$p/\rho = R\theta, \quad e = c_\nu \theta \quad \text{and} \quad p/\rho = A \exp(S/c_\nu) \rho^{\gamma-1}, \quad (5)$$

where $R > 0$ is the gas constant, A a positive constant of absolute value, $\gamma > 1$ the ratio of specific heats, $c_\nu = \frac{R}{\gamma-1}$ the specific heat at constant volume and S the entropy.

The blow-up of smooth solutions of compressible Euler equations has been studied by several mathematicians. In 1985[4], T. C. Sideris showed that the life span T of the C^1 solution of the compressible Euler equations is finite when the initial data is constant outside a bounded set and the initial flow velocity is sufficiently large (super-sonic) in some region. In 1998[5], Z. Xin showed, in a different way from [4], the blow-up result for the compressible Euler equations, when the initial density and initial velocity have compact

supports. In the paper, he also showed the blow-up of smooth solution for the compressible Navier-Stokes equations for polytropic gas with zero heat conduction (that is, $\kappa = 0$), when the initial density has compact support. His theorem was derived independently of the size of data, but his point of view cannot be applied for $\kappa > 0$, since in his argument the estimation for the lower bound of entropy or time decay of total pressure is strongly necessary, which seems hard to be obtained for the case $\kappa > 0$.

As for the positive result, one may refer to [1]. In the paper [1], the authors showed the local existence of the unique strong solutions of the compressible Navier-Stokes equations (1)-(3) with $n = 3$, $\kappa \geq 0$ and nonnegative density. In particular, they obtained for $\kappa > 0$ that there exists a finite time $T_* > 0$ such that for some $3 < q \leq 6$

$$\begin{aligned} \rho &\in C([0, T_*]; H^1 \cap W^{1, q}), & \rho_t &\in C([0, T_*]; L^2 \cap L^q), \\ (u, e) &\in C([0, T_*]; H_0^1 \cap H^2), & (u_t, e_t) &\in L^2(0, T_*; H_0^1). \end{aligned} \quad (6)$$

In this paper, we extend the Xin's blow-up result to the heat-conducting compressible Navier-Stokes equations, that is, for the case $\kappa > 0$ in view of the regularity (6) and hence we provide a sufficient condition for the local result of [1].

Before stating our main theorem, we introduce some notations. We denote by $B_R = B_R(0)$ the ball in \mathbb{R}^n of radius R centered at the origin. We will use several physical quantities:

$$\begin{aligned} m(t) &= \int_{\mathbb{R}^n} \rho(x, t) dx && \text{(total mass),} \\ M(t) &= \int_{\mathbb{R}^n} \rho(x, t) |x|^2 dx && \text{(second moment),} \\ A(t) &= \int_{\mathbb{R}^n} \rho(x, t) u(x, t) \cdot x dx && \text{(radial component of momentum),} \\ \mathcal{E}(t) &= \int_{\mathbb{R}^n} \rho(x, t) E(x, t) dx && \text{(total energy)} \\ P(t) &= \int_{\mathbb{R}^n} p(x, t) dx && \text{(total pressure).} \end{aligned}$$

We always assume that $m(0), M(0), |A(0)|, \mathcal{E}(0) < \infty$ and $m(0) > 0, \mathcal{E}(0) > 0$.

For the proof of blow-up, we have only to prove the following theorem.

Theorem 1 *We assume $\mu > 0$, $\lambda + \frac{2}{n}\mu > 0$, $1 \leq n \leq 3$ and $\kappa \geq 0$. Let $\gamma > 1$ and $T > 0$. Suppose that (ρ, u, e) is a solution to the cauchy problem (1), (2) and (3) with initial data (ρ_0, u_0, e_0) such that for some $q > \max(2, n)$*

$$\begin{aligned} \rho &\in C([0, T]; H^1 \cap W^{1, q}), & \rho_t &\in C([0, T]; L^2 \cap L^q), \\ (u, e) &\in C([0, T]; H_0^1 \cap H^2), & (u_t, e_t) &\in L^2. \end{aligned} \quad (7)$$

Furthermore, assume that the initial density ρ_0 is compactly supported in a ball B_{R_0} . Then we have

$$R_0^2 \geq \frac{M(0)}{m(0)} + 2\frac{A(0)}{m(0)}T + \min(2, n(\gamma - 1))\frac{\mathcal{E}(0)}{m(0)}T^2. \quad (8)$$

The restriction of dimension can be removed by an appropriate choice of Sobolev spaces guaranteeing the continuity of the solution. For example, we can take $C^1([0, T]; H^k)$ for $k > 2 + [\frac{n}{2}]$ as in [5].

Let T^* be the life span of the solution (ρ, u, e) . Then since $m(0)$ and $\mathcal{E}(0)$ are strictly positive, the theorem above implies that T^* should be finite for $\gamma > 1$. It also shows the exact relationship between the size of support and the life span. For example, the range of life span can be extended as the initial support of density become larger. Hence, from the relation, one can expect the global existence of smooth solution of compressible Navier-Stokes equations in case that the initial density is positive but has a decay at infinity in the sense of $M(0) < \infty$. However even in this case, we show that there is no global solution with u having a little bit fast decay as time goes on as follows:

Theorem 2 *Suppose that (ρ, u, e) is the solution of (1), (2), (3) satisfying (7), and initial density ρ_0 is not compactly supported but its second momentum is finite ($M(0) < \infty$). Then there is no global solution of regularity (7) with $T = \infty$ such that*

$$\limsup_{t \rightarrow \infty} \left\| \frac{t}{1 + |x|^2} u(x, t) \cdot x \right\|_{L^\infty} < 1. \quad (9)$$

In view of the parabolic scaling $\alpha u(\alpha x, \alpha^2 t)$, it is expected for the global solution with the density away from zero that

$$\limsup_{t \rightarrow \infty} \left\| \frac{(1+t)}{1+|x|} u(x, t) \right\|_{L^\infty} \leq c,$$

where the constant c can be chosen to be strictly smaller than 1 under a smallness assumption for initial data, rewriting the equation (1)-(3) with $(\tilde{u}, \tilde{e}) = (\frac{1+t}{1+|x|}u, \frac{1+t}{1+|x|}e)$ and using the usual energy estimate for (\tilde{u}, \tilde{e}) and elliptic regularity as in [3]. However Theorem 2 shows that even though the bound (9) seems to be reasonable for a density away from zero, the global existence satisfying (9) is impossible for the initial density having a decay at infinity in the sense of $M(0) < \infty$, no matter how small the data is.

Our proof is based on more elementary argument like integration by parts, energy estimate and Gronwall's inequality than in [5]. The key idea is to control the lower bound of the second moment of solution by the evolution of

total energy \mathcal{E} via the total radial component of momentum A . The control of second moment by total energy enables us not to rely on the lower bound of entropy nor on the time decay of the total pressure P . The argument can easily give another proof for the compressible Euler equations and also for the Korteweg type compressible fluid of non-isothermal case if the initial density is compactly supported (see [2] for the later). We leave the details of proof for the later two cases to the readers.

2 Proof of Theorem 1

Since we consider the case of compactly supported initial density, we can assume that there is a positive constant R_0 so that $\text{supp}\rho_0 \subset B_{R_0}$. We let (ρ, u, e) be a solution to the Cauchy problem (1), (2) and (3) satisfying the regularity (7). We denote by $X(\alpha, t)$ the particle trajectory starting at α when $t = 0$, that is,

$$\frac{d}{dt}X(\alpha, t) = u(X(\alpha, t), t) \quad \text{and} \quad X(\alpha, 0) = \alpha.$$

Since by Sobolev embedding, $u \in C(\mathbb{R}^n \times [0, T])$ for $1 \leq n \leq 3$, X is unique and differentiable. We set $\Omega(0) = \text{supp}\rho_0$ and

$$\Omega(t) = \{x = X(\alpha, t) | \alpha \in \Omega(0)\}.$$

From the transport equation (1), one can easily show that $\text{supp}\rho(x, t) = \Omega(t)$ and hence from the equation of state (5) that

$$p(x, t) = \theta(x, t) = 0 \quad \text{if} \quad x \in \Omega(t)^c.$$

Therefore, from the equation (2) and (3), we observe that

$$Lu = 0 \quad \text{and} \quad Q(\nabla u) = 0 \quad \text{a.e. in} \quad \Omega(t)^c.$$

The following lemma is a revisit of Z. Xin's in [5].

Lemma 3 *We assume $\mu > 0$, $\lambda + \frac{2}{n}\mu > 0$ and $\kappa \geq 0$. Suppose that (ρ, u, e) satisfying the regularity (7), is the solution of (1), (2) and (3). Then*

$$u(x, t) \equiv 0 \quad \text{in} \quad x \in B_{R(t)}^c$$

for some $B_{R(t)}$ containing $\Omega(t)$. Moreover, we can take $R(t) = R_0$ for all $0 < t < T$.

Proof. We observe that

$$Q(\nabla u) = 2\mu \sum_{i=1}^n (\partial_i u_i)^2 + \lambda (\operatorname{div} u)^2 + \mu \sum_{i \neq j}^n (\partial_i u_j)^2 + 2\mu \sum_{i > j} (\partial_i u_j)(\partial_j u_i).$$

Assume $\lambda \leq 0$. Then

$$\begin{aligned} Q(\nabla u) &\geq (2\mu + n\lambda) \sum_{i=1}^n (\partial_i u_i)^2 + \mu \sum_{i \neq j}^n (\partial_i u_j)^2 + 2\mu \sum_{i > j} (\partial_i u_j)(\partial_j u_i) \\ &= (2\mu + n\lambda) \sum_{i=1}^n (\partial_i u_i)^2 + \mu \sum_{i > j}^n (\partial_i u_j + \partial_j u_i)^2. \end{aligned}$$

Assume $\lambda > 0$. Then

$$\begin{aligned} Q(\nabla u) &\geq 2\mu \sum_{i=1}^n (\partial_i u_i)^2 + \mu \sum_{i \neq j}^n (\partial_i u_j)^2 + 2\mu \sum_{i > j} (\partial_i u_j)(\partial_j u_i) \\ &= 2\mu \sum_{i=1}^n (\partial_i u_i)^2 + \mu \sum_{i > j}^n (\partial_i u_j + \partial_j u_i)^2. \end{aligned}$$

Therefore, both of the cases imply that

$$\left. \begin{aligned} \partial_i u_i(x, t) &= 0 \\ \partial_i u_j(x, t) + \partial_j u_i(x, t) &= 0 \end{aligned} \right\} \text{ a.e. in } B_{R(t)}^c$$

for all $i, j = 1, \dots, n$. This again implies $\nabla^2 u(x, t) = 0$ a.e. in $B_{R(t)}^c$. Thus ∇u is constant except for a measure zero set. But since $u \in H_0^1$ and continuous in \mathbb{R}^n (this comes from the regularity (7)), we conclude that $u \equiv 0$ in $B_{R(t)}^c$. That is, $u(X(\alpha, t), t) = 0$ if $\alpha \in B_{R_0}^c$. Thus we observe that

$$X(\alpha, t) = \alpha + \int_0^t u(X(\alpha, s), s) ds = \alpha, \quad \text{if } \alpha \in B_{R_0}^c.$$

This implies that we can choose $R(t) = R_0$ for $0 \leq t \leq T$. \square

From now on, we assume that $\Omega(t) = \operatorname{supp} \rho(\cdot, t)$ is contained in a ball $B_{R(t)}$.

Multiplying $|x|^2$ to (1) and integrating it over \mathbb{R}^n , we get the identity

$$\frac{d}{dt} M(t) = 2A(t). \tag{10}$$

If we take inner product by x to (2) and integrate it over \mathbb{R}^n , then we also get the identity

$$\frac{d}{dt} A(t) = \int_{\mathbb{R}^n} \rho |u|^2 dx + nP(t). \tag{11}$$

Integrating (1) and (4) over \mathbb{R}^n , we finally get the identity

$$\frac{d}{dt}m(t) = 0, \quad \frac{d}{dt}\mathcal{E}(t) = 0. \quad (12)$$

The integration by parts applied for deriving the above identities can be justified by the regularity (7).

Integrating (10), (11) and (12) over $[0, t]$, respectively, we obtain the following identities:

$$M(t) = M(0) + 2 \int_0^t A(s) ds, \quad (13)$$

$$A(s) = A(0) + \int_0^s \int_{\mathbb{R}^n} \rho |u|^2(x, \tau) dx d\tau + n \int_0^s P(\tau) d\tau, \quad (14)$$

$$m(t) = m(0), \quad \mathcal{E}(t) = \mathcal{E}(0). \quad (15)$$

Using the definition of E , we have from (14) and (15)

$$A(s) = A(0) + 2 \int_0^s \mathcal{E}(0) d\tau + \left(n - \frac{2}{\gamma - 1} \right) \int_0^s P(\tau) d\tau. \quad (16)$$

Now we first assume $(n - \frac{2}{\gamma - 1}) \geq 0$. Then by (16) we obtain

$$A(s) \geq A(0) + 2\mathcal{E}(0)s. \quad (17)$$

Substituting (17) into (13), we get

$$M(t) \geq M(0) + 2A(0)t + 2\mathcal{E}(0)t^2. \quad (18)$$

Secondly, we consider the case $\gamma \in (1, 1 + \frac{2}{n})$. By the equation of state $p = (\gamma - 1)\rho e$ and the identity (14), we have

$$A(s) = A(0) + 2\mathcal{E}(0)s - (2 - n(\gamma - 1)) \int_0^s \int \rho e dx d\tau. \quad (19)$$

It follows from (15) and the definition of E that $\int \rho e dx \leq \mathcal{E}(0)$. Substituting this into (19), we have

$$A(s) \geq A(0) + n(\gamma - 1)\mathcal{E}(0)s. \quad (20)$$

and hence from (13) and (20), we have

$$M(t) \geq M(0) + 2A(0)t + n(\gamma - 1)\mathcal{E}(0)t^2. \quad (21)$$

On the other hand, since $\Omega(t) \subset B_{R(t)}$, from the mass conservation (15) we can estimate the upper bound of the second moment as follows:

$$\begin{aligned} M(t) &= \int_{\Omega(t)} \rho(x, t) |x|^2 dx = \int_{|x| \leq R(t)} \rho(x, t) |x|^2 dx \\ &\leq (R(t))^2 m(t) = (R(t))^2 m(0). \end{aligned} \quad (22)$$

Thus from (18) and (22), we conclude that

$$m(0)R(t)^2 \geq M(0) + 2A(0)t + 2\mathcal{E}(0)t^2$$

for $\gamma \geq 1 + \frac{2}{n}$, and from (21) and (22) that

$$m(0)R(t)^2 \geq M(0) + 2A(0)t + n(\gamma - 1)\mathcal{E}(0)t^2$$

for $1 < \gamma < 1 + \frac{2}{n}$.

Since the solution has strong regularity (7) in the time interval $[0, T]$, noting from Lemma 3 that $R(t) = R_0$ for $t \in [0, T]$, we get the inequality (8).

3 Proof of Theorem 2

Suppose that there is a global solution (ρ, u, e) satisfying (9). Then there exist constants $t_0 > 0$ and $c < 1$ such that for all $t \geq t_0$,

$$\left\| \frac{u(x, t) \cdot x}{1 + |x|^2} \right\|_{L^\infty} \leq \frac{c}{t}. \quad (23)$$

Let $\widetilde{M}(t) = \int \rho(1 + |x|^2) dx$. Then it follows from (10) and (23) that

$$\frac{d}{dt} \widetilde{M}(t) \leq 2\widetilde{M}(t) \left\| \frac{u(x, t) \cdot x}{1 + |x|^2} \right\|_{L^\infty} \leq 2c \frac{\widetilde{M}(t)}{t}$$

for all $t \geq t_0$. Integrating this over $[t_0, t]$, we have

$$\widetilde{M}(t) \leq \widetilde{M}(t_0) + 2c \int_{t_0}^t \frac{\widetilde{M}(s)}{s} ds.$$

By Gronwall's inequality, we finally have

$$\widetilde{M}(t) \leq \widetilde{M}(t_0) \exp(2c \log(t/t_0)) = \frac{\widetilde{M}(t_0)}{t_0^{2c}} t^{2c} = \frac{m(0) + M(t_0)}{t_0^{2c}} t^{2c}. \quad (24)$$

From (18), (20) and (24), it follows that

$$M(0) + 2A(0)t + n(\gamma - 1)\mathcal{E}(0)t^2 \leq \frac{m(0) + M(t_0)}{t_0^{2c}} t^{2c}$$

for all $t \geq t_0$. Thus the last inequality yields the contradiction to the hypothesis $c < 1$. This completes the proof of theorem.

References

- [1] Y. Cho, H. Kim, Existence results for viscous polytropic fluids with vacuum, to appear in J. Differential Equations, Hokkaido University preprint series in Mathematics #675.
- [2] R. Danchin, B. Desjardins, Existence of solutions for compressible fluid models of Korteweg type, Ann. Inst. Henri Poincaré Anal. nonlinear 18 (2001) 97-133.
- [3] A. Matsumura, T. Nishida, The initial value problem for the equations of motion of viscous and heat-conductive gases, J. Math. Kyoto Univ. 20 (1980) 67-104.
- [4] T. C. Sideris, Formation of singularity in three dimensional compressible fluids, Comm. Math. Phys. 101 (1985) 475-487.
- [5] Z. Xin, Blow up of smooth solutions to the compressible Navier-Stokes equations with compact density, Comm. Pure Appl. Math. 51 (1998) 229-240.