



HOKKAIDO UNIVERSITY

Title	First record of a Cretaceous cheilostome bryozoan from Hokkaido, Japan
Author(s)	Ostrovsky, Andrew N.; Takashima, Reishi; Dick, Matthew H. et al.
Citation	Cretaceous Research, 27(6), 859-862 https://doi.org/10.1016/j.cretres.2006.04.001
Issue Date	2006-12
Doc URL	https://hdl.handle.net/2115/17181
Type	journal article
File Information	CR27-6.pdf



**First record of a Cretaceous cheilostome bryozoan from Hokkaido,
Japan**

**Andrew N. Ostrovsky^a, Reishi Takashima^{b,*}, Matthew H. Dick^c,
Andrei V. Grischenko^c, Hiroshi Nishi^b, Shunshuke F. Mawatari^c**

^a*Department of Invertebrate Zoology, Faculty of Biology and Soil Science, St. Petersburg State University, Universitetskaja nab. 7/9, St. Petersburg 199034, Russia*

^b*Division of Earth and Planetary Sciences, Graduate School of Science, Hokkaido University, North 10 West 8, Kita-ku, Sapporo 060 0810, Japan*

^c*Division of Biological Sciences, Graduate School of Science, Hokkaido University, Sapporo 060 0810, Japan*

Received 20 September 2005; accepted in revised form 21 January 2006

Abstract

A small collection of recrystallised, encrusting colonies of a single species from the Mikasa Formation represents the first record of cheilostome (malacostegan or anascan) bryozoans of Lower–Middle Cenomanian age from Hokkaido, Japan.

Keywords: Cretaceous; Cenomanian; Hokkaido; Bryozoans

* Corresponding author.

E-mail address: rtaka@nature.sci.hokudai.ac.jp (R. Takashima).

1. Introduction

Cheilostome bryozoans are locally abundant in Upper Cretaceous deposits, having been described from numerous localities, in particular in Europe. They are also known from the USA, Argentina, South Africa, Madagascar, Saudi Arabia and Oman, Central Asia, India, and New Zealand (for details, see Voigt, 1972, 1981; Taylor and Badve, 1994; Horowitz, 1998). However, we are unaware of any records from China or, indeed, the whole of southeastern Asia. There is just one previous record of Cretaceous bryozoans from Japan, that of an anascan cheilostome of uncertain affinities, *Dysnoetocella? voighti* Nishizawa and Sakagami, 1997 of Campanian age (Izumi Group) from Shikoku.

Research on fossil bryozoans in Japan has focused mainly on Late Palaeozoic (Carboniferous-Permian) and Neogene (Miocene-Pleistocene) assemblages, as summarised in Sakakura (1935), Sakagami (1961, 1976), Hayami (1975), and Sakagami et al. (1980). With regard to Mesozoic occurrences, there are a number of papers on Triassic species (see e.g., Sakagami, 1972; Sakagami and Sakai, 1979). Of note is that Sakagami (1976, p. 30) observed that, ‘occurrences of the Jurassic and Cretaceous bryozoans are known [in Japan] without any paleontological studies as yet’.

In central Hokkaido, a continuous Aptian-Campanian marine succession is well exposed (Takashima et al., 2004); we searched for bryozoans in an outcrop near Mikasa (43°14'43'' N, 141°59'50'' E) (Fig. 1) that exposes strata assigned to the Tb Member of the Mikasa Formation (Fig. 2), middle Yezo Group (Matsuno et al., 1964;

Hirano et al., 1992). This member comprises hummocky, cross-stratified, fine-grained sandstones and bioturbated sandy siltstones with frequent intercalations of *Pterotrigonia* shell beds, and is considered to have formed in an inner- to outer-shelf setting (Hirano et al., 1992). Typically Early-Middle Cenomanian ammonites, such as *Mantelliceras japonicum* and *Calycoceras orientale*, occur within this member (Ando, 1991; Matsumoto et al., 1991; see Fig. 2).

Here we document the first record of a cheilostome bryozoan of Early–Middle Cenomanian age from this section in Hokkaido; the previous record of cheilostome colonies concerned material of Campanian age from Shikoku (Nishizawa and Sakagami, 1997). The efforts of four people working for three hours screening rubble of Mikasa Formation strata resulted in thirteen completely recrystallised colony fragments of a single bryozoan species, preserved in rocks containing abundant bivalve shells. All specimens are now housed in the Fossil Invertebrate Collection, Paleontology Department, Natural Sciences Museum, Hokkaido University (Sapporo), under registration numbers UHR32534A–M.

All of the colonies are found to encrust the convex outer surface of smooth valves of species of the trioniid bivalve genus *Pterotrigonia*. In some cases, only the lower side of the colony is visible, the bryozoan itself remaining on a concave cast left by the molluscan valve. The few colonies that show vertical walls are strongly abraded. The zooidal walls are preserved in white calcite, whereas the zooidal cavities are completely filled with cemented, dark greenish-grey sandy siltstone.

Colonies are encrusting, unilaminar, multiserial, more-or-less circular or oval, and even in outline (Fig. 3A). The largest colony fragment available measures 19 x 17 mm; zooids are arranged in a quincuncial pattern, hexagonal to oval in shape (Fig.

3B–D), widest in the centre, and 0.48–0.67 mm (0.56 ± 0.05 mm) in length, by 0.25–0.43 (0.32 ± 0.04 mm) in width (mean \pm SD; N = 30). The frontal zooidal surface has not been observed; it is worn away in some specimens and embedded in the sediment in others. Large distal basal pore chambers (Fig. 3D) are recognisable in a few instances, and smaller distolateral pore chambers are also present. Early growth stages are evident in two colonies (Fig. 3A). In the absence of any data on the frontal zooidal wall, the bryozoan species described may either represent a malacostegan (suborder Malacostegina, superfamily Membraniporoidea) or an anascan cheilostome (suborder Neocheilostomina, infraorder Flustrina). Poor preservation of the colonies does not allow a more precise identification, and additional sampling is necessary to clarify the taxonomic position of the species.

This new find of fossil bryozoans in Hokkaido substantiates Sakagami's (1976) view that there should be a more extensive record of Cretaceous cheilostomes in Japan, and also encourages future field work in the area in order to document more material and gain a better understanding of bryozoan biogeography and evolution.

Acknowledgements

For financial support, we are greatly indebted to the 21st Century Center of Excellence (COE) Program on 'Neo-Science of Natural History' (program leader, Dr H. Okada; co-ordinator, Dr P. Gautam) at Hokkaido University, funded by the Ministry of Education, Culture, Sports, Science and Technology of Japan. We thank Dr Paul D. Taylor (Natural History Museum, London), Dr Dennis P. Gordon (National Institute of

Water and Atmospheric Research, Wellington), Dr Frank K. McKinney (Appalachian State University, Boone), and an anonymous reviewer for useful discussion and criticism.

References

- Ando, H., 1991. Shallow-marine sedimentary facies distribution and progradational sequences of the Mikasa Formation, Middle Yezo Group (Upper Cretaceous). *Journal of the Geological Society of Japan* 96, 453-469 [In Japanese, English abstract].
- Hayami, T., 1975. Neogene Bryozoa from northern Japan. *Science Reports of the Tohoku University, Sendai, (2, Geology)* 45, 83-126.
- Hirano, H., Tanabe, K., Ando, H., Futakami, M., 1992. Cretaceous forearc basin of central Hokkaido: lithofacies and biofacies characteristics. In: Adachi, M., Suzuki, K. (Eds.), *Field Trip Guidebook, Paleozoic and Mesozoic Terranes: Basement of the Japanese Island Arcs*, 1 [29th International Geological Congress, Nagoya University], pp. 45-80.
- Horowitz, A., 1998. Checklist of Cretaceous Bryozoa.
<http://www.geology.iupui.edu/research/PaleoLab/Projects/Bryolists/CRET1AL.html> (also available through the International Bryozoology Association).
- Matsuno, H., Tanaka, K., Mizuno, A., Ishida, M., 1964. Geological sheet map, 'Iwamizawa', scale 1:50,000 and its explanatory text. Hokkaido Development Agency, Sapporo, 168 pp.
- Matsumoto, T., Noda, M., Maiya, S., 1991. Towards an integrated ammonoid-, inoceramid- and foraminiferal biostratigraphy of the Cenomanian and Turonian (Cretaceous) in Hokkaido. *Journal of Geography* 100, 378-398.
- Nishizawa, Y., Sakagami, S., 1997. Phylogenetic significance of a new cheilostome bryozoan

- species, *Dysnoetocella ? voigti* from the Upper Cretaceous Izumi Group in Shikoku, Japan. *Paleontological Research* 1, 267-273.
- Sakagami, S., 1961. Japanese Permian Bryozoa. *Special Papers of the Palaeontological Society of Japan* 7, 1-58.
- Sakagami, S., 1972. The Triassic Bryozoa from Kusaka, Sakawa Basin, Shikoku, Japan. *Transactions and Proceedings of the Palaeontological Society of Japan, new series* 85, 275-279.
- Sakagami, S., 1976. History of bryozoological research. *Transactions and Proceedings of the Palaeontological Society of Japan, new series* 100, 29-30.
- Sakagami, S., Arakawa, S., Hayami, T., 1980. Check list and bibliography of Japanese Cenozoic fossil Ectoprocta (Bryozoa), 1935-1978. In: Igo, H., Noda, H. (Eds.), *Professor Saburo Kanno Memorial volume, Ibaraki*, pp. 317-338.
- Sakagami, S., Sakai, A., 1979. Triassic bryozoans from the Hidaka Group in Hokkaido, Japan. *Transactions and Proceedings of the Palaeontological Society of Japan, new series* 114, 77-86.
- Sakakura, K., 1935. Pliocene and Pleistocene Bryozoa from the Boso Peninsula. *Journal of the Faculty of Science, Imperial University of Tokyo*, (2)4, 1-48.
- Takashima, R., Kawabe, F., Nishi, H., Moriya, K., Wani, R., Ando, H., 2004. Geology and stratigraphy of forearc basin sediments in Hokkaido, Japan: Cretaceous environmental events on the Northwest Pacific margin. *Cretaceous Research* 25, 365-390.
- Taylor, P.D., Badve, R.M. 1994. The mid-Cretaceous bryozoan fauna from the Bagh Beds of central India: composition and evolutionary significance. In: Hayward, P.J., Ryland, J.S., Taylor, P.D. (Eds.), *Biology and palaeobiology of bryozoans*. Olsen & Olsen, Fredensborg, pp. 33-36.
- Voigt, E., 1972. Les méthodes d'utilisation stratigraphique des bryozoaires du Crétacé supérieur. *Mémoires du Bureau de Recherches Géologiques et Minières de France* 77, 45-53.

Voigt, E., 1981. Répartition et utilisation stratigraphique des bryozoaires du Crétacé moyen (Aptien-Coniacien). *Cretaceous Research* 2, 439-462.

Captions

Fig. 1. Locality map of the study area, where the Mikasa Group of south-central Hokkaido is exposed.

Fig. 2. Composite column of the Cretaceous sequence in the section studied, showing lithology and ranges of age-diagnostic macrofossil taxa (ammonoids, inoceramid bivalves), modified from Ando (1991), Matsumoto et al. (1991), and Hirano et al. (1992). Abbreviation HCS: hummocky, cross-stratified.

Fig. 3. A, General view of specimen UHR32534H, the black arrow indicating the ancestrular zone; B-D, colony fragments (specimens UHR32534B, UHR32534C, and UHR32534A, respectively) shown at different magnifications; white arrows in D indicate distal basal pore chambers. Scale bars represent 1 mm (A–C) and 100 μm (D).




Figure 1; OSTROVSKY et al.




Figure 2; OSTROVSKY et al.




Figure 3; OSTROVSKY et al.