



HOKKAIDO UNIVERSITY

Title	STUDIES ON RESPONSES TO MORPHINE HYDROCHLORIDE IN DOGS AND SOME CLINICAL EVALUATIONS
Author(s)	MATSUHASHI, Akira
Citation	Japanese Journal of Veterinary Research, 13(4), 138-139
Issue Date	1965-12
Doc URL	https://hdl.handle.net/2115/1832
Type	departmental bulletin paper
File Information	KJ00003418288.pdf



In the second study gravid cows were found to show prominent sclerotic changes in the tunica intima of their arterinae uterinae mediae, i.e. prominent hyperplastic swelling of the tunica intima, hyperplastic smooth muscle fibers and interstitial ground substance. In the region of hyperplastic swelling of the tunica intima, one or more concentric layers of elastic membrane are clearly seen. This appearance is considered to represent gravidity sclerosis in the cow.

In many cases (35%) the number of layers of elastic membrane is exactly same as the parturition frequency. The appearance of sclerotic gestation layers ("sclerotic gestation rings", which TAKAHATA and others advocated) becomes more noticeable as parturition frequency increases.

STUDIES ON RESPONSES TO MORPHINE HYDROCHLORIDE IN DOGS AND SOME CLINICAL EVALUATIONS*

Akira MATSUHASHI

*Department of Veterinary Surgery
Faculty of Veterinary Medicine
Hokkaido University, Sapporo, Japan*

The experiment was conducted to establish criteria for the use of morphine in veterinary clinics. A one percent solution of morphine hydrochloride was administered to dogs, either intravenously (A_1) or subcutaneously (A_2) in doses of 1 mg (D_1), 10 mg (D_2) and 30 mg (D_3) per kg of body weight, and comparative biochemical studies and clinical examinations were carried out immediately, 24 hours following administration.

The results may be summarized as follows:

The difference between A_1 and A_2 was significant in the changes of eosinophil count, blood sugar, blood CO_2 , blood P_{CO_2} , blood HCO_3^- , blood H_2CO_3 and blood B_B^+ , and was insignificant in the changes of red cell count, white cell count, blood O_2 , blood pH and Ht. Comparison between A_1 and A_2 generally indicated that A_1 produced not only a more rapid effect, but smoother recovery.

Comparison between D_1 , D_2 and D_3 indicated that an increased dosage produced increased effect. Only slight changes were produced by D_1 , but considerable changes were observed by D_2 and D_3 . Changes of serum proteins and liver functions were slight, irrespective of the administered methods or doses.

In clinical symptoms, the considerable difference between A_1 and A_2 was the

* The original report of this work will appear in this Journal in the near future.

appearance of remarkable agitation, fall of blood pressure and respiratory depression in A₁ and the appearance of emesis in A₂. Observed clinical symptoms indicated that D₁ produced either no effect or only sedative effect while D₂ and D₃ produced surgical anesthesia in most cases, either A₁ or A₂.

It is concluded that morphine is unsuitable as general anesthetic for dogs because of its side effects, but it is felt that small doses of morphine (about 1 mg/kg) are very useful as pre-anesthetic, basal anesthetic or sedative, especially when administered intravenously.

STUDIES ON BOVINE ENTEROVIRUSES

Shinji YAMADA

*Chemo- Sero-Therapeutic Institute,
Kumamoto, Japan*

This article appeared in the Japanese Journal of Veterinary Science, **27**, 133, 171, 271 & 317 (1965) (in Japanese with English summary).

METHODOLOGISCH ORIENTIERTE QUANTITATIVE UNTERSUCHUNG DER SPERMATOGENESE BEIM BULLEN AN DREI FAELLEN VON ASTHENOSPERMIE, DIE MIT ASPERMIE EINHERGEHT

Tosiro TIBA

*Institut für Obstetrik
Tierärztliche Fakultät
Hokkaido-Universität, Sapporo, Japan*

This article appeared in this journal, Supplementum 2 (1965).