



HOKKAIDO UNIVERSITY

Title	A RADIOIMMUNOASSAY SYSTEM USING A SEROVAR-SPECIFIC LIPOPOLYSACCHARIDE ANTIGEN OF LEPTOSPIRA, AND STUDIES ON THE CHEMICAL NATURE OF THE ANTIGENIC SITES
Author(s)	KAWAOKA, Yoshihiro
Citation	Japanese Journal of Veterinary Research, 28(1-2), 45-45
Issue Date	1980-05-31
Doc URL	https://hdl.handle.net/2115/2190
Type	departmental bulletin paper
File Information	KJ00003407915.pdf



observed between the two lesions; however, there did appear to be some relation between their immunosuppressed condition and the incidence of MD lesions.

**A RADIOIMMUNOASSAY SYSTEM USING A SEROVAR-SPECIFIC
LIPOPOLYSACCHARIDE ANTIGEN OF *LEPTOSPIRA*,
AND STUDIES ON THE CHEMICAL NATURE OF
THE ANTIGENIC SITES**

Yoshihiro KAWAOKA

*Department of Hygiene and Microbiology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

A highly sensitive and serovar-specific radioimmunoassay (RIA) was established to implement studies on the chemical nature of leptospiral antigenic determinant. The serovar-specific lipopolysaccharide (TM) antigen extracted from *Leptospira interrogans* serovar *kremastos* strain Kyoto was labeled with tritium by a reduction using sodium boro [^3H]-hydride after 1 h oxidation with periodate for the RIA. When 50 ng (625 cpm) of the labeled compound was used in the RIA system, 50% inhibition of the labeled antigen-antibody binding was obtained by the addition of 9 ng of the homologous TM antigen, whereas 5,000 times as much as the TM antigen from *hebdomadis*, which belongs to the same serogroup, was required. The TM antigens from different serogroups, such as *icterohaemorrhagiae*, *copenhageni* and *pomona*, showed no inhibitions in amounts up to 3×10^3 , 2×10^5 , and 2×10^5 , respectively.

The mild acid hydrolysis of *kremastos* Kyoto TM antigen using 0.5 N sulfonic acid or 2 N formic acid produced an antigen-active fraction which was dialysable. The chemical, physicochemical, and immunochemical properties of the fraction obtained by formic acid hydrolysis were then characterized. This fraction was eluted in a gel filtration at a molecular weight which was calculated at approximately 10,000 dalton. As compared with the original TM compound containing 10.5% protein and 1.5% fatty acid, this fraction lacked almost all of these components and was composed mainly of carbohydrates. However, the antigenic potency of this fraction diminished to 1/350 in the RIA. The carbohydrate nature of the antigenic determinant was also discussed.