



# HOKKAIDO UNIVERSITY

Title	THE INHIBITORY EFFECTS OF OLIGONUCLEOTIDES, COMPLEMENTARY TO MAREK'S DISEASE VIRUS mRNA TRANSCRIBED FROM THE BamHI-H REGION, ON THE PROLIFERATION OF TRANSFORMED LYMPHOBLASTOID CELLS, MDCC-MSB1
Author(s)	KAWAMURA, Mikiko
Citation	Japanese Journal of Veterinary Research, 40(1), 37-37
Issue Date	1992-05-29
Doc URL	<a href="https://hdl.handle.net/2115/2370">https://hdl.handle.net/2115/2370</a>
Type	departmental bulletin paper
File Information	KJ00002377562.pdf



THE INHIBITORY EFFECTS OF OLIGONUCLEOTIDES,  
COMPLEMENTARY TO MAREK'S DISEASE VIRUS  
mRNA TRANSCRIBED FROM THE BamHI-H REGION,  
ON THE PROLIFERATION OF TRANSFORMED  
LYMPHOBLASTOID CELLS, MDCC-MSB1

Mikiko KAWAMURA

*Department of Laboratory Animal Science,  
Faculty of Veterinary Medicine,  
Hokkaido University, Sapporo 060, Japan*

An oligonucleotide complementary to the splice donor sequence of the 1.8-kb gene family produced from the BamHI-H region of Marek's disease virus (MDV) DNA inhibited the proliferation of the MDV-derived lymphoblastoid cell line, MDCC-MSB1 (MSB-1), but not that of the avian lymphoblastoid leukemia-derived lymphoblastoid cell line, LSCC-BK3. Colony formation in soft agar was also inhibited by treatment of MSB-1 cells with the antisense oligonucleotide.

When cellular RNA prepared from MSB-1 cells treated with antisense oligonucleotide was analyzed by Northern blot hybridization using an RNA probe synthesized from the BamHI-H region of MDV DNA, the 1.8-kb mRNA was not detected. Moreover, when cellular RNA prepared from MSB-1 cells treated with antisense oligonucleotide was analyzed by RT-PCR using two oligonucleotides complementary to sequences within the 1.8-kb mRNA as primers, the fragment could not be detected in the amplified product.

These results indicate that expression of the 1.8-kb gene family produced from the BamHI-H region is directly associated with the maintenance of the tumorigenic states of transformed Marek's disease-derived lymphoblastoid cells.