



# HOKKAIDO UNIVERSITY

Title	Effect of a Quaternary Ammonium Base on the Distribution Equilibrium of 8-Hydroxyquinoline
Author(s)	NORIKI, Shinichiro; 乗木, 新一郎
Citation	北海道大學水産學部研究彙報, 35(1), 35-42
Issue Date	1984-03
Doc URL	<a href="https://hdl.handle.net/2115/23846">https://hdl.handle.net/2115/23846</a>
Type	departmental bulletin paper
File Information	35(1)_P35-42.pdf



## Effect of a Quaternary Ammonium Base on the Distribution Equilibrium of 8-Hydroxyquinoline

Shinichiro NORIKI\*

### Abstract

Distribution equilibrium of 8-hydroxyquinoline (oxine, HA) between the aqueous and organic (1, 2-dichloroethane) phases was investigated in the presence of a quaternary ammonium base, so-called zephiramine ( $Z^+ \cdot Cl^-$ ). In acidic solution, zephiramine has no effect on the distribution of oxine. In alkaline solution, the formation of the ion-association complex of oxinate ion ( $A^-$ ) with zephiramine ion ( $Z^+$ ), AZ, was observed. And, the complex was distributed between the aqueous and 1, 2-dichloroethane phases. Constants obtained were  $pK_1=4.96$ ,  $pK_2=10.01$ ,  $\log D_{HA}=2.5$ ,  $\log K_{AZ}=3.0$ ,  $\log D_{AZ}=1.3$ ,  $\log K_{ZCl}=2.7$ , and  $\log D_{ZCl}=0.89$ , K and D being dissociation constant and distribution constant, respectively. The dissociation of oxine was apparently enhanced above pH 9.5. But, the enhancement of dissociation of ligand by a quaternary ammonium base is not reason for highly ligand-coordinated chelate formation.

### Introduction

The author has discussed the behavior of a valence-saturated but coordination-unsaturated chelate which can be extracted into an inactive solvent in the presence of a quaternary ammonium base.<sup>1-6)</sup> According to a series of their studies, the effects of a quaternary ammonium base on extraction systems have been grouped in the three categories: 1) Formation of an inactive solvent-soluble chelate by displacing the coordinated water molecule with the molecule of chelating agent.<sup>1)</sup> 2) Formation of a highly ligand-coordinated anionic chelate which results in the formation of an ion-associated complex with the quaternary ammonium base existing in the solution.<sup>1-4)</sup> 3) Formation of an inactive solvent-soluble chelate by displacing the water molecule with the quaternary ammonium base molecule itself.<sup>5)</sup>

For the detailed discussion on the second category, it is necessary to elucidate the equilibrium of species of a ligand in the presence of a quaternary ammonium base. In this paper, the equilibrium of 8-hydroxyquinoline (oxine) was investigated in the presence of tetradecyldimethylbenzylammonium chloride, so-called zephiramine ( $Z^+ \cdot Cl^-$ ).

The following symbols are used:

- [ ] ; concentration in the aqueous phase
- [ ]<sub>org</sub> ; concentration in the organic phase
- HA ; oxine molecule
- $H_2A^+$  ; oxinium ion
- $A^-$  ; oxinate ion

\* *Laboratory of Analytical Chemistry, Faculty of Fisheries, Hokkaido University*  
(北海道大学水産学部分析化学講座)

- $Z^+$  ; zephiramine ion  
 $AZ$  ; ion-association complex of oxinate ion and zephiramine ion  
 $K_1$  ; dissociation constant of  $H_2A^+$   
 $K_2$  ; dissociation constant of HA  
 $K_{AZ}$  ; dissociation constant of AZ  
 $D_{HA}$  ; distribution constant of HA  
 $D_{AZ}$  ; distribution constant of AZ

## Experimental

### Reagents

Oxine solution. (a)  $1.2 \times 10^{-3}$  M; dissolve 0.174 g of 8-hydroxyquinoline in 5 ml of hot acetic acid and diluted to 1 l with water. (b)  $3 \times 10^{-4}$  M; dissolve 0.0435 g of the reagent in 1 l of 1, 2-dichloroethane.

Zephiramine solution,  $2 \times 10^{-2}$  M. Dissolve 7.36 g of tetradecyldimethylbenzylammonium chloride in 1 l of water.

Buffer solution. The pH of a solution was adjusted with acetate buffer (pH < 6.2), borate buffer (6.2 < pH < 12) or sodium hydroxide solution (pH > 12).

### General procedure

#### Determination of dissociation constants

Each 5 ml of  $1.2 \times 10^{-3}$  M oxine solution, a buffer solution and the zephiramine solution was placed in a test tube and diluted to 20 ml with water. The absorption spectrum of the solution was measured against water as a reference. The dissociation constants,  $K_1$  and  $K_2$ , were determined from the change of absorption spectra caused by variation of pH.  $K_{AZ}$  was calculated from the change of absorption spectra with variation of zephiramine concentration.

#### Determination of distribution constants

Twenty ml of  $3 \times 10^{-4}$  M oxine-1, 2-dichloroethane solution and 20 ml of an aqueous solution containing buffer and zephiramine were shaken for a definite time in a separatory funnel. After each phase was filtered through a filter paper, the absorption spectra of the organic and the aqueous phases were measured against 1,2-dichloroethane and water as references, respectively. From the changes of the absorption spectra of the organic and the aqueous phases with variations of pH and zephiramine concentration,  $D_{HA}$  and  $D_{AZ}$  were determined.

## Results and Discussion

### Dissociation constants and distribution constants

Because of amphoteric nature of oxine,  $H_2A^+$  is formed in an acidic solution and  $A^-$  in an alkaline solution. The dissociation constants are expressed as follows:

$$K_1 = \frac{[HA][H^+]}{[H_2A^+]} \quad (1)$$

$$K_2 = \frac{[A^-][H^+]}{[HA]} \quad (2)$$

In acidic solution, the isobestic point at 335 nm indicated the equilibrium was between HA and  $H_2A^+$  (Table 1). The spectra did not change with the addition of

zephiramine whose concentrations were zero,  $1 \times 10^{-5}$  M,  $1 \times 10^{-4}$  M, and  $1 \times 10^{-3}$  M. Thus, the author concludes that zephiramine has no effect on the dissociation of oxine in acidic solution. As listed in Table 1, the increase of absorbance at 360 nm ( $H_2A$ ) and the decrease at 310 nm (HA) indicate the transition of oxine from HA to  $H_2A^+$ .  $K_1$  is determined by plotting  $\log ([HA]/[H_2A^+])$  against  $\log [H^+]$ . The results agree with the published data<sup>7-12)</sup> (Table 2).

When oxine was distributed between the aqueous phase and 1,2-dichloroethane, no changes in the absorption spectra were observed in either of the phases with the addition of zephiramine. Consequently, the author concludes that zephiramine has no effect on the distribution of HA.  $D_{HA}$  is expressed by

$$D_{HA} = \frac{[HA]_{org}}{[HA]} \quad (3)$$

Introduction of equation (1) into equation (3) gives:

Table 1. Spectrophotometric data on 8-hydroxyquinoline\* for the determination of  $K_1$  and  $K_2$ .

pH	Absorbance			
	310 nm	321 nm	335 nm	360 nm
1.72	0.459	—	0.370	0.485
3.40	0.465	—	0.370	0.480
3.71	0.468	—	0.370	0.450
4.20	0.499	—	0.370	0.414
5.05	0.605	—	0.370	0.259
6.12	0.711	—	0.370	0.085
7.40	0.740	0.624	—	0.050
9.10	0.696	0.624	—	0.189
9.93	0.595	0.624	—	0.400
10.30	0.539	0.624	—	0.521
11.31	0.440	0.624	—	0.786
13.00	0.430	0.624	—	0.810

\*  $[HA] = 3 \times 10^{-4}$  M.

Table 2. Values of the dissociation constants and the distribution constants.

$pK_1$	$pK_2$	$\log D_{HA}$	$\log K_{AZ}$	$\log D_{AZ}$	$\log K_{ZCl}$	$\log D_{ZCl}$	Ref.
4.5	9.7						7
5.09	9.82	2.86					8
4.93	9.71						9
4.88	9.89						10
4.910	9.814						11
5.004	9.658	2.66					12
4.96	10.01	2.5	3.0	1.3	2.7*	0.89*	This work

\* Ref. 5.

Table 3. Spectrophotometric data on 8-hydroxyquinoline\* for the determination of  $D_{HA}$ .

pH	Absorbance	
	360 nm of aqueous phase	310 nm of organic phase
0.1 N HCl	0.510	0.000
1.51	0.432	0.080
2.00	0.343	0.250
2.50	0.240	0.400
2.90	0.175	0.510
3.00	0.154	0.536
4.95	0.020	0.745
7.01	0.020	0.745

\*  $[HA] = 3 \times 10^{-4}$  M.

$$K_1 \cdot D_{HA} = \frac{[HA]_{org} [H^+]}{[H_2A^+]} \quad (4)$$

$[HA]_{org}$  and  $[H_2A^+]$  were determined by absorbances at 310 nm in the organic phase and 360 nm in the aqueous phase (Table 3), and  $D_{HA}$  is calculated (Table 2).





Fig. 1. Absorption spectra in alkaline solution in the absence and presence of zephiramine. -----  $[Z]=0$  and —  $[Z]=1 \times 10^{-3}$  M; pH=(a) 9.10, (b) 10.30 and (c) 12.01;  $[HA]=3 \times 10^{-4}$  M.

In alkaline solution, the increase of absorbance at 360 nm indicates the formation of  $A^-$  from HA with increase of pH in the absence of zephiramine (Table 1).  $K_2$  is determined by plotting  $\log ([A^-]/[HA])$  against  $\log [H^+]$ .

With the addition of zephiramine, no changes of the absorption spectra were observed in acidic solution, but changes of the absorption spectra occurred in an alkaline solution (Fig. 1). The shift of the absorption maximum to a longer wavelength indicates an interaction between  $A^-$  and zephiramine. Variations of absorbances depending on the concentration of zephiramine were investigated at pH 13 (Table 4). When the pH is over 12, total oxinate concentration is assumed to equal oxinate ion concentration, and an isosbestic point at 370 nm suggests the following equilibrium (Fig. 1-c):



In the presence of zephiramine over  $4 \times 10^{-3}$  M, all of the oxinate ion is supposed to combine with zephiramine. Molar absorption coefficients of the ion-association complex and oxinate ion are  $2.2 \times 10^3$  and  $2.7 \times 10^3$  at 360 nm, respectively.  $K_{AZ}$  is calculated by plotting  $\log ([A^-]/[AZ])$  against  $\log [Z^+]$  by using the spectrophotometric data. As shown in Fig. 2, the ion-association complex, AZ, is distributed between the two phases. The distribution constant of AZ is given by

$$D_{AZ} = \frac{[AZ]_{org}}{[AZ]} \quad (6)$$

At pH 14, the present species are  $A^-$ , AZ and  $Z^+$  in both phases. Therefore, introduction of equation (5) into equation (6) gives:

Table 4. Spectrophotometric data at pH 13 for the determination of  $K_{AZ}$ .

HA $10^{-4}$ M	Z $10^{-3}$ M	Absorbance	
		360 nm	370 nm
0.6	0	0.168	0.142
0.6	1	0.141	0.142
1.2	0	0.333	0.290
1.2	1	0.299	0.290
1.8	0	0.485	0.410
1.8	1	0.429	0.410
2.4	0	0.645	0.564
2.4	1	0.575	0.564
3.0	0	0.795	0.700
3.0	0.1	0.790	0.700
3.0	0.5	0.755	0.700
3.0	1	0.714	0.700
3.0	3	0.676	0.700
3.0	4	0.672	0.700
3.0	5	0.672	0.700




Fig. 2. Absorption spectra at pH 14 in the aqueous and the organic phases. ----- Aqueous phase and — organic phase;  $[Z]$ =(a) 0, (b)  $1 \times 10^{-5}$  M, (c)  $1 \times 10^{-4}$  M, (d)  $2 \times 10^{-4}$  M, and (e)  $5 \times 10^{-4}$  M and  $1 \times 10^{-3}$  M;  $[HA]=3 \times 10^{-4}$  M.




Fig. 3. Distribution of 8-hydroxyquinoline in the presence of zephiramine.  $[HA]=1 \times 10^{-2}$  M, and  $[Z]=1 \times 10^{-3}$  M; (a)  $[HA]_{org}$ , (b)  $[H_2A^+]_{aq}$ , (c)  $[A^-]_{aq}$ , and (d)  $[AZ]_{org}$ .



I am also very grateful to Dr. S. Tsunogai and the members of the Laboratory of Analytical Chemistry of this faculty for their valuable discussions.

### References

- 1) Nishimura, M., Noriki, S. and Muramoto, S. (1974). Effect of quaternary ammonium bases on valence-saturated but coordination-unsaturated chelates. Part I. Extraction of chelates of glyoxal bis-(2-hydroxyanil) and o-(salicylideneamino) phenol. *Anal. Chim. Acta* **70**, 121-126.
- 2) Noriki, S. and Nishimura, M. (1974). *Ditto*, Part II. Extraction of magnesium 8-hydroxyquinolate. *Ibid.* **72**, 339-343.
- 3) Noriki, S. (1975). *Ditto*, Part III. Extraction of nickel- and cobalt-thenoyltrifluoroacetone chelates. *Ibid.* **76**, 215-218.
- 4) Noriki, S. and Nishimura, M. (1977). *Ditto*, Part IV. Extraction of some divalent metal 8-hydroxyquinoline. *Ibid.* **94**, 57-62.
- 5) Noriki, S. (1983). Adductive effect of a quaternary ammonium base on solvent extraction of coordination-unsaturated chelates. *Bull. Fac. Fish. Hokkaido Univ.* **34**, 264-277. (In Japanese with English abstract).
- 6) Nishimura, M. and Noriki, S. (1975). Effect of a quaternary ammonium base on chelate formation. *Anal. Instruments* **13**, 24-31. (in Japanese).
- 7) Kolthoff, I.M. (1927). The detection and determination of metals by means of o-hydroxyquinoline (oxine). *Chem. Weekblad* **24**, 606-610.
- 8) Lacroix, S. (1947). Properties of the oxinates of aluminum, gallium, and indium. Separation of gallium by extraction with chloroform. *Anal. Chim. Acta* **1**, 260-290. (in French).
- 9) Phillips, J.P. and Merritt, L.L. (1948). Ionization constants of some substituted 8-hydroxyquinolines. *J. Amer. Chem. Soc.* **70**, 410-411.
- 10) Irving, H., Ewart, J.A.D. and Wilson, J.T. (1949). The dissociation constants of 8-hydroxyquinoline. *J. Chem. Soc.*, 2672-2674.
- 11) Nasanen, R., Lumme, P. and Mukula, A.-L. (1951). Potentiometric and spectrophotometric studies on 8-hydroxyquinolinol and its derivatives. I. Ionization of 8-quinolinol in aqueous solutions of potassium chloride. *Acta. Chem. Scand.* **5**, 1199-1208.
- 12) Dyrssen, D. (1952). Studies on the extraction of metal complex. IV. The dissociation constants and partition coefficients of 8-quinolinol (oxine) and N-nitroso-N-phenylhydroxyl-amine (cupferron). *Svensk Kem. Tidsskr.* **64**, 213-224.
- 13) Nishida, S. (1971). Spectrophotometric study of iron (III)-chromazurol S complex in the presence of zephiramine. *Bunseki Kagaku* **20**, 410-415. (In Japanese with English abstract).
- 14) Kohara, H. (1974). Mechanism of higher order complexation in the neighborhood of charge-cluster. *Ibid.* **23**, 39-46. (In Japanese with English abstract).