



# HOKKAIDO UNIVERSITY

Title	The in Vitro Effects of Various Steroid Hormones and Gonadotropin on Oocyte Maturation of the Viviparous Rockfish, <i>Sebastes taczanowskii</i>
Author(s)	TAKEMURA, Akihiro; 竹村, 明洋; TAKANO, Kazunori et al.
Citation	北海道大學水産學部研究彙報, 40(1), 1-7
Issue Date	1989-02
Doc URL	<a href="https://hdl.handle.net/2115/24014">https://hdl.handle.net/2115/24014</a>
Type	departmental bulletin paper
File Information	40(1)_P1-7.pdf



## The *in Vitro* Effects of Various Steroid Hormones and Gonadotropin on Oocyte Maturation of the Viviparous Rockfish, *Sebastes taczanowskii*

Akihiro TAKEMURA\*, Kazunori TAKANO\*  
and Kohei YAMAUCHI\*

### Abstract

The effects of seven steroid hormones (progesterone,  $17\alpha$ -hydroxyprogesterone,  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one, deoxycorticosterone, cortisol, testosterone and estradiol- $17\beta$ ) and human chorionic gonadotropin on the induction of *in vitro* oocyte maturation were examined in a viviparous rockfish, *Sebastes taczanowskii*. Among them,  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one was the most potent steroid in inducing *in vitro* germinal vesicle breakdown (GVBD), a criterion for oocyte maturation in follicle-enclosed oocytes. Progesterone,  $17\alpha$ -hydroxyprogesterone and deoxycorticosterone were moderately effective only at a high dose, while cortisol was slightly effective. Testosterone and estradiol- $17\beta$  did not have any effect at the concentrations tested. Human chorionic gonadotropin was very effective with rates of GVBD similar to those of  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one. Moreover, human chorionic gonadotropin was effective in stimulating  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one synthesis by follicle-enclosed oocytes. These results suggest that  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one is the maturation inducing steroid of this species and that synthesis of the steroid is regulated by gonadotropin.

### Introduction

Oocyte development and maturation in teleosts are regulated directly and indirectly by various hormones in the hypothalamus-pituitary-ovary axis. For the majority of fish species studied, oocyte maturation including germinal vesicle breakdown (GVBD) is particularly induced by either 11-deoxycorticoids or progestogens (cf. Goetz, 1983). Many studies have demonstrated that in various teleosts,  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one ( $17\alpha$ ,  $20\beta$ -diOHprog) is one of the most potent progestogens in inducing oocyte maturation, based on *in vitro* experiments (Duffey and Goetz, 1980; Nagahama et al., 1980; Young et al., 1982; Nagahama et al., 1983; Yamauchi et al., 1983; Upadhyaya and Haider, 1986; Adachi et al., 1988) and serum profiles of the steroid during the maturation period (Kagawa et al., 1983; Young et al., 1983; Ueda et al., 1984; Yamauchi et al., 1984; Shimizu et al., 1985; Levavi-Zermonsky and Yaron, 1986).

In viviparous teleosts, although a few studies on hormonal regulation of gestation and/or parturition have been carried out (Ishii, 1962 and 1963; Korsgaard and Petersen, 1979; deVlaming et al., 1983; Groves and Batten, 1986), there have been no reports on regulation of oocyte maturation. Fertilization, gestation and

---

\* Laboratory of Fresh-Water Fish-Culture, Faculty of Fisheries, Hokkaido University  
(北海道大学水産学部淡水増殖学講座)

parturition are a chain of events which follow on oocyte maturation in the ovary of viviparous species. To clarify these peculiar events associated with viviparity, it is very interesting to investigate the endocrine mechanisms which regulate oocyte maturation. The present study was designed to examine the effect of various hormones on oocyte maturation of a marine viviparous rockfish, *Sebastes taczanowskii*, by an *in vitro* incubation method. An experiment on *in vitro*  $17\alpha, 20\beta$ -diOHprog synthesis in the ovarian follicles by addition of gonadotropin was also carried out.

### Material and Methods

The white-edged rockfish, *Sebastes taczanowskii*, used in this study were collected by angling or with gill nets along the shores of Shikabe and Esan in the suburbs of Hakodate, southern Hokkaido from spring to fall in 1986. These were reared in indoor circular tanks (1 ton capacity) with seawater flowing through under natural water temperature and daylight condition until use in late March to early April of 1987. After anesthetization in ethyl 4-aminobenzoate, ovaries were dissected out and placed in cold Eagle's MEM solution buffered by 10 mM Hepes (pH 7.6). Follicle-enclosed oocytes were then teased from the overlaying connective tissues.

Effects on GVBD in oocytes were evaluated by seven steroid hormones: testosterone (T) and estradiol- $17\beta$  ( $E_2$ ),  $17\alpha, 20\beta$ -diOHprog,  $17\alpha$ -hydroxy-progesterone ( $17\alpha$ -OHprog), progesterone (Prog), deoxycorticosterone (DOC), cortisol (F) (Sigma, USA). Stock solutions of each steroid were dissolved in ethanol and then diluted with Eagle's MEM to give final concentrations of 0.001, 0.01, 0.1 and 1.0  $\mu\text{g/ml}$  which contained less than 1% ethanol. In addition, the effect of human chorionic gonadotropin (HCG; Teikoku zohki, Tokyo, Japan) on the induction of GVBD was examined. HCG (1, 10, 100 IU/ml) was directly dissolved in Eagle's MEM. Fifty oocytes were incubated in each hole of Nunclon (Nunc, Denmark) with or without hormones for 72 hours at 10°C. Three replicates were conducted for each treatment.

To determine *in vitro*  $17\alpha, 20\beta$ -diOHprog synthesis, 30 oocytes were incubated in three HCG concentrations (1, 10 and 100 IU/ml) for 72 hours at 10°C. After incubation, oocytes with induced GVBD were counted and the incubation media were kept at -40°C until analysis. The steroid was measured using the radioimmunoassay method of Kagawa et al. (1981).

All data are expressed as mean  $\pm$  SEM. Duncan's multiple range test was used for statistical analysis.

### Results and Discussion

As oocyte maturation progressed, the germinal vesicle became visible as it migrated from the inner portion of the cell to the animal pole (Fig. 1a). Subsequently, the germinal vesicle disappeared (GVBD) and the oocyte became translucent at the same time (Fig. 1b). During the course of maturation, small oil droplets dispersed through the ooplasm, and then gradually aggregated to form a large, prominent oil globule. Oocytes also increased significantly ( $p < 0.01$ ) in diameter during this process (Table 1). In this study, GVBD was used as a criterion for determining final maturation.




Fig. 1. Changes prior to (a) and after (b) germinal vesicle breakdown in *S. taczanowskii* oocytes. The bar represents 100  $\mu$ m. GV and OD show a germinal vesicle and oil droplet, respectively.

Table 1. Changes of oocyte diameter prior to and after oocyte maturation

	maturing	matured
Egg diameter (mm)	0.799 $\pm$ 0.003 (23) <sup>a</sup>	0.816 $\pm$ 0.004*(23)

<sup>a</sup> Number of oocytes measured. Each value represents the mean  $\pm$  standard error.

\* Significantly different from maturing oocytes at  $p < 0.01$ .

The effects of the seven steroid hormones (T, E<sub>2</sub>, DOC, F, Prog, 17 $\alpha$ -OHprog and 17 $\alpha$ , 20 $\beta$ -diOHprog) tested for *in vitro* induction of GVBD in isolated oocytes are shown in Fig. 2. Among these hormones, only T and E<sub>2</sub> did not have any effect at the concentrations tested.

DOC was moderately effective at 1.0  $\mu$ g/ml (53% GVBD), while F was slightly effective even at the same concentration (10% GVBD). In several species, 11-corticosteroids such as DOC are as potent as progestogens in inducing oocyte maturation (Nagahama et al., 1980; Duffey and Goetz, 1980; Young et al., 1982; Yamauchi et al., 1984; Upadhyaya and Haider, 1986; Adachi et al., 1988). A positive interpretation of the effect of DOC must be deferred, because physiological information such as the serum profile and gonadal synthesis of corticosteroids have not been obtained in *S. taczanowskii*.

Of the three progestogens used in this study, Prog and 17 $\alpha$ -OHprog, as well as DOC, were moderately effective only at a high dose (1.0  $\mu$ g), showing GVBD rates of 61% and 68%, respectively. The most effective steroid was 17 $\alpha$ , 20 $\beta$ -diOHprog which induced GVBD in 82% of oocytes tested at a concentration of 1.0  $\mu$ g/ml. At a level of 0.1  $\mu$ g/ml, this inducer also produced a 75% response, although the other six steroids were ineffective or much less effective at the same level. It has been reported that 17 $\alpha$ , 20 $\beta$ -diOHprog has the most potent effect in inducing *in vitro* oocyte maturation in fish (Duffey and Goetz, 1980; Nagahama et al., 1980; Young et al., 1982; Nagahama et al., 1983; Yamauchi et al., 1983; Upadhyaya and Haider, 1986; Adachi et al., 1988). In addition, the levels of this hormone in the blood reached a peak near the time of oocyte maturation in salmonids (Young et al.,




Fig. 2. *In vitro* effects of various steroid hormones on inducing germinal vesicle breakdown in *S. taczanowskii* oocytes.

1983; Yamauchi et al., 1984; Ueda et al., 1984), goldfish (Kagawa et al., 1983), carp (Levavi-Zermonsky and Yaron, 1986) and bitterling (Shimizu et al., 1985). The results described above show that 17 $\alpha$ , 20 $\beta$ -diOHprog is the maturation inducing steroid in many oviparous species. Based on the results of this study, it is suggested that 17 $\alpha$ , 20 $\beta$ -diOHprog is also the maturation inducing steroid in *S. taczanowskii*.

The present study also showed that HCG was considerably effective at 1, 10 and 100 IU/ml with GVBD of 11%, 77% and 86%, respectively (Fig. 3). Although HCG has been used extensively for the *in vivo* induction of final maturation and ovulation, it is less active *in vitro* in many fish (Goetz, 1983). Among those species examined, HCG has been shown to be very effective in inducing *in vitro* GVBD in killifish, *Fundulus heteroclitus* (Wallace and Selman, 1980) as well as in *S. taczanowskii*. The causes of these differences in biological activity of HCG among different fishes are still unknown. In a preliminary experiment, it has been ascertained that the pituitary preparation of *S. taczanowskii* was also very effective, inducing 91% and 52% responses at levels of 1% and 0.1% (w/v) fresh pituitary suspension, respectively (Takemura, unpublished). These facts suggest that the pituitary and gonadotropin preparations are also effective in inducing *in vitro* oocyte maturation in this fish.

Furthermore, the effect of HCG on *in vitro* 17 $\alpha$ , 20 $\beta$ -diOHprog synthesis by follicle-enclosed oocytes of *S. taczanowskii* was examined in this study. As shown in Fig. 4, HCG was significantly ( $p < 0.01$ ) effective in stimulating 17 $\alpha$ , 20 $\beta$ -diOH-




Fig. 3. An *in vitro* effect of HCG on inducing germinal vesicle breakdown in *S. taczanowskii* oocytes.

prog synthesis at concentrations of 10 and 100 IU/ml. In some fish species, the action of gonadotropin in inducing oocyte maturation appears to be dependent on the synthesis of a secondary steroidal effector, a maturation-inducing hormone (Goetz, 1983; Nagahama, 1987). Nevertheless definite evidence for the action of HCG is lacking in this study; the results obtained suggest that the pathway leading to the induction of final maturation in viviparous scorpaenids is similar to that in oviparous species already investigated (Goetz, 1983).

Groves and Batten (1986) suggested that in a viviparous teleost, *Poecilia latipinna*,  $17\alpha, 20\beta$ -diOHprog influenced the female reproductive cycle at ovulation/parturition rather than at oocyte maturation at the hypophyseal level, based on ultrastructural observations of *in vitro* effects of some gonadal steroids on the gonadotrophs. During the reproductive cycles of *S. taczanowskii* and a related species, *S. schlegeli*, however, the serum concentrations of  $17\alpha, 20\beta$ -diOHprog have been observed to increase around the estimated time of final oocyte maturation and subsequent gestation (Takemura, unpublished). Therefore, it is likely that  $17\alpha, 20\beta$ -diOHprog is at least partially involved in oocyte maturation and gestation maintenance in viviparous scorpaenids.




Fig. 4. *In vitro* synthesis of 17 $\alpha$ ,20 $\beta$ -diOHprog (bar graph) and the percentages of oocytes having induced germinal vesicle breakdown (line graph) at various HCG concentrations. \* Indicates significant difference ( $p < 0.01$ ) compared to control.

### Acknowledgments

We would like to express our gratitude to Professor Y. Nagahama, National Institute for Basic Biology, Okazaki, who kindly provided us with the antiserum against 17 $\alpha$ ,20 $\beta$ -diOHprog. We are also very grateful to Professor H. Takahashi, Hokkaido University, for his useful discussion of this work. This study was supported in part by a Grant-in Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

### References

- Adachi, S., Ouchi, K., Hirose, K. and Nagahama, Y. (1988). Induction of oocyte maturation *in vitro* by steroid hormones in the red sea bream *Pagrus major*. *Nippon Suisan Gakkaishi* **54**, 1665.
- deVlaming, V., Baltz, D., Anderson, S., Fitzgerald, R., Delahunty, G. and Barkley, M. (1983). Aspects of embryo nutrition and excretion among viviparous embiotocid teleosts: Potential endocrine involvements. *Comp. Biochem. Physiol.* **76A**, 189-198.
- Duffey, R.J. and Goetz, F.W. (1980). The *in vitro* effects of 17 $\alpha$ -hydroxy-20 $\beta$ -dihydroprogesterone on germinal vesicle breakdown in brook trout (*Salvelinus fontinalis*) oocytes. *Gen. Comp. Endocrinol.* **41**, 563-565.
- Goetz, F.W. (1983). Hormonal control of oocyte final maturation and ovulation in fishes. p. 117-170. In Hoar, W.S., Randall, D.J. and Donaldson, E.M. (eds.), *Fish Physiology*, Vol. IXB,

- 477 p, Academic Press, New York.
- Groves, D.J. and Batten, T.F.C. (1986). Direct control of the gonadotroph in a teleost, *Poecilia latipinna* : Gonadal steroids. *Gen. Comp. Endocrinol.* **61**, 402-416.
- Ishii, S. (1962). Effects of some hormones on the gestation of the top-minnow. *J. Fac. Sci. Univ. Tokyo. Sec. IV*, **9**, 279-290.
- Ishii, S. (1963). Some factors involved in the delivery of the young in the top-minnow, *Gambusia affinis*. *Ibid.* **10**, 181-187.
- Kagawa, H., Takano, K. and Nagahama, Y. (1981). Correlation of plasma estradiol-17 $\beta$  and progesterone levels with ultrastructure and histochemistry of ovarian follicles in the white-spotted char, *Salvelinus leucomaenis*. *Cell Tissue Res.* **218**, 315-329.
- Kagawa, H., Young, G. and Nagahama, Y. (1983). Changes in plasma steroid hormone levels during gonadal maturation in female goldfish *Carassius auratus*. *Bull. Jap. Soc. Sci. Fish.* **49**, 1783-1787.
- Korsgaard, B. and Petersen, I. (1979). Vitellogenin, lipid and carbohydrate metabolism during vitellogenesis and pregnancy, and after hormonal induction in the blenny *Zoarces viviparus* (L.). *Comp. Biochem. Physiol.* **63B**, 245-251.
- Levavi-Zermansky, B. and Yaron, Z. (1986). Changes in gonadotropin and ovarian steroids associated with oocytes maturation during spawning induction in the carp. *Gen. Comp. Endocrinol.* **62**, 89-98.
- Nagahama, Y. (1987). Gonadotropin action on gametogenesis and steroidogenesis in teleost gonads. *Zool. Sci.* **4**, 209-222.
- Nagahama, Y., Kagawa, H. and Tashiro, F. (1980). The *in vitro* effects of various gonadotropins and steroid hormones on oocyte maturation in amago salmon *Oncorhynchus rhodurus* and rainbow trout *Salmo gairdneri*. *Bull. Jap. Soc. Sci. Fish.* **46**, 1097-1102. (In Japanese with English abstract).
- Nagahama, Y., Hirose, K., Young, G., Adachi, S., Suzuki, K. and Tamaoki, B. (1983). Relative *in vitro* effectiveness of 17 $\alpha$ , 20 $\beta$ -dihydroxy-4-pregnen-3-one and other pregnene derivatives on germinal vesicle breakdown in oocytes of Ayu (*Plecoglossus altivelis*), amago salmon (*Oncorhynchus rhodurus*), rainbow trout (*Salmo gairdneri*), and goldfish (*Carassius auratus*). *Gen. Comp. Endocrinol.* **51**, 15-23.
- Shimizu, A., Aida, K. and Hanyu, I. (1985). Endocrine profiles during the short reproductive cycle of an autumn-spawning bitterling, *Acheilognathus rhombea*. *Ibid.* **60**, 361-371.
- Ueda, H., Hiroi, O., Hara, A., Yamauchi, K. and Nagahama, Y. (1984). Changes in serum concentrations of steroid hormones, thyroxine, and vitellogenin during spawning migration of the chum salmon, *Oncorhynchus keta*. *Ibid.* **53**, 203-211.
- Upadhyaya, N. and Haider, S. (1986). Germinal vesicle breakdown in oocytes of catfish, *Mystus vittatus* (Bloch) : Relative *in vitro* effectiveness of estradiol-17 $\beta$ , androgens, corticosteroids, progesterone, and other pregnene derivatives. *Ibid.* **63**, 70-76.
- Wallace, R.A. and Selman, K. (1980). Oogenesis in *Fundulus heteroclitus* II. The transition from vitellogenesis into maturation. *Ibid.* **42**, 345-354.
- Yamauchi, K., Ban, M., Tsuda, K., Kasahara, N. and Izumi, T. (1983). The *in vitro* effects of various steroid hormones and gonadotropin on oocyte maturation in masu salmon, *Oncorhynchus masou*. *Sci. Rec. Hokkaido Fish Hatch.* **38**, 85-91. (In Japanese with English abstract).
- Yamauchi, K., Kagawa, H., Ban, M., Kasahara, N. and Nagahama, Y. (1984). Changes in plasma estradiol-17 $\beta$  and 17 $\alpha$ , 20 $\beta$ -dihydroxy-4-pregnen-3-one levels during final oocyte maturation of the masu salmon *Oncorhynchus masou*. *Bull. Jap. Soc. Sci. Fish.* **50**, 2137.
- Young, G., Kagawa, H. and Nagahama, Y. (1982). Oocyte maturation in the amago salmon (*Oncorhynchus rhodurus*) : *In vitro* effects of salmon gonadotropin, steroids, and cyanoketone (an inhibitor of 3 $\beta$ -hydroxy- $\Delta^5$ -steroid dehydrogenase). *J. Exp. Zool.* **224**, 265-275.
- Young, G., Crim, L.W., Kagawa, H., Kambegawa, A. and Nagahama, Y. (1983). Plasma 17 $\alpha$ , 20 $\beta$ -dihydroxy-4-pregnen-3-one levels during sexual maturation of amago salmon (*Oncorhynchus rhodurus*) : Correlation with plasma gonadotropin and *in vitro* production by ovarian follicles. *Gen. Comp. Endocrinol.* **51**, 96-105.