



HOKKAIDO UNIVERSITY

Title	EVALUATION OF INTERFERON- α ANTI-VIRAL ACTIVITY AGAINST AUJESZKEY'S DISEASE VIRUS, AND THE MECHANISM(S) INVOLVED
Author(s)	TONOMURA, Noriko
Citation	Japanese Journal of Veterinary Research, 44(1), 29-29
Issue Date	1996-05-31
Doc URL	https://hdl.handle.net/2115/2529
Type	departmental bulletin paper
File Information	KJ00002398213.pdf



EVALUATION OF INTERFERON- α ANTI-VIRAL ACTIVITY
AGAINST AUJESZKEY'S DISEASE VIRUS,
AND THE MECHANISM(S) INVOLVED

Noriko TONOMURA

*Laboratory of Microbiology,
Department of Disease Control,
School of Veterinary Medicine,
Hokkaido University, Sapporo 060, Japan*

Interferon- α is a type I interferon, produced by neutrophils, macrophages, fibroblasts and so on. It is reported to inhibit Aujeszky's disease virus (ADV) replication in nervous tissue cells and porcine nasal mucosa *in vitro*. However the mechanism(s) by which interferon- α exerts its anti-viral activity against ADV is not known.

To evaluate the anti-viral activity of human natural interferon- α (IFN- α) against ADV, viral replication in IFN- α -treated Vero cells was examined by plaque assay. The replication of ADV was suppressed by treatment with IFN- α by 0.1~0.01 fold compared to that of the control until 24 hours post infection.

Effects of IFN- α on early steps of ADV infection, such as adhesion, penetration, uncoating, and transportation to the cell nucleus, were examined by dot blot analysis of the viral DNA in the IFN- α -treated cells. Equal amount of viral DNA was detected in IFN- α treated and control cell nuclei, indicating that the early steps of ADV infection were not impaired by IFN- α treatment.

Messenger RNA (mRNA) transcribed from the ADV IE gene was reduced in the IFN- α -treated cells when detected by Northern blot analysis, and transient expression assays showed that the transcription from the ADV IE promoter was selectively inhibited in IFN- α -treated Vero cells. Analysis of deletion mutants of the ADV IE promoter suggested that sequence between -90 and the transcription initiation site is needed for the inhibition.

These results indicate that IFN- α treatment inhibits ADV replication by altering the transcription of the IE gene, and suggest the involvement of ADV IE promoter in that event.