



# HOKKAIDO UNIVERSITY

Title	Magnetic resonance imaging of hepatocellular carcinoma in Long-Evans Cinnamon rats under a magnetic field of 7.05 T
Author(s)	ASANUMA, Taketoshi; HIRANO, Yoshiko; OHKURA, Kaori et al.
Citation	Japanese Journal of Veterinary Research, 45(3), 147-151
Issue Date	1997-11-28
DOI	<a href="https://doi.org/10.14943/jjvr.45.3.147">https://doi.org/10.14943/jjvr.45.3.147</a>
Doc URL	<a href="https://hdl.handle.net/2115/2601">https://hdl.handle.net/2115/2601</a>
Type	departmental bulletin paper
File Information	KJ00002398560.pdf



## Magnetic resonance imaging of hepatocellular carcinoma in Long-Evans Cinnamon rats under a magnetic field of 7.05 T

Taketoshi Asanuma, Yoshiko Hirano, Kaori Ohkura, Yasuhiro Kon<sup>1</sup>,  
Shigezou Shimokawa<sup>2</sup>, Noriyuki Kasai<sup>3</sup> and Mikinori Kuwabara

(Accepted for publication: Sep. 24, 1997)

### Abstract

Magnetic resonance (MR) images of livers in 3-, 12- and 29-month-old Long-Evans Cinnamon(LEC) rats (male) were taken under a magnetic field of 7.05 T. MR images of sagittal and transversal sections were obtained in 1-mm-thick slices by T<sub>1</sub>-weighted and two-dimensional Fourier transformation techniques. The data matrix size was 256 phase-encoded steps. Each image was obtained through four acquisitions. Three-month-old rats gave MR images with low signal intensity over the liver probably due to the shorting of its T<sub>1</sub> and T<sub>2</sub> relaxation times. However, 12-month-old rats gave hyperintense regions around hepatic veins in right hepatic lobe, which was assigned to hepatocellular carcinoma. In 29-month-old rats, MR images with hyperintensity throughout the hepatic lobe were observed. These MR images, therefore, suggested that hepatocellular carcinoma in LEC rats developed from the restricted regions surrounding hepatic veins. In the present study, T<sub>1</sub>-weighted MR imaging under a magnetic field of 7.05 T was shown to be applicable to the diagnosis of hepatic cancer in LEC rats.

Key words: hepatocellular carcinoma, Long-Evans Cinnamon rat, MRI

### Introduction

The theory of nuclear magnetic resonance (NMR) suggests that the higher magnetic field the stronger are NMR signals obtained<sup>2)3)8)</sup>. This means that MR imaging under a high magnetic field is favored over imaging under a low magnetic field for small animals like mice and rats. Our previous studies showed that MR imaging of rat brains under a magnetic field of 7.05 T, which

was higher than that of usual human MRI (0.5–2.0 T), was successful to visualize the age-dependent morphological changes in brains<sup>7)11)</sup>.

Recent MR imaging studies proved that this method was useful for the diagnosis of not only human brain cancer but also human hepatocellular carcinoma (HCC)<sup>9)</sup>. Since LEC rats are known as a good animal model for studying chronic hepatitis as well as hepatic cancer<sup>5)10)12)</sup>, it is natural to consider that the MR imaging method

---

Laboratory of Radiation Biology, Department of Environmental Veterinary Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060, Japan

<sup>1</sup> Laboratory of Experimental Animal Science, Department of Disease Control, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060, Japan

<sup>2</sup> Faculty of Engineering, Hokkaido University, Sapporo 060, Japan

<sup>3</sup> Institute for Animal Experimentation, Tohoku University School of Medicine, Sendai 980–77, Japan.  
FAX: 011–706–5236

E-Mail: teiou@vetmed.hokudai.ac.jp

should be applicable to the diagnosis of hepatic cancer of LEC rats. However, LEC rats contain several paramagnetic ions in the liver at high concentrations (40–50 times higher than those of normal rats). Generally, the higher the concentrations of paramagnetic ions and the higher magnetic field, the relaxation times ( $T_1$  and  $T_2$ ) of MRI become shorter<sup>1)14)</sup>. The shortening of  $T_1$  and  $T_2$  relaxation times results in the decrease of MR signal intensities and gives hypointense MR images. In the present study,  $T_1$ -weighted MR images of the livers of 3-, 12- and 29-month-old LEC rats were taken under a magnetic field of 7.05 T to know whether or not  $T_1$ -weighted MR imaging at high magnetic field was applicable to hepatic pathogenesis in LEC rats.

## Materials and Methods

### Animals

LEC rats which are maintained under conventional conditions at the Institute for Animal Experimentation, Hokkaido University School of Medicine were used. They fed a regular diet and water *ad libitum* in an air-conditioned animal room at  $22 \pm 3^\circ\text{C}$  with a relative humidity of  $55 \pm 5\%$ . Male rats aged at 3, 12 and 29 months old were used in the present experiments. Hepatitis with jaundice was once observed at 12- and 29-months of age<sup>15)</sup>.

### MRI

$T_1$ -weighted MR images were obtained on an Oxford 7.05 T superconducting magnet equipped with a Toshiba workstation with an SIS 300/183 imaging system using a two-dimensional Fourier transformation technique. LEC rats were anesthetized with pentobarbital sodium solution (40 mg/kg) diluted 2.5 times with a mixed solution of propylene glycol/ethanol/ $\text{H}_2\text{O}$  (2/1/7, w/w). The saddle-shaped coil (89 mm in diameter) designed for MR imaging of whole body of rat was used. Each rat was positioned supination in the probe running parallel to z-axis of the magnet, and fixed to a cradle using adhesive tape to minimize the

artifacts due to respiratory motions. After adjusting the coils, sagittal sections of body were roughly imaged under a  $T_1$ -weighted condition (repetition time (TR) / echo time (TE) = 500 ms / 20 ms) to determine the positions of transversal sections in the hepatic region. Then,  $T_1$ -weighted images were taken using a spin echo (SE) sequence with 4 acquisitions and pulse width of 4 kHz. In addition to these parameters, the followings were employed<sup>7)11)</sup>; the gradient magnetic field, 3 mT/cm; the field of view (FOV),  $10 \times 10 \text{ cm}^2$  for sagittal sections and  $6 \times 6 \text{ cm}^2$  for transversal section; the slice thickness, 1.0 mm; data matrix size,  $256 \times 256$ ; reconstructed images,  $512 \times 512$  pixel matrix.

## Results

Typical  $T_1$ -weighted MR image of the transversal section at about 10 mm from the top of xiphoid process to head in a 3-month-old male LEC rat is shown in Fig. 1a. Hepatic veins, muscle and spinal cord were clearly imaged, though whole liver showed hypointensity on  $T_1$ -weighted image. Shortening of  $T_1$  and/or  $T_2$  relaxation times of the liver due to the presence of excess metal ions is thought to be responsible for hypointensity (unpublished results).

MR image of the transversal section at about 5 mm from the top of xiphoid process to head in a 12-month-old rat is shown in Fig. 1b. The MR signal intensity in the normal hepatic region was the same as that of a 3-month-old rat, but the hyperintense regions (2–6 mm in diameter) were observed around hepatic veins in right hepatic lobe. These hyperintense regions were assigned to HCC regions because the hyperintense regions were proved to be HCC in  $T_1$ -weighted MR imaging of the livers in human<sup>4)6)13)</sup>.

MR image of the transversal section at about 15 mm from the top of xiphoid process to tail in 29-month-old rat is shown in Fig. 1c. A number of hyperintense regions (6–10 mm in diameter) with irregular shapes were observed all over




Fig. 1 (a)  $T_1$ -weighted MR image of the transversal section at about 10 mm from the top of xiphoid process to head of 3-month-old LEC rat abdomen. LL : left hepatic lobe, RL : right hepatic lobe, SC : spinal cord, DM : dorsal muscle, VM : ventral muscle, HV : hepatic vein. (b)  $T_1$ -weighted MR image of the transversal section at about 5 mm from the top of xiphoid process to head of 12-month-old LEC rat abdomen. S : stomach. Arrows show abnormal hyperintense regions around hepatic veins. These regions were assigned to HCC. (c)  $T_1$ -weighted MR image of the transversal section at about 15 mm from the top of xiphoid process to tail of 29-month-old LEC rat abdomen. Solid arrow heads show hyperintense regions (6–10 mm in diameter) with obscured borders gathering with one another. An open arrow head shows hyperintense region with egg-like structure consisting of the viscous liquid.

hepatic lobe. Additionally, several hyperintense regions with unclear border were observed to gather with one another. A large egg-like structure was observed as the hyperintense region in the left hepatic lobe. The pathological dissection after MR imaging revealed that a viscous liquid cyst was present at the surface of left hepatic lobe (Fig. 2). The egg-like structure was, therefore, assigned to the liquid cyst but not HCC.

#### Discussion

$T_1$ -weighted MR imaging technique was applied to visualize the livers of 3-, 12- and 29-month-old LEC rats under a high magnetic field of 7.05 T. Before experiments, there was a fear that  $T_1$ -weighted MR images of LEC rat livers under the high magnetic field, as a whole, become hypointense due to the shortening of  $T_1$  and  $T_2$  relaxation times arising from the presence of excess paramagnetic ions. However, anatomical and pathological information of the livers could be obtained in 1-mm-thick slices by MR images. A number of hyperintense regions in




Fig. 2 Macroscopic observation of 29-month-old LEC rat liver. The cystic hypertrophy of liver by HCC was observed in upper abdominal cavity (arrow). This cyst corresponded to the egg-like structure in Fig. 1c.

the 12- and 29-month-old rat livers were observed in  $T_1$ -weighted images. These regions were assigned to HCC because the hyperintense regions were identified as HCC in  $T_1$ -

weighted MR imaging of the livers in human<sup>4)6)13)</sup>. The interesting finding in 12-month-old rat livers was the fact that the hyperintense regions corresponding to HCC were distributed at the circumference of hepatic veins. Furthermore, it was found that hyperintense regions were distributed over the livers in 29-month-old rats. These results indicated that HCC in LEC rats developed from the regions surrounding the hepatic veins.

From these results, it was concluded that T<sub>1</sub>-weighted MR imaging at high magnetic field was applicable to hepatic pathogenesis in LEC rats.

#### Acknowledgements

This was supported in part by a Grant-in-Aid for Basic Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan (No. 09556064). In conducting this work, the investigators adhered to The Guide for the Care and Use of Laboratory Animals, Hokkaido University School of Medicine.

#### References

- 1) Bloembergen, N. and Morgan, L. O. 1961. Proton relaxation times in paramagnetic solutions. effects of electron spin relaxation. *J. Chem. Phys.*, 34 : 842–850.
- 2) Crooks, L. E., Mills, C. and Davis, P. L. 1982. Visualization of cerebral and vascular abnormalities by NMR imaging. The effects of imaging parameters on contrast. *Radiology*, 144 : 843–852.
- 3) Dixen, R. L. and Ekstrand, K. E. 1982. The physics of proton NMR. *Med. Phys.*, 9 : 807–818.
- 4) Ebara, M., Watanabe, S., Kita, K., Yoshikawa, M., Sugiura, N., Ohto, M., Kondo, F. and Kondo, Y. 1991. MR imaging of small hepatocellular carcinoma : effect of intratumoral copper content on signal intensity. *Radiology*, 180 : 617–621.
- 5) Enomoto, K., Sawaki, M., Takahashi, H., Nakajima, Y., Dempo, K. and Mori, M. 1991. The multistep nature of spontaneous liver cancer development in the LEC rat : Analysis of incidence and phenotype of preneoplastic and neoplastic liver lesions. In : *The LEC Rat*, pp.305–312, Mori, M., Yoshida, M., C., Takeichi, N. and Taniguchi, N. eds., Springer-Verlag, Tokyo.
- 6) Itoh, K., Nishimura, K., Togashi, K., Fujisawa, I., Noma, S., Minami, S., Sagoh, T., Nakano, Y., Itoh, H., Mori, K., Ozawa, K. and Torizuka, K. 1987. Hepatocellular carcinoma : MR imaging. *Radiology*, 164 : 21–25.
- 7) Inanami, O., Asanuma, T., Inukai, N., Jin, T., Shimokawa, S., Kasai, N., Nakano, M., Sato, F. and Kuwabara, M. 1995. The suppression of age-related accumulation of lipid peroxides in rat brain by administration of Rooibos tea (*Aspalathus linearis*). *Neuroscience Letters*, 196 : 85–88.
- 8) Johnson, G. A., Herfkens, R. J. and Brown, M. A. 1985. Tissue relaxation time : in vivo field dependence. *Radiology*, 156 : 805–810.
- 9) Kadoya, M., Matsui, O., Takashima, T. and Nonomura, A. 1992. Hepatocellular carcinoma : correlation of MR imaging and histopathologic findings. *Radiology*, 183 : 819–825.
- 10) Kawano, K., Hirashima, T., Mori, S., Bando, S., Yonemoto, K., Abe, F. and Natori, T. 1991. Neoplastic and non-neoplastic lesions in aging LEC/Otk rats. In : *The LEC Rat*, pp.305–312, Mori, M., Yoshida, M. C., Takeichi, N. and Taniguchi, N. eds., Springer-Verlag, Tokyo.
- 11) Kuwabara, M., Asanuma, T., Inanami, O., Jin, T., Shimokawa, S., Kasai, N., Kator, K. and Sato, F. 1994. Magnetic resonance imaging of young and aged rat brains under a magnetic field of 7.05 T. *J. Vet. Med. Sci.*, 55 : 933–938.
- 12) Li, Y., Togashi, Y. and Takeichi, N. 1991. Abnormal copper accumulation in the liver of LEC rats : a rat form of Wilson's disease. In : *The LEC Rat*, pp.305–312, Mori, M., Yoshida, M. C., Takeichi, N. and Taniguchi, N. eds., Springer-Verlag, Tokyo.
- 13) Mitchell, D. G., Palazzo, J., Hamm, Hie-Won, Y. L., Rifkin, M. D., Burk, D. L. Jr. and Rubin,

- R. 1991. Hepatocellular tumors with high signal on T1-weighted MR imagings : chemical shift MR imaging and histologic correlation. *J. Comput. Assis. Tomogr.*, 15 : 762-769.
- 14) Solomon, I. 1955. Relaxation processes in a system of two spins. *Phys. Rev.*, 99 : 559-565.
- 15) Yamashita, T., Ohshima, H., Asanuma, T., Inukai, N., Miyoshi, I., Kasai, N., Kon, Y., Watanabe, T., Sato, F. and Kuwabara, M. 1996. The effects of  $\alpha$ -phenyl-*tert*-butyl nitron (PBN) on copper-induced rat fulminant hepatitis with jaundice. *Free Radic. Biol. Med.*, 21 : 755-761.