HOKKAIDO UNIVERSITY

Title	MECHANISMS OF CATECHOLAMINE SECRETION BY Ca^<2+>-REINTRODUCTION AND Na^+ REMOVAL IN ADRENAL CHROMAFFIN CELLS OF THE GUINEA-PIG
Author(s)	KANEKO, Manami
Citation	Japanese Journal of Veterinary Research, 46(2-3), 132-133
Issue Date	1998-11-30
Doc URL	https://hdl.handle.net/2115/2681
Type	departmental bulletin paper
File Information	KJ00003408014.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

The voltage-current relationship of this current resembled that of T-type Ca^{2+} current but not R-type Ca^{2+} current. However, the rapid inactivation, one of the properties of T-type Ca^{2+} current, was not observed in the present experiment.

6. This residual current, insensitive to Ca^{2+} channel blockers, was inhibited by low concentrations of Ni^{2+} (0.01–0.1 mM), and was not suppressed by ATP (500 μM).

7. Catecholamine release induced by 60 mM KCl was significantly inhibited by nifedipine (3 μM), ω-conotoxin MVIIC (3 μM) and mixture of four blockers, but neither ω-conotoxin GVIA (1 μM) nor ω-agatoxin IVA (0.1 μM).

8. These results suggest that guinea pig adrenal chromaffin cells possess L- and N- and P- and Q-type Ca^{2+} channels and other subtypes of Ca^{2+} channel insensitive to these blockers, and that L- and Q-type channels and the unidentified Ca^{2+} channels mainly contribute to the catecholamine release by stimulation with high K^+. Some properties of unidentified Ca^{2+} channel currents were different from those of both T-type and R-type Ca^{2+} channel currents. As we can not identify the subtype of this Ca^{2+} channel, further studies are required to identify these Ca^{2+} channels.

MECHANISMS OF CATECHOLAMINE SECRETION BY Ca^{2+}-REINTRODUCTION AND Na^+ REMOVAL IN ADRENAL CHROMAFFIN CELLS OF THE GUINEA-PIG

Manami Kaneko

Laboratory of Pharmacology,
Department of Biomedical Sciences,
School of Veterinary Medicine,
Hokkaido University, Sapporo 060–0818, Japan

1. We investigated the mechanisms of increases in catecholamine (CA) secretion evoked by reintroduction of Ca^{2+} after removal of extracellular Ca^{2+} (Ca^{2+}-reintroduction) and removal of extracellular Na^+ in chromaffin cells of the guinea-pig.

2. CA secretion was increased with increasing the concentration of Ca^{2+} (0.1–10 mM) reintroduced. This secretory response was not affected by 1 μM atropine, 100 μM hexamethonium and 1 μM tetrodotoxin.

3. In the presence of 1 mM $CoCl_2$ or 1 mM $MgCl_2$, Ca^{2+}-reintroduction failed to increase CA secretion. The secretory response to Ca^{2+}-reintroduction was significantly inhibited by 10 μM methoxyverapamil (D600) and 1 μM nifedipine, but not by 1 μM ω-conotoxin GVIA and 0.1 μM ω-agatoxin IVA, and was greatly potentiated by 1 μM Bay K 8644.

4. Ca^{2+}-reintroduction caused an increase in the intracellular free Ca^{2+} concentration ($[Ca^{2+}]_i$) which was inhibited by D600 in isolated chromaffin cells loaded with fura-2.

5. CA secretion evoked by high K^+ (56 mM) which directly depolarizes the cell membranes was also inhibited only by nifedipine, and was greatly potentiated by Bay K 8644. However, the rate of inhibition in CA secretion evoked by high K^+ was much smaller than that by Ca^{2+}-reintroduction.

6. CA secretion was evoked by the removal of extracellular Na^+ in the absence of Ca^{2+} but not in its presence. This secretory responce was also significantly inhibited by D600, and was

greatly potentiated by Bay K 8644.
7. These results suggest that CA secretion evoked by Ca^{2+}-reintroduction, high K^+ and the removal of extracellular Na^+ and Ca^{2+} is mainly mediated by Ca^{2+} entered through L-type voltage dependent Ca^{2+} channels in adrenal chromaffin cells.

EFFECTS OF TYROSINE KINASE INHIBITORS AND NON-SELECTIVE CATION CHANNEL BLOCKERS ON CAPACITATIVE Ca^{2+} ENTRY IN RAT ILEAL SMOOTH MUSCLES

Wakana Yasuda

Laboratory of Pharmacology,
Department of Biochemical Siences,
School of Veterinary Medicine,
Hokkaido University, Sapporo 060–0818, Japan

1. The present experiment was performed to examine the involvement of tyrosine kinase and non-selective cation channels in capacitative Ca^{2+} entry (CCE) in the rat ileal smooth muscles. The effects of tyrosine kinase inhibitors (genistein and tyrphostin 47), an inactive analogue of genistein (daidzein), non-selective cation channel blocker (SK & F96365) and Ca^{2+} entry blocker (tetrandrine) were examined in the presence of methoxyverapamil.
2. In the presence of external Ca^{2+}, carbachol-induced sustained contractions were dose-dependently inhibited by genistein, daidzein, tyrphostin 47, SK & F96365 and tetrandrine.
3. Under Ca^{2+}-free conditions, after the depletion of stored Ca^{2+} by carbachol or caffeine, the application of Ca^{2+} evoked transient contractions due to CCE. These contractions were inhibited by genistein, daidzein, tyrphostin 47, SK & F96365 and tetrandrine. The inhibitory potency of genistein was greater than that of daidzein. The application of Ca^{2+} evoked sustained contractions due to CCE after the depletion of stored Ca^{2+} with the treatment of thapsigargin. These five drugs also inhibited the contraction, but the potency of daidzein was greater than that of genistein.
4. SK & F96365 produced no inhibitory effects on the carbachol- and caffeine-induced contractions due to Ca^{2+} released from Ca^{2+} store. Genistein, daidzein and tyrphostin 47 inhibited these contractions. However, these drugs were less effective in inhibiting the contraction evoked by Ca^{2+} release than that by CCE. Tetrandrine inhibited contraction induced by carbachol but not caffeine.
5. Genistein slightly suppressed Ca^{2+}-induced contractions in β-escine treated skinned fibers.
6. These results suggest that CCE induced by carbachol and caffeine may be mediated by tyrosine kinase and this pathway is sensitive to SK & F96365 and tetrandrine. However, the inhibitory effects on CCE were produced by not only genistein but also daidzein, indicating that further studies are nesessary to evaluate this hypothesis.