



HOKKAIDO UNIVERSITY

Title	Production and characterization of monoclonal antibodies against tick-borne encephalitis virus Hokkaido strain
Author(s)	KOMORO, Kimiyo
Citation	Japanese Journal of Veterinary Research, 47(1-2), 93-93
Issue Date	1999-08-31
Doc URL	https://hdl.handle.net/2115/2777
Type	departmental bulletin paper
File Information	KJ00003408109.pdf



Production and characterization of monoclonal antibodies
against tick-borne encephalitis virus Hokkaido strain

Kimiyo Komoro

*Laboratory of Public Health,
Department of Environmental Veterinary Sciences,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo 060-0818, Japan*

Tick-borne encephalitis (TBE) patient was found at Kamiiso, Hokkaido in 1993, and a TBE virus (Oshima5-10) was isolated from blood of a sentinel dog in 1995.

To develop a diagnostic reagent to identify TBE virus, monoclonal antibodies (MAbs) were produced against TBE virus Oshima5-10 strain. Reaction patterns of MAbs were examined against several flaviviruses by indirect fluorescent antibody (IFA) test, hemagglutination-inhibition (HI) test, and neutralization (NT) test.

Results were summarized as follows :

1. Seven hybridomas producing MAbs against TBE virus Oshima5-10 strain were obtained using inactivated virus by formaldehyde as the immunogen. These MAbs were confirmed to react with viral protein by IFA test.
2. All MAbs were confirmed to react with

envelope protein of Oshima5-10 strain by Western blot analysis after immunoprecipitation.

3. Two MAbs 4H8 and 4A2 showed high antibody titer to 4 strains of tick-borne encephalitis virus group and Japanese encephalitis virus by IFA test or HI test, which suggests 4H8 and 4A2 to be flavivirus genus-specific.
4. MAb 1H4 showed high antibody titer to 4 strains of tick-borne encephalitis virus group by IFA, HI and NT tests, suggesting 1H4 to be TBE group-specific.
5. MAb 2F9 showed high antibody titer to Sofjin strain and Oshima5-10 strain, suggesting 2F9 to be Russian spring-summer encephalitis type-specific.
6. The other MAbs, 2A5, 4H1 and 5D10 did not show consistent reaction patterns to each virus strain.

Development of highly sensitive hantavirus genome detection methods
and analysis of the viral replication

Hiroki Tanabe

*Laboratory of Public Health,
Department of Environmental Veterinary Sciences,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo, Japan*

To analyze the Hantavirus replication cycle of plus-strand and minus-strand RNA in infected Vero E6 cells and newborn rats, we developed a Northern blot hybridization technique (NB) using digoxigenin labeled RNA probes to quantitatively

analyse the Hantavirus genome and develop a highly sensitive and strand specific RT-nested-PCR (SS-PCR).

The results are summarized as follows :

1. NB and SS-PCR could specifically detect either