



HOKKAIDO UNIVERSITY

| | |
|------------------|---|
| Title | Spectrum and prevalence of autosomal dominant spinocerebellar ataxia in Hokkaido, the northern island of Japan : a study of 113 Japanese families |
| Author(s) | Basri, Rehana; Yabe, Ichiro; Soma, Hiroyuki et al. |
| Citation | Journal of Human Genetics, 52(10), 848-855 https://doi.org/10.1007/s10038-007-0182-x |
| Issue Date | 2007-10 |
| Doc URL | https://hdl.handle.net/2115/28746 |
| Rights | The original publication is available at www.springerlink.com |
| Type | journal article |
| File Information | JHG52-10.pdf |



Spectrum and prevalence of autosomal dominant spinocerebellar ataxia in Hokkaido, the northern island of Japan: a study of 113 Japanese families

Rehana Basri^{1*}, Ichiro Yabe^{1*}, Hiroyuki Soma^{1*}, Hidenao Sasaki¹

¹Department of Neurology, Hokkaido University, Graduate School of Medicine, N15W7, Kita-Ku, Sapporo, 060-8368, Japan.

* These authors contributed equally to this work.

Corresponding author: Ichiro Yabe Tel: 81 117066028; Fax: +81 117005356.

Email address: yabe@med.hokudai.ac.jp

Abstract

The autosomal dominant cerebellar ataxia (ADCA) is a genetically heterogeneous group of neurodegenerative disorders. To shed further light on the clinical and genetic spectrum of ADCA in Japan, we conducted a study to determine the frequency of a new variety of different subtypes of SCAs among ADCA patients. This current study was carried out from April 1999 to December 2006 on the basis of patients with symptoms and signs of ADCA disorders. PCR and/or direct sequencing were evaluated in a total of 113 families. Among them, 35 families were found to have the mutation associated with SCA6, 30 with SCA3, 11 with SCA1, 5 with SCA2, 5 with DRPLA, and 1 with SCA14. We also detected the heterozygous -16C→T single nucleotide substitution within the puratrophin-1 gene responsible for 16q22.1-linked ADCA in 10 families. In this study, unusual varieties of SCA, including 27, 13, 5, 7, 8, 12, 17, and 16 were not found. Of the 113 patients, 14% had as yet unidentified ADCA mutations. The present study validates the prevalence of genetically distinct ADCA subtypes based on ethnic origin and geographical variation, and that 16q-linked ADCA has strong hereditary effects in patients with ADCAs in Japan.

Key words Autosomal dominant cerebellar ataxia, SCA6, SCA3, 16q-linked ADCA,

Introduction

Autosomal dominant cerebellar ataxia (ADCAs) are a genetically heterogeneous group of neurodegenerative disorders affecting the cerebellum, brainstem, spinal cord and cranial nerve nuclei resulting in progressive cerebellar ataxia of gait and limbs and variably associated with nystagmus, dysarthria, intention tremor and ophthalmoparesis. Disease onset is usually between 30 and 50 years of age, although early onset in childhood and onset in later decades after 60 years have been reported. Spinocerebellar ataxia (SCAs) are also genetically heterogeneous and the clinical diagnosis of subtypes of SCAs is complicated by the salient overlap of the phenotype variations. Harding classified ADCA into three major categories based on clinical features and dominant mode of inheritance (Harding 1993). Classification of these ADCA was modified by Duenas on the basis of neuropathology and provides a guideline in clinical practice to prioritize genetic tests for diagnoses (Duenas et al. 2006). ADCA type I is variably associated with ataxia of the gait, ophthalmoplegia, pyramidal and extrapyramidal signs, cognitive impairment, optic atrophy or peripheral neuropathy; ADCA type II with retinopathy; and ADCA type III with late onset, pure cerebellar ataxia with genetic heterogeneity. To date, 29 different genetic loci have been reported to be responsible for ADCA (<http://www.gene.ucl.ac.uk/cgi-bin/nomenclature>): Spinocerebellar ataxia type 1 (SCA1) on chromosome 6p22-p23 (Orr et al. 1993); SCA2 on 12q23-q24.1 (Imbert et al. 1996; Sanpei et al. 1996); Machado-Joseph disease (MJD)/SCA3 on 14q24.3-q32.1 (Kawaguchi Y et al. 1994); SCA4 on 16q22.1 (Flanigan et al. 1996); SCA5 on 11q13.2 (Ranum et al. 1994); SCA6 on 19p13.1 (Zhuchenko et al. 1997); SCA7 on 3p12-p13 (David et al. 1997); SCA8 on 13p (Koob et al. 1999); SCA10 on 22q13-qter. (Matsuura et al. 2000); SCA11 on 15q14-q21.3 (Worth et al. 1999); SCA12 on 5q31-33 (Holmes et al. 1999); SCA13 on 19q13.3-q13.4 (Waters et al.

2006); SCA14 on 19q13.42 (Yamashita et al. 2000; Chen et al. 2003; Yabe et al. 2003); SCA15 on 3p24.2 (Gardner et al. 2005); SCA16 on 8q23-q24.1 (Miyoshi et al. 2001); SCA17 on 6q27 (Nakamura et al. 2001); SCA18 on 7q22 (Brkanac et al. 2002); SCA19 on 1p21-q21 (Verbeek et al. 2002); SCA20 on 11p11.2-q13.3 (Knight et al. 2004); SCA21 on 7p21.3-p15.1 (Vuillaume et al. 2002); SCA22 on 1p21-q23 (Chung et al. 2003); SCA23 on 20p13-p12 (Verbeek et al. 2004); SCA24 on 1p26 (Swartz et al. 2002); SCA25 on 2p21-p15 (Stevanin et al. 2004); SCA26 on 19p13.3 (Yu et al. 2005); SCA27 on 13q33.1 (Van Swieten et al. 2003); SCA28 on 18p11.22-q11.2 (Cagnoli et al. 2006); SCA 29 on 3p26 (Dudding et al. 2004) and dentatorubral pallidoluysian atrophy (DRPLA) on 12p13.31 (Koide et al. 1994; Nagafuchi et al. 1994). Among these loci, causative genes have been further identified containing trinucleotide repeats (CAG) in SCA1, SCA2, SCA3, SCA6, SCA7, SCA17, DRPLA (Orr et al. 1993; Sanpei et al. 1996; Kawaguchi Y et al. 1994; Zhuchenko et al. 1997; David et al. 1997; Nakamura et al. 2001; Koide et al. 1994), and in the promoter region of *PPP2R2B* related to SCA12 (Holmes et al. 1999); trinucleotide repeats (CTG) within an untranslated region of the SCA8 gene (Koob et al. 1999); and a pentanucleotide (ATTCT) repeat expansion within intron 9 of the SCA10 gene (Matsuura et al. 2000). While the exact mechanism for repeat expansion is still unknown, a number of hypotheses have evolved to explain how the consequences of these expansions cause disease. A prime example is the expansion of CAG repeats in specific genes that lead to long abnormal polyglutamine (polyQ) tracts in the encoded proteins (Zoghi and Orr 2000). Polyglutamine-containing protein tends to aggregate in diseased neurons, forming characteristic nuclear and cytoplasmic inclusions (ubiquitin, proteasome, HSP70, transcription factors), which are the neuropathological hallmarks of these diseases (Ross and Poirier 2004).

On the other hand, deletion and/or point mutations were identified in the *SPTBN2*, *KCNC3*, *PRKCG*, and *FGF14* genes, recently attributed to SCA5, SCA13, SCA14, and SCA27, respectively (Ikeda et al. 2006; Waters et al. 2006; Yamashita et al. 2000; Chen et al. 2003; Yabe et al. 2003; Van Swieten et al. 2003). Point mutations may result in decreased stability of the proteins, and frameshift mutations in truncated proteins (Van Swieten et al. 2003; Dalski et al. 2005). More recently, single nucleotide substitutions have been identified in the puratrophin-1 and contactin 4 (*CNTN4*) genes, classified as 16q-linked ADCA and SCA16, respectively (Takashima et al. 2001; Lie et al. 2003; Mizusawa et al. 2004; Ishikawa et al. 2005; Miura et al. 2006). These two varieties of ADCA have been seen only in Japan.

In Japan, the prevalence of spinocerebellar degeneration including multiple system atrophy (MSA) is 15.68 in 100,000 (Sasaki et al. 2003). The frequency of ADCA subtypes also varies among countries. SCA3, SCA6 and DRPLA were higher in Japanese populations than in Caucasian populations (Maruyama et al. 2002; Sasaki et al. 2003; Matsumura et al 2003), whereas SCA7, SCA8, SCA17, and SCA12 are less frequent in the Japanese population (Maruyama et al. 2002; Sasaki et al. 2000; Matsumura et al 2003; Onodera 2000). SCA14 phenotype has been also reported (Yamashita et al. 2000; Yabe et al. 2003; Morita et al. 2006), but there are no reported cases of SCA27, SCA13, or SCA5. And 16q22.1-linked ADCA has been identified only in the Japanese population (Wieczorek et al. 2006). Interestingly, the molecular basis of such differences in the prevalence of SCAs within and among populations is still unclear. Moreover, the causes of approximately 20-40% of dominant SCAs are still unknown (Sasaki et al. 2003).

The breadth of the clinical spectrum of ADCA has been reported previously, but the relative frequencies of new varieties of different subtypes of ADCA have not been considered. To shed light on this problem, we performed a study of patients with clinically and genetically confirmed ADCA. We investigated the relative prevalences of different subtypes of repeat expansions and of the other new varieties of 16q22.1-linked ADCA, SCA13, SCA5, SCA27, and SCA16 in Japanese patients on Hokkaido, the northern most island of Japan.

Subjects and Methods

The Department of Neurology at Hokkaido University Graduate School of Medicine is located in Sapporo, the capital of Hokkaido. Patients with SCAs were referred by physicians to the Department for DNA diagnoses. The laboratory serves as the major molecular genetic center for neurological diseases throughout Hokkaido. Thus, patients visiting our clinic can be regarded as reflecting the relative prevalence of hereditary SCAs in Hokkaido.

Subjects with ADCA

Genetic analyses of 113 unrelated Japanese families with ADCA from different areas of Hokkaido Island were carried out in our laboratory between April 1999 and the end of December 2006. Although we have published the similar work, the study was done for the patients who were diagnosed by genetic analysis until March 1999 (Sasaki et al. 2000). Therefore, there is no overlap of the subjects between this study and the previous study. In all cases, progressive cerebellar ataxia was the cardinal clinical manifestation. Nearly all patients presented with an adult onset phenotype. In addition to neurological examination, brain MRIs showed cerebellar atrophy or pontocerebellar

atrophy. All procedures of the study were approved by the Hokkaido University Ethics Committee and informed consent was obtained from all participants.

Genetic analyses

Genomic DNA was extracted from blood lymphocytes by standard methods (Sambrook, 1989, Cold Spring Harbor, NY). Expanded triplet repeats in the genes causing SCA1, SCA2, SCA3, SCA6, SCA7, SCA8, SCA12, SCA17, and DRPLA were amplified using the primer pairs, Rep-1/Rep-2 (Orr et al. 1993), F-1/F-2 (Sanpei et al. 1996), MJD25/MJD52 (Kawaguchi et al. 1994), S-5-F1/ S-5-R1 (Zhunchenko et al. 1997), 4U1024/4U716 (David et al. 1997), SCA8-F3/SCA8-R4 (Koob et al. 1999), PPP2R2B-A/PPP2R2B-B (Holmes et al. 1999), TBP-F/TBP-R (Nakamura et al. 2001) and CTG-B37(699)/ (840) (Koide et al. 1994), respectively. We did not test for the SCA10 mutation because its clinical manifestation is ataxia with seizure, which was not observed in our ADCA families. Conditions for polymerase chain reaction (PCR) amplification were based on the respective original reports. All the PCR products containing triplets repeats were separated in 4% polyacrylamide /6M urea gels at 1680 V in 1· TBE buffer for 2 hours. Genotypes were determined using Gene Scan Analysis Ver.2.0 computer software (Perkin Elmer), as described previously (Yabe et al. 1998). For the new varieties of ADCA subtypes, we also performed PCR and sequence analyses in undiagnosed ADCA families to detect mutations in all exons of the following genes: fibrous growth factor14 (*FGF14*) for SCA27, spectrin (*SPTBN2*) for SCA5, protein kinase C γ (*PRKCG*) for SCA14, and voltage-gated potassium channel (*KCNC3*) for SCA13. In puratrophin-1 (*PRTPHN1/PLEKHG4*) for 16q-linked ADCA phenotype and contactin 4 (*CNTN4*) for SCA16, each published specific single nucleotide polymorphism was analyzed. The PCR mixtures and fragment analyses

were basically the same as those reported previously (Ishikawa et al. 2005; van Sweeten et al. 2003; Ikeda et al. 2006; Miura et al. 2006; Yabe et al. 2003; Waters et al. 2006).

Results

A total of 113 ADCA families were examined for 15 different mutations, and 97 families (86%) were positive for various subtypes of SCA. The SCA6 mutation was found in 35 of the 113 families (31%), SCA3 was detected in 30 families (27%), SCA1 in 11 families (10%), SCA2 in 5 families (4%), DRPLA in 5 families (4%), and SCA14 in 1 family (1%). The screen for point mutations in the puratrophin-1, *FGF14*, *SPTBN2*, *CNTN4* and *KCNC3* genes, revealed only the -16C→T mutation in the puratrophin-1 gene in 10 families (9%). The remaining 16 ADCA families (14%) were genetically unidentified. The frequency of known SCA mutations in the consecutive series of 113 families is shown in Fig.1. Table 1 summarizes the clinical features of the ten 16q-linked ADCA families and the remaining 16 unidentified ADCA families. The two groups of patients with 16q-linked ADCA and unidentified ADCA differed in their mean age at onset, with a higher onset in the 16qADCA patients (mean±SD age: 58±8 years) than in those with unidentified ADCA (mean±SD age: 32±13 years).

Discussion

Among the 113 families with ADCA examined in our department in Hokkaido from April 1999 to the end of December 2006, SCA6 (31%) was the most prevalent subtype, followed by SCA3/MJD (27%), SCA1 (10%), 16q-ADCA (9%), SCA2 (4%), and DRPLA (4%). This is in good agreement with a previous study also reporting that SCA6 (29%) and SCA3 (23%) were the most prevalent in the Hokkaido region (Sasaki

et al. 2000). Earlier reports from other districts also demonstrated that the frequencies of SCA6 and SCA3 were the most prevalent in Japan (Sasaki et al. 2003; Matsumura et al. 2003; Maruyama et al. 2002) presumably, due to the founder effect. SCA3 is quite frequent in many countries with different ethnic backgrounds, such as Portugal (80%), Germany (40%), Japan (40%), and France 30% (Storey et al. 2000). In the Nagano district of Japan, DRPLA was reported to be the second most frequent of the ADCA subtypes (10%) (Shimizu et al. 2004). There have also been a few reports of DRPLA from Korea (Jin et al. 1999), America (Brunetti-Pierri et al. 2006) and Canada (Espay et al. 2006). The expanded repeat ranges and the clinical features of the SCA6, SCA3, SCA1, SCA2, and DRPLA families were similar to those reported previously (data not shown) (Zhuchenko et al. 1997; Kawaguchi Y et al. 1994; Orr et al. 1993; Sanpei et al. 1996; Koide et al. 1994).

That we did not find any cases of SCA17, SCA12, SCA7, and SCA8 is in agreement with previous reports showing that the frequencies of these subtypes were not common in most of the districts of Japan (Maruyama et al. 2002; Sasaki et al. 2000; Matsumura et al. 2003; Onodera 2000). The SCA10 expansion appears to be restricted to a few families of Mexican (Matsuura et al. 2000) and Brazilian (Teive et al. 2004) origins. The SCA12 mutation has been identified only in a large pedigree of German decent and in an Indian family (Holmes et al. 1999; Bahl et al. 2005). These data clearly indicate that ethnic origin and geographical variation play important roles in the prevalence of SCA types.

Recently, a novel form of cerebellar ataxia linked to the 16q22.1 locus was reported (Takashima et al. 2001; Li et al. 2003; Mizusawa et al. 2004; Ishikawa et al. 2005). Several of these Japanese ADCA families also mapped within the SCA4 candidate

region, and the disease was reported as “16q-linked ADCA type III” (Nagaoka et al. 2000; Li et al. 2003). 16q-linked ADCA is the third most prevalent subtype of ADCA in Japan, after SCA6 and SCA3, and is distributed nation wide (Mizusawa et al 2004). In our study, the -16C→T change in the 5'UTR of the puratrophin-1 gene causing 16q22-linked ADCA diseases was found in 10 families from the Hokkaido island. Ishikawa also previously reported on the existence of 16q22-linked ADCA families from this island (Ishikawa et al. 2005). Thus, an accumulation of 16q-linked ADCA families leads to a moderately high prevalence of SCD in Hokkaido. 16q-linked ADCA was also very prevalent in ADCA families in the Nagano districts of Japan (Ohata et al. 2006). Table 2 summarizes the high prevalence of 16q-linked ADCA in various localities throughout Japan.

Clinically, 16q-ADCA shows a slowly progressive pure cerebellar ataxia of late-onset (onset > 50 years old), increased deep tendon reflexes, hearing impairment, and normal sensation and behavior. 16q22-linked ADCA demonstrates a strong founder effect in Japan (Hirano et al. 2004; Ishikawa et al. 2005). The spectrum of these 16q linked ADCA symptoms appears more commonly in the Japanese population, but was not detected in 537 European patients with unknown ADCA (Wieczorek et al. 2006). However, the Japanese 16q22.1-linked ADCA ataxia and the European SCA4 have the same locus, but the European SCA4 gene has not yet been identified. In addition to the clinical phenotypes of the Japanese patients, 16q-linked ADCA is clearly distinguishable from the SCA4 disorder. The neuropathology of 16q-linked ADCA type III was characterized by abnormal Purkinje cells in the cerebellum, including shrunken cell bodies, abnormal dendrites, somatic sprouts, and amorphous materials surrounding the cells, all indicative of degeneration processes (Owada et al. 2005).

The more recently discovered SCA14, SCA27, SCA13, SCA16, and SCA5 are distinguished by alterations in amino acid compositions and/or frameshift mutations within the *PRKCG* (protein kinase C gamma), *FGF14*, *KCNC3*, *CNTN4* and *SPTBN2* genes, respectively. We detected 1 family of SCA14 displaying axial myoclonus followed by ataxia at early onset; this family has already been described (Yabe et al. 2003) and other Japanese patients with SCA14 displayed slowly progressive cerebellar syndromes with late onset that included gait and limb ataxia, dysarthria, and saccadic pursuit. Brain MRI of these patients revealed atrophy of the cerebellum (Hiramoto et al. 2006; Morita et al. 2006). More recently, a Dutch family with SCA14 patients was also described with early onset, mildly generalized myoclonus (Vlak et al. 2006; Verbeek et al. 2005). SCA14 is present in various different ethnic populations (Japan, German, Dutch, Portuguese and French) and displays a heterogeneous mutation spectrum (Yabe et al. 2003; Dalski et al. 2006; Verbeek et al. 2005; Alonso et al. 2005; Klebe et al. 2005).

Several pathways might be interrupted to cause neurodegenerations and neuronal function loss, which remains unclear. In addition, we detected 16 families (14%) with unidentifiable types of ADCA. Thus, further studies are needed to improve genetic diagnoses, and epidemiological surveys and genetic counseling are needed for awareness and prevention of ADCA disease progress.

Acknowledgements

We are very grateful to all the patients and their families for their willingness to participate in this study. This work was partly supported by a Grant-in-Aid for "the Research Committee for Ataxic Diseases" of the Research on Measures for Intractable Diseases from the Ministry of Health, Welfare and Labour, Japan, by a Grant-in-Aid

for Scientific Research from the Ministry of Education, Science, Sports, and Culture,
Japan and by a Grant from the Kanae Foundation

References

Alonso I, Costa C, Gomes A, Ferro A, Seixas AI, Silva S, Cruz VT, Coutinho P, Sequeiros J, Silveira I (2005) A novel H101Q mutation causes PKC gamma loss in spinocerebellar ataxia type 14. *J Hum Genet* 50:523-9

Bahl S, Viridi K, Mittal U, Sachdeva MP, Kalla AK, Holmes SE, O'Hearn E, Margolis RL, Jain S, Srivastava AK, Mukerji M (2005) Evidence of a common founder for SCA12 in the Indian population. *Ann Hum Genet* 69:528-34

Brkanac Z, Fernandez M, Matsushita M, Lipe H, Wolff J, Bird TD, Raskind WH (2002) Autosomal dominant sensory/motor neuropathy with ataxia (SMNA): linkage to chromosome 7q22- q32. *Am J Med Genet* 114:450-457

Brunetti-Pierrri N, Wilfong AA, Hunter JV, Craigen WJ (2006) A severe case of dentatorubro-pallidoluysian atrophy (DRPLA) with microcephaly, very early onset of seizures, and cerebral white matter involvement. *Neuropediatrics* 37:308-11

Burk K, Zuhlke C, Konig IR, Ziegler A, Schwinger E, Globas C, Dichgans J, Hellenbroich Y (2004) Spinocerebellar ataxia type 5: clinical and molecular genetic features of a German kindred. *Neurology* 62:327-9

Cagnoli C, Mariotti C, Taroni F, Seri M, Brussino A, Michielotto C, Grisoli M, Di Bella D, Migone N, Gellera C, Di Donato S, Brusco A (2006) SCA28, a novel form of autosomal dominant cerebellar ataxia on chromosome 18p11.22-q11.2. *Brain* 129:235-42

Chen DH, Brkanac Z, Verlinde CL, Tan XJ, Bylenok L, Nochlin D, Matsushita M, Lipe H, Wolff J, Fernandez M, Cimino PJ, Bird TD, Raskind WH (2003) Missense

mutations in the regulatory domain of PKC gamma: a new mechanism for dominant nonepisodic cerebellar ataxia. *Am J Hum Genet* 72:839-49

Chung MY, Lu YC, Cheng NC, Soong BW (2003) A novel autosomal dominant spinocerebellar ataxia (SCA22) linked to chromosome 1p21-q23. *Brain* 126:1293–1299

Dalski A, Atici J, Kreuz FR, Hellenbroich Y, Schwinger E, Zuhlke C (2005) Mutation analysis in the fibroblast growth factor 14 gene: frameshift mutation and polymorphisms in patients with inherited ataxias. *Eur J Hum Genet* 13:118-20

Dalski A, Mitulla B, Burk K, Schattenfroh C, Schwinger E, Zuhlke C (2006) Mutation of the highly conserved cysteine residue 131 of the SCA14 associated PRKCG gene in a family with slow progressive cerebellar ataxia. *J Neurol.* 253:1111-2

David G, Abbas N, Stevanin G, Durr A, Yvert G, Cancel G, Weber C, Imbert G, Saudou F, Antoniou E, Drabkin H, Gemmill R, Giunti P, Benomar A, Wood N, Ruberg M, Agid Y, Mandel JL, Brice A (1997) Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. *Nat Genet* 17:65–70

Dudding TE, Friend K, Schofield PW, Lee S, Wilkinson IA, Richards RI (2004) Autosomal dominant congenital non-progressive ataxia overlaps with the SCA15 locus. *Neurology* 63:2288-92

Duenas AM, Goold R, Giunti P (2006) Molecular pathogenesis of spinocerebellar ataxias. *Brain* 129: 1357-1370

Espay AJ, Bergeron C, Chen R, Lang AE (2006) Rapidly progressive sporadic dentatorubral pallidoluysian atrophy with intracytoplasmic inclusions and no CAG repeat expansion. *Mov Disord* 21:2251-4

Flanigan K, Gardner K, Alderson K, Galster B, Otterud B, Leppert MF, Kaplan C, Ptacek LJ (1996) Autosomal dominant spinocerebellar ataxia with sensory axonal neuropathy (SCA4): clinical description and genetic localization to chromosome 16q22.1. *Am J Hum Genet* 59:392-399

Gardner RJ, Knight MA, Hara K, Tsuji S, Forrest SM, Storey E (2005) Spinocerebellar ataxia type 15. *Cerebellum* 4:47-50

Harding AE (1993) Clinical features and classification of inherited ataxias. *Adv Neurol* 61:1-14

Herman-Bert A, Stevanin G, Netter JC, Rascol O, Brassat D, Calvas P, Camuzat A, Yuan Q, Schalling M, Durr A, Brice A (2000) Mapping of spinocerebellar ataxia 13 to chromosome 19q13.3-q13.4 in a family with autosomal dominant cerebellar ataxia and mental retardation. *Am J Hum Genet* 67:229-235

Hirano R, Takashima H, Okubo R, Tajima K, Okamoto Y, Ishida S, Tsuruta K, Arisato T, Arata H, Nakagawa M, Osame M, Arimura K (2004) Fine mapping of 16q-linked autosomal dominant cerebellar ataxia type III in Japanese families. *Neurogenetics* 5:215-21

Hiramoto K, Kawakami H, Inoue K, Seki T, Maruyama H, Morino H, Matsumoto M, Kurisu K, Sakai N (2006) Identification of a new family of spinocerebellar ataxia type

14 in the Japanese spinocerebellar ataxia population by the screening of PRKCG exon 4.
Mov Disord 21:1355-60

Holmes SE, O'Hearn EE, McInnis MG, Gorelick-Feldman DA, Kleiderlein JJ,
Callahan C, Kwak NG, Ingersoll-Ashworth RG, Sherr M, Sumner AJ, Sharp AH,
Ananth U, Seltzer WK, Boss MA, Viera-Saecker AM, Epplen JT, Riess O, Ross CA,
Margolis RL (1999) Expansion of a novel CAG trinucleotide repeat in the 5' region of
PPP2R2B is associated with SCA12. Nat Genet 23:391–392

Ikeda Y, Dick KA, Weatherspoon MR, Gincel D, Armbrust KR, Dalton JC, Stevanin G,
Durr A, Zuhlke C, Burk K, Clark HB, Brice A, Rothstein JD, Schut LJ, Day JW,
Ranum LP (2006) Spectrin mutations cause spinocerebellar ataxia type 5. Nat Genet
38:184-90

Imbert G, Saudou F, Yvert G, Devys D, Trottier Y, Garnier JM, Weber C, Mandel JL,
Cancel G, Abbas N, Durr A, Didierjean O, Stevanin G, Agid Y, Brice A (1996) Cloning
of the gene for spinocerebellar ataxia 2 reveals a locus with high sensitivity to expanded
CAG/glutamine repeats. Nat Genet 14:285–291

Ishikawa K, Toru S, Tsunemi T, Li M, Kobayashi K, Yokota T, Amino T, Owada K,
Fujigasaki H, Sakamoto M, Tomimitsu H, Takashima M, Kumagai J, Noguchi Y,
Kawashima Y, Ohkoshi N, Ishida G, Gomyoda M, Yoshida M, Hashizume Y, Saito Y,
Murayama S, Yamanouchi H, Mizutani T, Kondo I, Toda T, Mizusawa H (2005) An
autosomal dominant cerebellar ataxia linked to chromosome 16q22.1 is associated with
a single-nucleotide substitution in the 5' untranslated region of the gene encoding a
protein with spectrin repeat and Rho guanine-nucleotide exchange-factor domains. Am
J Hum Genet 77:280-96

Jin DK, Oh MR, Song SM, Koh SW, Lee M, Kim GM, Lee WY, Chung CS, Lee KH, Im JH, Lee MJ, Kim JW, Lee MS (1999) Frequency of spinocerebellar ataxia types 1,2,3,6,7 and dentatorubral pallidoluysian atrophy mutations in Korean patients with spinocerebellar ataxia. *J Neurol* 246:207-10

Kawaguchi Y, Okamoto T, Taniwaki M, Aizawa M, Inoue M, Katayama S, Kawakami H, Nakamura S, Nishimura M, Akiguchi I, Kimura J, Narumia S, Kakizuka A (1994) CAG expansions in a novel gene for Machado–Joseph disease at chromosome 14q32.1. *Nat Genet* 8:221–228

Klebe S, Durr A, Rentschler A, Hahn-Barma V, Abele M, Bouslam N, Schols L, Jedynek P, Forlani S, Denis E, Dussert C, Agid Y, Bauer P, Globas C, Wullner U, Brice A, Riess O, Stevanin G (2005) New mutations in protein kinase Cgamma associated with spinocerebellar ataxia type 14. *Ann Neurol* 58:720-9

Knight MA, Gardner RJ, Bahlo M, Matsuura T, Dixon JA, Forrest SM, Storey E (2004) Dominantly inherited ataxia and dysphonia with dentate calcification: spinocerebellar ataxia type 20. *Brain* 127:1172–1181

Koide R, Ikeuchi T, Onodera O, Tanaka H, Igarashi S, Endo K, Takahashi H, Kondo R, Ishikawa A, Hayashi T, Saito M, Tomoda A, Miike T, Naito H, Ikuta F, Tsuji S. (1994) Unstable expansion of CAG repeat in hereditary dentatorubral pallidoluysian atrophy (DRPLA). *Nat Genet* 6:9–13

Koob MD, Moseley ML, Schut LJ, Benzow KA, Bird TD, Day JW, Ranum LP (1999) An untranslated CTG expansion causes a novel form of spinocerebellar ataxia (SCA8) *Nat Genet* 21:379–384

Li M, Ishikawa K, Toru S, Tomimitsu H, Takashima M, Goto J, Takiyama Y, Sasaki H, Imoto I, Inazawa J, Toda T, Kanazawa I, Mizusawa H (2003) Physical map and haplotype analysis of 16q-linked autosomal dominant cerebellar ataxia (ADCA) type III in Japan. *Hum Genet* 48:111-8

Maruyama H, Izumi Y, Morino H, Oda M, Toji H, Nakamura S, Kawakami H (2002) Difference in disease-free survival curve and regional distribution according to subtype of spinocerebellar ataxia: a study of 1,286 Japanese patients. *Am J Med Genet* 114:578-83

Matsumura R, Futamura N, Ando N, Ueno S (2003) Frequency of spinocerebellar ataxia mutations in the Kinki district of Japan. *Acta Neurol Scand* 107:38-41

Matsuura T, Yamagata T, Burgess DL, Rasmussen A, Grewal RP, Watase K, Khajavi M, McCall AE, Davis CF, Zu L, Achari M, Pulst SM, Alonso E, Noebels JL, Nelson DL, Zoghbi HY, Ashizawa T (2000) Large expansion of the ATTCT pentanucleotide repeat in spinocerebellar ataxia type 10. *Nat Genet* 26:191–194

Miura S, Shibata H, Furuya H, Ohyagi Y, Osoegawa M, Miyoshi Y, Matsunaga H, Shibata A, Matsumoto N, Iwaki A, Taniwaki T, Kikuchi H, Kira J, Fukumaki Y (2006) The *CNTN4* gene locus at 3p26 is a candidate gene of SCA16. *Neurology* 67:1236-41

Miyoshi Y, Yamada T, Tanimura M, Taniwaki T, Arakawa K, Ohyagi Y, Furuya H, Yamamoto K, Sakai K, Sasazuki T, Kira J (2001) A novel autosomal dominant spinocerebellar ataxia (SCA16) linked to chromosome 8q22.1–24.1. *Neurology* 57: 96–100

Mizusawa H, Ishikawa K, Toru S, Li M (2004) The prevalence and clinical features of 16q-linked autosomal dominant cerebellar ataxia (ADCA) type III in Japan. Annual report of the research committee of ataxic disease, Research on specific disease. The Ministry of Health, Labour, and Welfare of Japan, pp 67–70 (In Japanese)

Morita H, Yoshida K, Suzuki K, Ikeda S (2006) A Japanese case of SCA14 with the Gly128Asp mutation. *J Hum Genet* 51:1118-21

Nagafuchi S, Yanagisawa H, Ohsaki E, Shirayama T, Tadokoro K, Inoue T, Yamada M (1994) Structure and expression of the gene responsible for the triplet repeat disorder, dentatorubral and pallidolusian atrophy (DRPLA). *Nat Genet* 8:177–182

Nagaoka U, Takashima M, Ishikawa K, Yoshizawa K, Yoshizawa T, Ishikawa M, Yamawaki T, Shoji S, Mizusawa H (2000) A gene on SCA4 locus causes dominantly inherited pure cerebellar ataxia. *Neurology* 54:1971-5

Nakamura K, Jeong SY, Uchihara T, Anno M, Nagashima K, Nagashima T, Ikeda S, Tsuji S, Kanazawa I (2001) SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. *Hum Mol Genet* 10:1441–8

Nozaki H, Ikeuchi T, Kawakami A, Kimura A, Koide R, Tsuchiya M, Nakamura Y, Mutoh T, Yamamoto H, Nakao N, Sahashi K, Nishizawa M, Onodera O (2007) Clinical and genetic characterizations of 16q-linked autosomal dominant spinocerebellar ataxia (AD-SCA) and frequency analysis of AD-SCA in the Japanese population. *Mov Disord* 22:857-62

Ohata T, Yoshida K, Sakai H, Hamanoue H, Mizuguchi T, Shimizu Y, Okano T, Takada F, Ishikawa K, Mizusawa H, Yoshiura K, Fukushima Y, Ikeda S, Matsumoto N (2006) A -16C>T substitution in the 5' UTR of the puratrophin-1 gene is prevalent in autosomal dominant cerebellar ataxia in Nagano. *J Hum Genet* 51:461-6

Onodera Y, Aoki M, Mizuno H, Warita H, Shiga Y, Itoyama Y (2006) Clinical features of chromosome 16q22.1 linked autosomal dominant cerebellar ataxia in Japanese. *Neurology* 67:1300-2

Onodera Y, Aoki M, Tsuda T, Kato H, Nagata T, Kameya T, Abe K, Itoyama Y (2000) High prevalence of spinocerebellar ataxia type 1 (SCA1) in an isolated region of Japan. *J Neurol Sci* 178:153-8

Orr HT, Chung MY, Banfi S, Kwiatkowski TJ Jr, Servadio A, Beaudet AL, McCall AE, Duvick LA, Ranum LP, Zoghbi HY (1993) Expansion of an unstable trinucleotide CAG repeat in spinocerebellar ataxia type 1 *Nat Genet* 4:221–226

Ouyang Y, Sakoe K, Shimazaki H, Namekawa M, Ogawa T, Ando Y, Kawakami T, Kaneko J, Hasegawa Y, Yoshizawa K, Amino T, Ishikawa K, Mizusawa H, Nakano I, Takiyama Y (2006) 16q-linked autosomal dominant cerebellar ataxia: a clinical and genetic study. *J Neurol Sci* 247:180-6

Owada K, Ishikawa K, Toru S, Ishida G, Gomyoda M, Tao O, Noguchi Y, Kitamura K, Kondo I, Noguchi E, Arinami T, Mizusawa H (2005) A clinical, genetic, and neuropathologic study in a family with 16q-linked ADCA type III. *Neurology* 65:629-32

Ranum LP, Schut LJ, Lundgren JK, Orr HT, Livingston DM (1994) Spinocerebellar ataxia type 5 in a family descended from the grandparents of President Lincoln maps to chromosome 11. *Nat Genet* 8:280–284

Ross CA, Poirier MA (2004) Protein aggregation and neurodegenerative disease. *Nat Med* 10: 10–7

Sanpei K, Takano H, Igarashi S, Sato T, Oyake M, Sasaki H, Wakisaka A, Tashiro K, Ishida Y, Ikeuchi T, Koide R, Saito M, Sato A, Tanaka T, Hanyu S, Takiyama Y, Nishizawa M, Shimizu N, Nomura Y, Segawa M, Iwabuchi K, Eguchi I, Tanaka H, Takahashi H, Tsuji S (1996) Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT. *Nat Genet* 14:277–284

Sambrook J, Fritsh EF, Maniatis T (1989) *Molecular cloning-a laboratory manual*, 2nd edn. Cold Spring Harbor laboratory press, Cold Spring Harbor NY

Sasaki H, Yabe I, Tashiro K (2003) The hereditary spinocerebellar ataxias in Japan. *Cytogenet Genome Res* 100:198–205

Sasaki H, Yabe I, Yamashita I, Tashiro K (2000) Prevalence of triplet repeat expansion in ataxia patients from Hokkaido, the northernmost island of Japan. *J Neurol Sci* 175:45-51

Shimizu Y, Yoshida K, Okano T, Ohara S, Hashimoto T, Fukushima Y, Ikeda S (2004) Regional features of autosomal-dominant cerebellar ataxia in Nagano: clinical and molecular genetic analysis of 86 families. *J Hum Genet* 49:610-6

- Stevanin G, Herman A, Brice A, Durr A (1999) Clinical and MRI findings in spinocerebellar ataxia type 5. *Neurology* 53:1355-7
- Stevanin G, Bouslam N, Thobois S, Azzedine H, Ravaux L, Boland A, Schalling M, Broussolle E, Durr A, Brice A (2004) Spinocerebellar ataxia with sensory neuropathy (SCA25) maps to chromosome 2p. *Ann Neurol* 55:97–104
- Storey E, du Sart D, Shaw JH, Lorentzos P, Kelly L, McKinley Gardner RJ, Forrest SM, Biros I, Nicholson GA (2000) Frequency of spinocerebellar ataxia types 1, 2, 3, 6, and 7 in Australian patients with spinocerebellar ataxia. *Am J Med Genet* 95:351-7
- Swartz BE, Burmeister M, Somers JT, Rottach KG, Besselova IN, Leigh RJ (2002) A form of inherited cerebellar ataxia with saccadic intrusions, increased saccadic speed, sensory neuropathy, and myoclonus. *Ann N Y Acad Sci* 956:441–444
- Takashima M, Ishikawa K, Nagaoka U, Shoji S, Mizusawa H (2001) A linkage disequilibrium at the candidate gene locus for 16q-linked autosomal dominant cerebellar ataxia type III in Japan. *J Hum Genet* 46:167–171
- Teive HA, Roa BB, Raskin S, Fang P, Arruda WO, Neto YC, Gao R, Werneck LC, Ashizawa T (2004) Clinical phenotype of Brazilian families with spinocerebellar ataxia 10. *Neurology* 63:1509-12
- van Swieten JC, Brusse E, de Graaf BM, Krieger E, van de Graaf R, de Koning I, Maat-Kievit A, Leegwater P, Dooijes D, Oostra BA, Heutink P (2003) A mutation in the fibroblast growth factor 14 gene is associated with autosomal dominant cerebellar ataxia *Am J Hum Genet* 72:191-9

Verbeek DS, Schelhaas JH, Ippel EF, Beemer FA, Pearson PL, Sinke RJ (2002) Identification of a novel SCA locus (SCA19) in a Dutch autosomal dominant cerebellar ataxia family on chromosome region 1p21-q21. *Hum Genet* 111:388–393

Verbeek DS, Warrenburg BP, Hennekam FA, Dooijes D, Ippel PF, Verschuuren-Bemelmans CC, Kremer HP, Sinke RJ (2005) Gly118Asp is a SCA14 founder mutation in the Dutch ataxia population. *Hum Genet* 117:88-91

Verbeek DS, van de Warrenburg BP, Wesseling P, Pearson PL, Kremer HP, Sinke RJ (2004) Mapping of the SCA23 locus involved in autosomal dominant cerebellar ataxia to chromosome region 20p13-12.3. *Brain* 127:2551-7

Vlak MH, Sinke RJ, Rabelink GM, Kremer BP, van de Warrenburg BP (2006) Novel PRKCG/SCA14 mutation in a Dutch spinocerebellar ataxia family: expanding the phenotype. *Mov Disord* 21:1025-8

Vuillaume I, Devos D, Schraen-Maschke S, Dina C, Lemainque A, Vasseur F, Bocquillon G, Devos P, Kocinski C, Marzys C, Destee A, Sablonniere B (2002) A new locus for spinocerebellar ataxia (SCA21) maps to chromosome 7p21.3-p15.1. *Ann Neurol* 52:666–670

Waters MF, Minassian NA, Stevanin G, Figueroa KP, Bannister JP, Nolte D, Mock AF, Evidente VG, Fee DB, Muller U, Durr A, Brice A, Papazian DM, Pulst SM (2006) Mutations in voltage-gated potassium channel KCNC3 cause degenerative and developmental central Nervous system phenotypes. *Nat Genet* 38:447-51

Waters MF, Fee D, Figueroa KP, Nolte D, Muller U, Advincula J, Coon H, Evidente VG, Pulst SM (2005) An autosomal dominant ataxia maps to 19q13: Allelic heterogeneity of SCA13 or novel locus? *Neurology* 65:1111-3

Wieczorek S, Arning L, Alheite I, Epplen JT (2006) Mutations of the puratrophin-1 (PLEKHG4) gene on chromosome 16q22.1 are not a common genetic cause of cerebellar ataxia in a European population. *J Hum Genet* 51:363-7

Worth PF, Giunti P, Gardner-Thorpe C, Dixon PH, Davis MB, Wood NW (1999) Autosomal dominant cerebellar ataxia type III: linkage in a large British family to a 7.6-cM region on chromosome 15q14–21.3. *Am J Hum Genet* 65:420–426

Yabe I, Sasaki H, Chen DH, Raskind WH, Bird TD, Yamashita I, Tsuji S, Kikuchi S, Tashiro K (2003) Spinocerebellar ataxia type 14 caused by a mutation in protein kinase C gamma. *Arch Neurol* 12:1749-51

Yabe I, Sasaki H, Matsuura T, Takada A, Wakisaka A, Suzuki Y, Fukazawa T, Hamada T, Oda T, Ohnishi A, Tashiro K (1998) SCA6 mutation analysis in a large cohort of the Japanese patients with late-onset pure cerebellar ataxia. *J Neurol Sci* 156:89–95

Yamashita I, Sasaki H, Yabe I, Fukazawa T, Nogoshi S, Komeichi K, Takada A, Shiraishi K, Takiyama Y, Nishizawa M, Kaneko J, Tanaka H, Tsuji S, Tashiro K (2000) A novel locus for dominant cerebellar ataxia (SCA14) maps to a 10.2-cM interval flanked by D19S206 and D19S605 on chromosome 19q13.4-qter. *Ann Neurol* 48:156-63

Yu GY, Howell MJ, Roller MJ, Xie TD, Gomez CM (2005) Spinocerebellar ataxia type 26 maps to chromosome 19p13.3 adjacent to SCA6. *Ann Neurol* 57:349-54

Zhuchenko O, Bailey J, Bonnen P, Ashizawa T, Stockton DW, Amos C, Dobyns WB, Subramony SH, Zoghbi HY, Lee CC (1997) Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the alpha 1A voltage-dependent calcium channel. *Nat Genet* 15:62–69

Zoghbi HY, Orr HT (2000) Glutamine repeats and neurodegeneration. *Annu Rev Neurosci* 23: 217–47

Legends

Figure 1. The Frequency of different subtypes of ADCA in 113 families from April 1999 to December 2006

Table 1. Clinical features of patients with 16q-ADCA and other unknown ADCAs

ADCA: Autosomal dominant cerebellar ataxia

Table 2. Frequencies of 16q-linked ADCA type III in Japan

Fig. 1




Table 1

| | 16q-ADCA | Unknown ADCA |
|---------------------------------|--------------|--------------|
| Number of families | 10 (M=4,F=6) | 16 (M=9,F=7) |
| Family history | AD | AD |
| Mean age at examination (range) | 66.9 (52-76) | 53.0 (30-80) |
| Mean age at onset (range) | 58.0 (45-66) | 32.4 (3-55) |
| Disease duration (years) | 9.1 (2-19) | 20.2 (7-60) |
| Neurological findings | | |
| Dysarthria | 100% | 100% |
| Gaze nystagmus | 100% | 38% |
| Limb ataxia | 100% | 100% |
| Truncal ataxia | 100% | 100% |
| Hyporeflexia | 10% | 26% |
| Hyperreflexia | 0% | 43% |
| Spasticity | 0% | 38% |
| Babinski/chaddock reflexes | 0% | 43% |
| Hearing impairment | 20% | 0% |
| Decrease vibration sence | 20% | 43% |
| Dystonia | 0% | 17% |
| Dementia | 0% | 13% |

Table 2

| Location | Number of SCA families | 16q-linked ADCA type III | References |
|--|-------------------------------------|---|-----------------------|
| Kagoshima and southern Miyazaki, the Southern Kyushu | 30 families | 20 case/ 4 families | Hirano R et al. 2004 |
| Throughout Japan | 52 families | 109 case/52 families | Ishikawa K et al.2005 |
| Tokyo | 1 family | 12case/ 1 family | Owada k et al.2005 |
| Tohoku, Northernmost part of Honshu Island | 218 families plus 128 sporadic case | 31 case,7 case/ 26 families | Onodera Y et al. 2006 |
| Nagano | 67 families | 63 case/51 family | Ohata T et al.2006 |
| Tochigi, a northern region of Kanto Plane | 24 families | 20cases/9 families, plus 2 sporadic cases | Ouyang Y et al.2006 |
| Honshu Island | 686 families | 65 cases/ 57 families | Nozaki H et al.2007 |
| Hokkaido | 113 families | 10cases/ 10 families | Present study |