



# HOKKAIDO UNIVERSITY

Title	Genetic and biological comparision of tick-borne encephalitis viruses from Hokkaido and Far-Eastern Russia
Author(s)	GOTO, Akiko; HAYASAKA, Daisuke; YOSHII, Kentarou et al.
Citation	Japanese Journal of Veterinary Research, 49(4), 297-307
Issue Date	2002-02-28
DOI	<a href="https://doi.org/10.14943/jjvr.49.4.297">https://doi.org/10.14943/jjvr.49.4.297</a>
Doc URL	<a href="https://hdl.handle.net/2115/2927">https://hdl.handle.net/2115/2927</a>
Type	departmental bulletin paper
File Information	KJ00002400405.pdf



## Genetic and biological comparison of tick-borne encephalitis viruses from Hokkaido and Far-Eastern Russia

Akiko Goto, Daisuke Hayasaka, Kentarou Yoshii, Tetsuya Mizutani, Hiroaki Kariwa and Ikuo Takashima\*

(Accepted for publication : January 31, 2002)

### Abstract

We compared the biological properties of Oshima 5-10 (tick-borne encephalitis [TBE] virus isolated in Hokkaido, Japan) and Sofjin-HO (Far-Eastern subtype TBE virus) including plaque formation, virus replication and virus protein synthesis in BHK-21 cell cultures to reveal strain differences. We also determined the complete nucleotide sequences of both strains and compared the deduced amino acid sequences. Plaques of Oshima 5-10 were smaller than those of Sofjin-HO. Virus titers in culture fluid of Oshima 5-10 were 1/100 of those of Sofjin-HO at 9 and 12 hr after infection. Less viral protein and RNA syntheses of strain Oshima 5-10 was observed than with Sofjin-HO. Genetic analysis revealed 1.4% of amino acids to differ with Sofjin-HO. No difference between the two strains was detected in the motif sequence of the viral enzyme, cleavage sites of viral protein or glycosylation sites of NS1.

Key words : Tick-borne encephalitis virus, flavivirus, genome sequence, biological property

### Introduction

Tick-borne encephalitis (TBE) virus is a member of the genus *Flavivirus* in the family *Flaviviridae* and causes a fatal encephalitis in humans. TBE virus is a single-stranded

positive-polarity enveloped RNA virus. The RNA genome is about 11kb in length. A single large open reading frame encodes three structural proteins (core [C], membrane[prM] and envelope [E]) and seven nonstructural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4

---

Laboratory of Public Health, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, 060-0818, Japan

\*Corresponding author :

Ikuo Takashima, D. V. M., Ph. D

Graduate School of Veterinary Medicine, Hokkaido University, Kita-18 Nishi-9, Kita-ku, Sapporo, Hokkaido 060-0818, Japan.

Fax : +81-1-11-706-5211, E-mail : takashima@vetmed.hokudai.ac.jp

B and NS5). The 5' and 3' end of the viral genome is a non-coding region (NCR)<sup>6,29</sup>.

E protein is associated with the virus envelope and mediates both receptor binding and fusion, therefore, E protein is assumed to be an important determinant of virulence and pathogenicity<sup>16-19,27</sup>. NS3 protein contains motifs for serine protease<sup>3,10,25</sup>, nucleotide triphosphatase<sup>41</sup>, and helicase<sup>11</sup>. NS5 protein contains motifs for methyltransferase<sup>24</sup> and RNA-dependent RNA polymerase<sup>23,32</sup>. NS2B protein makes a complex with NS3 protein, and NS1, NS2A and NS4A proteins are assumed to be involved in RNA replication complex<sup>21,22</sup>.

Based on phylogenetic analysis, TBE virus can be divided into three subtypes, Far-Eastern, European and Siberian<sup>2,8,13,14</sup>. The Far-Eastern subtype is known as the Russian spring summer encephalitis (RSSE) virus and prevalent in Far Eastern Russia. It is known to be highly pathogenic in humans, with a mortality rate of 5 to 20%<sup>8</sup>.

In October 1993, a human case of TBE was reported in Kamiiso, Hokkaido, where TBE had not been reported as endemic. The virus strain, Oshima 5-10, was isolated from sentinel dogs in the same area in 1995, and identified as a TBE virus of the Far-Eastern subtype from an analysis of antigen and viral envelope (E) protein gene nucleotide sequence<sup>38</sup>. In Far East Russia, the annual number of TBE patients is more than 500 and fatality rates are around 15% each year<sup>8</sup>. In contrast, very few cases of TBE and no fatal cases had been reported in Hokkaido. Heterogenous strains with different levels of virulence are reportedly distributed in endemic areas. Our previous study showed that Oshima 5-10 strain was less virulent and replicated slower than another TBE virus, strain Sofjin-HO (the prototype of the Far-Eastern subtype) in a mouse model<sup>7</sup>.

In this study, we compared the biological properties of the two strains including plaque formation, virus replication, and virus genome RNA and protein syntheses in BHK - 21 cell cultures to clarify strain differences. We also determined the complete nucleotide sequences of both strains and compared the deduced amino acid sequences. We discuss the correlation between biological and genetic properties.

## Materials and Methods

### *Cells lines and viral strains*

The baby hamster kidney (BHK - 21 ; ICN Biomedicals, Aurora, OH) cell line was grown at 37°C in Eagle's minimum essential medium (MEM ; Nissui Pharmaceutical, Tokyo, Japan) containing fetal calf serum (FCS ; ICN Biomedicals). The strain of TBE virus used were Oshima 5-10 (Far-Eastern subtype TBE virus isolated from dog's blood in Hokkaido, Japan)<sup>38</sup>, and Sofjin-HO (RSSE virus ; the prototype of the Far-Eastern subtype of TBE virus)<sup>42</sup>. Langat virus TP-21 strain, tick-borne flavivirus isolated from ticks in Malaysia in 1956<sup>12</sup> was used to obtain immune mouse ascites. The brains of virus-inoculated one-day old suckling mice were homogenized and diluted into a 1% suspension in phosphate-buffered saline (PBS ; pH7.6) containing 10% FCS and stored at - 80°C for use as virus stock. The infectious virus titer was assayed by the focus count assay described previously<sup>20</sup>. Briefly, monolayers of BHK - 21 cells grown in 96 - well plates were inoculated with serially diluted virus. After 38 hours at 37°C, foci of virus on the monolayer were visualized by immunohistochemical staining by using the peroxidase-antiperoxidase procedure.

### *Plaque forming assay*

Monolayers of BHK - 21 cells grown in 6 -

well plates were infected with serially diluted virus in culture medium. After 90 minutes at 37°C, cells were washed twice in PBS and were overlaid with 2 ml of MEM containing 10% FCS and 7% agarose (SEA KEM ME : FMC BioProducts, Rockland, ME). At 96 hr post-infection (p. i.), the cells were stained with 2 ml of MEM containing 10% FCS, 7% agarose and 5% neutral red. After incubation for 24 hr, the plaque morphology was observed.

#### *Growth curve*

Monolayers of BHK-21 cells grown in 24-well plates were infected with virus in culture medium at a multiplicity of infection (m. o. i.) of 1 focus forming unit (FFU). After 90 min at 37°C, cells were washed twice with PBS and 1 ml of MEM containing 4% FCS was added. At 6, 9, 12, 18, 24 and 30 hr p. i., samples were harvested from the cell culture medium, stored at -80°C and titrated by focus count assay for quantification of the infectious virus titer.

#### *Western blotting*

Viral protein production was measured by Western blotting. Monolayers of BHK-21 cells grown in 35-mm-dishes were infected with virus in culture medium at an m. o. i. of 1 FFU. After 90 min at 37°C, cells were washed twice with PBS and 1 ml of MEM containing 4% FCS was added. At 6, 9 and 12 hr p. i., the cells were harvested and lysed with 200 µl of sodium dodecyl sulfate (SDS) buffer (62.5 mM Tris-HCl buffer (pH 6.8), 2% SDS, 5% 2-mercaptoethanol, 10% glycerol and 0.005% bromo-phenol blue). After heating for 2 min at 100°C, the samples (10 µl) were separated by 10% SDS-polyacrylamide gel electrophoresis (SDS-PAGE). For calculation of the molecular weights of proteins, a molecular standard was included in each gel.

Samples were transferred to nitrocellu-

lose membranes by electroblotting, the membranes were blocked with Block Ace (Dainippon Pharmaceutical Co., Ltd., Osaka, Japan), and the blotted membranes were incubated with anti-Langat virus immune mouse ascites (diluted of 1 : 200) in PBS. Bound antibodies were reacted with horseradish peroxidase-conjugated anti-mouse IgG goat antibody (Zymed Laboratories, Inc., South San Francisco, CA) as a second step, and bands were visualized with 2.5 mg/ml of 4-chloro-1-naphthol (SIGMA-Aldrich Co., St Louis, MO), 17% methanol and 0.25% H<sub>2</sub>O<sub>2</sub> in PBS.

#### *Northern blotting*

Digoxigenin (DIG) - labeled positive- or negative-sense RNA probes, encompassing the prM and E region of the Oshima-5-10 genome, were produced by *in vitro* transcription using a DIG RNA labeling kit (Roche Diagnostics, Basel, Switzerland).

Monolayers of BHK-21 cells grown in 35-mm dishes were infected with virus in culture medium at an m. o. i. of 1 FFU. At 5, 6, 7, 8 and 9 hr p. i., the cells were harvested and RNA was extracted using an IsoGen Kit (Nippon Gene, Tokyo, Japan). The RNA samples (10 µg) were denatured with glyoxal and dimethyl sulfoxide and used in Northern blot hybridization reactions which were performed as described previously<sup>35</sup>.

Briefly, pre-hybridization and hybridization were carried out in hybridization buffer (7% SDS, 50 mM sodium phosphate buffer (pH 7.2), 2% Blocking Reagent [Roche Diagnostics], 50% deionized formamide, 5×SSC (pH 7.0), 1 mM EDTA and 50 µg/ml yeast tRNA [Roche Diagnostics]) at 70°C. Both positive- and negative-sense DIG-labeled RNA probes were diluted to 0.2 µl/ml for use.

After hybridization, the membranes were blocked with blocking buffer (1% Blocking Reagent [Roche Diagnostics], 0.15 M NaCl and

0.1 M Tris-HCl [pH 7.5]), the bound probes were reacted with alkaline phosphatase-conjugated anti-DIG Fab fragments (Roche Diagnostics), and the bands were visualized with 2.5  $\mu$ M CDP-Star (Roche Diagnostics) in buffer containing 0.1 M Tris-HCl (pH 9.5), 0.1 M NaCl and 50 mM MgCl<sub>2</sub>.

#### *Genome sequence analysis*

The nucleotide sequences of the viral genome were determined by directly sequencing the RT-PCR products. Viral RNA was extracted from brains of virus-inoculated suckling mice with the Isogen Kit. For cDNA synthesis, RT-PCR was performed with a THERMOSCRIPT™ RT-PCR System Kit (Invitrogen, Carlsbad, CA). The cycle sequencing reaction was performed with the DNA Sequence Kit (Applied Biosystems, Foster city, CA), and the sequence was determined by fluorescence autosequencer (ABI PRISM 310 Genetic Analyser, Applied Biosystems). Nucleotide sequences and deduced amino acid sequences were aligned out with GENETYX-MAC ver. 10 (Software Development, Tokyo, Japan).

## Results

#### *Plaque forming assay*

To distinguish Oshima 5-10 from Sofjin-HO based on biological characteristics, we examined their plaque morphology in BHK - 21 cells (Fig. 1). BHK - 21 cells in 6 - well plates were infected with 50 FFU/well of each TBE virus and the plaques were observed at 120 hr p. i.. The plaques of Oshima 5-10 were smaller than those of Sofjin-HO.

#### *Growth curve*

To identify differences in viral replication between the two strains, the infectious virus titer of culture fluid harvested at various time points was assayed by the focus count method (Fig. 2). Virus titers in the culture fluid of Sofjin-HO increased at 9 hr p. i.. In contrast, those of Oshima 5-10 remained low until 9 hr p. i. and then increased at 12 hr p. i. to be 1/100 the level of Sofjin-HO. This result reveals that strain Oshima 5-10 replicates slower than strain Sofjin-HO.




Fig. 1 Morphology of plaques at 120 hr p. i. . BHK-21 cells were infected with Oshima 5-10(A) and Sofjin-HO (B).




Fig. 2. Growth curve of TBE virus. BHK-21 cells were infected at an m. o. i. of 1 FFU with either Oshima 5-10 (●) or Sofjin-HO (◇). The titer of virus released during each time period was determined by focus counting assay in BHK-21 cells.

#### Western blotting

Viral protein synthesis levels were compared between the two strains by Western blotting (Fig. 3). No viral protein was detected in samples from cells infected with either Oshima 5-10 or Sofjin-HO at 6 hr p. i.. At 9 hr p. i., parts of various viral proteins (E, NS1 and NS5) were detected from Sofjin-HO-infected cells, but not from Oshima 5-10-infected cells (Fig. 3). Furthermore, at 12 hr p. i., the viral proteins of Oshima 5-10 was less abundant than those of Sofjin-HO. The results show that the viral proteins of strain Oshima 5-10 in BHK-21 cells were produced later than those of strain Sofjin-HO.

#### Northern blotting

Northern blotting analysis was carried out to compare the viral RNA synthesis of Oshima 5-10 and Sofjin-HO (Fig. 4). To make


sure that DIG-labeled positive- or negative-sense RNA probes detect only complementary viral RNA of both senses, we carried out Northern blotting with synthesized non-labeled complementary RNA. The results showed that both probes could detect only the corresponding RNA (data not shown). We used both positive- and negative-sense RNA probes to detect viral RNA in infected BHK-21 cells, but found no negative-strand RNA (data not shown). Positive-strand viral RNA was detected in samples from cells infected with either strain from 5 to 9 hr p. i.. Less viral RNA of Oshima 5-10 than Sofjin-HO was produced in the cells (Fig. 4).

The result shows that the RNA synthesis of Oshima 5-10 was delayed in BHK-21 cells at an early stage, as compared with that of Sofjin-HO.


#### Genome sequence analysis

Nucleotide sequence analysis revealed that there were 11,100 nucleotides in the genome of Oshima 5-10 and 10,894 nucleotides in that of Sofjin-HO. This difference is caused by the shorter length of 3' NCR in Sofjin-HO as compared with Oshima 5-10<sup>15)</sup>. On the other hands, numbers of deduced amino acid were same (3414 a. a.) in both strains. Oshima 5-10 differed by a total of 46 amino acids (1.4%) from Sofjin-HO (Table 1). No difference between the two strains was detected in the motif of the NS3 serine protease<sup>3,10,25)</sup>/helicase<sup>11)</sup>, NS5 RNA-dependent RNA polymerase<sup>23,32)</sup>/methyltransferase<sup>24)</sup>, cleavage sites of viral protein<sup>1,6,9,36,37)</sup> and glycosylation sites of NS1<sup>6)</sup>.

Amino acid differences included 15 non-conservative changes which affect the characteristics of the amino acid such as polarity or hydrophobicity. NS3 protein has the most non-conservative changes (Table 1). In the 5' NCR, Oshima 5-10 and Sofjin-HO differed by



**Fig. 3** Viral protein synthesis. Samples were separated by 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Lane N: Negative Control (BHK-21 cells), P: Positive Control (BHK-21 cells infected with Oshima 5-10 48 hr p. i.), M: Marker. Samples were transferred to nitrocellulose membranes by electroblotting, and the blotted membranes were incubated with anti-Langat virus immune mouse ascites. Bound antibodies were reacted with horseradish peroxidase-conjugated anti-mouse IgG goat antibody as a second step, and bands were visualized with 2.5 mg/ml of 4-chloro-1-naphthol.



**Fig. 4** Analysis of viral RNA replication by detection of positive-strand viral RNA. RNA samples (10 µg) were denatured and used in Northern blot hybridization reactions which were performed as described previously<sup>35</sup>. Pre-hybridization and hybridization were carried out in hybridization buffer at 70°C. After the hybridization, the membranes were blocked with blocking buffer, the bound DIG-labeled probes were reacted with alkaline phosphatase-conjugated anti-DIG Fab fragments and bands were visualized with CDP-Star.

Oshima 5-10 5'	1	AGATTTTCTTGACGTCATGCGTTTGCTTCGGATAGCAACAGCAGCGAC	50
Sofjin-HO 5'	1	.....G.....	50
		*****	
Oshima 5-10 5'	51	AGGCTTGAGAGAGACAATCTTTTGTGGATCAGTCGTGAACGTGTTGAGA	100
Sofjin-HO 5'	51	...T.....CAC.....	100
		***.....	
Oshima 5-10 5'	101	AAAAGACAGCTTAGGAGAACAAGAGCTGGGG	131
Sofjin-HO 5'	101	.....	131
		*****	

**Fig. 5** Comparison of the 5' NCR of Oshima 5-10 and Sofjin-HO.

5nucleotides (Fig. 5).

**Discussion**

In this study, we compared the biological and genetic characteristics of the viral strains Oshima 5-10 and Sofjin-HO to reveal differences between TBE virus strains of the Far-Eastern subtype.

Oshima 5-10 strain replicated more slowly in BHK-21 cell cultures than did Sofjin-HO. Plaques of Oshima 5-10 were also smaller than those of Sofjin-HO (Fig. 1) and virus titers in culture fluid were 1/100 of those of Sofjin-HO at 9 and 12 hr after infection (Fig. 2).

Next, we examined viral protein and

Table 1 . Amino acid differences between Sofjin-HO and Oshima 5 - 10. \*

Compared protein	Amino acid differences (Oshima 5-10→Sofji-HO)	Total	Percentage identity
C	Arg-32→Gln* Met-43→Leu Ala-54→Val Asn-64→Lys* Val-111→Leu	5	95.5
prM	Ile-134→Met Val-145→Ala	2	98.8
E	Ala-153→Val Val-308→Met Met-462→Val Ala-463→Val	4	99.2
NS1	Leu-58→Val Val-72→Ala Phe-93→Leu Ile-138→Thr* Val-346→Ile	5	98.6
NS2A	Arg-52→Lys Ile-168→Thr* Ala-169→Val Val-225→Ile	4	98.3
NS2B		0	100
NS3	Asp-2→Gly* Phe-45→Ser* Ser-62→Tyr Val-63→Ile Val-68→Ala Gly-124→Arg* Ile-184→Thr* Thr-380→Ala* Gly-85→Ser*	9	98.6
NS4A	Ala-63→Gly	1	99.3
NS4B	Leu-24→Val Phe-88→Leu Ser-198→Ala* Ser-227→Gly*	4	98.4
NS5	Lys-15→Arg Arg-101→Lys Met-130→Val Tyr-519→His* Thr-634→Ser Phe-642→Leu Arg-677→Gly* Val-692→Ile Phe-778→Leu Gly-827→Ser* Val-831→Ala Arg-862→Lys	12	98.7
polyprotein		46	98.6

\*The sequences used in this analysis were as follows (GenBank numbers in parentheses) : Oshima 5-10 (AB062063), Sofjin-HO (AB062064).

RNA levels of both strains early in the infection when the difference was most evident. Protein production was delayed in Oshima 5-10 infected cells as compared with in Sofjin-HO infected cells (Fig. 3). Viral RNA was also detected later in strain Oshima 5-10 than Sofjin-HO (Fig. 4). The later onset of protein and RNA production in Oshima 5-10 may be related with the slower viral replication in BHK - 21 cells.

We also determined the complete nucleotide sequences of both strains and compared their deduced amino acid sequences. There were 5 nucleotides differences in the 5' NCR gene between Oshima 5-10 and Sofjin-HO (Fig. 5). There are reports that attenuation of flavivirus was caused by changes of nucleotides in the 5' or 3' NCR<sup>5, 28, 31, 33)</sup>. The 5' NCR is capped<sup>6)</sup> and the 5' and 3' NCR make secondary structure and have the inverted complementary repeat sequences which are considered to play an important role in viral transcription<sup>13)</sup>. But in this study, there was no difference in those regions between the two strains and we could not conclude that above mentioned 5 nucleotide differences had influence on virus replication.

Oshima 5-10 possessed a total of 46 amino acids (1.4%) that differed from Sofjin-HO (Table 1). There are reports that attenuation of flavivirus was caused by changes of amino acids in the motif sequence of the NS proteins<sup>4, 27, 40)</sup>, cleavage sites of viral protein<sup>1, 9, 16, 36)</sup> and glycosylation sites of NS1<sup>30)</sup>. Particularly, a lack of motif sequence of NS3 serine protease<sup>3, 10, 25)</sup> / helicase<sup>11)</sup>, NS5 RNA-dependent RNA polymerase<sup>23, 32)</sup> / methyltransferase<sup>24)</sup> is lethal. But in this study, there was no difference in those regions between the two strains.

The 46 amino acid changes included 15 non-conservative types. In flavivirus, E protein is assumed to be an important determinant of virulence and pathogenicity, because

it is associated with the viral envelope and mediates both receptor binding and fusion<sup>16-19, 27, 29, 34)</sup>. However, sequence analyses of the E protein gene of Oshima 5-10 showed high identity with Sofjin-HO and the amino acid differences between the two strains did not include non-conservative changes<sup>15)</sup>. Therefore, other proteins may influence the biological characteristics and virulence. But we could not correlate these amino acid differences with the biological differences between the strains.

We demonstrated that in BHK - 21 cells, Oshima 5-10 replicated at a more slowly pace and produced protein and RNA later than Sofjin-HO. The results encourage further study to correlate the biological, genetic and pathogenetic characteristics of the two strains for understanding the virulence of the TBE virus.

#### References

- 1) Amberg, S. M. and Rice, C. M. 1999. Mutagenesis of the NS 2 B-NS 3-mediated cleavage site in the flavivirus capsid protein demonstrates a requirement for coordinated processing. *J. Virol.*, 73 : 8083-8094.
- 2) Bakhvalova, V. N., Rar, V. A., Tkachev, S. E., Matveev, V. A., Matveev, L. E., Karavanov, A. S., Dobrotvorsky, A. K. and Morozova, O. V. 2000. Tick-borne encephalitis virus strains of Western Siberia. *Virus Res.*, 70 : 1-12.
- 3) Bazan, J. F. and Fletterick, R. J. 1989. Detection of a trypsin-like serine protease domain in flaviviruses and pestiviruses. *Virology* 171 : 637-639.
- 4) Butrapet, S., Huang, C. Y. H., Pierro, D. J., Bhamarapravati, N., Gubler, D. J. and Kinney, R. M. 2000. Attenuation markers of a candidate dengue type 2 vaccine virus, strain 16681 (PDK - 53),

- are defined by mutations in the 5' non-coding region and nonstructural proteins 1 and 3. *J. Virol.*, 74 : 3011-3019.
- 5) Cahour, A., Pletnev, A., Vazeille-Falcoz, M., Rosen, L. L. and Lai, C. 1995. Growth-restricted dengue virus mutants containing deletions in the 5' non-coding region of the RNA genome. *Virology*, 207 : 68-76.
  - 6) Chambers, T. J., Hahn, C. S., Galler, R. and Rice, C. M. 1990. Flavivirus genome organization, expression, and replication. *Annu. Rev. Microbiol.*, 44 : 649-688.
  - 7) Chiba, N., Iwasaki, T., Mizutani, T., Kariwa, H., Kurata, T. and Takashima, I. 1999. Pathogenicity of tick-borne encephalitis virus isolated in Hokkaido, Japan in mouse model. *Vaccine*, 17 : 779-787.
  - 8) Dumpis, U., Crook, D. and Oksi, J. 1999. Tick-borne encephalitis (Review). *Clin. Infect. Dis.*, 28 : 882-890.
  - 9) Falgout, B. and Markoff, L. 1995. Evidence that flavivirus NS1 - NS2A cleavage is mediated by a membrane-bound host protease in the endoplasmic reticulum. *J. Virol.*, 69 : 7232-7243.
  - 10) Gorbalenya, A. E., Donchenko, A. P., Koonin, E. V. and Blinov, V. M. 1989. N-terminal domains of putative helicases of flavi- and pestiviruses may be serine proteases. *Nuc. Acids Res.*, 17 : 3889-3897.
  - 11) Gorbalenya, A. E., Koonin, E. V., Donchenko, A. P. and Blinov, V. M. 1989. Two related superfamilies of putative helicases involved in replication, recombination, repair and expression of DNA and RNA genomes. *Nuc. Acids Res.*, 17 : 4713-4729.
  - 12) Gordon Smith, C. E. 1956. A virus resembling Russian spring-summer encephalitis virus from an *Ixodid* in Malaya. *Nature*, 178 : 581-582.
  - 13) Gritsun, T. S., Venugopal, K., Zanotto, P. M. de A., Mikhailov, M. V., Sall, A. A., Holmes, E. C., Polkinghorne, I., Frolova, T. V., Pogodina, V. V., Lashkevich, V. A. and Gould, E. A. 1997. Complete sequence of tick-borne flaviviruses isolated from Siberia and the UK: analysis and significance of the 5' and 3' -UTRs. *Virus Res.*, 49 : 27-39.
  - 14) Hayasaka, D., Ivanov, L., Leonova, G., Goto, A., Yoshii, K., Mizutani, T., Kariwa, H. and Takashima, I. 2000. Distribution and characterization of tick-borne encephalitis viruses from Siberia and far-eastern Asia. *J. Gen. Virol.*, 82 : 1319-28.
  - 15) Hayasaka, D., Suzuki, Y., Kariwa, H., Ivanov, L., Volkov, V., Dmenev, V., Mizutani, T., Gojobori, T. and Takashima, I. 1999. Phylogenetic and virulence analysis of tick-borne encephalitis viruses from Japan and far-eastern Russia. *J. Gen. Virol.*, 80 : 3127-3135.
  - 16) Heinz, F. X. and Allison, S. L. 2000. Structures and mechanisms in flavivirus fusion. *Adv. Virus Res.*, 55 : 231-269.
  - 17) Heinz, F. X. and Mandl, C. W. 1993. The molecular biology of tick-borne encephalitis virus. *APMIS*, 101 : 735-745.
  - 18) Holzmann, H., Heinz, F. X., Mandl, C. W., Guirakhoo, F. and Kunz, C. 1990. A single amino acid substitution in envelope protein E of tick-borne encephalitis virus leads to attenuation in the mouse model. *J. Virol.*, 64 : 5156-5159.
  - 19) Jiang, W. R., Lowe, A., Higgs, S., Reid, H. and Gould, E. A. 1993. Single amino acid codon changes detected in louping ill virus antibody-resistant mutants with reduced neurovirulence. *J. Gen.*

- Viol.*, 74 : 931-935.
- 20) Kariwa, H., Yoshizumi, S., Arikawa, J., Yoshimatu, K., Takahashi, K., Takashima, I. and Hashimoto, N. 1995. Evidence for the existence of Puumala-related virus among *Clethrionomys rufocanus* in Hokkaido, Japan. *Am. J. Trop. Med. Hyg.*, 53 : 222-227.
  - 21) Khromykh, A. A., Sedlak, P. L. and Westaway, E. G. 1999. *trans*-Complementation analysis of the flavivirus Kunjin ns 5 gene reveals an essential role for translation of its N-terminal half in RNA replication. *J. Virol.*, 73 : 9247-9255.
  - 22) Khromykh, A. A., Sedlak, P. L. and Westaway, E. G. 2000. *cis*- and *trans*-acting elements in flavivirus RNA replication. *J. Virol.*, 74 : 3253-3263.
  - 23) Koonin, E. V. 1991. The phylogeny of RNA-dependent RNA polymerases of positive-strand RNA viruses. *J. Gen. Virol.*, 72 : 2197-2206.
  - 24) Koonin, E. V. 1993. Computer-assisted identification of a putative methyltransferase domain in NS 5 protein of flaviviruses and  $\lambda$  2 protein of reovirus. *J. Gen. Virol.*, 74 : 733-740
  - 25) Krishna Murthy, H. M., Clum, S. and Padmanabhan, R. 1999. Dengue virus NS 3 serine protease. *J. Biol. Chem.*, 274 : 5573-5580.
  - 26) Lindenbach, B. D. and Rice, C. M. 1999. Genetic interaction of flavivirus non-structural proteins NS 1 and NS 4 A as a determinant of replicase function. *J. Virol.*, 73 : 4611-4621.
  - 27) Mandl, C. W., Allison, S. L., Holzmann, H., Meixner, T. and Heinz, F.X. 2000. Attenuation of tick-borne encephalitis virus by structure-based site-specific mutagenesis of a putative flavivirus receptor binding site. *J. Virol.*, 74 : 9601-9609.
  - 28) Mandl, C. W., Holzmann, H., Meixner, T., Rauscher, S., Stadler, P. F., Allison, S. L. and Heinz, F. X. 1998. Spontaneous and engineered deletions in the 3' noncoding region of tick-borne encephalitis virus : construction of highly attenuated mutants of a flavivirus. *J. Virol.*, 72 : 2132-2140.
  - 29) McMinin, P. C. 1997. The molecular basis of virulence of the encephalitogenic flaviviruses. *J. Gen. Virol.*, 78 : 2711-2722.
  - 30) Muylaert, I. R., Chambers, T. J., Galler, R. and Rice, C. M. 1996. Mutagenesis of the N-linked glycosylation sites of the yellow fever virus NS 1 protein : effects on virus replication and mouse neurovirulence. *Virology*, 222 : 159-168.
  - 31) Ni, H., Burns, N. J., Chang, G. J., Zhang, M., Wills, M. R., Trent, D. W., Sanders, P. G. and Barrett, A. D. T. 1994. Comparison of nucleotide and deduced amino acid sequence of the 5' noncoding region and structural protein genes of the wild-type Japanese encephalitis virus strain SA 14 and its attenuated vaccine derivatives. *J. Gen. Virol.*, 75 : 1505-1510.
  - 32) O'Reilly, E. K. and Kao, C. C. 1998. Analysis of RNA-dependent RNA polymerase structure and function as guided by known polymerase structures and computer predictions of secondary structure. *Virology*, 252 : 287-303.
  - 33) Proutski, V., Gritsun, T. S., Gould, E. A., Holmes, E. C. 1999. Biological consequences of deletions within the 3'-untranslated region of flaviviruses may be due to rearrangements of RNA secondary structure. *Virus Res.*, 64 : 107-123.
  - 34) Rey, F. A., Heinz, F. X., Mandl, C., Kunz,

- C. and Harrison, S. C. 1995. The envelope glycoprotein from tick-borne encephalitis virus at 2 Å resolution. *Nature*, 375 : 291-298.
- 35) Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. *Molecular Cloning : A Laboratory Manual*, 2nd edn., Cold Spring Harbor Laboratory, New York
- 36) Stadler, K., Allison, S. L., Schalich, J. and Heinz, F. X. 1997. Proteolytic activation of tick-borne encephalitis virus by furin. *J. Virol.*, 71 : 8475-8481.
- 37) Stocks, C. E. and Lobigs, M. 1998. Signal peptidase cleavage at the flavivirus C-prM junction : dependence on the viral NS2B-3 protease for efficient processing requires determinants in C, the signal peptide, and prM. *J. Virol.*, 72 : 2141-2149.
- 38) Takashima, I., Morita, K., Chiba, M., Hayasaka, D., Sato, T., Takezawa, C., Igarashi, A., Kariwa, H., Yoshimatsu, K., Arikawa, J. and Hashimoto, N. 1997. A case of tick-borne encephalitis in Japan and isolation of the virus. *J. Clin. Microbiol.*, 35 : 1943-1947.
- 39) Wallner, G., Mandl, C. W., Ecker, M., Holzmann, H., Stiasny, K., Kunz, C. and Heinz, F. X. 1996. Characterization and complete genome sequences of high- and low-virulence variants of tick-borne encephalitis virus. *J. Gen. Virol.*, 77 : 1035-1042.
- 40) Wallner, G., Mandl, C. W., Kunz, C. and Heinz, F. X. 1995. The flavivirus 3' - noncoding region : Extensive size heterogeneity independent of evolutionary relationships among strains of tick-borne encephalitis virus. *Virology*, 213 : 169-178.
- 41) Wengler, G., Wengler, G. 1993. The NS3 nonstructural protein of flavivirus contains an RNA triphosphatase activity. *Virology*, 197 : 265-273.
- 42) Zilber, L. A. and Soloviev, V. D. 1946. Far Eastern tick-borne spring-summer (spring) encephalitis. *Am. Rev. Soviet Med.*, 5 (special suppl.)