



HOKKAIDO UNIVERSITY

Title	COMPARISON OF ANTIGENS AND PROTEIN PROFILES OF VIRULENT AND AVIRULENT CLONES OF LEPTOSPIRA INTERROGANS SEROVAR COPENHAGENI STRAIN SHIBAURA
Author(s)	NIIKURA, Masahiro
Citation	Japanese Journal of Veterinary Research, 34(2), 144-144
Issue Date	1986-04-30
Doc URL	https://hdl.handle.net/2115/2991
Type	departmental bulletin paper
File Information	KJ00002374398.pdf



COMPARISON OF ANTIGENS AND PROTEIN PROFILES OF VIRULENT AND
AVIRULENT CLONES OF *LEPTOSPIRA INTERROGANS* SEROVAR
COPENHAGENI STRAIN SHIBAURA

Masahiro NIKURA

*Department of Hygiene and Microbiology
Faculty of Veterinary Medicine
Hokkaido University, Sapporo 060, Japan*

The antigens and protein profiles of virulent and avirulent clones of *Leptospira interrogans* serovar *copenhageni* strain Shibaura were compared by SDS-polyacrylamide gel electrophoresis (SDS-PAGE), Western blotting and radioimmunoprecipitation.

A virulent clone specific antigen was detected by Western blotting. The molecular weight of the antigen was estimated to be approximately 32000 daltons by SDS-PAGE, and the antigen was found to be thermostable.

A 43000 dalton antigenic molecule and a 41000 dalton antigenic molecule that were specific, respectively, for the virulent clone and the avirulent clone were detected by Western blotting. However, these antigenic molecules contained common antigenic determinants between the virulent and avirulent clones.

No difference between the virulent and avirulent clones was observed in the comparison of protein profiles on SDS-PAGE and of the surface-exposed protein antigens detected by radioimmunoprecipitation.

Production of virulent clone specific monoclonal antibodies was attempted. Although 12 hybridoma cell lines secreting monoclonal antibodies were produced, none of them was specific for the virulent clone of leptospira. However, one of these monoclonal antibodies showed over 10 times higher agglutination titer for the virulent clone than for the avirulent clone of leptospira.